Reasons underlying the intention to vaccinate children aged 5-11 against COVID-19: A

cross-sectional study of parents in Israel, November 2021

Authors:

Nicole G. Morozov[1], Amiel A. Dror[2,3*], Amani Daoud[2,3]; Netanel Eisenbach[2,3]; Edward

Kaykov[3,4]; Masad Barhoum[5]; Tsvi Sheleg[5]; Eyal Sela[2,3]; Michael Edelstein[3]

*Morozov and Dror are equal first authors

[1]Tel Aviv University, Tel Aviv, Israel

[2]Department of Otolaryngology, Head and Neck Surgery, Galilee Medical Center, Nahariya,

Israel

[3]Azrieli Faculty of Medicine, Bar-Ilan University, Safed, Israel

[4]Geriatric Medicine Department, Galilee Medical Center, Nahariya, Israel

[5]Galilee Medical Center, Nahariya, Israel

*Corresponding author: Amiel A. Dror, Department of Otolaryngology, Head and Neck Surgery,

Galilee Medical Center, Nahariya, Israel; Azrieli Faculty of Medicine, Bar-Ilan University,

Safed, Israel Tel: +972544530534; email: amieldror@gmail.com

Abstract

Background

Vaccination is a key tool to mitigate impacts of the COVID-19 pandemic. In Israel, COVID-19

vaccines became available to adults in December 2020 and to 5–11-year-old children in

November 2021. Ahead of the vaccine roll-out in children, we aimed to determine whether

parents intended to vaccinate their children and describe reasons for their intentions.

Methods

We recruited parents on social media and collected information on parental socio-demographic

characteristics, COVID-19 vaccine history, intention to vaccinate their children against COVID-

19, and reasons for parental decisions, using an anonymous online survey. We identified

associations between parental characteristics and intention to vaccinate children using a logistic

regression model and described reasons for intentions to vaccinate or not using proportions

together with 95% confidence intervals (CI).

Results

1837 parents participated. Parental non-vaccination and having experienced major vaccination

side effects were strongly associated with non-intention to vaccinate their children (OR 0.09 and

0.18 respectively, p<0.001). Compared with others, parents who were younger, lived in the

socio-economically deprived periphery, and belonged to the Arab population had lower

intentions to vaccinate their children. Commonly stated reasons for non-intention to vaccinate

included vaccine safety and efficacy (53%, 95%CI 50-56) and the belief that COVID-19 was a

mild disease (73%, 95%CI 73-79). The most frequently mentioned motives for intending to

vaccine children was returning to normal social and educational life (89%, 95%CI 87-91).

Conclusion

Parental socio-demographic background and their own vaccination experience was associated

with intention to vaccinate their children aged 5-11. Intention to vaccinate was mainly for social

and economic reasons rather than health, whereas non-intention to vaccinate mainly stemmed

from health concerns. Understanding rationales for COVID-19 vaccine rejection or acceptance,

as well as parental demographic data, can pave the way for intentional educational campaigns to

encourage not only vaccination against COVID-19, but also regular childhood vaccine

programming.

Keywords: COVID-19; vaccine hesitancy; childhood vaccines; parental vaccine hesitancy

Highlights

• Parental intention to vaccinate 5-11 children is much lower than vaccine coverage in

parental age groups

• Being unvaccinated and having experienced side effects following vaccination were the

greatest negative predictors in parents of intention to vaccinate their children

• Parents were more likely to accept a COVID-19 vaccine for their children to allow them

to return to daily social life and to ensure economic security in the family

• Parents were more likely to reject a COVID-19 vaccination for health reasons such as

safety concerns or due the belief that COVID-19 was a mild disease in children

Introduction

The rapid, global spread of COVID-19, an infection caused by the SARS-CoV-2 virus, spurred

the expeditious development of several vaccines and the eventual implementation of vaccine

programs at record speeds. In Israel, COVID-19 vaccines became available in December 2020

for adults and eligibility gradually extended to younger ages. Teenagers aged 12-15 became

eligible in June 2021 and children aged 5-11 in November 2021. Most high-income countries

now offer teenage vaccination, but vaccination of younger children is less common despite

evidence of its safety and effectiveness [1]. Outside of Israel, the United States' (USA) Centers

for Disease Control and Prevention (CDC) recommends vaccination in the 5-11 age group while

the European Centre for Disease prevention and Control has also recommended considering

universal vaccination in this demographic [2, 3].

The short period between the pandemic's onset and vaccine availability has added additional

challenges for vaccine acceptance among the public [4]. Almost two years after the onset of the

pandemic and one year after the first administration of COVID-19 vaccines in adults, the USA's

Food and Drug Administration (FDA) approved the Pfizer/BioNTech mRNA vaccine for 5- to

11-year-old children [1, 5]. Studies conducted in the UK, USA, and Canada suggest that many

parents remain concerned about the vaccine's side effects and safety, leading parents to be

hesitant to administer the COVID-19 vaccine to their children [6-8]. Research has not yet

examined the reasons why parents would be motivated or hesitant to vaccinate their children in

Israel.

Demographic criteria which have been associated with higher rates of vaccine refusal for

children include younger parental age, non-white or minority ethnicity, lower education level,

history of vaccination non-compliance to recommended childhood vaccinations, and lower

income [7, 9-11]. Parents concerned about future finances were more likely to accept a

vaccination for their offspring [11]. Attitudes and behaviors towards vaccination also tend to be

specific to the vaccine in question and to each country [12]. Compared with most high-income

countries, Israel typically achieves high coverage for routine childhood vaccines. The 2020

national coverage for the first MMR dose among 24-month-olds was 99% in Israel vs 91% in the

United States and in the UK [13]. In addition, Israel's largest ethnic minority typically achieves

higher coverage for routine childhood vaccines compared with the general population [14]. It is,

therefore, important to understand and document attitudes to COVID-19 vaccination in different

contexts due to the implications for public health interventions.

We aimed to investigate the intentions of parents in Israel to vaccinate their 5-11-year-old

children against COVID-19 and the underlying reasons for their intentions. We sought to

determine what common reasons parents listed for accepting or rejecting vaccinating their

children. Our survey was initiated two weeks before children in this age group became eligible

for COVID-19 vaccination.

Methods

Study design

This study was designed as a cross sectional survey to understand the general populations'

willingness to vaccinate their children against COVID-19 and the reasons for their acceptance or

refusal of the vaccine using an online questionnaire in Hebrew and Arabic and distributed

through social media channels (Facebook and Twitter). Informed consent was requested on the

introductory page of the online questionnaire, prior to survey enrollment. The anonymous web-

based survey followed the American Association for Public Opinion Research (AAPOR)

reporting guidelines [15]. The survey was anonymous and no identifiable information was

collected. Participants were permitted to terminate participation at will. Approval of this study

was granted by the Research Ethics Committee of the Galilee Medical Center. The questionnaire

was designed and distributed using Qualtrics (Qualtrics, Provo, UT, United States of America).

Data collection

The survey had two sections: The first part included nine questions about socio-demographic

characteristics of the participating parents including parental age, gender, area of residence,

household composition, number of children, ethnicity, parental education, parental COVID-19

vaccination status, and self-reported side effects from the vaccine (major, mild, or no symptoms),

and one question on intention to vaccinate their child(ren) against COVID-19.

In the second section, responders who indicated they were not intending to vaccinate their

children against COVID-19 were asked to state their reasons from a list of predefined

explanations, and those who indicated intention to vaccinate were also asked to affirm why from

a list of potential reasons. A free text answer was also provided.

The full questionnaire is available in Appendix 1. The estimated answering time for completing

the survey was under two minutes.

Statistical analysis

We calculated the proportion of parents intending to vaccinate their children together with 95%

confidence intervals, overall and in each stratum. We compared parents who intended to

vaccinate their children to those who did not in terms of all socio-demographic factors recorded

in the questionnaire using a univariate logistic regression model. In order to determine the extent

to which each factor was an independent predictor, we conducted a multivariate logistic

regression analysis adjusting for all other factors found to be statistically significant (at the

alpha=0.05 level) in the univariate analysis. We also calculated the proportion of parents giving

each reason for intending or not intending to vaccinate together with 95% confidence intervals.

Data were prepared using Microsoft Excel and analyzed using STATA 15.

Results

We received 1837 responses to our survey between November 1-14, 2021. Of those, 1514

(82.4%) completed the survey in its entirety. The distribution of responders according to age,

sex, ethnicity, education level, and other demographic criteria is described in Table 1. Overall

intention to vaccinate was 44% (95% CI 41.3-45.9).

Female parents and parents above the age of 35 were more likely to vaccinate than male parents

and parents aged 35 and younger (47% vs. 40%, p<0.002; 45% vs. 39%, p<0.02, Table 1).

Compared with non-Arab Israeli parents, Israeli Arab parents reported a lower intention to

vaccinate their children (34% vs. 47%, p<0.001, Table 1). Other parental factors significantly

associated with a lower intention to vaccinate included having a single child, living in the

periphery (Northern or Southern, socioeconomically depressed regions of the country), having

received fewer than three doses of a COVID-19 vaccine and having experienced major side

effects of COVID-19 vaccination. After adjusting for other factors, being male, living in the

periphery, belonging to the Arab population, reception of fewer than three doses of the COVID-

19 vaccine and having experienced major side effects following COVID-19 vaccination all

remained significantly associated with a lower intention to vaccinate children (Table 1). Being

completely unvaccinated and having suffered major side effects were associated with the largest

reduction in intention to vaccinate children (-91% and -81% respectively, p<0.001).

Of 801 parents intending to vaccinate their children, 93% were COVID-19 vaccinated at least

once. Intention to vaccinate their children increased with the number of doses the parents had

received; while only 13% of unvaccinated parents intended to vaccinate their children against

COVID-19, 32% of parents vaccinated with one dose, 48% of parents vaccinated with two doses,

and 62% of parents vaccinated with three doses were willing to vaccinate their children

(p<0.001, Table 1).

The most often cited reason for choosing to vaccinate their children was to allow their children to

return to everyday social, educational, and financial life, followed by protecting other family

members from COVID-19 infection (89 and 77.9% respectively, Table 2). The protection of the

children against viral COVID-19 infection followed by vaccine safety were the least often

mentioned explanations for parents who intended to vaccinate their children (37.9%).

The most frequently mentioned explanation for not vaccinating their children is the belief that

COVID-19 leads to mild or asymptomatic disease in children (76.1%, Table 2). Concerns of the

parents regarding vaccine safety were ordered second in their consideration not to vaccinate their

children (53.1%, Table 2).

Discussion

Routine childhood vaccination offers the prospect of protection against a transmissible and

infectious, but preventable, threat to health. In most settings, there is no ongoing endemic

transmission of poliovirus, measles, or diphtheria infections due to effective global vaccination

campaigns [16-18]. However, COVID-19 is unique in several ways: first, the pandemic is

continuing to develop and therefore vaccine protection is against an immediate and tangible

threat; second, the COVID-19 pandemic is the only instance in recent history that a mass

vaccination campaign concurrently targets children and their parents for initiating vaccination;

third, the volume and availability of information around COVID-19 vaccines, offering arguments

both for and against vaccination with very few verification pathways for parents to assess the

accuracy of the information they are accessing, is an unprecedent phenomenon; fourth, the speed

at which COVID-19 vaccines were developed have raised some concerns about safety among the

general public [19, 20]

We found that the intention to vaccinate children aged 5-11 was low, overall. Less than half of

parents intended to vaccinate their 5-11 children, and as of February 2022 just over 20% had

vaccinated their offspring[21]. Even among parents who had received three doses of the vaccine,

intention to vaccinate was only 62%. This is consistent with experience from other countries

where vaccinated parents did not intend to vaccination their children [22]. Parental experience

was clearly a key factor in the decision, with unvaccinated parents and those who experienced

perceived severe side effects very unlikely to vaccinate their children. Parents with higher

education were not more likely to vaccinate than others.

Our study has highlighted that the main reasons for intending to vaccinate their children were not

biomedical. Our study suggests that, for COVID-19, the traditional perception of vaccination

benefits such as protection against illness and abolishing the spread of infection has been

superseded by indirect benefits such as returning to regular societal life and education

institutions, as well as assuring financial resilience for the family. While this finding is not

surprising considering the severe disruption to normal life caused by the COVID-19 pandemic, it

is not clear whether this perception of the societal utility of vaccines will continue to impact the

perception of other vaccines beyond the pandemic.

Groups who may be targets for intersectional outreach include minorities, parents with a lower

educational level, and families in lower financial brackets as these parents have been highlighted

230 as more likely to be vaccine hesitant [10, 23]. Our research found that Arab parents were less

231 likely to vaccinate their children as compared to other parents, consistent with research finding a

232 higher rate of COVID-19 vaccine hesitancy among minority communities in other countries, as

233 well as in Israel's Arab and Ultra-Orthodox communities [7, 24, 25].

235 Our study has identified that key factors within Israeli society associated with an increased

236 intention to vaccinate children against COVID-19 include the parent's desire for the child to

237 return to normal social and educational life along with secondary motives such as the parent's

238 desire to maintain job security. The main drivers against intention to vaccinate are much more

239 biomedical, encompassing beliefs that COVID-19 causes a mild disease course in children and

240 concerns about vaccine safety. These findings have implications for health promotion and

241 communication campaigns; ensuring parents' confidence in the safety of the vaccine and

242 highlighting the sociocultural benefits of vaccination may be key in guiding communities to

243 higher levels of vaccine uptake.

245 Limitations of our investigation include the self-selection of participants in answering the

246 questionnaire and the participants' self-identification of socio-demographic characteristics.

247 Potentially, individuals who have stronger opinions either for or against vaccinating their

248 children for COVID-19 may be more likely to complete a questionnaire about the topic. It must

249 be noted that the age, sex, and geographical distribution of participants suggest that all segments

250 of the population were represented.

It is not clear to what extent our findings are generalizable to other countries. Globally, there are

wide discrepancies in vaccine uptake from one country to another with some countries enjoying

remarkably low levels of vaccine hesitancy among parents [9, 11]. Understanding what factors

enable childhood COVID-19 vaccination in countries with high parental intent and transferring

this knowledge to other contexts could help countries struggling with COVID-19 childhood

vaccination programs. Because a growing number of individuals, including parents, are seeking

health and vaccine information online or through peers, the availability of trustworthy sources

and early educational campaigns cannot be overstated [26].

Conclusion

The availability of COVID-19 vaccines and licensing of these vaccinations for use in children

aged five to 11 is not a guarantee that parents will inoculate their children. Our study showed that

intention to vaccinate 5- to 11-year-old children was low even among fully vaccinated parents,

with socio-cultural factors, as well as vaccination experience in parents, being associated with

intention to vaccinate their children. Our study showed that indirect benefits of vaccination such

as returning to regular societal activities, including educational institutions and personal

economic stability, were bigger motivators than direct protection against the virus, while the

most common reasons for intending not to vaccinate children against COVID-19 were classical

biomedical concerns such disease severity and vaccine safety. These findings are important for

health authorities to shape educational and communication campaigns: focusing on the message

that vaccines protect against COVID-19 may not be as effective as messages stating that

vaccines will allow children to go to school and adults to work. Different messaging campaigns

should be implemented and evaluated to test this possibility.

Conflicts of interest: The authors declare no conflicts of interest.

Attestation: All authors attest they meet the ICMJE criteria for authorship

Citations

1.   Walter EB, Talaat KR, Sabharwal C, Gurtman A, Lockhart S, Paulsen GC, et al.

Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age. New England

Journal of Medicine. 2021.10.1056/NEJMoa2116298

2.   CDC. COVID-19 Vaccination for Children 5-11 Years Old [Internet]. Centers for

Disease Control and Prevention. 2022 [cited 2022 Feb 19]. Available from:

https://www.cdc.gov/vaccines/covid-19/planning/children.html

3.   Interim public health considerations for COVID-19 vaccination of children aged 5-11

years [Internet]. European Centre for Disease Prevention and Control. 2021 [cited 2022 Feb 19].

Available from: https://www.ecdc.europa.eu/en/publications-data/interim-public-health-

considerations-covid-19-vaccination-children-aged-5-1

4.   Soares P, Rocha JV, Moniz M, Gama A, Laires PA, Pedro AR, et al. Factors Associated

with COVID-19 Vaccine Hesitancy. 2021;9(3):300

5.   Woodworth KR, Moulia D, Collins JP, Hadler SC, Jones JM, Reddy SC, et al. The

Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-

BioNTech COVID-19 Vaccine in Children Aged 5-11 Years - United States, November 2021.

MMWR Morb Mortal Wkly Rep. 2021;70(45):1579-83.10.15585/mmwr.mm7045e1

6.   Ruggiero KM, Wong J, Sweeney CF, Avola A, Auger A, Macaluso M, et al. Parents'

Intentions to Vaccinate Their Children Against COVID-19. Journal of Pediatric Health Care.

2021;35(5):509-17.https://doi.org/10.1016/j.pedhc.2021.04.005

7.   Bell S, Clarke R, Mounier-Jack S, Walker JL, Paterson P. Parents' and guardians' views

on the acceptability of a future COVID-19 vaccine: A multi-methods study in England. Vaccine.

2020;38(49):7789-98.https://doi.org/10.1016/j.vaccine.2020.10.027

8.   Humble RM, Sell H, Dubé E, MacDonald NE, Robinson J, Driedger SM, et al. Canadian

parents' perceptions of COVID-19 vaccination and intention to vaccinate their children: Results

from a cross-sectional national survey. Vaccine. 2021;39(52):7669-

76.https://doi.org/10.1016/j.vaccine.2021.10.002

9.   Bagateli LE, Saeki EY, Fadda M, Agostoni C, Marchisio P, Milani GP. COVID-19

Vaccine Hesitancy among Parents of Children and Adolescents Living in Brazil. Vaccines.

2021;9(10).10.3390/vaccines9101115

10.   McKinnon B, Quach C, Dubé È, Tuong Nguyen C, Zinszer K. Social inequalities in

COVID-19 vaccine acceptance and uptake for children and adolescents in Montreal, Canada.

Vaccine. 2021;39(49):7140-5.https://doi.org/10.1016/j.vaccine.2021.10.077

11.   Goldman RD, Krupik D, Ali S, Mater A, Hall JE, Bone JN, et al. Caregiver Willingness

to Vaccinate Their Children against COVID-19 after Adult Vaccine Approval. International

Journal of Environmental Research and Public Health. 2021;18(19).10.3390/ijerph181910224

12.   Larson HJ, de Figueiredo A, Xiahong Z, Schulz WS, Verger P, Johnston IG, et al. The

State of Vaccine Confidence 2016: Global Insights Through a 67-Country Survey.

EBioMedicine. 2016;12:295-301.10.1016/j.ebiom.2016.08.042

13.   Measles vaccination coverage [Internet]. World Health Organization Available from:

https://immunizationdata.who.int/pages/coverage/mcv.html?CODE=ISR+GBR+USA&ANTIGE

N=MCV1&YEAR=.

14.   Stein-Zamir C, Israeli A. Timeliness and completeness of routine childhood vaccinations

in young children residing in a district with recurrent vaccine-preventable disease outbreaks,

Jerusalem, Israel. Euro surveillance : bulletin Europeen sur les maladies transmissibles =

European communicable disease bulletin. 2019;24(6).10.2807/1560-7917.Es.2019.24.6.1800004

15.   Transparency Initiative Disclosure Elements. American Assocation for Public Opinion

Research; 2021

16.   Obregón R, Chitnis K, Morry C, Feek W, Bates J, Galway M, et al. Achieving polio

eradication: a review of health communication evidence and lessons learned in India and

Pakistan. Bulletin of the World Health Organization. 2009;87(8):624-30.10.2471/blt.08.060863

17.   Clarke KEN, MacNeil A, Hadler S, Scott C, Tiwari TSP, Cherian T. Global

Epidemiology of Diphtheria, 2000-2017(1). Emerging infectious diseases. 2019;25(10):1834-

42.10.3201/eid2510.190271

18.   Holzmann H, Hengel H, Tenbusch M, Doerr HW. Eradication of measles: remaining

challenges. Medical microbiology and immunology. 2016;205(3):201-8.10.1007/s00430-016-

0451-4

19.   The Lancet Infectious D. The COVID-19 infodemic. The Lancet Infectious diseases.

2020;20(8):875.10.1016/s1473-3099(20)30565-x

20.   Salerno L, Craxì L, Amodio E, Lo Coco G. Factors Affecting Hesitancy to mRNA and

Viral Vector COVID-19 Vaccines among College Students in Italy. Vaccines.

2021;9(8).10.3390/vaccines9080927

21.   Health IMo. Coronavirus in Israel - general situation 2022. p.

https://datadashboard.health.gov.il/COVID-19/general

22.   Horiuchi S, Sakamoto H, Abe SK, Shinohara R, Kushima M, Otawa S, et al. Factors of

parental COVID-19 vaccine hesitancy: A cross sectional study in Japan. PloS one.

2021;16(12):e0261121.10.1371/journal.pone.0261121

23.   Alfieri NL, Kusma JD, Heard-Garris N, Davis MM, Golbeck E, Barrera L, et al. Parental

COVID-19 vaccine hesitancy for children: vulnerability in an urban hotspot. BMC Public

Health. 2021;21(1):1662.10.1186/s12889-021-11725-5

24.   Robinson E, Jones A, Lesser I, Daly M. International estimates of intended uptake and

refusal of COVID-19 vaccines: A rapid systematic review and meta-analysis of large nationally

representative samples. Vaccine. 2021;39(15):2024-

34.https://doi.org/10.1016/j.vaccine.2021.02.005

25.   Gorelik Y, Anis E, Edelstein M. Inequalities in initiation of COVID19 vaccination by age

and population group in Israel- December 2020-July 2021. The Lancet regional health Europe.

2022;12:100234.10.1016/j.lanepe.2021.100234

26.   Rhodes ME, Sundstrom B, Ritter E, McKeever BW, McKeever R. Preparing for A

COVID-19 Vaccine: A Mixed Methods Study of Vaccine Hesitant Parents. Journal of Health

Communication. 2020;25(10):831-7.10.1080/10810730.2021.1871986

Table 1. Parental socio-demographic background and vaccination history and intention to vaccinate											
		Number of respondents	Number intending to vaccinate their 5–11-year-old child	Percent intending to vaccinate their 5-11-year-old child	95% CI	Unadjusted OR	Unadjusted OR 95%CI	P value	Adjusted OR*	Adjusted OR 95%CI	P value
Total		1837	801	43.6	41.3-45.9	-	-		-		-
Parent sex	Female	891	422	47.4	44-50.7	baseline					
	Male	946	379	40.1	36.9-43.2	0.74	0.62-0.89	0.002	0.75	0.6-0.94	0.01
Parent age	>35	1309	594	45.4	42.6-48.1	baseline					
	≤35	528	207	39.2	35-43.5	0.78	0.74-0.93	0.02	0.85	0.67-1.1	0.2
Number of children	One child	421	161	38.2	33.6-43.1	0.75	0.6-0.94	0.01	0.74	0.57-0.96	0.02
	> one child	1416	640	45.2	42.6-47.8	baseline					
Household composition	Two parents	1568	669	42.7	40.2-45.1	0.77	0.6-1	0.05	0.8	0.59-1.1	0.2
	Not two parents	269	132	49.1	42.9-55.2	baseline					
Parent education level	High school or under	379	161	42.5	37.4-47.6	baseline					
	Bachelor's degree	1155	525	45.5	42.5-48.4	1.13	0.89-1.4	0.3	-	-	-
	Higher degree	303	115	38.0	32.5-43.7	0.83	0.61-1.13	0.2	-	-	-
Resident area	Center	1103	508	46.1	43.1-49	baseline					
	Periphery	734	293	39.9	36.3-43.6	0.78	0.64-0.94	0.01	0.73	0.59-0.92	0.006
Parent ethnicity	Non-Arab Israeli	1298	611	47.1	44.3-49.8	baseline					
	Arab Israeli	469	159	33.9	29.6-38.4	0.58	0.46-0.72	<0.001	0.65	0.51-0.84	0.001
	Other	70	31	44.3	32.4-56.7	0.89	0.55-1.45	0.6	1.3	0.71-2.3	0.4
Parent COVID-19 vaccination status	3 vaccine doses	578	357	61.8	57.7-65.7	baseline					
	2 vaccine doses	728	351	48.2	44.5-51.9	0.57	0.46-0.72	<0.001	0.61	0.49-0.77	<0.001
	1 vaccine doses	125	40	32.0	23.9-40.9	0.29	0.19-0.43	<0.001	0.29	0.19-0.45	<0.001
	Not vaccinated	406	53	13.1	9.9-16.7	0.09	0.07-0.13	<0.001	0.09	0.07-0.13	<0.001
Parent COVID-19 vaccination side effects	Yes: major	112	22	19.6	12.7-28.2	0.18	0.11-0.29	<0.001	0.19	0.11-0.31	<0.001
	Yes: minor	546	279	51.1	46.8-55.3	0.76	0.61-0.95	0.02	0.76	0.60-0.95	0.02
	No	773	447	57.8	54.2-61.3	baseline					
	Missing data	406									
	For variable significance in the univariate analysis, adjusted for other significant variables										

Table 2. Stated reasons given by parents for intention or refusal to vaccinate their children, Israel, November 2021				
		n	%	95% CI
I want to vaccinate my child…(n=801)	to return to normal life and activities	713	89	86.6- 91.1
	to loss and secure income	624	77.9	74.8-80.7
	to protect others	560	69.9	66.6-73.1
	to protect my child	448	55.9	52.4-59.4
	because the vaccine is safe	304	37.9	34.6-41.4
I don't want to vaccinate my child… (n=1036)	because COVID-19 is a mild illness	788	76.1	73.3-78.6
	because of vaccine safety concerns	550	53.1	50-56.2
	because I don't trust the government	435	42	39-45
	because I don't trust the Ministry of Health	217	21	18.5-23.5
	because I think COVID-19 is a conspiracy	145	14	11.9-16.2