Burden of Disease from Contaminated Drinking Water in Countries with High Access to Safely Managed Water: A Systematic Review

Debbie Lee[1,2], Jacqueline MacDonald Gibson[3], Joe Brown[4], Jemaneh Habtewold[5], Heather M. Murphy[1,5#]

[1] Water, Health and Applied Microbiology (WHAM) Lab, Department of Epidemiology and Biostatistics, College of Public Health, Temple University, Philadelphia, PA, USA
[2] Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO, USA
[3] Department of Civil, Construction, and Environmental Engineering, North Carolina State University, Raleigh, NC, USA
[4] Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
[5] Water, Health and Applied Microbiology (WHAM) Lab, Department of Pathobiology, Ontario Veterinary College, University of Guelph, Guelph, ON, CAN

#Corresponding author:
Dr. Heather M. Murphy
50 Stone Road East
Guelph, ON, CAN N1G 2W1
heather.murphy@uoguelph.ca

Abstract
The vast majority of residents of high-income countries (≥90%) reportedly have high access to safely managed drinking water. Owing perhaps to the widely held perception of near universal access to high-quality water services in these countries, the burden of waterborne disease in these contexts is understudied. This systematic review aimed to: identify population-scale estimates of waterborne disease in countries with high access to safely managed drinking water, compare methods to quantify disease burden, and identify gaps in available burden estimates. We conducted a systematic review of population-scale disease burden estimates attributed to drinking water in countries where ≥90% of the population has access to safely managed drinking water per official United Nations monitoring. We identified 24 studies reporting estimates for disease burden attributable to microbial contaminants. Across these studies, the population-weighted average burden of gastrointestinal illness risks attributed to drinking water was ~3,529 annual cases per 100,000 people. Beyond exposure to infectious agents, we identified 10 studies reporting disease burden—predominantly, cancer risks—associated with chemical contaminants. Across these studies, the pooled population-weighted average of excess cancer cases attributable to drinking water was 1.8 annual cancer cases per 100,000 people. These estimates exceed WHO-recommended normative targets for disease burden attributable to drinking water and highlight that there remains important preventable disease burden in these contexts. However, the available literature was scant and limited in geographic scope, disease outcomes, range of microbial and chemical contaminants, and inclusion of subpopulations (rural, low-income communities; Indigenous or Aboriginal peoples; and populations marginalized due to discrimination by race, ethnicity, or socioeconomic status) that could most benefit from water infrastructure investments. Studies quantifying drinking water-associated disease burden in

47 countries with reportedly high access to safe drinking water, with a focus on specific
48 subpopulations and promoting environmental justice, are needed.
49
50 **Keywords: burden of disease; drinking water; waterborne disease; high-income countries;**
51 **water access**
52
53 **1. Introduction**
54
Since 1990, the WHO/UNICEF Joint Monitoring Programme for Water Supply, Sanitation and
Hygiene has reported on access to safe drinking water worldwide. WHO/UNICEF defines
drinking water as safely managed if it comes from an improved source that is accessible on
premises, available when needed, and free of fecal and priority chemical contamination.
WHO/UNICEF data indicate that in countries classified by the World Bank as low- or middle-
income (per-capita incomes <$12,696), major gaps in safe water access remain, with 49.7% of
the population unserved by piped water as of 2017 (Deshpande et al., 2020). These gaps manifest
in high disease burdens. Recent estimates attribute unsafe drinking water as the cause of 36% of
the 1.4 million annual diarrheal deaths worldwide in low- and middle-income countries (Prüss-
Ustün et al., 2019).
65
In contrast to the situation in low- and middle-income countries, WHO/UNICEF data suggest
that in high-income countries, most of the population has safe water, with an average of <3%
lacking access as of 2017 (Table S1). However, the WHO/UNICEF's national-scale data and
data collection methods can mask local-scale inequities in high-income countries, such as in
rural, low-income, and minority communities (Anderson, 2008; Balazs and Ray, 2014). They can
also lead to a misconception that water problems of high-income countries have been solved,
resulting in underinvestment in maintaining and extending water infrastructure. For example, a
recent commentary on infrastructure investment needs in the United States (US) noted
"Americans expect the water flowing from their kitchen faucets to be clean and safe," yet "water
infrastructure is particularly vulnerable to neglect" because it "is buried underground or removed
from public view, and is thus easily ignored" (Morris, 2017). Given the focus of the global water,
sanitation, and hygiene (WASH) sector on low- and middle-income countries, the present work
was undertaken to better understand the burden of waterborne disease in high-income countries
and to inform a more global understanding of WASH.
80
Improved understanding of the burden of waterborne disease in countries with high access to
safely managed drinking water (most high-income and some middle-income countries) can help
identify remaining contamination issues and gaps for water infrastructure investments. To
support such efforts, we conducted a systematic review to identify studies estimating the disease
burden attributed to contaminated drinking water in the 64 countries/territories with ≥90% safe
water access per the WHO/UNICEF definition (Table S1, Figure S1). Among these
countries/territories, 54 are classified as high-income; 7 are upper middle income (per-capita
incomes between $4,096 and $12,695); and 3 are lower middle income (per-capita incomes
between $1,046 and $4,095). The review was intended to:
90
91   1. identify available estimates of waterborne disease burden in countries with reportedly
92   high access to safely managed drinking water;

93   2. evaluate and compare methods to quantify disease burden; and
94   3. identify gaps in available disease burden estimates.
95
96   **2. Methods**
97
98   2.1 Review Method
99   We followed the Cochrane Handbook for Systematic Reviews (Higgins et al., 2019) and the
100 Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) when
101 conducting our review (Page et al., 2021). Table S2 documents study adherence to PRISMA
102 guidelines for reporting.
103
2.2 Search Strategy
We searched PubMed, Web of Science, and SCOPUS for relevant articles published before
September 10, 2021. We also searched for grey literature on burden estimates by country on
Google and Google Scholar. The objective of this systematic review was to identify articles with
explicit population-scale estimates of disease burden, so we limited our search terms
accordingly. We searched for articles that included "burden of disease" OR "disease burden" OR
"gastrointestinal illness" AND the terms "drinking water" OR "tap water" in text. Last,
additional articles were identified from consultations with subject-matter experts and from hand
searching reference lists of included articles. Text S1 provides additional details on the
development of search criteria.
114
115 2.3 Inclusion Criteria
116 Articles in English meeting these inclusion criteria were considered:
117
118   a) population-scale burden estimate is for a country where ≥90% of the population has
119   access to safely managed water supplies (as defined by the WHO; Table S1, Figure S1)
120   and
121   b) estimates cases, hospitalizations, or deaths that could be prevented if ingestion of one or
122   more contaminants in drinking water were prevented.
123
124 Articles focusing exclusively on exposure to contaminated water via recreation (e.g., swimming
125 in contaminated water), inhalation, or dermal uptake were excluded. At least two independent
126 reviewers screened the articles against these criteria.
127
2.4 Data Extraction & Synthesis
Disease burden estimates were extracted by co-authors (DL; HM; JMG) using a custom
spreadsheet to record estimates, study design, population, disease type(s), contaminant type(s)
(microbial or chemical), and drinking water source. Given the limited availability of burden
estimates and the wide variation in metrics for characterizing the disease burden, a formal meta-
analysis was not possible. However, to enable comparison of estimates across studies, the
reported disease burden was converted to incidence per 100,000 population. For some studies,
conversion factors were needed to translate the metrics used to report the burden estimate to
cases of illness or deaths per 100,000 people. For example, some studies estimated the number of
hospitalizations for gastrointestinal illness (GI) attributable to contamination of drinking water.
To express these results as total GI cases attributable to drinking water contamination, estimates

of the proportion of GI cases resulting in hospitalization were obtained using data for the
location where the study was carried out. Table S3 provides details. Disease burden estimates
also were compared by country or region. Text S2 outlines approaches for generating these
geographic comparisons.
143
3. Results
A total of 514 articles were recovered from the databases, and 14 articles were added by subject-
matter experts. After removal of duplicates, 528 articles remained for screening (Figure S2).
Thirty-three articles met the inclusion criteria. All but one of the articles focused on a single
country or union of countries. Of these 32 articles, most (68.8%) burden estimates were from the
United States or Canada; 12.5% were from Asia; 9.4% reported on European countries; and
9.4% were from Australasia (Figure S3, Table S4). Two articles provided estimates from a
middle-income country (Iran); the rest focused on high-income countries.
152
3.1 Infectious Diseases
Twenty-four articles estimated GI risks (Table 1). Estimates are difficult to compare across
studies because some estimated total GI burden, whereas others quantified the burden only for
specific pathogens or specific outcomes (e.g., hospitalization). When sufficient information was
available, we converted estimates to units of total cases per 100,000 people to enable cross-study
comparisons (Table S3).
159
The population-weighted average burden of GI attributable to drinking water was approximately
3,529 cases per 100,000 population (range 40–18,250 per 100,000) across studies that estimated
total GI burden or that could be adjusted to estimate this burden. The highest estimate was from a
study in Iowa (Eisenberg et al., 2006), using data from a drinking water intervention trial
comparing GI cases among participants with and without household water filters (Figure 1). The
lowest was from a North Carolina (US) study (DeFelice et al., 2015) that compared three
different estimation methods; two resulted in estimates of ~40 annual cases per 100,000, while
the third estimate was ~6,300 per 100,000 people. Across studies, the population-weighted
standard deviation (2,535 per 100,000) was relatively low, just over half the mean.
169
170 Of the 24 studies, five directly estimated the number of deaths attributable to microbial
171 contaminants. These estimates were very low, averaging 1 per 100,000 people annually (range =
172 7.3×10^{-3} to 2.1 per 100,000).
173
174 3.2 Non-infectious Diseases
175 Ten articles estimated the disease burden from chemical contaminants in drinking water (Table
176 2).
177
178 Cancers were the most common health outcome considered, included in 7 of the 10 studies
179 (Figure 2). Estimated cancer risks varied depending on the cancer type and contaminants
180 considered. Across all studies with cancer risk estimates, the population-weighted average
181 disease burden was 1.8 excess cases per 100,000 people annually (range = 0.09–2.91).
182
183 Two studies, both in the United States, estimated adverse reproductive outcomes from nitrate in
184 drinking water (Mathewson et al., 2020; Temkin et al., 2019). In both studies, the disease burden

was low on a per-capita basis. The highest burden was 1.6 low-birthweight babies per 100,000 people annually.

Surprisingly, despite the recent attention to lead in drinking water resulting from the Flint water crisis (US) and similar events elsewhere, only one study (from the Netherlands) quantified the health burden from lead (de Hollander et al., 1999). That study, published in 1999, estimated that 11 in 100,000 children were losing 1-3 IQ points as a result of neurocognitive damage from lead in drinking water.

The largest reported fatality risks from chemical contaminants were attributed to arsenic and disinfection byproducts (Figure 2b; Figure 3b), though these risks were very low. Across studies, about 0.1–1.3 premature annual deaths per 100,000 people were attributed to these contaminants. A 2017 study from North Carolina (US) attributed fewer than 0.0015 excess annual deaths per 100,000 people to gross alpha radiation, volatile organic compounds, and pesticides in drinking water (DeFelice et al., 2017).

4. Discussion

We found 23 studies that estimated the burden of infectious diseases and 9 that estimated the chemical disease burden—and one study that estimated both—from drinking water contamination in countries with reportedly high access to safely managed drinking water. Among the infectious disease burden studies, nearly all focused exclusively on GI; the average burden across studies was approximately 3,529 excess GI cases per 100,000 people yearly (Table 1; Figure 1). Of note, the range in burden estimates was large (40-18,250 cases per 100,000 population), which may be attributed to variability in drinking water sources, study populations, and/or estimation methods. Among studies of non-infectious diseases attributable to chemical contamination, the majority focused on carcinogens. The excess cancer risk across studies averaged 1.8 cases per 100,000 people yearly (ranging from less than one in one billion to two cases per 100,000 people yearly) (Table 2; Figure 2).

These results suggest that risks of GI from drinking water contamination are low in many high-income countries relative to those in developing nations, but that the risks are still higher than recommended targets established by the WHO's Guidelines for Drinking Water Quality (World Health Organisation, 2017). The Institute of Health Metrics and Evaluation estimates that globally, the burden of disease from unsafe water is 922 annual disability-adjusted life years (DALYs) per 100,000 people, with the highest burden in sub-Saharan Africa, at 2,813 annual DALYs per 100,000 people ("GBD 2019 Resources | Institute for Health Metrics and Evaluation," n.d.). Using a disability weight of 0.074 for mild diarrhoea as in recent global studies, (Salomon et al., 2015) the population-weighted average 3,529 annual GI cases per 100,000 people equates to approximately 260 annual DALYs per 100,000 people, or less than one-third the global average and about 90% less than in the most affected regions. However, the WHO defines "the tolerable burden of disease . . . as an upper limit of 10^{-6} DALY per person per year," equivalent to 0.1 per 100,000 people. By this measure, the burden of infectious diseases associated with drinking water contamination is still much higher than desired, even in high-income nations and even for the lowest risk estimates identified in this review (40 annual GI cases per 100,000 people, equivalent to 3 DALYs per 100,000).

231
Cancer risk estimates also suggest the magnitude of the disease burden associated with drinking
water contamination is relatively low but still exceeds the WHO's recommendation of one
excess annual cancer case per 100,000 people attributed to drinking water. Across studies, the
average cancer risk was about twice this amount.

3.3 Comparison of Approaches Across Studies
Broadly, study methods can be classified as "top-down" or "bottom up" (Tables 1-2).

3.3.1. Top-Down Approach
Unlike foodborne illness surveillance, a robust system of waterborne illness surveillance does
not exist. In the absence of such a system, some studies use overall disease surveillance data.
They estimate the fraction of diseases that may be attributable to water and multiply this fraction
by the total number of relevant diseases in the population. The WHO uses this method for its
periodic global assessments of the contribution of various risk factors to population health
(Steenland and Armstrong, 2006).

Top-down studies have used a range of methods to estimate the attributable fraction. Some use
previous epidemiologic studies, combined with estimates of the population fraction exposed to
various contamination levels. For example, Colford et al. (Colford et al., 2006) and Murphy et al.
(H.M. Murphy et al., 2016b) used data from epidemiologic studies in which participants were
randomized to receive either an active or sham water treatment system. DeFelice et al. (DeFelice
et al., 2017) obtained information on the distribution of occurrence and concentration of all
regulated chemical contaminants in public water systems from state and federal water quality
surveillance data. Fractions of illnesses attributable to these exposures were estimated by
matching exposure concentrations to relative risks of illness from previous epidemiologic
studies.

An alternative top-down method, used by Gibney et al. (Gibney et al., 2017), relies on elicitation
of expert judgments. Gibney et al. relied on 13 expert opinions on the fractions of illness from
five different waterborne pathogens. There was wide variation in expert views. For example, for
noroviruses, experts' median assessments of the attributable fraction ranged from 2% to 50%.

Another top-down approach is population intervention modeling, as illustrated in two studies by
DeFelice et al. (DeFelice et al., 2015, 2016). In this approach, disease surveillance data are
combined with water source and water quality monitoring data, and a regression model
estimating the disease incidence rate as a function of these variables (along with additional
controls) is fitted. The number of cases attributable to water contamination is estimated from the
regression model by comparing the baseline (observed) rate to the expected rate if water quality
were improved.

3.3.2. Bottom-Up Approach
The bottom-up approach combines data on microbial or chemical contaminant exposure
concentrations with dose-response information from epidemiologic or toxicologic studies. The
dose-response functions are used to predict the estimated probability of illnesses for specific
exposure concentrations. Those probabilities are then multiplied by the size of the exposed

277 population. A drawback of this approach is that the estimated number of attributable cases could
278 exceed the total number of actual cases—a potential error that the top-down approach avoids. On
279 the other hand, in the absence of sufficiently comprehensive surveillance data, this is the only
280 alternative (Murphy et al., 2014).
281
Among studies of the disease burden associated with microbial contamination of drinking water,
the bottom-up approach was illustrated by Murphy et al. (H.M. Murphy et al., 2016c), Burch et
al. (Burch et al., 2021a), Reynolds et al. (Reynolds et al., 2008), and DeFelice et al. (DeFelice et
al., 2015) For example, Murphy et al. used quantitative microbial risk assessment to estimate the
disease burden from five pathogens in private wells and small community systems in Canada.
Burch *et al.* used well water monitoring data for eight pathogens, along with published dose-
response information, to characterize the burden from these pathogens among residents of
Kewaunee County, Wisconsin (US), using private wells. Reynolds *et al.* estimated the disease
burden attributable to microbial contaminants in US drinking water by making assumptions
about the prevalence and concentrations of selected pathogens in different water source types
and then using literature-derived dose-response information to estimate illness probabilities from
these assumed exposures. DeFelice et al. estimated the burden of GI from three reference
pathogens in North Carolina community water systems using monitoring data on the occurrence
of *E. coli* and a literature review of the ratio of concentrations of each reference pathogen to *E.*
coli.
297
The bottom-up approach is also common in studies quantifying the disease burden from
chemical contaminants. Examples include studies of the disease burden from fluoride (Abtahi et
al., 2019) and disinfection byproducts (Dobaradaran et al., 2020) in drinking water in Iran and
characterization of the disease burden from arsenic in drinking water in the US (Greco et al.,
2019). These studies begin by estimating the probability distribution of contaminant exposure
across the study population, and they use a dose-response function derived from previous animal
or human studies to predict the number of adverse health outcomes associated with each
exposure dose.
306
Overall, this review indicates that the lack of robust surveillance systems and epidemiological
data linking drinking water to adverse health outcomes, particularly for chemical contaminants.
The lack of data hinders estimation of the burden of disease attributable to drinking water
contamination. A variety of approaches has been used to develop estimates in the face of limited
data, but there is no single, agreed-upon approach. This methodological uncertainty complicates
comparisons across geographic regions and study years.
313
314 4.1. Limitations of the Available Studies
315
This review identified critical gaps in efforts to characterize the disease burden attributable to
drinking water contamination in countries with high access to safely managed drinking water.
These gaps include a lack of country representation, insufficient spatial resolution in countries
with available studies, a limited scope of contaminants considered, and insufficient information
to support decision-making about the best ways to decrease the disease burden. Although we
attempted to recover estimates from grey literature, only two from New Zealand were found.
There may be other unpublished estimates that this review did not recover.

4.1.1. Limited Geographic Representation

The published disease burden studies covered only 10 of 64 countries/territories reporting ≥90% access to safely managed drinking water. One study covered disease burden in the European Union. More than half of studies are from the United States or Canada. As such, the results of this review may be heavily biased toward these countries.

4.1.2. Lack of estimates focusing on specific sub-populations

Even in countries with available disease burden estimates, the estimates are produced at a large scale (i.e., country-wide) and therefore are insufficient to support decision-making about allocation of resources for infrastructure improvements. Such allocation decisions typically begin at the local level and therefore require local-level estimates highlighting populations most in need of assistance. Most available studies provide estimates for large geographic regions (such as an entire country or a large political subdivision within the country), not for smaller spatial units or special subpopulations. The high risks that some populations may face are subsumed when they are averaged in with the general population receiving high-quality water services. Among studies identified in this review, only Haass et al. (Haass et al., 1996) and Beaudeau et al. (Beaudeau et al., 2014) looked at specific vulnerable populations. Haass et al. examined the burden associated with communities in El Paso, Texas (US), inadequately served by water and sanitation services. Beaudeau et al. estimated burden in elderly populations in Massachusetts (US).

As an example, small water systems experience water quality and service delivery problems more often than larger systems due to the lack of economies of scale (Allaire et al., 2018; Cretikos et al., 2009; McFarlane and Harris, 2018). This is of particular concern when considering that small water systems frequently serve rural, low-income communities. Allaire et al. found that violations of health-based drinking water standards in the US often occur in low-income rural areas (Allaire et al., 2018). A cross-sectional study of water quality in small and medium water systems in rural Alabama found an association between self-reported GI symptoms and the following water system characteristics: respondent-reported water supply interruption, low water pressure, lack of total chlorine in water, and detection of *E. coli* in water (Stauber et al., 2016).

Water systems serving communities of color also are more likely to experience water quality problems, with a consequent disease burden that can be masked in studies lacking sufficient spatial resolution. In Canada, boil-water advisories for systems serving Indigenous communities are common—typically a result of inadequate disinfection, mechanical failure, and insufficient technical expertise of system operators (Lui, 2015; Murphy et al., 2015; H.M. Murphy et al., 2016a). A scoping review of drinking water quality and health outcomes of Canadian Indigenous communities found various reports of adverse health outcomes (mostly GI) associated with drinking water but highlighted the paucity of relevant research (Bradford et al., 2016). The remote, Indigenous communities of Australia may also be at higher risk of infection with waterborne pathogens. The water supply in many of these communities often fails to meet microbiological water quality standards (OAG, 2015) and Ng-Hublin et al. found that the notification rate of cryptosporidiosis in Aboriginal people was 50 times that among non-Aboriginal people in Western Australia (Ng-Hublin et al., 2017). Schaider et al. found a positive

association across the United States between the proportion of Hispanic residents and nitrate
levels in drinking water (Schaider et al., 2019). Balasz et al. found a similar association in the
San Joaquin Valley of California for small community water systems serving a large proportion
of Latino residents (Balazs et al., 2011). A study by Nigra and Navas-Acien found that
incarcerated people in the southwestern US were disproportionately at risk of exposure to arsenic
in drinking water (Nigra and Navas-Acien, 2020). MacDonald Gibson et al. found that in Wake
County, North Carolina (US), children in households relying on private wells had higher blood
lead levels than those with community water service and that these differences were especially
acute in peri-urban minority communities (MacDonald Gibson et al., 2020).

4.1.3. Limited Scope of Disease Types and Contaminants

Nearly all the studies of the waterborne infectious disease burden consider only GI, overlooking
other critical diseases (such as respiratory illnesses) that can be transmitted through drinking
water. For example, only three studies identified in this review (Collier et al., 2020;
Verhougstraete et al., 2020; Vinson, 2012) characterized the respiratory disease burden
attributable to *Legionella*, even though it caused all deaths from waterborne disease outbreaks in
the United States in 2013–2014 (Benedict et al., 2017).

More information on the comparative disease burden from chemical contaminants is especially
important for developed nations since these contaminants tend to be the major focus of drinking
water regulations in developed nations (DeFelice et al., 2017; Roberson, 2011). Especially
notable is the dearth of studies estimating the disease burden from lead in drinking water, despite
recent water crises brought about by lead (for example, in Flint, Michigan, US), along with
mounting evidence that exposure to lead in private well water may be much more prevalent than
previously recognized and, in some cases, can be a source of health disparities (Hanna-Attisha et
al., 2016; MacDonald Gibson et al., 2020). This review located only one study that quantified the
burden of disease due to lead in drinking water in a developed nation, and it was completed more
than 20 years ago. This finding is surprising, since lead is ubiquitous in plumbing (even in
plumbing branded as "lead-free") (Katner et al., 2016). Updated estimates of lead exposure risks
from drinking water and the associated disease burden are needed across developed nations.

Also needed are studies of the disease burden attributable to "emerging" chemical
contaminants—that is, those for which knowledge on their potential prevalence in drinking water
is relatively new. One example is per- and polyfluoroalkyl substances (PFAS), which have been
reported in drinking water sources and systems globally in recent years (Domingo and Nadal,
2019). Another example is toxins formed by aquatic fungi and algae, such as mycotoxins and
cyanotoxins (Székács, 2021). The most recently published waterborne disease surveillance data
from the US Centers for Disease Control and Prevention indicate that cyanotoxins caused 12% of
reported US waterborne disease cases (Benedict et al., 2017).

4.1.4. Insufficient Consideration of Causal Factors

Also lacking is information that would enable estimates of specific water system deficiencies
causing preventable illness. Available estimates focus on total disease burden, not on
contributing factors. These might include source water contamination, treatment system
deficiencies or failures, distribution system breaches, disinfectant loss in the distribution system,

water shutoffs and low-pressure events, or release or formation of contaminants as water travels
through plumbing within buildings.
417
418 *4.1.5 Strengths and Limitations*
419
This review is a first attempt to summarize the global evidence for waterborne microbial and
chemical disease burdens in high-income countries. Current estimates (Vos et al., 2020) using
alternative methods have assumed substantially lower attributable burdens where water
infrastructure coverage is high. Our review shows that these estimates are likely to be too
conservative. Where burden estimates exist, they are typically much higher, even for the very
limited range of contaminants that have been the focus of burden estimates. Our review
additionally reveals that there are major gaps in burden estimation for high-income countries,
suggesting that it is a lack of data – not lack of burden – that has led to a perception that
economically advanced countries no longer face waterborne diseases. This perception may lead
to underinvestment in infrastructure.
430
Despite our review providing a useful summary of evidence, our analysis comes with clear
limitations. First, we were limited by our reliance on the WHO/UNICEF definition of safely
managed drinking water. These estimates of the proportion of the population with safely
managed water may not be accurate, though they are the only global estimates of infrastructure
coverage and are widely used to track progress in meeting the Sustainable Development Goals.
As one example, according to the WHO/UNICEF, 99% of the US population has access to safely
managed drinking water but nearly 13% of the US population relies on federally unregulated
private wells for drinking water (Dieter et al., 2018). Households in the US are responsible for
monitoring and maintaining their own wells, and thus, the safety of these water supplies remains
unknown. One major issue identified by this review was the disparity between the perception of
near universal access to safely managed drinking water in high-income countries (as indicated by
the WHO/UNICEF definition) and the blind spots in our understanding of water quality for a
sizeable proportion of the population within these high-income countries. This disconnect may
be perpetuating the information gaps about the burden of disease attributed to drinking water in
these contexts.
446
Second, this review focused on areas where ≥90% of the population had access to safely
managed drinking water, which meant that, up to 10% of the population in countries/territories
included in the review did not have access to safely managed drinking water. Many of the burden
estimates did not differentiate by water source (surface water vs. groundwater) or system (public
vs. private) while others did. Comparisons of burden are complicated by the potential variations
in water quality within a country depending on drinking water source. This further highlights the
need for higher resolution disease burden data for high-income countries.
454
455 A third limitation includes the low number of available studies, and their narrow range of
456 contaminants and the limited diversity of their methods, which makes synthesis impossible. We
457 also limited our review to studies in English, which may have eliminated burden estimates from
458 many countries.
459

Limitations notwithstanding, we conducted a rigorous systematic review of the literature to
identify and compare estimates of burden attributed to drinking water contamination in countries
perceived to provide near universal access to safely managed drinking water. The major finding
and takeaway of this review is precisely the lack of burden estimates available. The dearth of
these estimates in countries with robust water infrastructure coupled with the evidence of water
quality issues for subpopulations of these countries and the limitations of relying on
WHO/UNICEF definitions of safely managed drinking water may suggest that the true burden of
disease attributed to drinking water contamination may be greater in these countries than
previously recognized.
469
470
5. Conclusions
This review sheds light on the dearth of research on the disease burden attributable to unsafe
drinking water in countries with reportedly high access to safe drinking water. Estimating this
disease burden remains difficult given the lack of a comprehensive surveillance system and gaps
in drinking water quality data. WHO/UNICEF data suggesting the vast majority of the
population has access to safely managed water can be misleading because these data mask
variation in the quality of water service delivery that can have important public health
consequences. Further research on the drinking water-attributed disease burden in these countries
remains vital to identifying the main drivers of waterborne disease, the populations at greatest
risk, and strategies for improving health outcomes related to drinking water.
481

482 **Contributors**
483 HMM, JMG, and JB designed the study. DL, JH, JMG, and HMM recovered articles, assessed
484 article eligibility, and extracted and analyzed data from articles. DL, JMG, and HMM wrote the
485 initial drafts of the manuscript. DL, JH, JMG, HMM, and JB contributed to edits of the
486 manuscript.
487

488 **Data sharing**
489 Extracted data are available on request to the corresponding author.
490

491 **Acknowledgements**
492 We would like to thank Jessica Evans, Ilya Law (IL), Tala Alahdab (TA) for their help screening
493 titles and abstracts for this review.
494
Funding
JMG was supported in part by the U.S. National Science Foundation (CMMI-2017207). HMM,
JH and IL were supported in part by the Canada Research Chairs Program (Grant # 950-232787).
DL was supported in part by Pennsylvania Department of Health's Pennsylvania Commonwealth
Universal Research Enhancement Program (PA CURE) Formula Funds 2019 (Grant #
4100083099 to Temple University). TA was supported in part by the University of Guelph's
Undergraduate Research Assistant Program. Publication costs were covered by the UK Foreign
Commonwealth and Development Office.
503

504 **Protocol Registration**
505 A protocol for this review was not formally registered.

Competing Interests

The authors have no competing interests to declare.

References

Abtahi, M., Dobaradaran, S., Jorfi, S., Koolivand, A., Khaloo, S.S., Spitz, J., Saeedi, H., Golchinpour, N., Saeedi, R., 2019. Age-sex specific disability-adjusted life years (DALYs) attributable to elevated levels of fluoride in drinking water: A national and subnational study in Iran, 2017. WATER Res. 157, 94–105. https://doi.org/10.1016/j.watres.2019.03.087

Allaire, M., Wu, H., Lall, U., 2018. National trends in drinking water quality violations. Proc. Natl. Acad. Sci. U. S. A. 115, 2078–2083. https://doi.org/10.1073/pnas.1719805115

Anderson, M.W., 2008. Cities inside out: race, poverty, and exclusion at the urban fringe. UCLA Law Rev. 55, 1095–1160. https://doi.org/10.1111/j.1467-8276.2009.01305.x

Balazs, C., Morello-Frosch, R., Hubbard, A., Ray, I., 2011. Social disparities in nitrate-contaminated drinking water in California's San Joaquin Valley. Environ. Health Perspect. 119, 1272–1278. https://doi.org/10.1289/ehp.1002878

Balazs, C.L., Ray, I., 2014. The drinking water disparities framework: on the origins and persistence of inequities in exposure. Am. J. Public Health 104, 603–611. https://doi.org/10.2105/AJPH.2013.301664

Ball, A., 2006. Estimation of the burden of water-borne disease in New Zealand: preliminary report. Christchurch Environ. Sci. Res. Ltd.

Beaudeau, P., Schwartz, J., Levin, R., 2014. Drinking water quality. and hospital admissions of elderly people for gastrointestinal illness in Eastern Massachusetts, 1998-2008. WATER Res. 52, 188–198. https://doi.org/10.1016/j.watres.2014.01.005

Benedict, K.M., Reses, H., Vigar, M., Roth, D.M., Roberts, V.A., Mattioli, M., Cooley, L.A., Hilborn, E.D., Wade, T.J., Fullerton, K.E., Yoder, J.S., Hill, V.R., 2017. Surveillance for Waterborne Disease Outbreaks Associated with Drinking Water — United States, 2013–2014. MMWR. Morb. Mortal. Wkly. Rep. 66, 1216–1221. https://doi.org/10.15585/mmwr.mm6644a3

Bradford, L.E.A., Bharadwaj, L.A., Okpalauwaekwe, U., Waldner, C.L., 2016. Drinking water quality in Indigenous communities in Canada and health outcomes: a scoping review. Int. J. Circumpolar Health 75. https://doi.org/10.3402/ijch.v75.32336

Burch, T.R., Stokdyk, J.P., Spencer, S.K., Kieke, B.A., Firnstahl, A.D., Muldoon, M.A., Borchardt, M.A., 2021a. Quantitative Microbial Risk Assessment for Contaminated Private Wells in the Fractured Dolomite Aquifer of Kewaunee County, Wisconsin. Environ. Health Perspect. 129. https://doi.org/10.1289/EHP7815

Burch, T.R., Stokdyk, J.P., Spencer, S.K., Kieke, B.A., Firnstahl, A.D., Muldoon, M.A., Borchardt, M.A., 2021b. Quantitative Microbial Risk Assessment for Contaminated Private Wells in the Fractured Dolomite Aquifer of Kewaunee County, Wisconsin. Environ. Health Perspect. 129. https://doi.org/10.1289/EHP7815

Butler, A.J., Pintar, K.D.M., Thomas, M.K., 2016. Estimating the Relative Role of Various Subcategories of Food, Water, and Animal Contact Transmission of 28 Enteric Diseases in Canada. Foodborne Pathog. Dis. 13, 57–64. https://doi.org/10.1089/fpd.2015.1957

Colford, J.M., Roy, S., Beach, M.J., Hightower, A., Shaw, S.E., Wade, T.J., 2006. A review of household drinking water intervention trials and an approach to the estimation of endemic

waterborne gastroenteritis in the United States. J. Water Health 4, 71–88.

Collier, S.A., Deng, L., Adam, E.A., Benedict, K.M., Beshearse, E.M., Blackstock, A.J., Bruce,
B.B., Derado, G., Edens, C., Fullerton, K.E., 2020. Estimate of burden and direct healthcare
cost of infectious waterborne disease in the United States.

Cretikos, M., Byleveld, P., Durrheim, D.N., Porigneaux, P., Merritt, T., Leask, S., 2009. Supply
system factors associated with microbiological drinking water safety in regional New South
Wales, Australia, 2001–2007. J. Water Health 8, 257–268.
https://doi.org/10.2166/wh.2009.203

de Hollander, A.E.M., Melse, J.M., Lebret, E., Kramers, P.G.N., 1999. An aggregate public
health indicator to represent the impact of multiple environmental exposures. Epidemiology
10, 606–617. https://doi.org/10.1097/00001648-199909000-00030

DeFelice, N.B., Johnston, J.E., MacDonald Gibson, J., 2016. Reducing Emergency Department
Visits for Acute Gastrointestinal Illnesses in North Carolina (USA) by Extending
Community Water Service. Environ. Health Perspect. 124, 1583–1591.
https://doi.org/10.1289/EHP160

DeFelice, N.B., Johnston, J.E., MacDonald Gibson, J., 2015. Acute gastrointestinal illness risks
in North Carolina community water Systems: a methodological comparison. Environ. Sci.
Technol. 49, 10019–10027. https://doi.org/10.1021/acs.est.5b01898

DeFelice, N.B., Leker, H.G., MacDonald Gibson, J., 2017. Annual cancer risks from chemicals
in North Carolina community water systems. Hum. Ecol. Risk Assess. 23, 974–991.
https://doi.org/10.1080/10807039.2017.1292842

Deshpande, A., Miller-Petrie, M.K., Lindstedt, P.A., Baumann, M.M., Johnson, K.B., Blacker,
B.F., Abbastabar, H., Abd-Allah, F., Abdelalim, A., Abdollahpour, I., 2020. Mapping
geographical inequalities in access to drinking water and sanitation facilities in low-income
and middle-income countries, 2000–17. Lancet Glob. Heal. 8, e1162–e1185.

Dieter, C.A., Maupin, M.A., Caldwell, R.R., Harris, M.A., Ivahnenko, T.I., Lovelace, J.K.,
Barber, N.L., Linsey, K.S., 2018. Estimated use of water in the United States in 2015,
Circular. Reston, VA. https://doi.org/10.3133/cir1441

Dobaradaran, S., Fard, E.S., Tekle-Rottering, A., Keshtkar, M., Karbasdehi, V.N., Abtahi, M.,
Gholamnia, R., Saeedi, R., 2020. Age-sex specific and cause-specific health risk and burden
of disease induced by exposure to trihalomethanes (THMs) and haloacetic acids (HAAs)
from drinking water: An assessment in four urban communities of Bushehr Province, Iran,
2017. Environ. Res. 182. https://doi.org/10.1016/j.envres.2019.109062

Domingo, J.L., Nadal, M., 2019. Human exposure to per- and polyfluoroalkyl substances (PFAS)
through drinking water: A review of the recent scientific literature. Environ. Res. 177,
108648. https://doi.org/https://doi.org/10.1016/j.envres.2019.108648

Eisenberg, J.N.S., Hubbard, A., Wade, T.J., Sylvester, M.F., LeChevallier, M.W., Levy, D.A.,
Colford, J.M., 2006. Inferences Drawn from a Risk Assessment Compared Directly with a
Randomized Trial of a Home Drinking Water Intervention. Environ. Health Perspect. 114,
1199–1204. https://doi.org/10.1289/ehp.8682

Evlampidou, I., Font-Ribera, L., Rojas-Rueda, D., Gracia-Lavedan, E., Costet, N., Pearce, N.,
Vineis, P., Jaakkola, J.J.K., Delloye, F., Makris, K.C., Stephanou, E.G., Kargaki, S.,
Kozisek, F., Sigsgaard, T., Hansen, B., Schullehner, J., Nahkur, R., Galey, C., Zwiener, C.,
Vargha, M., Righi, E., Aggazzotti, G., Kalnina, G., Grazuleviciene, R., Polanska, K.,
Gubkova, D., Bitenc, K., Goslan, E.H., Kogevinas, M., Villanueva, C.M., 2020.
Trihalomethanes in Drinking Water and Bladder Cancer Burden in the European Union.

Environ. Health Perspect. 128. https://doi.org/10.1289/EHP4495

Fewtrell, L., Smith, S., Kay, D., Bartram, J., 2006. An attempt to estimate the global burden of
disease due to fluoride in drinking water. J. Water Health 4, 533–542.

GBD 2019 Resources | Institute for Health Metrics and Evaluation [WWW Document], n.d.

Gibney, K.B., O'Toole, J., Sinclair, M., Leder, K., 2017. Burden of Disease Attributed to
Waterborne Transmission of Selected Enteric Pathogens, Australia, 2010. Am. J. Trop.
Med. Hyg. 96, 1400–1403. https://doi.org/10.4269/ajtmh.16-0907

Greco, S.L., Belova, A., Haskell, J., Backer, L., 2019. Estimated burden of disease from arsenic
in drinking water supplied by domestic wells in the United States. J. Water Health 17, 801–
812. https://doi.org/10.2166/wh.2019.216

Haass, J.A., Miller, G.L., Haddix, A.C., Nickey, L.N., Sinks, T., 1996. An economic analysis of
water and sanitation infrastructure improvements in the Colonias of EI Paso County, Texas.
Int. J. Occup. Environ. Health 2, 211–221.

Hanna-Attisha, M., LaChance, J., Sadler, R.C., Schnepp, A.C., 2016. Elevated blood lead levels
in children associated with the Flint drinking water crisis: A spatial analysis of risk and
public health response. Am. J. Public Health 106, 283–290.
https://doi.org/10.2105/AJPH.2015.303003

Higgins, J., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M., Welch, V., 2019. Cochrane
Handbook for Systematic Reviews of Interventions, 2nd ed. John Wiley & Sons,
Chichester, UK.

Katner, A., Pieper, K.J., Lambrinidou, Y., Brown, K., Hu, C.-Y., Mielke, H.W., Edwards, M.A.,
2016. Weaknesses in Federal Drinking Water Regulations and Public Health Policies that
Impede Lead Poisoning Prevention and Environmental Justice. Environ. Justice 9, 109–117.
https://doi.org/10.1089/env.2016.0012

Kim, J.H., Cheong, H.K., Jeon, B.H., 2018. Burden of Disease Attributable to Inadequate
Drinking Water, Sanitation, and Hygiene in Korea. J. KOREAN Med. Sci. 33.
https://doi.org/10.3346/jkms.2018.33.e288

Lui, E., 2015. On notice for a drinking water crisis in Canada. Council of Canadians Ottawa.

MacDonald Gibson, J., Fisher, M., Clonch, A., MacDonald, J., Cook, P., 2020. Children drinking
private well water have higher blood lead than those with city water. Proc. Natl. Acad. Sci.
117, 16898–16907. https://doi.org/10.1073/pnas.2002729117

MacDonald Gibson, J., Thomsen, J., Launay, F., Harder, E., DeFelice, N.B., 2013. Deaths and
medical visits attributable to environmental pollution in the United Arab Emirates. PLoS
One 8, e57536–e57536. https://doi.org/10.1371/journal.pone.0057536

Masciopinto, C., De Giglio, O., Scrascia, M., Fortunato, F., La Rosa, G., Suffredini, E., Pazzani,
C., Prato, R., Montagna, M.T., 2019. Human health risk assessment for the occurrence of
enteric viruses in drinking water from wells: Role of flood runoff injections. Sci. Total
Environ. 666, 559–571. https://doi.org/10.1016/j.scitotenv.2019.02.107

Mathewson, P.D., Evans, S., Byrnes, T., Joos, A., Naidenko, O. V, 2020. Health and economic
impact of nitrate pollution in drinking water: a Wisconsin case study. Environ. Monit.
Assess. 192, 1–18.

McFarlane, K., Harris, L.M., 2018. Small systems, big challenges: review of small drinking
water system governance. Environ. Rev. 26, 378+.

Messner, M., Shaw, S., Regli, S., Rotert, K., Blank, V., Soller, J., 2006. An approach for
developing a national estimate of waterborne disease due to drinking water and a national
estimate model application. J. Water Health 4, 201–240.

Moore, D., Black, M., Valji, Y., Tooth, R., 2010. Cost Benefit Analysis of Raising the Quality of
New Zealand Networked Drinking Water.
Morris, J.C., 2017. Planning for Water Infrastructure: Challenges and Opportunities. Public
Work. Manag. Policy 22, 24–30. https://doi.org/10.1177/1087724X16668182
Morris, R.D., Levin, R., 1995. Estimating the incidence of waterborne infectious disease related
to drinking water in the United States. IAHS Publ. Proc. Reports-Intern Assoc Hydrol. Sci.
233, 75–88.
Murphy, H.M., Bhatti, M., Harvey, R., McBean, E.A., 2016a. Using Decision Trees to Predict
Drinking Water Advisories in Small Water Systems. J. AWWA 108, E109–E118.
https://doi.org/https://doi.org/10.5942/jawwa.2016.108.0008
Murphy, H.M., Corston-Pine, E., Post, Y., McBean, E.A., 2015. Insights and opportunities:
challenges of Canadian First Nations drinking water operators. Int. Indig. Policy J. 6.
Murphy, H.M., Pintar, K.D.M., McBean, E.A., Thomas, M.K., 2014. A systematic review of
waterborne disease burden methodologies from developed countries. J. Water Health 12,
634–655. https://doi.org/http://dx.doi.org/10.2166/wh.2014.049
Murphy, H.M., Thomas, M.K., Medeiros, D.T., McFadyen, S., Pintar, K.D.M., 2016b.
Estimating the number of cases of acute gastrointestinal illness (AGI) associated with
Canadian municipal drinking water systems. Epidemiol. Infect. 144, 1371–1385.
https://doi.org/10.1017/S0950268815002083
Murphy, H.M., Thomas, M.K., Schmidt, P.J., Medeiros, D.T., McFadyen, S., Pintar, K.D.M.,
2016c. Estimating the burden of acute gastrointestinal illness due to Giardia,
Cryptosporidium, Campylobacter, E. coli O157 and norovirus associated with private wells
and small water systems in Canada. Epidemiol. Infect. 144, 1355–1370.
https://doi.org/10.1017/S0950268815002071
Ng-Hublin, J.S.Y., Combs, B., Reid, S., Ryan, U., 2017. Differences in the occurrence and
epidemiology of cryptosporidiosis in Aboriginal and non-Aboriginal people in Western
Australia (2002−2012). Infect. Genet. Evol. 53, 100–106.
https://doi.org/https://doi.org/10.1016/j.meegid.2017.05.018
Nigra, A.E., Navas-Acien, A., 2020. Arsenic in US correctional facility drinking water, 2006–
2011. Environ. Res. 188, 109768.
https://doi.org/https://doi.org/10.1016/j.envres.2020.109768
OAG, 2015. Delivering Essential Services to Remote Aboriginal Communities.
Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D.,
Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw,
J.M., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S.,
McGuinness, L.A., Stewart, L.A., Thomas, J., Tricco, A.C., Welch, V.A., Whiting, P.,
Moher, D., 2021. The PRISMA 2020 statement: An updated guideline for reporting
systematic reviews. BMJ 372. https://doi.org/10.1136/BMJ.N71
Payment, P., 1997. Epidemiology of endemic gastrointestinal and respiratory diseases:
Incidence, fraction attributable to tap water and costs to society. WATER Sci. Technol. 35,
7–10. https://doi.org/10.1016/S0273-1223(97)00226-6
Perz, J.F., Ennever, F.K., Le Blancq, S.M., 1998. Cryptosporidium in Tap Water: Comparison of
Predicted Risks with Observed Levels of Disease. Am. J. Epidemiol. 147, 289–301.
https://doi.org/10.1093/oxfordjournals.aje.a009449
Prüss-Ustün, A., Wolf, J., Bartram, J., Clasen, T., Cumming, O., Freeman, M.C., Gordon, B.,
Hunter, P.R., Medlicott, K., Johnston, R., 2019. Burden of disease from inadequate water,

sanitation and hygiene for selected adverse health outcomes: An updated analysis with a focus on low- and middle-income countries. Int. J. Hyg. Environ. Health 222, 765–777. https://doi.org/10.1016/j.ijheh.2019.05.004

Reynolds, K., D Mena, K., Gerba, C., 2008. Risk of Waterborne Illness Via Drinking Water in the United States, Reviews of environmental contamination and toxicology. https://doi.org/10.1007/978-0-387-71724-1_4

Roberson, J.A., 2011. What's next after 40 years of drinking water regulations? Environ. Sci. Technol. 45, 154–60. https://doi.org/10.1021/es101410v

Salomon, J.A., Haagsma, J.A., Davis, A., de Noordhout, C.M., Polinder, S., Havelaar, A.H., Cassini, A., Devleesschauwer, B., Kretzschmar, M., Speybroeck, N., Murray, C.J.L., Vos, T., 2015. Disability weights for the Global Burden of Disease 2013 study. Lancet Glob. Heal. 3, e712–e723. https://doi.org/10.1016/S2214-109X(15)00069-8

Schaider, L.A., Swetschinski, L., Campbell, C., Rudel, R.A., 2019. Environmental justice and drinking water quality: are there socioeconomic disparities in nitrate levels in U.S. drinking water? Environ. Heal. 18, 3. https://doi.org/10.1186/s12940-018-0442-6

Stauber, C.E., Wedgworth, J.C., Johnson, P., Olson, J.B., Ayers, T., Elliott, M., Brown, J., 2016. Associations between Self-Reported Gastrointestinal Illness and Water System Characteristics in Community Water Supplies in Rural Alabama: A Cross-Sectional Study. PLoS One 11. https://doi.org/10.1371/journal.pone.0148102

Steenland, K., Armstrong, B., 2006. An overview of methods for calculating the burden of disease due to specific risk factors. Epidemiology 17, 512–9. https://doi.org/10.1097/01.ede.0000229155.05644.43

Székács, A., 2021. Mycotoxins as emerging contaminants. Introduction to the special issue "rapid detection of mycotoxin contamination." Toxins (Basel). https://doi.org/10.3390/toxins13070475

Temkin, A., Evans, S., Manidis, T., Campbell, C., Naidenko, O. V, 2019. Exposure-based assessment and economic valuation of adverse birth outcomes and cancer risk due to nitrate in United States drinking water. Environ. Res. 176, 108442. https://doi.org/https://doi.org/10.1016/j.envres.2019.04.009

Verhougstraete, M., Reynolds, K.A., Pearce-Walker, J., Gerba, C., 2020. Cost-benefit analysis of point-of-use devices for health risks reduction from pathogens in drinking water. J. Water Health 18, 968–982. https://doi.org/10.2166/wh.2020.111

Vinson, N.G., 2012. Towards estimating the economic burden of waterborne illness in Canada: what do we know, where do we go, in: Poster Presented at The Ontario Public Health Convention (TOPHC), Toronto, ON, Canada. pp. 2–4.

Vos, T., Lim, S.S., Abbafati, C., Abbas, K.M., Abbasi, M., Abbasifard, M., Abbasi-Kangevari, M., Abbastabar, H., Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abolhassani, H., Aboyans, V., Abrams, E.M., Abreu, L.G., Abrigo, M.R.M., Abu-Raddad, L.J., Abushouk, A.I., Acebedo, A., Ackerman, I.N., Adabi, M., Adamu, A.A., Adebayo, O.M., Adekanmbi, V., Adelson, J.D., Adetokunboh, O.O., Adham, D., Afshari, M., Afshin, A., Agardh, E.E., Agarwal, G., Agesa, K.M., Aghaali, M., Aghamir, S.M.K., Agrawal, A., Ahmad, T., Ahmadi, A., Ahmadi, M., Ahmadieh, H., Ahmadpour, E., Akalu, T.Y., Akinyemi, R.O., Akinyemiju, T., Akombi, B., Al-Aly, Z., Alam, K., Alam, N., Alam, S., Alam, T., Alanzi, T.M., Albertson, S.B., Alcalde-Rabanal, J.E., Alema, N.M., Ali, M., Ali, S., Alicandro, G., Alijanzadeh, M., Alinia, C., Alipour, V., Aljunid, S.M., Alla, F., Allebeck, P., Almasi-Hashiani, A., Alonso, J., Al-Raddadi, R.M., Altirkawi, K.A., Alvis-

Guzman, N., Alvis-Zakzuk, N.J., Amini, S., Amini-Rarani, M., Aminorroaya, A., Amiri, F.,

Amit, A.M.L., Amugsi, D.A., Amul, G.G.H., Anderlini, D., Andrei, C.L., Andrei, T.,

Anjomshoa, M., Ansari, F., Ansari, I., Ansari-Moghaddam, A., Antonio, C.A.T., Antony,

C.M., Antriyandarti, E., Anvari, D., Anwer, R., Arabloo, J., Arab-Zozani, M., Aravkin,

A.Y., Ariani, F., Ärnlöv, J., Aryal, K.K., Arzani, A., Asadi-Aliabadi, M., Asadi-Pooya,

A.A., Asghari, B., Ashbaugh, C., Atnafu, D.D., Atre, S.R., Ausloos, F., Ausloos, M., Ayala

Quintanilla, B.P., Ayano, G., Ayanore, M.A., Aynalem, Y.A., Azari, S., Azarian, G., Azene,

Z.N., Babaee, E., Badawi, A., Bagherzadeh, M., Bakhshaei, M.H., Bakhtiari, A.,

Balakrishnan, S., Balalla, S., Balassyano, S., Banach, M., Banik, P.C., Bannick, M.S.,

Bante, A.B., Baraki, A.G., Barboza, M.A., Barker-Collo, S.L., Barthelemy, C.M., Barua, L.,

Barzegar, A., Basu, S., Baune, B.T., Bayati, M., Bazmandegan, G., Bedi, N., Beghi, E.,

Béjot, Y., Bello, A.K., Bender, R.G., Bennett, D.A., Bennitt, F.B., Bensenor, I.M.,

Benziger, C.P., Berhe, K., Bernabe, E., Bertolacci, G.J., Bhageerathy, R., Bhala, N.,

Bhandari, D., Bhardwaj, P., Bhattacharyya, K., Bhutta, Z.A., Bibi, S., Biehl, M.H., Bikbov,

B., Bin Sayeed, M.S., Biondi, A., Birihane, B.M., Bisanzio, D., Bisignano, C., Biswas,

R.K., Bohlouli, S., Bohluli, M., Bolla, S.R.R., Boloor, A., Boon-Dooley, A.S., Borges, G.,

Borzì, A.M., Bourne, R., Brady, O.J., Brauer, M., Brayne, C., Breitborde, N.J.K., Brenner,

H., Briant, P.S., Briggs, A.M., Briko, N.I., Britton, G.B., Bryazka, D., Buchbinder, R.,

Bumgarner, B.R., Busse, R., Butt, Z.A., Caetano dos Santos, F.L., Cámera, L.L.A.A.,

Campos-Nonato, I.R., Car, J., Cárdenas, R., Carreras, G., Carrero, J.J., Carvalho, F.,

Castaldelli-Maia, J.M., Castañeda-Orjuela, C.A., Castelpietra, G., Castle, C.D., Castro, F.,

Catalá-López, F., Causey, K., Cederroth, C.R., Cercy, K.M., Cerin, E., Chandan, J.S.,

Chang, A.R., Charlson, F.J., Chattu, V.K., Chaturvedi, S., Chimed-Ochir, O., Chin, K.L.,

Cho, D.Y., Christensen, H., Chu, D.-T., Chung, M.T., Cicuttini, F.M., Ciobanu, L.G.,

Cirillo, M., Collins, E.L., Compton, K., Conti, S., Cortesi, P.A., Costa, V.M., Cousin, E.,

Cowden, R.G., Cowie, B.C., Cromwell, E.A., Cross, D.H., Crowe, C.S., Cruz, J.A.,

Cunningham, M., Dahlawi, S.M.A., Damiani, G., Dandona, L., Dandona, R., Darwesh,

A.M., Daryani, A., Das, J.K., Das Gupta, Rajat, das Neves, J., Dávila-Cervantes, C.A.,

Davletov, K., De Leo, D., Dean, F.E., DeCleene, N.K., Deen, A., Degenhardt, L.,

Dellavalle, R.P., Demeke, F.M., Demsie, D.G., Denova-Gutiérrez, E., Dereje, N.D.,

Dervenis, N., Desai, R., Desalew, A., Dessie, G.A., Dharmaratne, S.D., Dhungana, G.P.,

Dianatinasab, M., Diaz, D., Dibaji Forooshani, Z.S., Dingels, Z. V, Dirac, M.A., Djalalinia,

S., Do, H.T., Dokova, K., Dorostkar, F., Doshi, C.P., Doshmangir, L., Douiri, A., Doxey,

M.C., Driscoll, T.R., Dunachie, S.J., Duncan, B.B., Duraes, A.R., Eagan, A.W., Ebrahimi

Kalan, M., Edvardsson, D., Ehrlich, J.R., El Nahas, N., El Sayed, I., El Tantawi, M.,

Elbarazi, I., Elgendy, I.Y., Elhabashy, H.R., El-Jaafary, S.I., Elyazar, I.R.F., Emamian,

M.H., Emmons-Bell, S., Erskine, H.E., Eshrati, B., Eskandarieh, S., Esmaeilnejad, S.,

Esmaeilzadeh, F., Esteghamati, A., Estep, K., Etemadi, A., Etisso, A.E., Farahmand, M.,

Faraj, A., Fareed, M., Faridnia, R., Farinha, C.S. e S., Farioli, A., Faro, A., Faruque, M.,

Farzadfar, F., Fattahi, N., Fazlzadeh, M., Feigin, V.L., Feldman, R., Fereshtehnejad, S.-M.,

Fernandes, E., Ferrari, A.J., Ferreira, M.L., Filip, I., Fischer, F., Fisher, J.L., Fitzgerald, R.,

Flohr, C., Flor, L.S., Foigt, N.A., Folayan, M.O., Force, L.M., Fornari, C., Foroutan, M.,

Fox, J.T., Freitas, M., Fu, W., Fukumoto, T., Furtado, J.M., Gad, M.M., Gakidou, E.,

Galles, N.C., Gallus, S., Gamkrelidze, A., Garcia-Basteiro, A.L., Gardner, W.M.,

Geberemariyam, B.S., Gebrehiwot, A.M., Gebremedhin, K.B., Gebreslassie, A.A.A.A.,

Gershberg Hayoon, A., Gething, P.W., Ghadimi, M., Ghadiri, K., Ghafourifard, M., Ghajar,

A., Ghamari, F., Ghashghaee, A., Ghiasvand, H., Ghith, N., Gholamian, A., Gilani, S.A.,

Gill, P.S., Gitimoghaddam, M., Giussani, G., Goli, S., Gomez, R.S., Gopalani, S.V., Gorini,

G., Gorman, T.M., Gottlich, H.C., Goudarzi, H., Goulart, A.C., Goulart, B.N.G., Grada, A.,

Grivna, M., Grosso, G., Gubari, M.I.M., Gugnani, H.C., Guimaraes, A.L.S., Guimarães,

R.A., Guled, R.A., Guo, G., Guo, Y., Gupta, Rajeev, Haagsma, J.A., Haddock, B., Hafezi-

Nejad, N., Hafiz, A., Hagins, H., Haile, L.M., Hall, B.J., Halvaei, I., Hamadeh, R.R.,

Hamagharib Abdullah, K., Hamilton, E.B., Han, C., Han, H., Hankey, G.J., Haro, J.M.,

Harvey, J.D., Hasaballah, A.I., Hasanzadeh, A., Hashemian, M., Hassanipour, S.,

Hassankhani, H., Havmoeller, R.J., Hay, R.J., Hay, S.I., Hayat, K., Heidari, B., Heidari, G.,

Heidari-Soureshjani, R., Hendrie, D., Henrikson, H.J., Henry, N.J., Herteliu, C.,

Heydarpour, F., Hird, T.R., Hoek, H.W., Hole, M.K., Holla, R., Hoogar, P., Hosgood, H.D.,

Hosseinzadeh, M., Hostiuc, M., Hostiuc, S., Househ, M., Hoy, D.G., Hsairi, M., Hsieh,

V.C., Hu, G., Huda, T.M., Hugo, F.N., Huynh, C.K., Hwang, B.-F., Iannucci, V.C., Ibitoye,

S.E., Ikuta, K.S., Ilesanmi, O.S., Ilic, I.M., Ilic, M.D., Inbaraj, L.R., Ippolito, H., Irvani,

S.S.N., Islam, M.M., Islam, M., Islam, S.M.S., Islami, F., Iso, H., Ivers, R.Q., Iwu, C.C.D.,

Iyamu, I.O., Jaafari, J., Jacobsen, K.H., Jadidi-Niaragh, F., Jafari, H., Jafarinia, M.,

Jahagirdar, D., Jahani, M.A., Jahanmehr, N., Jakovljevic, M., Jalali, A., Jalilian, F., James,

S.L., Janjani, H., Janodia, M.D., Jayatilleke, A.U., Jeemon, P., Jenabi, E., Jha, R.P., Jha, V.,

Ji, J.S., Jia, P., John, O., John-Akinola, Y.O., Johnson, C.O., Johnson, S.C., Jonas, J.B., Joo,

T., Joshi, A., Jozwiak, J.J., Jürisson, M., Kabir, A., Kabir, Z., Kalani, H., Kalani, R.,

Kalankesh, L.R., Kalhor, R., Kamiab, Z., Kanchan, T., Karami Matin, B., Karch, A., Karim,

M.A., Karimi, S.E., Kassa, G.M., Kassebaum, N.J., Katikireddi, S.V., Kawakami, N.,

Kayode, G.A., Keddie, S.H., Keller, C., Kereselidze, M., Khafaie, M.A., Khalid, N., Khan,

M., Khatab, K., Khater, M.M., Khatib, M.N., Khayamzadeh, M., Khodayari, M.T.,

Khundkar, R., Kianipour, N., Kieling, C., Kim, D., Kim, Y.-E., Kim, Y.J., Kimokoti, R.W.,

Kisa, A., Kisa, S., Kissimova-Skarbek, K., Kivimäki, M., Kneib, C.J., Knudsen, A.K.S.,

Kocarnik, J.M., Kolola, T., Kopec, J.A., Kosen, S., Koul, P.A., Koyanagi, A., Kravchenko,

M.A., Krishan, K., Krohn, K.J., Kuate Defo, B., Kucuk Bicer, B., Kumar, G.A., Kumar, M.,

Kumar, P., Kumar, V., Kumaresh, G., Kurmi, O.P., Kusuma, D., Kyu, H.H., La Vecchia,

C., Lacey, B., Lal, D.K., Lalloo, R., Lam, J.O., Lami, F.H., Landires, I., Lang, J.J.,

Lansingh, V.C., Larson, S.L., Larsson, A.O., Lasrado, S., Lassi, Z.S., Lau, K.M.-M.,

Lavados, P.M., Lazarus, J. V, Ledesma, J.R., Lee, P.H., Lee, S.W.H., LeGrand, K.E., Leigh,

J., Leonardi, M., Lescinsky, H., Leung, J., Levi, M., Lewington, S., Li, S., Lim, L.-L., Lin,

C., Lin, R.-T., Linehan, C., Linn, S., Liu, H.-C., Liu, S., Liu, Z., Looker, K.J., Lopez, A.D.,

Lopukhov, P.D., Lorkowski, S., Lotufo, P.A., Lucas, T.C.D., Lugo, A., Lunevicius, R.,

Lyons, R.A., Ma, J., MacLachlan, J.H., Maddison, E.R., Maddison, R., Madotto, F.,

Mahasha, P.W., Mai, H.T., Majeed, A., Maled, V., Maleki, S., Malekzadeh, R., Malta, D.C.,

Mamun, A.A., Manafi, A., Manafi, N., Manguerra, H., Mansouri, B., Mansournia, M.A.,

Mantilla Herrera, A.M., Maravilla, J.C., Marks, A., Martins-Melo, F.R., Martopullo, I.,

Masoumi, S.Z., Massano, J., Massenburg, B.B., Mathur, M.R., Maulik, P.K., McAlinden,

C., McGrath, J.J., McKee, M., Mehndiratta, M.M., Mehri, F., Mehta, K.M., Meitei, W.B.,

Memiah, P.T.N., Mendoza, W., Menezes, R.G., Mengesha, E.W., Mengesha, M.B., Mereke,

A., Meretoja, A., Meretoja, T.J., Mestrovic, T., Miazgowski, B., Miazgowski, T., Michalek,

I.M., Mihretie, K.M., Miller, T.R., Mills, E.J., Mirica, A., Mirrakhimov, E.M., Mirzaei, H.,

Mirzaei, M., Mirzaei-Alavijeh, M., Misganaw, A.T., Mithra, P., Moazen, B.,

Moghadaszadeh, M., Mohamadi, E., Mohammad, D.K., Mohammad, Y., Mohammad Gholi

Mezerji, N., Mohammadian-Hafshejani, A., Mohammadifard, N., Mohammadpourhodki,
R., Mohammed, S., Mokdad, A.H., Molokhia, M., Momen, N.C., Monasta, L., Mondello,
S., Mooney, M.D., Moosazadeh, M., Moradi, G., Moradi, M., Moradi-Lakeh, M.,
Moradzadeh, R., Moraga, P., Morales, L., Morawska, L., Moreno Velásquez, I., Morgado-
da-Costa, J., Morrison, S.D., Mosser, J.F., Mouodi, S., Mousavi, S.M., Mousavi
Khaneghah, A., Mueller, U.O., Munro, S.B., Muriithi, M.K., Musa, K.I., Muthupandian, S.,
Naderi, M., Nagarajan, A.J., Nagel, G., Naghshtabrizi, B., Nair, S., Nandi, A.K., Nangia,
V., Nansseu, J.R., Nayak, V.C., Nazari, J., Negoi, I., Negoi, R.I., Netsere, H.B.N., Ngunjiri,
J.W., Nguyen, C.T., Nguyen, J., Nguyen, Michele, Nguyen, Minh, Nichols, E., Nigatu, D.,
Nigatu, Y.T., Nikbakhsh, R., Nixon, M.R., Nnaji, C.A., Nomura, S., Norrving, B., Noubiap,
J.J., Nowak, C., Nunez-Samudio, V., Oţoiu, A., Oancea, B., Odell, C.M., Ogbo, F.A., Oh,
I.-H., Okunga, E.W., Oladnabi, M., Olagunju, A.T., Olusanya, B.O., Olusanya, J.O.,
Oluwasanu, M.M., Omar Bali, A., Omer, M.O., Ong, K.L., Onwujekwe, O.E., Orji, A.U.,
Orpana, H.M., Ortiz, A., Ostroff, S.M., Otstavnov, N., Otstavnov, S.S., Øverland, S.,
Owolabi, M.O., P A, M., Padubidri, J.R., Pakhare, A.P., Palladino, R., Pana, A., Panda-
Jonas, S., Pandey, A., Park, E.-K., Parmar, P.G.K., Pasupula, D.K., Patel, S.K., Paternina-
Caicedo, A.J., Pathak, A., Pathak, M., Patten, S.B., Patton, G.C., Paudel, D., Pazoki
Toroudi, H., Peden, A.E., Pennini, A., Pepito, V.C.F., Peprah, E.K., Pereira, A., Pereira,
D.M., Perico, N., Pham, H.Q., Phillips, M.R., Pigott, D.M., Pilgrim, T., Pilz, T.M.,
Pirsaheb, M., Plana-Ripoll, O., Plass, D., Pokhrel, K.N., Polibin, R. V, Polinder, S.,
Polkinghorne, K.R., Postma, M.J., Pourjafar, H., Pourmalek, F., Pourmirza Kalhori, R.,
Pourshams, A., Poznańska, A., Prada, S.I., Prakash, V., Pribadi, D.R.A., Pupillo, E., Quazi
Syed, Z., Rabiee, M., Rabiee, N., Radfar, A., Rafiee, A., Rafiei, A., Raggi, A., Rahimi-
Movaghar, A., Rahman, M.A., Rajabpour-Sanati, A., Rajati, F., Ramezanzadeh, K.,
Ranabhat, C.L., Rao, P.C., Rao, S.J., Rasella, D., Rastogi, P., Rathi, P., Rawaf, D.L., Rawaf,
S., Rawal, L., Razo, C., Redford, S.B., Reiner Jr, R.C., Reinig, N., Reitsma, M.B.,
Remuzzi, G., Renjith, V., Renzaho, A.M.N., Resnikoff, S., Rezaei, N., Rezai, M. sadegh,
Rezapour, A., Rhinehart, P.-A., Riahi, S.M., Ribeiro, A.L.P., Ribeiro, D.C., Ribeiro, D.,
Rickard, J., Roberts, N.L.S., Roberts, S., Robinson, S.R., Roever, L., Rolfe, S., Ronfani, L.,
Roshandel, G., Roth, G.A., Rubagotti, E., Rumisha, S.F., Sabour, S., Sachdev, P.S., Saddik,
B., Sadeghi, E., Sadeghi, M., Saeidi, S., Safi, S., Safiri, S., Sagar, R., Sahebkar, A.,
Sahraian, M.A., Sajadi, S.M., Salahshoor, M.R., Salamati, P., Salehi Zahabi, S., Salem, H.,
Salem, M.R.R., Salimzadeh, H., Salomon, J.A., Salz, I., Samad, Z., Samy, A.M., Sanabria,
J., Santomauro, D.F., Santos, I.S., Santos, J.V., Santric-Milicevic, M.M., Saraswathy,
S.Y.I., Sarmiento-Suárez, R., Sarrafzadegan, N., Sartorius, B., Sarveazad, A., Sathian, B.,
Sathish, T., Sattin, D., Sbarra, A.N., Schaeffer, L.E., Schiavolin, S., Schmidt, M.I., Schutte,
A.E., Schwebel, D.C., Schwendicke, F., Senbeta, A.M., Senthilkumaran, S., Sepanlou, S.G.,
Shackelford, K.A., Shadid, J., Shahabi, S., Shaheen, A.A., Shaikh, M.A., Shalash, A.S.,
Shams-Beyranvand, M., Shamsizadeh, M., Shannawaz, M., Sharafi, K., Sharara, F., Sheena,
B.S., Sheikhtaheri, A., Shetty, R.S., Shibuya, K., Shiferaw, W.S., Shigematsu, M., Shin, J.
Il, Shiri, R., Shirkoohi, R., Shrime, M.G., Shuval, K., Siabani, S., Sigfusdottir, I.D.,
Sigurvinsdottir, R., Silva, J.P., Simpson, K.E., Singh, A., Singh, J.A., Skiadaresi, E., Skou,
S.T.S., Skryabin, V.Y., Sobngwi, E., Sokhan, A., Soltani, S., Sorensen, R.J.D., Soriano,
J.B., Sorrie, M.B., Soyiri, I.N., Sreeramareddy, C.T., Stanaway, J.D., Stark, B.A., Ştefan,
S.C., Stein, C., Steiner, C., Steiner, T.J., Stokes, M.A., Stovner, L.J., Stubbs, J.L.,
Sudaryanto, A., Sufiyan, M.B., Sulo, G., Sultan, I., Sykes, B.L., Sylte, D.O., Szócska, M.,

Tabarés-Seisdedos, R., Tabb, K.M., Tadakamadla, S.K., Taherkhani, A., Tajdini, M.,
Takahashi, K., Taveira, N., Teagle, W.L., Teame, H., Tehrani-Banihashemi, A.,
Teklehaimanot, B.F., Terrason, S., Tessema, Z.T., Thankappan, K.R., Thomson, A.M.,
Tohidinik, H.R., Tonelli, M., Topor-Madry, R., Torre, A.E., Touvier, M., Tovani-Palone,
M.R.R., Tran, B.X., Travillian, R., Troeger, C.E., Truelsen, T.C., Tsai, A.C., Tsatsakis, A.,
Tudor Car, L., Tyrovolas, S., Uddin, R., Ullah, S., Undurraga, E.A., Unnikrishnan, B.,
Vacante, M., Vakilian, A., Valdez, P.R., Varughese, S., Vasankari, T.J., Vasseghian, Y.,
Venketasubramanian, N., Violante, F.S., Vlassov, V., Vollset, S.E., Vongpradith, A.,
Vukovic, A., Vukovic, R., Waheed, Y., Walters, M.K., Wang, J., Wang, Y., Wang, Y.-P.,
Ward, J.L., Watson, A., Wei, J., Weintraub, R.G., Weiss, D.J., Weiss, J., Westerman, R.,
Whisnant, J.L., Whiteford, H.A., Wiangkham, T., Wiens, K.E., Wijeratne, T., Wilner, L.B.,
Wilson, S., Wojtyniak, B., Wolfe, C.D.A., Wool, E.E., Wu, A.-M., Wulf Hanson, S.,
Wunrow, H.Y., Xu, G., Xu, R., Yadgir, S., Yahyazadeh Jabbari, S.H., Yamagishi, K.,
Yaminfirooz, M., Yano, Y., Yaya, S., Yazdi-Feyzabadi, V., Yearwood, J.A., Yeheyis, T.Y.,
Yeshitila, Y.G., Yip, P., Yonemoto, N., Yoon, S.-J., Yoosefi Lebni, J., Younis, M.Z.,
Younker, T.P., Yousefi, Z., Yousefifard, M., Yousefinezhadi, T., Yousuf, A.Y., Yu, C.,
Yusefzadeh, H., Zahirian Moghadam, T., Zaki, L., Zaman, S. Bin, Zamani, M., Zamanian,
M., Zandian, H., Zangeneh, A., Zastrozhin, M.S., Zewdie, K.A., Zhang, Y., Zhang, Z.-J.,
Zhao, J.T., Zhao, Y., Zheng, P., Zhou, M., Ziapour, A., Zimsen, S.R.M., Naghavi, M.,
Murray, C.J.L., 2020. Global burden of 369 diseases and injuries in 204 countries and
territories, 1990–2019: a systematic analysis for the Global Burden of Disease
Study 2019. Lancet 396, 1204–1222. https://doi.org/10.1016/S0140-6736(20)30925-9
World Health Organisation, 2017. Guidelines for drinking-water quality, 4th edition: 1st
addendum, World Health Organization. WHO.
898

Table 1. Estimated disease burden from waterborne microbial contaminants in high-income countries

Author	Date	Location	Health Outcomes	Exposure Source	Pathogens	Approach	Burden Estimate (Per Year)
Ball et al. (Ball, 2006)	2006	New Zealand	GI cases	Drinking	Unspecified	Top down	33,743 GI cases
Beaudeau et al. (Beaudeau et al., 2014)	2014	Boston, Massachusetts, USA	GI hospital admissions	Drinking (public systems)	Unspecified	Top down	153 GI hospital admissions
Burch et al. (Burch et al., 2021b)	2021	Kewaunee County, Wisconsin, USA	GI cases	Drinking (private wells)	Adenovirus group A, *Campylobacter*, *Cryptosporidium*, enteropathogenic *E. coli*, *Giardia*, *Salmonella*	Bottom up	301 GI cases
Butler et al. (Butler et al., 2016)	2016	Canada	Enteric illness cases	Food, drinking, recreation, animal contact	*Campylobacter*, *Cryptosporidium*, *Giardia*, verotoxin-producing *E. coli* (non-O157 and O157)	Top down	447,459 enteric illness cases attributed to only drinking water (private wells, public water systems, bottled)
Colford et al. (Colford et al., 2006)	2006	USA	GI cases	Drinking (public systems)	Unspecified	Top down	4.26-11.69 million GI cases
Collier et al. (Collier et al., 2020)	2021	USA	GI and respiratory cases, ED visits, hospitalizations, deaths	Drinking, recreation, inhalation	*Campylobacter*, *Cryptosporidium*, *Giardia*, *Legionella*, nontuberculous mycobacteria*,* norovirus*,* *Pseudomonas*, *Salmonella*, Shiga toxin-producing *E. coli*, *Shigella*, *Vibrio*, unspecified otitis externa	Top down	7.15 million waterborne illnesses, 601,000 ED visits, 118,000 hospitalizations, 6,630 deaths
DeFelice et al. (DeFelice et al., 2016)	2016	North Carolina, USA	ED visits for GI	Drinking (public systems and private wells)	Unspecified	Top down	29,400 ED visits
DeFelice et al. (DeFelice et al., 2015)	2015	North Carolina, USA	ED visits for GI	Drinking	*Giardia*, *Campylobacter*, rotavirus	Top down, bottom up comparison	Varied by approach: 190, 190, and 32,000 ED visits
Eisenberg et al. (Eisenberg et al., 2006)	2006	Iowa, USA	GI cases	Drinking (public systems)	Burden from trial data: unspecified	Top down, bottom up comparison	Burden from trial data: 1,825 cases per 10,000 population

Author	Date	Location	Health Outcomes	Exposure Source	Pathogens	Approach	Burden Estimate (Per Year)
					Burden from risk model: *Giardia*, *Cryptosporidium*, unspecified viruses		Burden from risk model: 13.9 cases per 10,000 population
Gibney et al. (Gibney et al., 2017)	2017	Australia	GI cases, deaths, DALYs	Drinking, recreation	*Campylobacter*, *Cryptosporidium*, *Giardia*, nontyphoidal *Salmonella*, norovirus	Top down	820,003 GI cases; 11 deaths
Haass et al. (Haass et al., 1996)	1996	San Elizario, El Paso County, Texas, USA	GI cases, Hepatitis A cases	Drinking, sanitation	*Entamoeba histolytica*, *Campylobacter*, hepatitis A virus, nontyphoidal *Salmonella*, typhoidal *Salmonella, Shigella*	Top down	199 GI cases, 6 hepatitis A cases
Kim et al. (Kim et al., 2018)	2018	Korea	Deaths, DALYs	Drinking, sanitation, hygiene	Unspecified GI pathogens (includes hepatitis A virus and helminths)	Top down	6 deaths
MacDonald Gibson et al. (MacDonald Gibson et al., 2013)	2013	United Arab Emirates	Gastroenteritis cases	Drinking	Unspecified	Top down	46,200 cases (4 deaths)
Masciopinto et al. (Masciopinto et al., 2019)	2019	Salento Region, Italy	GI cases, hepatitis A, DALYs	Drinking (groundwater)	Adenovirus, enterovirus, hepatitis A virus, norovirus, rotavirus	Bottom up	348 GI cases; 2 hepatitis A cases (summer only)
Messner et al. (Messner et al., 2006)	2006	USA	AGI cases	Drinking	Unspecified	Top down	16.4 million cases
Moore et al. (Moore et al., 2010)	2010	New Zealand	AGI cases	Drinking (noncompliant public systems)	Unspecified	Top down	35,000 cases
Morris and Levin (Morris and Levin, 1995)	1995	USA	GI cases, deaths	Drinking	*Campylobacter*, *Cryptosporidium*, *E. coli*, *Giardia*, norovirus *Salmonella*, *Shigella*	Top down (bacteria); bottom up (parasites & viruses)	7.66 million cases; 1,210 deaths
Murphy et al. (H.M. Murphy et al., 2016b)	2016	Canada	AGI cases	Drinking (public systems serving >1,000)	Unspecified	Top down	334,966 cases

Author	Date	Location	Health Outcomes	Exposure Source	Pathogens	Approach	Burden Estimate (Per Year)
Murphy et al. (H.M. Murphy et al., 2016c)	2016	Canada	AGI cases	Drinking (private wells, small public systems)	*Campylobacter, Cryptosporidium, E. coli* O157:H7, *Giardia*, norovirus	Bottom up	103,230 cases
Payment (Payment, 1997)	1997	Canada	HCGI cases	Drinking (public systems)	Unspecified	Top down	4,200,000
Perz et al. (Perz et al., 1998)	1998	New York City, USA	GI cases	Drinking (public systems)	*Cryptosporidium*	Bottom up	42 cases
Reynolds et al. (Reynolds et al., 2008)	2008	USA	GI cases	Drinking (public systems)	*Campylobacter, Cryptosporidium,* and rotavirus (and other unspecified viruses)	Bottom up	19.5 million cases
Verhougstraete et al. (Verhougstraete et al., 2020)	2020	USA	GI cases, Legionnaires' disease, sequela and mortality	Drinking	Adenovirus, *Campylobacter, Cryptosporidium Salmonella, E. coli* O157:H7, *Giardia*, hepatitis A virus, *Legionella,* norovirus, and rotavirus	Top down	9 million GI cases; 13,000 Legionnaire's disease cases; 32,531 Hepatitis A cases; 0.6 million sequela; 1,470 GI deaths
Vinson (Vinson, 2012)	2012	Canada	GI cases, respiratory diseases, toxoplasmosis, ear infections	Drinking, recreation, inhalation	*Legionella pneumophila, Mycoplasma pneumoniae, Toxoplasma gondii* and unspecified pathogens causing GI/otitis externa,	Top down	1.32 million GI cases

GI- Gastrointestinal illness
AGI- Acute gastrointestinal illness
HCGI- Highly credible gastrointestinal illness
ED- Emergency department
DALYs- Disability Adjusted Life Years

Table 2. Estimated disease burden from waterborne chemical contaminants in high-income countries

Author	Date	Location	Health Outcomes	Exposure Source	Chemical Contaminant	Approach	Burden Estimate (Per Year)
Abtahi et al. (Abtahi et al., 2019)	2019	Iran	Dental fluorosis	Drinking	Fluoride	Bottom up	60 cases/100,000 people (95% CI 48-69)
de Hollander et al. (de Hollander et al., 1999)	1999	The Netherlands	Loss of 1-3 IQ points	Drinking	Lead	Bottom up	1,764 cases
DeFelice et al. (DeFelice et al., 2017)	2017	North Carolina, USA	Cancer	Drinking (public systems)	20 carcinogens	Top down	53.9 (95% CI 27.8-79.4) deaths statewide; individual risk 7.2×10^{-6}
Dobaradaran et al. (Dobaradaran et al., 2020)	2020	Bushehr Province, Iran	Cancer	Drinking, dermal, inhalation	Disinfection byproducts	Bottom up	1 (95% CI 0.32-2.8) death in Bushehr Province; 95.0 DALYs (94.7–95.2).
Evlampidou et al. (Evlampidou et al., 2020)	2020	28 European Union countries	Cancer	Drinking	Disinfection byproducts	Top down	6,561 (95% CI 3,389, 9,537) cases across the EU
Fewtrell et al. (Fewtrell et al., 2006)	2006	USA, United Kingdom, New Zealand	Dental and skeletal fluorosis	Drinking	Fluoride	Bottom up	Zero
Greco et al. (Greco et al., 2019)	2019	USA	Ischemic heart disease; cancer	Drinking (private wells)	Arsenic	Bottom up	500 ischemic heart disease deaths; 496 fatal cancer cases; 578 nonfatal cancer cases
MacDonald Gibson et al. (MacDonald Gibson et al., 2013)	2013	United Arab Emirates	Bladder and colorectal cancers	Drinking	Trihalomethanes (THMs)	Top down	154 healthcare facility visits (and 3 deaths) for bladder cancers and 328 visits (and 9 deaths) for colorectal cancers
Mathewson et al. (Mathewson et al., 2020)	2020	Wisconsin, USA	Cancer; neural tube defects; low birth weight; preterm birth	Drinking	Nitrate	Top down	95 (range 46-149 very low birth weight babies; 51 (range 26-79) very preterm births; 1-2 neural tube defect births; 169 (range 111-297) cancer cases
Temkin et al. (Temkin et al., 2019)	2019	USA	Cancer; neural tube defects; low birth weight; preterm birth	Drinking	Nitrate	Top down	2,939 very low birth weight cases; 1,725 very preterm births; 41 neural tube defect births; 6,537 (range 2,300-12,594) cancer cases

Figure 1. Summary of burden estimates (incidence per 100,000 population) for a) cases of GI illness attributed to microbiological contamination of drinking water and b) deaths from illness attributed to microbiological contamination of drinking water.

Countries: AUS=Australia, CAN=Canada, ITA=Italy, KOR=Republic of Korea, NZL=New Zealand, UAE=United Arab Emirates, USA=United States of America

Figure 2. Summary of burden estimates (incidence per 100,000 population) for a) cases of illness attributed to chemical contamination of drinking water, and b) deaths from illness attributed to chemical contamination of drinking water.

Note: Estimates of zero cases (per 100,000 population) of fluorosis in USA, GBR (UK), NZL by Fewtrell et al. were excluded

Countries: EU=European Union, GBR=United Kingdom, IRN=Iran, NLD=The Netherlands, UAE=United Arab Emirates, USA=United States of America
Contaminants: DBPs=Disinfection by-products, VOCs=Volatile Organic Compounds
Outcomes: VLBW=Very low birth weight, VPTB=Very preterm birth, NTD=Neural tube defects

Figure 3. Summary of gaps in the literature on a) cases and b) deaths by chemical contaminant and country. For countries where disease burden estimates were available through multiple studies, disease outcome estimates from different studies were averaged to produce country-specific estimates. For cancers, specific cancer estimates (e.g., bladder cancer, liver cancer) were summed to produce an overall cancer estimate.

Note: The burden of neural tube defects in the United States is not visible due to its relatively low level (0.02 cases per 100,000)

Countries: EU=European Union, GBR=United Kingdom, IRN=Iran, NLD=The Netherlands, UAE=United Arab Emirates, USA=United States of America
Contaminant: DBPs=Disinfection by-products, VOCs=Volatile Organic Compounds