

Determinants of Neonatal Sepsis Admitted In Neonatal Intensive Care Unit At Public Hospitals Of Kaffa Zone, South West Ethiopia.

Melesse Belayneh^{1*}, Gebremariam Getaneh¹, Alemaw Gebretsadik²

^{1,2}Department of Health System Management and Health Economics College of Medicine and Health Science Bahir Dar University, Ethiopia,

❖ Current address: 1Department of Health System Management and Health Economics College of Medicine and Health Science Bahir Dar University, Ethiopia * meleseyayeh@gmail.com

Abstract

Background: Neonatal sepsis is a systemic inflammatory response syndrome in the presence of infection during the first 28 days of life. Globally every year about 4 million children die in the first 4 weeks of life, of which 99% of the deaths occur in low and middle income countries and the most common causes of neonatal death in Ethiopia. Identification of the determinants for neonatal sepsis and treatment of newborns with sepsis is not adequate in low income countries like Ethiopia especially in southern part of the country.

Objective: To identify determinants of neonatal sepsis admitted in neonatal intensive care unit at public hospitals of Kaffa zone, south west Ethiopia 2021.

Methods: Institutional based unmatched case control study was conducted on a total sample of 248 (62 cases and 186 controls) in public Hospitals of kaffa zone from March to April 2021. The collected data were entered, coded and cleaned by Epidata version 3.1 and it was exported to SPSS version 25. Bi-variable and multivariable logistic regression was conducted. Variables with ($p < 0.25$) in bi-variable logistic regression analysis, were entered to multivariable logistic regression and then determinants which is statistical significant will be declared at $P < 0.05$.

Result: A total of 248 (62 cases and 186 controls) were included in the study. variables like prolonged rupture of membrane ≥ 18 hours [AOR =5.13, 95%CI=1.38-19.05], meconium stained amniotic fluid [AOR =6.03, 95%CI=2.16-16.90], intra-partum fever [AOR =8.26, 95%CI=3.12-21.97], urinary tract infections [AOR=14.55, 95%CI=4.91-43.10], breast feeding after a hour [AOR =3.9, 95%CI=1.27-12.02], resuscitation [AOR =13.25, 95%CI=3.44-51.01], no chlorohexidine application [AOR =4.27, 95%CI=1.65-11.08] were significantly associated with neonatal sepsis.

Conclusion and Recommendation: Among the variables prolonged rupture of membranes, meconium stained amniotic fluid, intra-partum fever, UTI/STI, and not breast feeding within an hour were maternal variables and resuscitation at birth and not application of chlorohexidine ointment on the umbilicus were neonatal variables that were found to be neonatal-related risk factors of neonatal sepsis. Infection prevention strategies and clinical management need to be strengthening and/or implementing by providing especial attention for the specified determinants.

Keywords: Neonatal, sepsis, Infection, Kaffa zone, NICU Care

Introduction

Neonatal sepsis is a systemic inflammatory response syndrome in the presence of infection during the first 28 days of life(1, 2). It could be of bacterial, viral, fungal or rickettsia origin(2). It is estimated to cause twenty-six percent of all neonatal deaths worldwide(3).

Neonatal sepsis can be classified in to two subtypes depending upon whether the onset of symptoms is before 72 hours of life (early onset) or after 72 hours (late onset). Early onset neonatal sepsis (EONS) is caused by organisms prevalent in the maternal genital tract or in the delivery area(4, 5). Late onset neonatal sepsis (LONS) is caused by the organisms thriving in the external environment of the home or hospital. The infection is often transmitted through the hands of the care-providers(5, 6). It comprises abundant systemic infections of the neonate like septicemia, meningitis, pneumonia, arthritis, urinary tract infection but, it does not embrace mucocutaneous infections like conjunctivitis and oral thrush. Neonatal infections in health care facilities are still a reason for inflated death and illness in Neonatal Intensive Care Unit (NICU) (5).

World health organization (WHO) has recognized seven clinical indicators [difficulty feeding, convulsions, movement only while stimulated /lethargy, respiratory rate of >60 breaths in a minute, chest in drawing, or axillary temperature of >37.5 °C or <35.5 °C and respiratory distress(7). Another study has additionally incorporated cyanosis and grunting(8).

Risk factors for EONS include prematurity, low birth weight, premature and prolonged rupture of membranes, maternal fever, uroinfection and chorioamnionitis whereas for LONS are invasive procedures such as resuscitation in delivery room, intubation, mechanical ventilation, central venous catheters, surgical procedures (5). As shown in different literatures, neonatal sepsis is caused by factors related to both maternal and neonatal factors (9-11). Delivery by caesarian section, male sex and prematurity have been showed as determinants of neonatal sepsis(11, 12). Maternal factors such as prolonged rupture of membrane, urinary tract infection, intra-partum fever, instrumental delivery and place delivery are significant predictors of neonatal sepsis (10, 13).

In other studies, use of endotracheal intubation, resuscitation at birth, surgery are significant determinants (11). Neonates born from women with meconium stained amniotic fluid, more than three times digital per vaginal examination and never attend antenatal care (ANC) are at higher risk for neonatal sepsis(14-16).

Neonatal sepsis is the commonest cause of neonatal morbidity and mortality. It is responsible for about thirty up to fifty percent of total neonatal deaths in developing countries(10). Globally every year about 4 million children die in the first 4 weeks of life, of which 99% of the deaths occur in low- and middle-income countries and of which 75% are considered as avoidable. Ethiopia continuous to struggle with a prevalence of about 42% or 81,000 newborn deaths every year(17).

Although several novel diagnostic markers have been discovered to detect onset of disease, at this moment, there is no proper diagnostic tool developed that can accurately diagnose sepsis. Promising results are emerging from recently identified cytokines, chemokines, cell surface antigens, physiomarkers, and metabolomics(18). The gold standard for establishing a diagnosis of neonatal sepsis is through culture. However, several factors ,including the small blood volumes obtained from neonates as well as maternal intra-partum antimicrobial exposure, can make the confirmation of sepsis in a neonate a diagnostic challenge(19)

Although recent medical advances have improved neonatal care many challenges remain in the diagnosis and management of neonatal infections. The diagnosis of neonatal sepsis is complicated by the frequent presence of non-infectious conditions that resemble sepsis, especially in preterm infants, and by the absence of optimal diagnostic tests. Efforts to reduce the rates of infection in this vulnerable population are one of the most important interventions in neonatal care(20).

Even though the presence of few studies regarding risk factors and etiology of neonatal sepsis, there are some contradicting or inconsistent findings on some predictors for neonatal sepsis, like prematurity, low birth weight and residence (10, 27). A study in Mexico showed, prematurity, low birth weight and residence as significant determinant of neonatal sepsis(11). On the contrary a study in Ethiopia showed that, all the three variables were not determinants of neonatal sepsis(10). Also a study in Wolayta Sodo showed that ANC visit, maternal age, maternal income as independent determinants of neonatal sepsis but in a study conducted in Jinka the above variables are not determinants(32, 33).

As far as literature searching showed, there is no study conducted in the study area regarding determinants of neonatal sepsis. Therefore, the aim of this study is to identify the determinants of neonatal sepsis among neonate admitted to the Neonatal Intensive Care Unit (NICU) in public hospitals of Kaffa zone, Southwest Ethiopia, 2021.

Objective

To identify determinants of neonatal sepsis admitted in neonatal intensive care unit at public hospitals of Kaffa zone, south west Ethiopia 2021

Methods and Materials

Study design

Institutional based unmatched case control study design was conducted to assess the determinants of neonatal sepsis among neonates admitted at neonatal intensive care unit in public hospitals of Kaffa zone south west region of Ethiopia.

Study Area and Period

The study was conducted from March 01 to April 10/2021 in Kafa zone south west region of Ethiopia. Kafa zone is one among 15 zones and 4 special woredas in SNNPRS with an elevation of 1450-2700 meters above sea level. The capital of kaffa is Bonga town which is located 450Km south west of Addis Ababa and 559 Km from regional city Hawassa. Kaffa zone is divided in to 12 woredas and 2 administrative towns. It has 1 General hospital which is Gebretsadik shawo General Hospital serving for more than 1,200,000 populations and 2 primary hospital (Wacha Primary Hospital and Telo Primary Hospital) serving for 211,000 and 84,000 populations respectively. According to information obtained from administrative offices of these hospitals, they provide different services in outpatient department, inpatient department and operation room theatre department. GSGH had 1 GP and 5 neonatal nurses with 10 functional beds while CPH had three nurses while TPH has two clinical nurses with three beds each in NICU with total annual neonatal admission of more than 2500 of which more than 850 was by neonatal sepsis in all hospitals of the catchment area. As a report from kaffa zone health department indicates the monthly neonatal sepsis new admission in March 2020 was 32 in GSGH, 13 in WPH, and 10 in TPH(55).

According to the 2020/21 annual report of Zonal health department, currently the population is 1,224,684 (male: 600,095 and female: 624,589 and from this 42,374 and 39,067 are live births and surviving infants respectively. The zone has up and down topography which is not suitable to get their health needs of the community. It is bordered by Jima zone in East and North, Konta special woreda and Basketo woreda in South, Benchimaji and Sheka zone in west.

Population

Source population: All neonates admitted in neonatal intensive care unit in public hospitals of kaffa zone

Study population: All neonates admitted in neonatal intensive care unit in public hospitals of kaffa zone during the study period

Sample population: All randomly selected neonates for controls and consecutively selected neonates for cases in the neonatal intensive care unit.

Variables

Dependent variable: Neonatal sepsis.

Independent Variables: Socio-demographic characteristics of neonates and parents (Age of mother, Ethnicity, Religion, Residence, Marital status, Educational status of mother, Occupation of mother, Neonatal age, Neonatal sex, Monthly income and Family size), maternal factors (Gravidity, Parity, ANC visit, Frequency of ANC, Place of delivery, Mode of delivery, Attendant of delivery, PROM, MSAF, Number of PV exam, Intrapartum fever, Foul smelling vaginal discharge, PIH /eclapsia, APH, UTI or STI and Breast feeding) and neonatal factors (Gestational age, Apgar score at 1st minute, Apgar score at 5th minute, Birth weight, Immediate cry, Birth asphyxia, Resuscitation, Birth trauma, Cord cut, Chlorohexidine application, Congenital anomalies and Invasive procedure) were the hypothesized determinants

Inclusion and Exclusion criteria

Inclusion criteria

Neonates whose age less than one month(at the time of admission) with their mothers and admitted to neonatal intensive care unit of public hospitals of kaffa zone and fulfills the neonatal sepsis criteria for cases and neonates who admitted in NICU other than sepsis for controls.

Exclusion criteria

Neonates whose mothers had hearing impairments or unable to talk, abandoned neonates who were admitted in the hospital, and neonates whose cards had incomplete information, neonates whose mother is died, was excluded from the study in both controls and cases.

Sample size estimation

The sample size was determined by using double population proportion formula in STATCALC program of the EPI INFO version 7.4.2 statistical software by considering double population proportion formula Proportion of maternal UTI/STI among controls is 14.9% and proportion of UTI/STI among cases is 33.8%; then by taking

95% CI, power of the study is 80%, control to case ratio of 3:1 the sample size becomes 227. After adding 10% non-respondent rate the final required sample will be 248 (62 cases and 186 controls) participants. Cases were neonates with sepsis and controls were neonates without sepsis who were admitted to NICU of the three hospitals during the study period.

Sampling procedure

Cases (diagnosed with sepsis) were selected through consecutive sampling technique from the neonates who were admitted in NICU until the required sample size achieved. The next immediate three corresponding controls were selected by simple random sampling (lottery method) at the same day and in the same neonatal intensive care unit. Proportional allocation of the sample was applied for the hospitals based on their case load (36 cases and 108 controls from GSGH, 15 cases and 45 controls from WPH, 11 cases and 33 controls from TPH).

Operational and term definitions

Case (Neonatal sepsis): The established Integrated Management of Neonatal and Childhood Illness (IMNCI) clinical features, includes the presence of two or more of persistent fever (≥ 37.5 °C) or persistent hypothermia (≤ 35.5 °C) for more than one hour, fast breathing (≥ 60 breath per minute), severe chest in drawing, grunting, not feeding well, movement only when stimulated, bulged fontanel, convulsion, lethargic or unconsciousness along with ≥ 2 of the hematological criteria such as total leukocyte count (< 4000 or $> 12,000$ cells/mm³), absolute neutrophil count (< 1500 cells/mm³ or > 7500 cells/mm³), platelet count (< 150 or > 450 cells/mm³), and random blood sugar (< 40 mg/dl or > 125 mg/dl) will be used to diagnose neonatal sepsis cases(10).

Control: neonates with the diagnosis of non-sepsis case with their index mother(10).

Meconium stained amniotic fluid (MSAF): will be considered if the amniotic fluid was green /brown in color or mixed with meconium, or appears meconium stained on the baby(54).

Prolonged rupture of membrane (PROM): the time from membranes' rupture to delivery more than 18 hours(54).

Invasive procedures: Procedures which have either of the following like Surgical procedures, Umbilical catheterization, Urinary catheterization, Nasogastric/oropharynx tube insertion, Endotracheal tube(32).

Data Collection materials and procedure

The questionnaire and checklist were developed after reviewing different literature which was related to determinants of neonatal sepsis. Then the designed questionnaires was changed from English to local

language(kafinoono) and back to English to check the consistency of the questionnaire.. Training for supervisors and data collectors about data collection procedure was given for two days. Data were collected through a face to face interview of the index mothers with pretested structured interviewer administered questionnaires and reviewing neonates' medical records (laboratory results:-CBC, CRP, ESR,) using checklist by trained experienced health professionals. They were interviewed about their socio-demographic characteristics, maternal factors and neonatal factors in each NICU in two primary hospitals and one general hospital. The data were collected from March to April 10/2021 by 4 BSc nurses and 1 MPH supervisor.

Data Processing and Analysis

The filled questionnaires were checked for completeness and entered to Epidata version 3.1 then exported to SPSS version 25 for farther analysis. Descriptive statistics like frequency tables and percentages were used to describe the study participants in relation to relevant variables. The goodness of the model was assessed whether the required assumptions for the application of multivariate logistic regression will be fulfilled and the model will adequately fits the data (Hosmer and Lemeshow test was assessed). Both bi-variable and multivariable logistic regression models were used to identify factors which determinate neonatal sepsis. In order to minimize factors in multivariable analysis only variables that had showed $p\text{-value} < 0.25$ on the Bi-variable logistic analysis were entered in to the multivariable logistic regression models. Odds Ratio and their 95% confidence intervals were computed and variables with $p\text{-value}$ less than 0.05 in multivariable analysis were considered as statistically significant. Backward stepwise technique method will be used.

Data Quality Assurance

Data quality assurance was implemented in all stage of the study starting from questionnaire designing, training, data collection and data entry. Pretest was done at Bonga health center which had NICU by taking 5% of sample size (3 cases and 9 controls) before the study period and the consistency of the questionnaire was checked. The questionnaire was objective based, logically sequenced and pre tested. The data collector and supervisor had received training on the objective of the study. The collected data was checked by the principal investigator on daily basis for any incompleteness. Data entry was conducted by principal investigator and other skilled personnel to minimize error during data entry.

Ethical Consideration

Ethical clearance was obtained from Institutional Review Board (IRB) of Bahir Dar University College of Medicine and health science. Letter of permission was obtained from kaffa zone health department and each

hospital Chief executive office. The confidentiality of information had maintained by excluding personal identifiers and interview privately. Data were collected after securing informed written consent from every respondent.

Result

Socio-demographic characteristics of the respondent

This study involved a total of 248 neonates with their index mother among which 62 had sepsis (cases) and 186 was controls with 100% response rate. The study was intended to assess determinants of neonatal sepsis among neonates admitted in neonatal intensive care unit in kaffa zone hospitals. The mean age of index mothers was 28.06 years (SD \pm 5.82) with minimum age of 18 and maximum age of 42. The mean age of neonates was 5.04 days (SD \pm 6.09) with the age range of 1-27 days. Most of the participants were kaffa by ethnicity 43(69.4%) of cases and 139(74.7%) controls and 42(67.7%) of cases and 119 (64%) of controls were orthodox Christian. Among participants most of them were rural residents 38 (61.3%) of cases and 120 (64.5%) of controls). 82.3% cases and 83.9% of controls were married while 11(17.7%) of cases and 30(16.1%) of controls were educated above college level. Concerning maternal occupation, 75.8% of cases and 73.7% of controls were house wives while 19.4% of cases and of controls were go vernment employee. More than half of cases 36(58.1%) and controls 95(51%) were male neonate. The mean monthly income of cases and controls were 1970(SD \pm 1628) and 2197 (SD \pm 1191) ETB respectively(Table 1).

Table 1: Socio-demographic characteristics of mothers and their neonates for determinants of neonatal sepsis admitted in Neonatal Intensive Care unit of public hospitals in Kaffa zone southwest Ethiopia 2021.

Variables	Categories	Case n =62(%)	Control =186(%)	Total 248(%)
Age of mother	15-20	8(12.9)	14(7.5)	22(8.9)
	21-34	44(71.0)	139(74.1)	183(73.8)
	\geq 35	10(16.1)	33(17.7)	43(17.3)
Ethnicity	Kafa	43(69.4)	139(74.7)	182(73.4)
	Oromo	6(9.7)	33(17.7)	39(15.7)
	Amhara	11(17.7)	14(7.5)	25(10.1)
	Konta	2(3.2)	0(0.0)	2(0.8)
	others	0(0.0)	0(0.0)	0(0.0)
Religion	Orthodox	42(67.7)	119(64.0)	161(64.9)
	Protestant	16(28.8)	55(29.6)	71(28.6)
	Muslim	4(6.5)	12(6.5)	16(6.5)
Residence	Urban	24(38.7)	66(35.5)	90(36.3)
	Rural	38(61.3)	120(64.5)	158(63.7)

Marital status	Single	8(12.9)	17(9.1)	25(10.1)
	Married	51(82.3)	156(83.9)	207(83.5)
	Divorced	2(3.2)	7(3.8)	9(3.6)
	Widowed	1(1.6)	6(3.2)	7(2.8)
Educational status of mother	Illiterate	31(50.0)	94(50.5)	125(50.4)
	Read and write	10(16.1)	40(21.5)	50(20.2)
	Elementary school	5(8.1)	11(5.9)	16(6.5)
	Secondary school	5(8.1)	11(5.9)	16(6.5)
	College/above	11(17.7)	30(16.1)	41(16.5)
Occupation of mother	House wife	47(75.8)	137(73.7)	184(74.2)
	Gov't employee	12(19.4)	36(19.4)	48(19.4)
	Private institution	2(3.2)	5(2.7)	7(2.8)
	Self-employee	0(0.0)	4(2.2)	4(1.6)
	Daily laborer	1(1.6)	4(2.2)	5(2.0)
Neonatal age	1-3	43(69.3)	107(57.5)	150(60.5)
	>3	19(30.6)	79(42.5)	98(39.5)
Neonatal sex	Male	36(58.1)	95(51.1)	131(52.8)
	Female	26(41.9)	91(48.9)	117(47.2)
Monthly income	<3500	49(79.0)	147(79.0)	196(79.0)
	>=3500	13(21.0)	39(21.0)	52(21.0)
Family size	<5	39(62.9)	107(57.5)	146(58.9)
	>=5	23(37.1)	79(42.5)	102(41.1)

Maternal factors

According to this study 5(8.1%) of cases and 21(11.3%) of controls were Primigravida while 67.7% of cases and 64.5% of controls were multipara index mothers. Nearly two-third (63%) of cases and 166(89.2%) of controls were visited health facility for ANC service among them who visited for ANC 66% of cases and 70.4% of controls were visited more than three times. Among the participants, 23 (37.1%) of cases and 40.9% of controls, 30(40.4%) of cases and 43.5% of controls, 9(14.5%) of cases and 15.6% of controls were delivered at hospital, health center and home respectively. Three fourth (75.8%) of cases and 87.6% of controls were delivered through SVD while 8.1% of cases and 4.8% of controls were CS delivery and most deliveries were attended by health professionals. Twenty two (35.5%) of cases and 7% of controls had rupture of membranes greater than or equal to 18 hours. Two third (65.6%) of cases and nearly one-fourth (24.2%) controls had amniotic fluid discoloration while nearly half 48.4% of cases and 22.6 % controls had foul smelling amniotic fluid. Thirty nine (62.9%) of cases and 139(74.7%) of controls had per-vaginal examination of three times or

less. Among respondents 46(74.2%) of cases and 41(22%) of controls had history of fever during the index pregnancy. Twenty two (35.5%) of cases and 27.4% of cases had PIH/eclapsia and APH respectively. One-fourth (19.9%) of controls 74.2% of case had UTI/STI during their index pregnancy while 40(64.5%) of cases 165(88.7%) of controls start breast feeding within one hour(Table 2).

Table 2: Maternal factors for determinants of neonatal sepsis admitted in Neonatal Intensive Care Unit of public hospitals in Kaffa zone southwest Ethiopia 2021.

Variables	Categories	Case n=62(%)	Control n=186(%)	Total n=248(%)
Gravidity	Primigravida	5(8.1)	21(11.3)	26(10.5)
	Multigravida	57(91.9)	165(88.7)	222(89.5)
Parity	Nulipara	6(9.7)	21(11.3)	27(10.9)
	Paraone	14(22.6)	45(24.2)	59(23.8)
	Multipara	42(67.7)	120(64.5)	162(65.3)
ANC visit	Yes	39(62.9)	166(89.2)	205(82.7)
	No	23(37.1)	20(10.8)	43(17.3)
Frequency of ANC	<=3	21(33.9)	55(29.6)	76(30.6)
	>3	41(66.1)	131(70.4)	172(69.4)
Place of delivery	Home	9(14.5)	29(15.6)	38(15.3)
	Health center	30(48.4)	81(43.5)	111(44.8)
	Hospital	23(37.1)	76(40.9)	99(39.9)
Mode of delivery	SVD	47(75.8)	163(87.6)	210(84.7)
	Instrumental	10(16.1)	14(7.5)	24(9.7)
	CS	5(8.1)	9(4.8)	14(5.6)
Attendant of delivery	TBA	7(11.3)	28(15.1)	35(14.1)
	HEW	0(0.0)	0(0.0)	0(0.0)
	Health professional	55(88.7)	158(84.9)	213(85.9)
PROM	<12	38(61.3)	171(91.9)	209(84.3)
	12-18	2(3.2)	2(1.1)	4(1.6)
	>=18	22(35.5)	13(7.0)	35(14.1)
MSAF	Yes	40(65.6)	45(24.2)	85(34.3)
	No	21(34.4)	141(75.8)	162(65.3)
Number of PV exam	<=3	39(62.9)	139(74.7)	178(71.8)
	>3	23(37.1)	47(25.3)	70(28.2)
Intra-partum fever	Yes	46(74.2)	41(22.0)	87(35.1)
	No	16(25.8)	145(78.0)	161(64.9)
Foul smelling vaginal discharge	Yes	30(48.4)	42(22.6)	72(29.0)
	No	32(51.6)	144(77.4)	176(71.0)
PIH /eclapsia	Yes	22(35.5)	25(13.4)	47(19.0)
	No	40(64.5)	161(86.6)	201(81.0)

APH	Yes	17(27.4)	41(22.0)	58(23.4)
	No	45(72.6)	145(78.0)	190(76.6)
UTI or STI	Yes	46(74.2)	37(19.9)	83(33.5)
	No	16(25.8)	149(80.1)	165(66.5)
Breast feeding	<60 minute	40(64.5)	165(88.7)	205(82.7)
	>60 minute	22(35.5)	21(11.3)	43(17.3)

APH ante partum hemorrhage, PPH postpartum hemorrhage, GDM gestational diabetes mellitus, PIH pregnancy induced hypertension, PROM premature rupture of membrane and UTI Urinary Tract Infection

Neonatal factors

According to this study 35(56.5%) of cases and 115(61.8%) of controls were in between 37-42 completed weeks of gestational age. The 1st minute and 5th minute apgar score <7 among case neonates were 24.2% and 9.2% respectively while the 1st and 5th minute apgar score <7 in controls were 9.7% and 1.6% respectively. More than 90% of cases and controls were in the normal range of birth weight. Forty one 66.1% of cases and 166(89.2%) of controls cries immediately after birth where as 34% of cases and 9.7% of controls developed birth asphyxia. One third of cases and 8.1% of controls were resuscitated after birth and more than 92% of case and controls had cord cut with new blade. Twenty two (35.5%) of cases 134(72%) of controls had history of chlorohexidine application for neonates where as 4.8% and 6% cases and controls had had invasive procedure(Table 3).

Table 3. Neonatal factors for the study of determinants of neonatal sepsis admitted in Neonatal Intensive Care Unit of public hospitals in Kaffa zone southwest Ethiopia 2021.

Variables	Categories	Case n=62(%)	Control n=186(%)	Total n=248(%)
Gestational age	<37	27(43.5)	71(38.2)	98(39.5)
	37-42	35(56.5)	115(61.8)	150(60.5)
	>42	0(0.0)	0(0.0)	0(0.0)
Apgar score at 1 st minute	<7	15(24.2)	18(9.7)	33(13.3)
	≥7	47(75.8)	168(90.3)	215(86.7)
Apgar score at 5 th minute	<7	6(9.7)	3(1.6)	9(3.6)
	≥7	56(90.3)	183(98.4)	239(96.4)
Birth weight	<2500	6(9.7)	11(5.9)	17(6.9)
	≥2500	56(90.3)	175(94.1)	231(93.1)
Immediate cry	Yes	41(66.1)	166(89.2)	207(83.5)
	No	21(33.9)	20(10.8)	41(16.5)
Birth asphyxia	Yes	21(33.9)	18(9.7)	39(15.7)
	No	41(66.1)	168(90.3)	209(84.3)
Resuscitation	Yes	21(33.9)	15(8.1)	36(14.5)
	No	41(66.1)	171(91.9)	212(85.5)

Birth trauma	Yes	4(6.5)	8(4.3)	12(4.8)
	No	58(93.5)	178(95.7)	236(95.2)
Cord cut	New blade	57(91.9)	173(93.0)	230(92.7)
	Old blade	5(8.1)	13(7.0)	18(7.3)
Chlorohexidine application	Yes	22(35.5)	134(72.0)	156(62.9)
	No	40(64.5)	52(28.0)	92(37.1)
Congenital anomalies	Yes	2(3.2)	13(7.0)	15(6.0)
	No	60(96.8)	173(93.0)	233(94.0)
Invasive procedure	Yes	3(4.8)	11(5.9)	14(5.6)
	No	59(95.2)	175(94.1)	234(94.4)

Health facility related factors

All neonates admitted in which all hospitals implemented infection prevention practice in the neonatal intensive care unit while 82.3% of case and control neonates were admitted in hospitals which had trained professional on infection prevention and control practice. Thirty nine (62.9%) of cases and 102(54.8%) of controls lives with in 5km or less and their common means of transportation for rural residents were hoarse(Table 4).

Table 4:Health facility related factors for the study of determinants of neonatal sepsis admitted in Neonatal Intensive Care Unit of public hospitals in Kaffa zone southwest Ethiopia 2021.

Variables	Categories	Case n=62(%)	Control n= 186(%)	Total n=248(%)
Infection prevention implemented	Yes	62(100.0)	186(100.0)	248(100.0)
	No	0(0.0)	0(0.0)	0(0.0)
Trained professional on IPC	Yes	51(82.3)	153(82.3)	204(82.3)
	No	11(17.7)	33(17.7)	44(17.7)
Distance of HF	≤5	39(62.9)	102(54.8)	141(56.9)
	>5	23(37.1)	84(45.2)	107(43.1)
Means of transport	Car	32(51.6)	85(45.7)	117(47.2)
	Hoarse	25(40.3)	85(45.7)	110(44.4)
	Foot	5(8.1)	16(8.6)	21(8.5)

Determinants of Neonatal Sepsis Admitted In Neonatal Intensive Care

In bi-variable logistic regression analysis result ANC visit, mode of delivery, duration of PROM, meconium stained amniotic fluid, frequency of PV exam, intra-partum fever, foul smelling amniotic fluid, PIH, UTI, breast feeding, gestational age, apgar score <7 at 1st and 5th minute, immediate cry, asphyxia, resuscitation, chlorohexidine application were showed association with neonatal sepsis(P<0.25). then applying multivariable logistic regression maternal PROM ≥ 18 hours, MSAF, intra-partum fever, history of UTI/STI, breast feeding after 60 minute, neonatal resuscitation at birth and chlorohexidine application was significantly associated with neonatal sepsis.

In multivariable logistic regression analysis, neonates born from women who had prolonged rupture of membrane ≥ 18 hours had five times more likely to develop neonatal sepsis than that of women who had PROM <12 hours [AOR =5.13; 95% CI(1.38, 19.05)]. Those neonates born from mothers who had MSAF were six times more likely to suffer from neonatal sepsis than their counter parts [AOR =6.03; 95%CI(2.16, 16.90)]. Intra-partum fever was also had a significant association with neonatal sepsis, mothers who had history of intra-partum fever were eight times more risk for their neonates to develop neonatal sepsis than mothers who had no fever [AOR =8.26; 95%CI(3.12, 21.97)].

In this study neonatal sepsis was strongly associated with maternal history of UTI/STI, those neonates born from mothers who had history of UTI/STI were fourteen times more likely to have neonatal sepsis than their counter parts [AOR =14.55; 95%CI(4.91, 43.10)]. Mothers who breast feed their neonate after 1 hour were nearly four times more likely to suffer from neonatal sepsis than that of neonates who breast feed within 1 hour [AOR =3.9; 95%CI(1.27, 12.02)]. Resuscitation of the newborn was also had significant association with neonatal sepsis, resuscitated neonates were thirteen times more likely to have neonatal sepsis than their counter parts [AOR =13.25; 95%CI(3.44, 51.01)]. Those neonates who were not got chlorohexidine ointment on their umbilicus had four times more likely to develop neonatal sepsis than neonates who had got chlorohexidine application [AOR =4.27; 95%CI(1.65, 11.08)] (Table 5).

Table 5: multivariable logistic regression analysis result determinants of neonatal sepsis admitted in Neonatal Intensive Care Unit of public hospitals in Kaffa zone southwest Ethiopia 2021.

Variables	Categories	Cases n=62(%)	Control n=186(%)	AOR [95%CI]	P-Value
Mode of delivery	SVD	47(75.8)	163(87.6)	1	
	instrumental	10(16.1)	14(7.5)	6.130[0.759, 49.486]	0.089
	CS	5(8.1)	9(4.8)	1.520[0.110, 21.110]	0.755
PROM	<12	38(61.3)	171(91.9)	1	
	12-18	2(3.2)	2(1.1)	1.663[0.108, 25.511]	0.715
	>=18	22(35.5)	13(7.0)	5.129[1.381,19.055]**	0.015
Chlorohexidine application	yes	22(35.5)	134(72.0)	1	
	no	40(64.5)	52(28.0)	4.275[1.649, 11.083]**	0.003
MSAF	yes	40(65.6)	45(24.2)	6.038[2.157, 16.905]**	0.001
	no	21(34.4)	141(75.8)	1	
Number of PV exam	<=3	39(62.9)	139(74.7)	1	
	>3	23(37.1)	47(25.3)	1.977[0.608, 6.431]	0.257
Intra-partum fever	yes	46(74.2)	41(22.0)	8.267[3.119, 21.916]**	0.000
	no	16(25.8)	145(78.0)	1	
MSAF	yes	30(48.4)	42(22.6)	0.712[0.217,2.333]	0.575
	no	32(51.6)	144(77.4)	1	
PIH eclapsia	yes	22(35.5)	25(13.4)	1.889[0.511, 6.981]	0.340
	no	40(64.5)	161(86.6)	1	
UTI or STI	yes	46(74.2)	37(19.9)	14.548[4.911, 43.100]**	0.000
	no	16(25.8)	149(80.1)	1	
Breast feeding	<60 minute	40(64.5)	165(88.7)	1	
	>60 minute	22(35.5)	21(11.3)	3.910[1.271, 12.023]**	0.017
Resuscitation	yes	21(33.9)	15(8.1)	13.256[3.445, 51.010]**	0.000
	no	41(66.1)	171(91.9)	1	
Apgar score at 5th minute	<7	6(9.1)	3(1.6)	3.831[0.033, 47.243]	0.580
	≥7	56(90.3)	183(98.4)	1	
Gestational age	<37	27(43.5)	71(38.2)	1.274[0.409, 3.969]	0.677
	37-42	35(56.5)	115(61.8)	1	
Apgar score at 1st minute	<7	15(24.2)	18(9.7)	0.230[0.024, 2.171]	0.200
	≥7	47(75.8)	168(90.3)	1	
ANC visit	yes	39(62.9)	166(89.2)	1	
	no	23(37.1)	20(10.8)	0.644[0.166, 2.496]	0.524

**significant Variable

Discussion

The finding of this study showed that the duration of rupture of membrane ≥ 18 hours, meconium stained amniotic fluid, intra-partum maternal fever, history of UTI/STI, and breast feeding after 1 hour were independent maternal factors for neonatal sepsis whereas neonatal resuscitation and chlorohexidine application were identified as neonatal determinants of neonatal sepsis.

In this study prolonged rupture of membrane ≥ 18 hours was significantly associated with neonatal sepsis, neonates born from women who had prolonged rupture of membrane ≥ 18 hours had five times more likely to develop neonatal sepsis than that of women who had PROM < 12 hours. This finding is in line with studies conducted in Mekele, Mexico, Jinka, Nepal, USA, China, Pakistan, Northwest Ethiopia (10, 11, 33, 35, 47, 48, 49, 53). These may be due to birth canal is colonized with aerobic and anaerobic pathogens that might cause in ascending amniotic fluid infection and colonization of the neonate at birth. Early rupture of membrane increases the chance of ascending microorganisms from the birth canal into the amniotic sac causing chorioamnionitis and fetal compromise which frequently leads the new born to infection in utero. Mother to fetus transmission of bacterial agents that infect the amniotic fluid and birth canal may occur in uterus more commonly during labor and delivery which results in neonatal sepsis.

Meconium stained amniotic fluid had also a significant association with neonatal sepsis. Those neonates born from mothers who had MSAF were six times more likely to suffer from neonatal sepsis than neonates who had no MSAF. These finding is comparable with a study conducted in Mexico, south Africa, Uganda, Nepal, Ghana (11, 15, 16, 35, 45). This may be due to that, neonates delivered from women with meconium stained amniotic fluid are more liable to aspirate it and fill smaller air ways and alveoli in the lung. And it increases the multiplication of microbes that cause sepsis.

Intra-partum fever was also had a significant association with neonatal sepsis, mothers who had history of intra-partum fever were eight times more risk for their neonates to develop neonatal sepsis than mothers who had no fever. These finding is consistent with studies conducted in Mekele, Wolayta sodo, and Pakistan (32). This might be because of the reason that fever is a sign of local or systemic infections such as chorioamnionitis and urinary tract infection, which are frequently transmitted to the baby in the uterus or during passage through the canal This results in hematogenous spread and vertical transmission of pathogens to the newborn before or during labor and delivery which further results in neonatal sepsis.

In this study neonatal sepsis was strongly associated with maternal history of UTI/ STI, those neonates born from mothers who had history of UTI/STI were fourteen times more likely to have neonatal sepsis than mothers who had no UTI/STI. This is in line with Mekele, Uganda, Jinka, central Ethiopia, and Ghana which showed UTI was significant determinants of neonatal sepsis (10, 16, 33, 34,45). This might be due to, the wall of birth canal in women with urinary tract infection are colonized with pathogens. The most common microorganisms that cause neonatal sepsis are found across the birth canal and possibly increase the risk while the newborn was born and pass through the vaginal wall.

Breast feeding was also had association with neonatal sepsis, Mothers who breast feed their neonate after 1 hour were nearly four times more likely to suffer from neonatal sepsis than that of neonates who breast feed within 1 hour. These finding is consistent with a study conducted in Wolayta sodo (32). Early initiation of breast feeding has different health benefit like increase ability to defense infection and increase the survival rate of children because colostrum is the first milk that is very important for newborns in protecting infection since rich in immunoglobulin G that has great role in disease resistance.

Resuscitation of the newborn had significant association with neonatal sepsis; resuscitated neonates were thirteen times more likely to have neonatal sepsis than their counter parts. This finding is comparable with studies done in Tanzania, Ghana, NW Ethiopia, and another Northwest Ethiopia (3, 45, 53, 54). These findings might be due to the fact that, if procedure of resuscitation is done forcefully, it may cause laceration to the susceptible and easily breakable mucous membrane of the neonate and serve as a route of entry for pathogens from unsterile equipment. It may also lead microbes into the lower air way of the newborn with an immature immune system. This is due to the lumen of airways of the neonate is too narrow, and respiratory secretions are copious compared to older children who could predispose to easily destruction of smaller air sacs and sepsis. Moreover, resuscitation might be done with unsterile equipment, which could introduce microbes into the lungs of the neonate whose immune system is not yet well developed.

Chlorohexidine application had also significant association with the occurrence of neonatal sepsis. Neonates who had no application of chlorohexidine were having a significant risk for neonatal sepsis. This may be due to that umbilical cord is the main portal of entry for microorganisms that cause neonatal sepsis(56, 57).

Limitation of the study

Since the study was done on admitted neonates, thus results might lack generalizability to the entire population.

Conclusion

Among the variables prolonged rupture of membranes, meconium stained amniotic fluid, intra-partum fever, UTI/STI, and not breast feeding within an hour were maternal variables and resuscitation at birth and not application of chlorohexidine ointment on the umbilicus were neonatal variables that were found to be neonatal-related risk factors of neonatal sepsis. Infection prevention strategies and clinical management need to be strengthening and/or implementing by providing special attention for the specified determinants.

Recommendation

Health professionals who are working in NICU and delivery room should use aseptic technique, clean sterilize or disinfect instruments and equipment, and routinely clean the neonatal intensive care unit. Special attention and follow up should be given for neonates who born from mothers who had history of PROM, MSAF, UTI/STI and intra-partum fever. Screen mothers for UTI and STI and treat accordingly. Mobilize and support early health-seeking behavior for health problems during pregnancy, like seeking care for UTI/STI. Encourage mothers to breastfeed their newborns and initiate as early as possible. Give regular infection prevention, health education and promotion on maternal and neonatal health.

Availability of data and materials

All the data are available in the manuscript.

Abbreviations

ANC: Antenatal Care, AOR: Adjusted Odds Ratio, APGAR: Activity, Pulse, Grimace, Appearance, Respiration, APH : Antepartum Hemorrhage, BW: Birth Weight, CBC: Complete Blood Count, CI: Confidence Interval, COR: Crude Odds Ratio, CRP: C-Reactive Protein, EDHS: Ethiopian Demographic and Health Survey, EONS: Early Onset Neonatal Sepsis; Integrated Management of Neonatal and Childhood Illness, GSGH: Gebretsadik Shawo General Hospital, LONS: Late onset neonatal sepsis, MSAF: Meconium Stained Amniotic Fluid, NGT: Nasogastric Tube

NMR: Neonatal Mortality Rate, NICU: Neonatal intensive care unit, PIH: Pregnancy Induced Hypertension PROM: Premature Rupture of Membrane, PSBI: Possible Serious Severe Bacterial Infection, SDG: -Sustainable Development Goal, SPSS: Statistical package for Social Science SSA: Sub-Saharan Africa, UTI: Urinary Tract Infection, TPH: Telo Primary Hospital, WHO: World Health Organization and WPH: Wacha Primary Hospital

Acknowledgments

The authors acknowledged Bahir Dar University, Wacha Primary Hospital, Telo Primary Hospital, Gebretsadik Shawo General Hospital, and data collectors.

Funding

There is no funding for this work.

Contributions

MB is the pioneer to draft the manuscript. GG & AG are research assistant who has been actively engaged in each step in the research process. MB, GG & AG were involved in the conception and design of the whole research. MB, GG & AG is involved in the analyzed and interpreted of findings. All authors have read and approved their ownership the manuscript.

Corresponding authors

*Correspondence to Melesse Belayneh; meleseayeh@gmail.com

Ethics declarations

Ethics approval and consent to participate

Ethical clearance was obtained from Bahir Dar University, College of Medicine and Health sciences, Institutional Ethics Review Board (IRB) with letter refrence number (IRB 2345/2021). Informed consent was taken from each participant.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no any competing interests.

Reference

1. Chaurasia S, Sivanandan S, Agarwal R, Ellis S, Sharland M, Sankar MJ. Neonatal sepsis in South Asia: huge burden and spiralling antimicrobial resistance. *bmj*. 2019;364.
2. Goldstein B, Giroir B, Randolph A. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatric critical care medicine*. 2005;6(1):2-8.
3. Jabiri A, Wella HL, Semiono A, Saria A, Protas J. Prevalence and factors associated with neonatal sepsis among neonates in Temeke and Mwananyamala Hospitals in Dar es Salaam, Tanzania. *Tanzania Journal of Health Research*. 2016;18(4).
4. Palatnik A, Liu LY, Lee A, Yee LM. Predictors of early-onset neonatal sepsis or death among newborns born at < 32 weeks of gestation. *Journal of Perinatology*. 2019;39(7):949-55.
5. Stefanovic IM. Neonatal sepsis. *Biochemia medica*. 2011;21(3):276-81.
6. Dong Y, Speer CP. Late-onset neonatal sepsis: recent developments. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2015;100(3):F257-F63.

7. Lancet. The Young Infants Clinical signs that predict severe illness in children under age 2 months: a multicentre study. *Lancet* (London, England). 2008:135-42.
8. Opiyo N EM. What clinical signs best identify severe illness in young infants aged 0-59 days in developing countries? A systematic review. *Archives of disease in childhood*. 2011:1052-9.
9. Moges F, Eshetie S, Yeshitela B, Abate E. Bacterial etiologic agents causing neonatal sepsis and associated risk factors in Gondar, Northwest Ethiopia. *BMC pediatrics*. 2017;17(1):137.
10. Gebremedhin D, Berhe H, Gebrekirstos K. Risk Factors for Neonatal Sepsis in Public Hospitals of Mekelle City, North Ethiopia, 2015: Unmatched Case Control Study. *PLoS One*. 2016;11(5):e0154798.
11. Leal YA, Álvarez-Nemegyei J, Velázquez JR, Rosado-Quiab U, Diego-Rodríguez N, Paz-Baeza E, et al. Risk factors and prognosis for neonatal sepsis in southeastern Mexico: analysis of a four-year historic cohort follow-up. *BMC pregnancy and childbirth*. 2012;12(1):48.
12. Wang ME, Patel AB, Hansen NI, Arlington L, Prakash A, Hibberd PL. Risk factors for possible serious bacterial infection in a rural cohort of young infants in central India. *BMC public health*. 2016;16(1):1097.
13. Weldearegawi B MY, Abera SF, Ashebir Y, Haile F, Mulugeta A, Eshetu F, Spigt M. Infant mortality and causes of infant deaths in rural Ethiopia: a population-based cohort of 3684 births. *BMC public health* 2015.
14. John B, David M, Mathias L, Elizabeth N. Risk factors and practices contributing to newborn sepsis in a rural district of Eastern Uganda, August 2013: a cross sectional study. *BMC research notes*. 2015;8(1):339.
15. Schrag SJ, Cutland CL, Zell ER, Kuwanda L, Buchmann EJ, Velaphi SC, et al. Risk factors for neonatal sepsis and perinatal death among infants enrolled in the prevention of perinatal sepsis trial, Soweto, South Africa. *The Pediatric infectious disease journal*. 2012;31(8):821-6.
16. Santhanam S, Arun S, Rebekah G, Ponmudi NJ, Chandran J, Jose R, et al. Perinatal risk factors for neonatal early-onset group B streptococcal sepsis after initiation of risk-based maternal intrapartum antibiotic prophylaxis—a case control study. *Journal of tropical pediatrics*. 2018;64(4):312-6.
17. Assemie MA, Alene M, Yismaw L, Ketema DB, Lamore Y, Petrucka P, et al. Prevalence of Neonatal Sepsis in Ethiopia: A Systematic Review and Meta-Analysis. *International Journal of Pediatrics*. 2020;2020.
18. Chryssoula Tzialla, Paolo Manzoni, Cristian Achille, Lina Bollani, Mauro Stronati, Borghesi A. New Diagnostic Possibilities for Neonatal Sepsis. *American Journal of Perinatology*. 2018;Vol. 35.
19. Tam P-YI, Bendel CM. Diagnostics for neonatal sepsis: current approaches and future directions. *Pediatric research*. 2017;82(4):574-83.
20. Zea-Vera A, Ochoa TJ. Challenges in the diagnosis and management of neonatal sepsis. *Journal of tropical pediatrics*. 2015;61(1):1-13.
21. Sorsa A. Diagnostic significance of white blood cell count and c-reactive protein in neonatal sepsis; Asella Referral Hospital, South East Ethiopia. *The Open Microbiology Journal*. 2018;12:209.
22. Yamey G, Horváth H, Schmidt L, Myers J, Brindis CD. Reducing the global burden of Preterm Birth through knowledge transfer and exchange: a research agenda for engaging effectively with policymakers. *Reproductive health*. 2016;13(1):26.
23. Aijaz N, Huda N, Kausar S. Disease burden of NICU, at a tertiary care hospital, Karachi. *Journal of the Dow University of Health Sciences (JDUHS)*. 2012;6(1):32-5.
24. Edmond K, Zaidi A. New approaches to preventing, diagnosing, and treating neonatal sepsis. *PLoS Med*. 2010;7(3):e1000213.
25. Unicef. Committing to child survival: a promise renewed. *eSocialSciences*; 2015.
26. Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kisssoon N. The global burden of paediatric and neonatal sepsis: a systematic review. *The Lancet Respiratory Medicine*. 2018;6(3):223-30.

27. Tewabe T, Mohammed S, Tilahun Y, Melaku B, Fenta M, Dagnaw T, et al. Clinical outcome and risk factors of neonatal sepsis among neonates in Felege Hiwot referral Hospital, Bahir Dar, Amhara Regional State, North West Ethiopia 2016: a retrospective chart review. *BMC research notes*. 2017;10(1):265.
28. Organization WH. Every newborn: an action plan to end preventable deaths. 2014.
29. Ethiopia E. Health Survey. Key Indicators Report. Central Statistical Agency Addis Ababa, Ethiopia. Rockville: The DHS program, ICF; 2016. 2018.
30. Organization WH. World health statistics 2016: monitoring health for the SDGs sustainable development goals: World Health Organization; 2016.
31. survey. EDah. Key indicator report. CSA Addis Ababa, Ethiopia. 2019.
32. Alemayehu A, Alemayehu M, Arba A, Abebe H, Goa A, Paulos K, et al. Predictors of Neonatal Sepsis in Hospitals at Wolaita Sodo Town, Southern Ethiopia: Institution-Based Unmatched Case-Control Study, 2019. *International Journal of Pediatrics*. 2020;2020.
33. Ketema E, Mamo M, Miskir D, Hussen S, Boti N. Determinants of neonatal sepsis among neonates admitted in a neonatal intensive care unit at Jinka General Hospital, Southern Ethiopia. *International Journal of Nursing and Midwifery*. 2019;11(3):18-24.
34. Chan GJ, Lee AC, Baqui AH, Tan J, Black RE. Prevalence of early-onset neonatal infection among newborns of mothers with bacterial infection or colonization: a systematic review and meta-analysis. *BMC infectious diseases*. 2015;15(1):118.
35. Thapa B, Thapa A, Aryal DR, Thapa K, Pun A, Khanal S, et al. Neonatal sepsis as a major cause of morbidity in a tertiary center in Kathmandu. *Journal of Nepal Medical Association*. 2013;52(192).
36. Shehab El-Din EMR, El-Sokkary MMA, Bassiouny MR, Hassan R. Epidemiology of neonatal sepsis and implicated pathogens: a study from Egypt. *BioMed research international*. 2015;2015.
37. Dramowski A, Aucamp M, Bekker A, Mehtar S. Infectious disease exposures and outbreaks at a South African neonatal unit with review of neonatal outbreak epidemiology in Africa. *International Journal of Infectious Diseases*. 2017;57:79-85.
38. Velaphi SC, Westercamp M, Moleleki M, Pondo T, Dangor Z, Wolter N, et al. Surveillance for incidence and etiology of early-onset neonatal sepsis in Soweto, South Africa. *PloS one*. 2019;14(4):e0214077.
39. Abate BB, Kasie AM, Reta MA, Kassaw MW. Neonatal sepsis and its associated factors in East Africa: a systematic review and meta-analysis. *International Journal of Public Health*. 2020:1-11.
40. Getabelew A, Aman M, Fantaye E, Yeheyis T. Prevalence of neonatal sepsis and associated factors among neonates in neonatal intensive care unit at selected governmental hospitals in Shashemene Town, Oromia Regional State, Ethiopia, 2017. *International journal of pediatrics*. 2018;2018.
41. Agnche Z, Yeshita HY, Gonete KA. Neonatal Sepsis and Its Associated Factors Among Neonates Admitted to Neonatal Intensive Care Units in Primary Hospitals in Central Gondar Zone, Northwest Ethiopia, 2019. *Infection and Drug Resistance*. 2020;13:3957.
42. Ahmed M, Magzoub O. Risk factors for neonatal sepsis in paediatric ward at Khartoum North Teaching Hospital, Sudan. *Basic Research Journal of Medicine and Clinical Science*. 2015;4(1):37-43.
43. Ashraf MN, ul Haq MZ, Ashraf MW, Sajjad R, Ahmed F. Early onset neonatal sepsis in preterm premature rupture of membranes. *Pakistan Armed Forces Medical Journal*. 2015;65(2):226-30.
44. Salem SY, Sheiner E, Zmora E, Vardi H, Shoham-Vardi I, Mazor M. Risk factors for early neonatal sepsis. *Archives of gynecology and obstetrics*. 2006;274(4):198.
45. Siakwa M, Kpikpitse D, Mupepi SC, Semuatu M. Neonatal sepsis in rural Ghana: A case control study of risk factors in a birth cohort. 2014.

46. Qiu X, Zhang L, Tong Y, Qu Y, Wang H, Mu D. Interleukin-6 for early diagnosis of neonatal sepsis with premature rupture of the membranes: A meta-analysis. *Medicine*. 2018;97(47).
47. Chan GJ, Lee AC, Baqui AH, Tan J, Black RE. Risk of early-onset neonatal infection with maternal infection or colonization: a global systematic review and meta-analysis. *PLoS Med*. 2013;10(8):e1001502.
48. Jiang Z, Ye G. 1: 4 matched case-control study on influential factor of early onset neonatal sepsis. *Eur Rev Med Pharmacol Sci*. 2013;17(18):2460-6.
49. Alam MM, Saleem AF, Shaikh AS, Munir O, Qadir M. Neonatal sepsis following prolonged rupture of membranes in a tertiary care hospital in Karachi, Pakistan. *The Journal of Infection in Developing Countries*. 2014;8(01):067-73.
50. Hayun M, Alasiry E, Daud D, Febriani DB, Madjid D. The risk factors of early onset neonatal sepsis. *American Journal of clinical and experimental medicine*. 2015;3(3):78-82.
51. Tran H, Doyle L, Lee K, Dang N, Graham S. A high burden of late-onset sepsis among newborns admitted to the largest neonatal unit in central Vietnam. *Journal of Perinatology*. 2015;35(10):846-51.
52. Kung Y-H, Hsieh Y-F, Weng Y-H, Lien R-I, Luo J, Wang Y, et al. Risk factors of late-onset neonatal sepsis in Taiwan: A matched case-control study. *Journal of Microbiology, Immunology and Infection*. 2016;49(3):430-5.
53. Alemu M, Ayana M, Abiy H, Minuye B, Alebachew W, Endalamaw A. Determinants of neonatal sepsis among neonates in the northwest part of Ethiopia: case-control study. *Ital J Pediatr*. 2019;45(1):150.
54. Akalu TY, Gebremichael B, Desta KW, Aynalem YA, Shiferaw WS, Alamneh YM. Predictors of neonatal sepsis in public referral hospitals, Northwest Ethiopia: A case control study. *Plos one*. 2020;15(6):e0234472.
55. department Kzh. Annual report of MCH in kaffa zone health institution. 2020.
56. Kalathia MB SP, Parmer PNParkih YN. study of umbilical cord blood culture in diagnosis of neonatal sepsis. 2013;2:169-72.
57. Tilayie FG YD, Begu A. effect of chlorohexidine cord application on prevention of neonatal sepsis in developing countries: systematic review and meta analysis. 2019.