Extending non-targeted exposure discovery of environmental chemical exposures during pregnancy and their association with pregnancy complications—a cross-sectional study

Jessica Trowbridge[1], Dimitri Abrahamsson[1], Ting Jiang[2], Miaomiao Wang[2], June-Soo Park[2], Rachel Morello-Frosch[3], Marina Sirota[4], Dana E. Goin[1], Marya Zlatnick[1] and Tracey J. Woodruff[1]

[1]Department of Obstetrics, Gynecology and Reproductive Sciences, Program on Reproductive Health and the Environment, University of California San Francisco, San Francisco 94143 California
[2]Department of Toxic Substances Control, Environmental Chemistry Laboratory, California Environmental Protection Agency, Berkeley 94710, California
[3]Department of Environmental Science and Policy Management and School of Public Health, University of California, Berkeley, Berkeley 94720 California
[4]Bakar Computational Health Sciences Institute and Department of Pediatrics, University of California San Francisco, San Francisco 94158, California

*Corresponding author: Tracey J. Woodruff, tracey.woodruff@ucsf.edu

Abstract

Background: Non-targeted Analysis (NTA) methods identify novel exposures; however, few chemicals have been quantified and interrogated with pregnancy complications.

Objectives: We characterize levels of nine exogenous and endogenous chemicals in maternal and cord blood identified, selected, and confirmed in prior NTA steps including: linear and branched isomers perfluorooctane sulfonate (PFOS); perfluorohexane sulfonate (PFHxS); monoethylhexyl phthalate; 4-nitrophenol; tetraethylene glycol; tridecanedioic acid, octadecanedioic acid; and deoxycholic acid. We evaluate relationships between maternal and cord levels and the relationship gestational diabetes mellitus (GDM) and hypertensive disorders of pregnancy in a diverse pregnancy cohort in San Francisco.

Methods: We collected matched maternal and cord serum samples from 302 pregnant people at delivery from the Chemicals in Our Bodies cohort in San Francisco. Chemicals were identified via NTA and quantified using targeted approaches. We calculate distributions and Spearman correlation coefficients testing the relationship of chemicals within and between the maternal and cord blood matrices. We used logistic regression to calculate the odds of GDM and hypertensive disorders of pregnancy associated with an interquartile range increase in maternal chemical exposures.

Results: We detected linear PFOS, PFHxS, octadecanedioic acid, and deoxycholic acid in at least 97% of maternal samples. Correlations ranged between -0.1 and 0.9. We observed strong correlations between cord and maternal levels of PFHxS (coefficient = 0.9), linear PFOS (0.8), and branched PFOS (0.8). An IQR increase in linear PFOS, branched PFOS, and octadecanedioic acid is associated with increased odds of GDM [OR (95%CI): 1.43 (0.96, 2.14), 1.56 (1.00, 2.44), and 1.26 (0.83, 1.92) respectively] and tridecanedioic acid positively associated with hypertensive disorders of pregnancy [1.28 (0.90, 1.86)].

Discussion: We identified both exogenous and endogenous chemicals, two of which (octadecanedioic acid and tridecanedioic acid) have both endogenous and exogenous sources, and which have seldom been quantified in pregnant people or related to pregnancy complications.

Keywords: Non-targeted analysis, linear and branched PFOS, tridecanedioic acid, fatty acids, gestational diabetes mellitus, pregnancy hypertension

1. Introduction

Prenatal exposures to environmental chemicals are ubiquitous in the United States (U.S.)[1] and studies have demonstrated they can have lifelong consequences for maternal and child health outcomes,[2] such as neurodevelopment, cardiovascular disease, and reproduction. However, fewer than 1% of over 80,000 chemicals produced and used in the U.S. are regularly biomonitored[3] and fewer still have been assessed for adverse health outcomes during pregnancy. Non-targeted analysis (NTA) coupled with a prioritization step and quantification using targeted methods can facilitate the identification and quantitation of previously unmeasured chemical exposures.[4] Additionally, by identifying and measuring novel chemical exposures we can investigate their relationship with pregnancy complications, facilitating the identification of opportunities for prevention.

Increasingly, studies are applying NTA to better characterize exposure to toxic environmental chemicals seldom measured in pregnant people.[5,6] NTA studies in pregnant and non-pregnant populations have identified both known and novel chemical exposures, including PFAS congeners, pesticides, flame retardants, and other industrial chemicals, which have been linked with pregnancy complications.[6,7] For example, PFAS has been linked with increased risk of miscarriage and preeclampsia,[8] and chemicals with fatty acid-like structures, including PFAS and other long-chain hydrocarbons, have been shown to interact with endogenous fatty acid metabolism impacting placental development and may contribute to pregnancy complications and adverse birth outcomes.[6,9–11] Similarly, NTA studies have been able to link novel chemical exposures and metabolism biomarkers.[6,12] However, the vast majority of novel chemicals identified through NTA have not been evaluated for associations with pregnancy related complications.

This study builds on previous NTA work done on the Chemicals in Our Bodies-2 (CIOB2) cohort, in which we first identified chemical features using high resolution mass spectrometry (HRMS)

and confirmed chemical identities via comparison to a standard.[5,6,13] The aim of the present study is to

describe the quantification of a subset of chemicals identified and prioritized through this prior NTA and

to evaluate relationships between chemical levels and pregnancy complications. Supplementing the non-

targeted discovery of chemicals with targeted quantification described in this analysis addresses some of

the limitations of NTA; namely, targeted approaches increase the sensitivity to measure and compare

chemical concentrations and facilitate assessment of relationships between chemicals and pregnancy

complications.

2. Methods

2.1 Data and Sample Collection

We enrolled pregnant participants in the CIOB2 cohort between 2014 and 2018 from the prenatal

clinics at the Zuckerberg San Francisco General Hospital (SFGH) and Mission Bay or Moffit Long

(MB/ML) hospitals. Participant recruitment and sample collection is described elsewhere.[4,5,14] Briefly,

participants were recruited during the second trimester of pregnancy, with eligibility criteria including

18-40 years of age, English or Spanish speaking, with singleton pregnancies, and no diagnosed

pregnancy complications at recruitment. Paired maternal and umbilical cord blood were collected from

participants who agreed to bank their samples and be contacted for participation in future studies.

Maternal samples were collected at delivery and cord blood was collected after delivery but before

clamping whenever possible. Blood was collected in BD Vacutainer Plus serum tubes, spun at 3000 rpm

to collect the serum, and stored at -80°C until analysis. Study protocols were approved by the

Institutional Review Boards at the University of California, San Francisco and Berkeley (# 13-12160).

Demographic information was collected via interviews conducted during the second trimester of

pregnancy and included maternal age, race and ethnicity (white, Latinx, and Non-Hispanic Asian/Pacific

Islander, other race—due to low numbers of Black, Native American, and multiracial participants, these

groups were collapsed into "other" for model specification), maternal education (some college or less,

completed a college degree, >college degree), household income (<40k per year, 40 to 100k, >100k),

nativity (born in the U.S. yes or no). Pre-pregnancy body mass index (BMI kg/m2; <18.5, 18.5 to <25,

25 to <35, >=35)[15] and hospital of delivery [Zuckerberg San Francisco General Hospital (SFGH) or

Mission Bay/ Moffit Long (MB/ML)] were abstracted from the medical record.

Pregnancy complications were identified by clinical diagnoses and abstracted from the medical

record. These included the binary outcomes of: gestational diabetes mellitus (GDM), preeclampsia, and

gestational hypertension. Preeclampsia and pregnancy related hypertension were combined into one

outcome called hypertensive disorder of pregnancy because they make up part of a larger spectrum of

pregnancy complications related to hypertension.[16,17]

2.2 Chemical selection

Nine environmental chemicals were measured in maternal and cord samples using targeted

approaches described below. Linear and branched isomers of perfluorooctane sulfonate (PFOS);

perfluorohexane sulfonate (PFHxS); monoethylhexyl phthalate (MEHP), a metabolite of diethylhexyl

phthalate, a common plasticizer[18]; 4-nitrophenol, a known endocrine disruptor used in pesticides, dyes,

and pharmaceuticals[19]; tetraethylene glycol, a plasticizer and solvent used in metal lubricants, printing

inks, and cosmetics;[20] tridecanedioic acid and octadecanedioic acid which are endogenous fatty acids

that also have exogenous sources primarily from plastics synthesis;[21–23] and deoxycholic acid, a bile acid

and endogenous biomarker associated with GDM **(Table 1)**.[24]

The selection and confirmation procedures for selection of these chemicals is described

elsewhere.[5,6,13] Briefly, in previous NTA studies, 30 matched maternal and cord samples were scanned

using HRMS in positive and negative ionization modes identifying suspect features. Detected features

were matched to an in-house database and features were then prioritized for MS/MS fragmentation

based on the ubiquity of exposure, feature intensity differences across demographic variables, and

correlation between maternal and cord samples. Fifteen chemicals were confirmed at level 1 confidence

via comparison to analytical standards.[28] Level 1 confirmed features included monoethylhexyl phthalate,

4-nitrophenol, tridecanedioic acid, and octadecanedioic acid. Subsequent generalized suspect screening

analysis of this cohort was expanded to include 295 matched maternal and cord samples and a more

comprehensive database including both endogenous and exogenous compounds. This resulted in

nineteen chemicals confirmed with analytical standards to level 1 confidence, including those previously

identified, and adding PFHxS, PFOS, and deoxycholic acid which were tentatively identified through

MS/MS fragmentation (confidence level 2). We selected chemicals for targeted quantification if the

chemical was identified in previous NTA with level 1 or level 2 confidence, is a widely used industrial

compound with high potential for exposure, is associated with pregnancy complications, or is not

regularly biomonitored by the Center for Disease Control or other large biomonitoring studies.[5]

2.3 Targeted chemical analysis

Quantitation analysis of the nine selected chemicals was carried out using Agilent MassHunter

Quantitative Analysis software package (Agilent Technologies Inc., Version 10.0). Analytical grade

standards were purchased from suppliers (Sigma Aldrich and Thermo Fisher) and serial dilutions of the

standards were prepared in methanol. Calibration curve of each of the specific standard was acquired

under the same condition as serum sample analysis, using the LC/QTOF-MS (Agilent 1290 UPLC

interfaced with Agilent iFunnel 6550 QTOF-MS system) as described previously.[5,6,13] M2-PFOA

(Wellington Laboratories, Ontario, Canada) was used as internal standard for negative ionization

chemicals, while D15-TPP (Cambridge Isotope Laboratories, Tewksbury, MA) for positive ionization

chemicals. Particularly, PFOS in serum samples were separated chromatically, thus both total and linear

PFOS quantitation were calculated, and branched PFOS concentrations were obtained by subtracting

linear portion from the total PFOS.

2.4 Statistical analysis

To address chemical levels below the limit of detection we used machine reported values when

available or, if not available, we imputed levels below the LOD with LOD/sqrt(2). We assessed the

detection frequency (percent above the limit of detection) and limited our statistical analysis to

chemicals with a detection frequency of 67% or higher. We described the distribution of chemicals in

maternal and cord blood samples, and calculated Spearman correlation coefficients for chemicals within

and between maternal and cord samples. We natural log-transformed the data for use in statistical

models and applied logistic regression models to assess the relationship between maternal chemical

exposure and pregnancy complications. Potential confounders were identified *a priori* (**Supplemental**

Figure 1), and our final models were adjusted for maternal age, hospital of delivery, and race/ethnicity.

We considered hospital of delivery as a cumulative indicator of socioeconomic status (including income,

educational attainment, and insurance status). Patients in both hospitals are racially and ethnically

diverse. However, SFGH is a safety net hospital that serves a large proportion of patients without

private health insurance, while MB/ML serves patients of higher socioeconomic status and with private

health insurance. Race and ethnicity are included in models as an indicator of the exposure to structural

racism, that is independently associated with chemical exposure and adverse pregnancy complications.

We calculated the interquartile range (IQR) of each natural log transformed chemical. Beta coefficients

and 95% confidence intervals (CI) were multiplied IQR and exponentiated to identify the odds ratio

(OR) of each pregnancy complication for each IQR increase in exposure level.

3. Results

Participants, demographics, and medical record data

The analysis included 302 participants (**Table 2**). The average age at delivery was 33 years of age. Forty-three percent of participants identified as white and 32% as Latinx or Hispanic. Most participants completed college or graduate degrees (>65%), were born in the U.S. (54%), and had a household income of over $100,000 per year (56%). For the outcomes, 21% of participants had a diagnosis of GDM, 13% gestational hypertension, and 7% preeclampsia.

3.1 Distribution and correlation of chemical levels in maternal samples

Participants had at least four chemicals in maternal samples and three chemicals in cord samples. PFHxS, linear PFOS, deoxycholic acid, and octadecanedioic acid were all detected in at least 97% of maternal samples. Deoxycholic acid, tridecanedioic acid, PFHxS, were detected in at least 87% of cord blood samples (Table 3). Since we estimated branched PFOS levels by subtracting linear PFOS from the total PFOS we did not calculate a detection frequency for this chemical. Chemical detection frequency varied between maternal and cord blood samples. Linear PFOS, and 4-nitrophenol were detected more often maternal samples while monoethylhexyl phthalate, and tridecanedioic acid were detected more frequently in cord blood. Similarly, the average chemical levels varied across the different matrices with branched PFOS, linear PFOS, deoxycholic acid, and octadecanedioic acid found at higher levels in maternal samples compared to cord samples, while tridecanedioic acid was higher in cord than in maternal samples.

Monoethylhexyl phthalate, tetraethylene glycol, and 4-nitrophenol were each detected in fewer than 50% of maternal samples, however 4-nitrophenol was detected in more maternal serum samples than cord samples, while monoethylhexyl phthalate was found in more cord blood samples. Due to their

low detection frequency, monoethylhexyl phthalate, tetraethylene glycol, and 4-nitrophenol, were

excluded from further analyses.

When assessing the Spearman correlation coefficients (**Figure 2**) of the same chemical across

maternal and cord samples, we found that linear PFOS, and branched PFOS and PFHxS were highly

correlated in the maternal and cord samples (correlation coefficients of 0.8, 0.8, and 0.9 respectively).

When looking at the correlations of different chemicals in the maternal and cord samples we found

moderate correlations between cord-linear PFOS and maternal-branched PFOS, as well as between cord-

linear PFOS and maternal-PFHxS (coefficients 0.7 and 0.5 respectively). In the cord blood samples,

linear PFOS was highly correlated with branched PFOS and moderately with PFHxS (coefficients of 0.8

to 0.6 respectively), and octadecanedioic acid was moderately correlated with tridecanedioic acid

(coefficient of 0.4). In the maternal samples, the relationship between linear and branched PFOS had a

correlation coefficient of 0.8, branched PFOS and PFHxS a correlation of 0.6, and between linear PFOS

and PFHxS of 0.6. The PFAS chemicals were weakly correlated with the fatty acids—octadecanedioic

acid, tridecanedioic acid—and neither PFAS, nor the fatty acids were correlated with deoxycholic acid,

within or between the maternal serum and cord blood samples.

3.2 Association between chemical exposure with pregnancy complications

From our logistic models we observed a trend of increased odds of GDM for each IQR increase in levels

of exposure to environmental chemicals and increased odds of pregnancy hypertensive disorders with

tridecanedioic acid (Table 4). For the relationship between chemical exposures and GDM, all the

chemicals included in this analysis had an OR above 1. Stronger associations were observed for

branched PFOS, linear PFOS, and octadecanedioic acid; The unadjusted relationship between linear

PFOS and branched PFOS with GDM was statistically significant. The direction and measures of

association we observed between chemical levels and hypertensive disorders of pregnancy was less

consistent than for GDM. Exposure to tridecanedioic acid was associated with an increase in the odds of

hypertensive disorders of pregnancy [OR (95%CI): 1.28(0.90, 1.86)]. Exposure to tridecanedioic acid

was also associated in increased odds of the individual outcomes that made up hypertensive disorders of

pregnancy—preeclampsia and gestational hypertension (**Supplemental Table 1**). The ORs for PFHxS,

octadecanedioic acid, and deoxycholic acid with gestational hypertension approximated 1. Linear and

branched PFOS had an OR of <1 but the confidence intervals included the null value.

4. Discussion

Building on NTA steps from prior studies[4,6,13], we confirmed and quantified the presence of nine

chemicals, some of which have not been previously measured in pregnant people, such as tridecanedioic

acid, octadecanedioic acid, and tetraethylene glycol, or rarely measured in pregnant people like 4-

nitrophenol. In addition, a subset of these were moderately to strongly associated with GDM and

hypertensive disorders of pregnancy. Linear and Branched PFOS were associated with increased odds of

GDM and tridecanedioic acid was associated with hypertensive disorders of pregnancy. This study adds

to our understanding of the associations between environmental exposures during pregnancy and

pregnancy complications.

This study provides further evidence of prenatal exposure to environmental chemicals, some of

which have positive associations with pregnancy complications. A 2020 study suggests that PFAS

exposure during pregnancy may also affect metabolic pathways that adversely impact placental

development[9] and while evidence remains mixed, PFAS has been shown to be associated with

hypertensive disorders of pregnancy, preeclampsia, and GDM.[9,29–31] The linear and branched forms of

PFOS have not previously been well characterized during pregnancy. Furthermore, metabolism,

excretion, and placental transfer may vary by the linear and branched forms of this ubiquitous chemical

and lead to different health outcomes and pregnancy complications.[25]

Significant positive correlations between branched PFOS, linear PFOS and PFHxS in the

maternal and cord samples suggest that they may share a common source of exposure. Indeed,

traditional PFOS production yielded a mixture of branched and linear isomers with approximately 30%

of isomers in the branched form.[25] From linear models we found that the linear and branched forms of

PFOS may have a relationship with pregnancy complications. Branched and Linear PFOS were

significantly associated with GDM in unadjusted models and although the relationship was attenuated in

adjusted models, the direction and magnitude of the association were maintained. Scientific evidence

indicates transplacental transfer of PFAS chemicals[6,32,33] and gestational diabetes may facilitate

transplacental transfer of PFOS.[34] Linear PFOS makes up a larger proportion of the measured PFOS,[25] is

hypothesized to remain longer in maternal serum and is therefore more available in maternal samples

than branched PFOS, which may contribute to the differences in association we observed with

GDM.[35,36]

Previous studies have found that increased exposure to fatty acids may be associated with an

increased risk of hypertensive disorders of pregnancy due to disruption of fatty acid regulation and

metabolism.[37] We saw an association between tridecanedioic acid and hypertensive disorders of

pregnancy, but not for octadecanedioic acid. Similarly, although the bile acid, deoxycholic acid, is

associated GDM in other studies,[24] we did not see elevated risk of these pregnancy complications

associated with deoxycholic acid in our study. The endogenous chemicals in our study have seldom been

studied in pregnant women and the fatty acids are also known to have exogenous sources and uses.

Tetraethylene glycol, 4-nitrophenol and monoethylhexyl phthalate were present in a small subset

of our participants but were not evaluated for an association with pregnancy complications because their

detection frequency in maternal samples was below 65%. In addition, these chemicals have short half-

lives and are quickly metabolized and excreted in urine contributing to their low detection frequency in

serum samples.[19,26,38] These are high production volume chemicals that may interact with endocrine and

metabolic pathways; however, their relationship with adverse reproductive or pregnancy outcomes has

not been previously assessed, warranting further study. Despite their low detection frequency and that

they were measured in serum rather than in urine our study demonstrates that novel exposures to these

environmental chemicals can still be identified through non-targeted methods and confirmed through

targeted approaches.

There are several limitations that may influence the associations we observed in our study. First

is the timing of the collection of samples and assessment of health outcomes. This is a cross-sectional

study with samples collected at delivery and pregnancy complications were identified from clinical

diagnosis abstracted from medical record. Because of this we cannot assess temporality of our exposures

and outcome. Nevertheless, the chemicals included in this study are industrial chemicals, many of which

are high production volume chemicals that have a high potential for exposure. For example, PFAS

chemicals are ubiquitous in biological matrices and the environment, and tridecanedioic acid and

octadecanedioic acid are used in the synthesis of plastics and are considered high production volume

chemicals.[21] While the chemical measurements reflect a snapshot of an individual's exposure history

during pregnancy the sources and uses of these environmental chemicals imply that there is a potential

for prolonged exposure throughout the pregnancy. Studies show that PFAS levels decrease in later

trimesters of pregnancy,[39,40] thus the PFAS levels we measured at delivery may be lower than in other

trimesters of pregnancy biasing our results toward to the null. Another potential limitation is that levels

of branched PFOS were estimated and not quantified with a standard and thus the levels of branched

PFOS may reduce precision of calculated measures of association. Likewise, the low detection

frequency of 4-nitrophenol, tetraethylene glycol, or monoethylhexyl phthalate precluded our ability to

assess the chemical-outcome relationship. However, these chemicals remain understudied in pregnant

people and are of high concern due to their widespread use as industrial chemicals and limited

understanding of their health effects. Finally, because our recruitment criteria indicated that participants

with pregnancy complications by the second trimester we may have missed earlier manifestations of

pregnancy complications. This could undercount the number of GDM and hypertensive disorders of

pregnancy biasing our results toward the null. However, diagnosis of hypertensive disorders of

pregnancy and testing for GDM usually occurs in the third trimester of pregnancy while our participants

were recruited in the second.[41,42]

In this analysis we identified and quantified exogenous and endogenous environmental

chemicals, some of which have not been previously measured during pregnancy nor evaluated for

associations with pregnancy complications. Our study supports prior reports of associations between the

branched and linear forms of PFOS and GDM and adds to the limited research on exposure to

tridecanedioic acid and octadecanedioic acid, endogenous fatty acids that also have industrial sources.

Finally, this study further illustrates the benefit of pairing non-targeted and targeted approaches to

identify, confirm, and quantify novel chemical exposures. Combining NTA methods with targeted

chemicals measurement can be further applied to identifying, quantifying, and evaluating metabolism

biomarkers that are on the pathway between exposure and downstream adverse health effects.

5. References

Woodruff TJ, Zota AR, Schwartz JM. Environmental chemicals in pregnant women in the United States: NHANES 2003-2004. *Environ Health Perspect* 2011; **119**: 878–885.

American College of Obstetricians and Gynecologists' Committee on Obstetric Practice. Reducing Prenatal Exposure to Toxic Environmental Agents: ACOG Committee Opinion, Number 832. *Obstet Gynecol* 2021; **138**: e40–e54.

CDC. National Report on Human Exposure to Environmental Chemicals | CDC. Center for Disease Control and Prevention. 2019.https://www.cdc.gov/exposurereport/index.html (accessed 18 Nov2018).

Wang A, Gerona RR, Schwartz JM, Lin T, Sirota M, Morello-Frosch R *et al.* A Suspect Screening Method for Characterizing Multiple Chemical Exposures among a Demographically Diverse Population of Pregnant Women in San Francisco. *Environ Health Perspect* 2018; **126**: 077009.

Wang A, Abrahamsson DP, Jiang T, Wang M, Morello-Frosch R, Park J-S *et al.* Suspect Screening, Prioritization, and Confirmation of Environmental Chemicals in Maternal-Newborn Pairs from San Francisco. *Environ Sci Technol* 2021; **55**: 5037–5049.

Panagopoulos Abrahamsson D, Wang A, Jiang T, Wang M, Siddharth A, Morello-Frosch R *et al.* A Comprehensive Non-targeted Analysis Study of the Prenatal Exposome. *Environ Sci Technol* 2021; **55**: 10542–10557.

Grashow R, Bessonneau V, Gerona RR, Wang A, Trowbridge J, Lin T *et al.* Integrating Exposure Knowledge and Serum Suspect Screening as a New Approach to Biomonitoring: An Application in Firefighters and Office Workers. *Environ Sci Technol* 2020; **54**: 4344–4355.

Gao X, Ni W, Zhu S, Wu Y, Cui Y, Ma J *et al.* Per- and polyfluoroalkyl substances exposure during pregnancy and adverse pregnancy and birth outcomes: A systematic review and meta-analysis. *Environ Res* 2021; **201**: 111632.

Szilagyi JT, Avula V, Fry RC. Perfluoroalkyl Substances (PFAS) and Their Effects on the Placenta, Pregnancy, and Child Development: a Potential Mechanistic Role for Placental Peroxisome Proliferator-Activated Receptors (PPARs). *Curr Environ Health Rep* 2020; **7**: 222–230.

Ledda C, Fiore M, Santarelli L, Bracci M, Mascali G, D'Agati MG *et al.* Gestational Hypertension and Organophosphorus Pesticide Exposure: A Cross-Sectional Study. *Biomed Res Int* 2015; **2015**: 280891.

Kahn LG, Trasande L. Environmental Toxicant Exposure and Hypertensive Disorders of Pregnancy: Recent Findings. *Curr Hypertens Rep* 2018; **20**: 87.

Bessonneau V, Gerona RR, Trowbridge J, Grashow R, Lin T, Buren H *et al.* Gaussian graphical modeling of the serum exposome and metabolome reveals interactions between environmental chemicals and endogenous metabolites. *Sci Rep* 2021; **11**: 7607.

Jiang T, Wang M, Wang A, Abrahamsson D, Kuang W, Morello-Frosch R *et al.* Large-Scale Implementation and Flaw Investigation of Human Serum Suspect Screening Analysis for Industrial Chemicals. *J Am Soc Mass Spectrom* 2021; **32**: 2425–2435.

Eick SM, Hom Thepaksorn EK, Izano MA, Cushing LJ, Wang Y, Smith SC *et al.* Associations between prenatal maternal exposure to per- and polyfluoroalkyl substances (PFAS) and polybrominated diphenyl ethers (PBDEs) and birth outcomes among pregnant women in San Francisco. *Environ Health* 2020; **19**: 100.

CDC. Defining Adult Overweight and Obesity. Centers for Disease Control and Prevention. 2021.https://www.cdc.gov/obesity/adult/defining.html (accessed 12 Feb2022).

Pinheiro TV, Brunetto S, Ramos JGL, Bernardi JR, Goldani MZ. Hypertensive disorders during pregnancy and health outcomes in the offspring: a systematic review. *J Dev Orig Health Dis* 2016; **7**: 391–407.

Folk DM. Hypertensive Disorders of Pregnancy: Overview and Current Recommendations. *J Midwifery Womens Health* 2018; **63**: 289–300.

Martínez-Razo LD, Martínez-Ibarra A, Vázquez-Martínez ER, Cerbón M. The impact of Di-(2-ethylhexyl) Phthalate and Mono(2-ethylhexyl) Phthalate in placental development, function, and pathophysiology. *Environ Int* 2021; **146**: 106228.

Li R, Song M, Li Z, Li Y, Watanabe G, Nagaoka K *et al.* 4-Nitrophenol exposure alters the AhR signaling pathway and related gene expression in the rat liver. *Journal of Applied Toxicology* 2017; **37**: 150–158.

Final report on the safety assessment of Triethylene Glycol and PEG-4. *Int J Toxicol* 2006; **25 Suppl 2**: 121–138.

Tridecanedioic acid - Chemical Details. https://comptox.epa.gov/dashboard/chemical/details/DTXSID9021683 (accessed 24 Dec2021).

Cui X, Yan D, Xiao D. Synthesis and characterization of novel polyamides based on tridecanedioic acid: Nylons 3 13, 5 13, 6 13, 7 13, 9 13, 10 13, 11 13. *e-Polymers* 2004; **4**. doi:10.1515/epoly.2004.4.1.784.

PubChem. Octadecanedioic acid. https://pubchem.ncbi.nlm.nih.gov/compound/70095 (accessed 1 Mar2022).

Liu J, Li J, Li S, Leng J, Li W, Yang W *et al.* Circulating Lysophosphatidylcholines in Early Pregnancy and Risk of Gestational Diabetes in Chinese Women. *J Clin Endocrinol Metab* 2020; **105**: dgaa058.

Schulz K, Silva MR, Klaper R. Distribution and effects of branched versus linear isomers of PFOA, PFOS, and PFHxS: A review of recent literature. *Science of The Total Environment* 2020; **733**: 139186.

Koch HM, Bolt HM, Preuss R, Angerer J. New metabolites of di(2-ethylhexyl)phthalate (DEHP) in human urine and serum after single oral doses of deuterium-labelled DEHP. *Arch Toxicol* 2005; **79**: 367–376.

Jaacks LM, Diao N, Calafat AM, Ospina M, Mazumdar M, Ibne Hasan MOS *et al.* Association of prenatal pesticide exposures with adverse pregnancy outcomes and stunting in rural Bangladesh. *Environ Int* 2019; **133**: 105243.

Schymanski EL, Jeon J, Gulde R, Fenner K, Ruff M, Singer HP *et al.* Identifying small molecules via high resolution mass spectrometry: communicating confidence. *Environ Sci Technol* 2014; **48**: 2097–2098.

C8 Science Panel. C8 Science Panel Website. 2019.http://www.c8sciencepanel.org/ (accessed 30 Jul2019).

Liu X, Zhang L, Chen L, Li J, Wang Y, Wang J *et al.* Structure-based investigation on the association between perfluoroalkyl acids exposure and both gestational diabetes mellitus and glucose homeostasis in pregnant women. *Environment International* 2019; **127**: 85–93.

Gao Y, Chen L, Wang C, Zhou Y, Wang Y, Zhang Y *et al.* Exposure to polybrominated diphenyl ethers and female reproductive function: A study in the production area of Shandong, China. *Sci Total Environ* 2016; **572**: 9–15.

Li A, Zhuang T, Shi W, Liang Y, Liao C, Song M *et al.* Serum concentration of bisphenol analogues in pregnant women in China. *Sci Total Environ* 2020; **707**: 136100.

Blake BE, Fenton SE. Early life exposure to per- and polyfluoroalkyl substances (PFAS) and latent health outcomes: A review including the placenta as a target tissue and possible driver of peri- and postnatal effects. *Toxicology* 2020; **443**: 152565.

Eryasa B, Grandjean P, Nielsen F, Valvi D, Zmirou-Navier D, Sunderland E *et al.* Physico-chemical properties and gestational diabetes predict transplacental transfer and partitioning of perfluoroalkyl substances. *Environment International* 2019; **130**: 104874.

Beesoon S, Martin JW. Isomer-Specific Binding Affinity of Perfluorooctanesulfonate (PFOS) and Perfluorooctanoate (PFOA) to Serum Proteins. *Environ Sci Technol* 2015; **49**: 5722–5731.

Watkins DJ, Vélez-Vega CM, Rosario Z, Cordero JF, Alshawabkeh AN, Meeker JD. Preliminary assessment of exposure to persistent organic pollutants among pregnant women in Puerto Rico. *International Journal of Hygiene and Environmental Health* 2019; **222**: 327–331.

Varga T, Czimmerer Z, Nagy L. PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 2011; **1812**: 1007–1022.

Fowles J, Banton M, Klapacz J, Shen H. A toxicological review of the ethylene glycol series: Commonalities and differences in toxicity and modes of action. *Toxicol Lett* 2017; **278**: 66–83.

Jain RB. Effect of Pregnancy on The Levels of Selected Perfluoroalkyl Compounds for Females Aged 17–39 Years: Data From National Health and Nutrition Examination Survey 2003–2008. *Journal of Toxicology and Environmental Health, Part A* 2013; **76**: 409–421.

Fisher M, Arbuckle TE, Liang CL, LeBlanc A, Gaudreau E, Foster WG *et al.* Concentrations of persistent organic pollutants in maternal and cord blood from the maternal-infant research on environmental chemicals (MIREC) cohort study. *Environmental Health* 2016; **15**: 59.

Ohkuchi A, Hirashima C, Takahashi K, Suzuki H, Matsubara S. Prediction and prevention of hypertensive disorders of pregnancy. *Hypertens Res* 2017; **40**: 5–14.

Rodriguez BSQ, Mahdy H. *Gestational Diabetes*. StatPearls Publishing, 2021https://www.ncbi.nlm.nih.gov/books/NBK545196/ (accessed 2 Mar2022).

6. Tables

Table 1. Chemical uses and sources of the nine chemicals identified from non-targeted analyses (NTA) and selected for confirmation and quantification

Name	Sources and Uses
Branched perfluorooctane sulfonate (PFOS)	Industrial Chemicals used for stain, water, and fire resistance. In fabrics and food packaging.[25]
Linear PFOS	
Perfluorohexane sulfonate (PFHxS)	
Deoxycholic acid	A naturally occurring bile acid and endogenous biomarkers.
Tridecanedioic acid	Endogenous fatty acid, also used in plastic and resin manufacturing. [21,22]
Octadecanedioic acid	Endogenous fatty acid also used in plastic and resin manufacturing. [23] CDR database and estimates use for 3 companies at <1,000,000 lbs/year each.
Monoethylhexyl phthalate (MEHP)	Secondary diethylhexyl phthalate (DEHP) metabolite. DEHP is a common plasticizer. Studies suggest that secondary oxidized metabolites, not DEHP metabolites, are the developmental toxicants[18,26]
Tetraethylene glycol (TEG)	Industrial solvent used in cosmetics, fragrances, manufacturing, food contact, industrial fluids, lubricants.[20]. CDR from 11 companies each reporting 10,000 -50,000 pounds per year
4-nitrophenol	Xenobiotic metabolite of the pesticides parathion and methyl parathion, 4-nitroanisole. [19,27]
Chemical Data Reporting (CDR) a USEPA program where companies/manufacturers report quantities of chemicals being used and imported. Tetraethylene Glycol and octadecanedioic acid were the only ones found on the CDR database. Basic Information about Chemical Data Reporting	US EPA	

Table 2. Demographic variables and outcomes in Chemicals in Our Bodies 2 participants (n= 302) recruited between 2014 and 2018 in San Francisco

	Mean(SD) or N(%)
COVARIATES	
Maternal age at delivery (years)	33 (±5)
Race/ethnicity	
White	130 (43%)
Latinx	96 (32%)
Asian	47 (16%)
Other or Unknown [a]	29 (10%)
Education	
Some college or less	94 (31%)
College degree	73 (24%)
Graduate degree	125 (41%)
Missing	10 (3%)
Household income	
<40k	62 (21%)
40k-100k	36 (12%)
>100k	170 (56%)
Missing	34 (11%)
Born in U.S.	
No	110 (36%)
Yes	147 (49%)
Missing	45 (15%)
Pre-pregnancy BMI [b]	
<18.5	6 (2%)
18.5 to <25	147 (49%)
25 to <30	73 (24%)
30 or higher	41 (14%)
Missing	35 (12%)
Hospital of Delivery	
San Francisco General Hospital	88 (29%)
Mission Bay	213 (71%)
Missing	1 (<1%)
OUTCOMES	
Gestational diabetes mellitus	
No	237 (78%)
Yes	63 (21%)
Missing	2 (1%)
Preeclampsia	
No	268 (89%)
Yes	20 (7%)
Missing	14 (5%)
Gestational hypertension	
No	248 (82%)
Yes	39 (13%)
Missing	15 (5%)
Hypertensive disorders of pregnancy [c]	
None	238 (79%)
Any	50 (17%)
Missing	14 (5%)

[a] Race and ethnicity categories were collapsed for use in linear models due to small numbers in some categories. Other/unknown includes Black, Native Hawaiian or Pacific Islander, Native American, multiracial and individuals of unknown race; [b] BMI is calculated as kg/m^2 and cut-offs are based on the CDC guidelines;[15]
[c] Combines preeclampsia and gestational hypertension.

Table 3. Distribution of chemical levels in maternal serum (n= 302) and cord blood[a] (n= 299) (ng/ml)

Chemical	LOD	DF%	GM	GSD	min	max	Percentile 25	50	75	95
Branched PFOS[b] (Ratio GM cord/maternal: 0.7)										
cord	NR	NR	0.2	2.4	0	1.3	0.1	0.2	0.3	0.6
mat	NR	NR	0.3	2	0	3	0.2	0.3	0.5	1
Linear PFOS (Ratio: 0.5)										
cord	0.43	67	0.6	1.6	<LOD	3.8	<LOD	0.5	0.8	1.4
mat	0.43	97	1.1	1.8	<LOD	6.7	0.7	1	1.6	3.1
PFHxS (Ratio: 0.8)										
cord	0.11	98	0.3	1.6	<LOD	2.6	0.2	0.3	0.4	0.7
mat	0.11	100	0.4	1.8	<LOD	4.2	0.3	0.4	0.5	1.1
Tridecanedioic acid (Ratio: 1.5)										
cord	6.79	86	11.2	1.6	<LOD	47.7	8.1	11.2	14.9	23.7
mat	6.79	67	7.7	1.5	<LOD	26.2	6.3	8.1	9.9	14
Octadecanedioic acid (Ratio: 0.2)										
cord	0.48	100	3.1	1.8	0.5	35.7	2.2	3	4.3	9.2
mat	0.48	100	13.6	2	1.9	102.8	8.6	13.6	21.8	40.3
Deoxycholic acid (Ratio: 0.2)										
cord	0.81	99	3.7	2.2	<LOD	54.4	2.2	3.2	5.7	15.3
mat	0.81	99	16.1	3.4	<LOD	432.6	7.5	16.5	37.1	104.8
Monoethylhexyl phthalate (Ratio: 1.3)										
cord	5.19	58	6.7	2.7	<LOD	136.1	<LOD	6	10.7	50.4
mat	5.19	44	<LOD	<LOD	<LOD	452.6	<LOD	<LOD	16	60.2
Tetraethylene glycol (Ratio: 1.0)										
cord	2.1	43	3	3.1	<LOD	358	<LOD	<LOD	4.7	35.3
mat	2.1	43	2.9	2.9	<LOD	226.2	<LOD	<LOD	5	24.3
4-nitrophenol (Ratio: 0.8)										
cord	0.67	17	<LOD	1.5	<LOD	1.6	<LOD	<LOD	<LOD	0.9
mat	0.67	27	<LOD	1.6	<LOD	2.5	<LOD	<LOD	0.7	1

Abbreviations: GM geometric mean, GSD geometric standard deviation, LOD limit of detection, DF detection frequency, NR not reportable. [a] N is smaller for cord blood samples because we were not able to collect cord blood from a small number of participants. [b] The LOD and DF for branched PFOS were not reportable because they were estimated by subtracting the levels of linear PFOS from the total PFOS. Ratio illustrates transfer efficiency between maternal and cord samples and was calculated by GM(Cord)/GM(Mat)

Table 4. Adjusted and unadjusted odds ratios and 95% confidence intervals from linear models of the relationship between an interquartile range (IQR) chemical exposure at delivery and pregnancy complications (n = 302)

	OR (95% CI)	
Chemical	**gestational diabetes**	**hypertensive disorders** [b]
Number missing	2	14
N(%) with outcome	63 (21%)	50 (17%)
Branched PFOS		
Model0	1.65 (1.10 , 2.52)	1.05 (0.68 , 1.62)
Model1	1.56 (1.00 , 2.46)	1.06 (0.65 , 1.71)
Linear PFOS		
Model0	1.50 (1.05 , 2.17)	0.85 (0.56 , 1.27)
Model1	1.43 (0.96 , 2.14)	0.89 (0.56 , 1.39)
PFHxS		
Model0	1.15 (0.86 , 1.51)	1.00 (0.72 , 1.35)
Model1	1.09 (0.80 , 1.48)	0.98 (0.67 , 1.38)
Octadecanedioic acid		
Model0	1.20 (0.82 , 1.77)	1.05 (0.69 , 1.63)
Model1	1.26 (0.83 , 1.92)	1.00 (0.65 , 1.54)
Tridecanedioic acid		
Model0	1.08 (0.79 , 1.49)	1.25 (0.88 , 1.80)
Model1	1.03 (0.74 , 1.45)	1.28 (0.90 , 1.86)
Deoxycholic acid		
Model0	1.16 (0.81 , 1.69)	0.96 (0.65 , 1.43)
Model1	1.17 (0.80 , 1.72)	0.97 (0.65 , 1.45)

[a] CI = Confidence Interval; Model0 is the unadjusted model; Model1 is adjusted for maternal age, hospital of delivery, and race/ethnicity; [b] includes both preeclampsia and gestational hypertension

7. Figures

Study	HRMS	Database match	Feature selection from *a priori* criteria	Confirmed with analytical standard	Selected for quantification
Wang *et al* 2020	17,512 ESI+ 17219 ESI-	557 unique compounds	73 from MS/MS spectra	15	Level 1 PFHxS, PFOS, Octadecanedioic acid, Tridecanedioic acid, 4-nitrophenol, Monoethylhexyl phthalate, Tetraethylene glycol
Abrahamsson *et al* 2021	712 features ESI+ and ESI-	543 exogenous 142 endogenous compounds	73 from MS/MS spectra 98 via annotation algorithm	19	Level 2 Deoxycholic Acid

Figure 1. Summary of NTA procedures leading to the selection of nine chemical for quantification.

Figure 2: Spearman correlation coefficients for chemicals detected in maternal and cord samples, chemical have at least 65% DF in maternal and cord samples (n = 302).