

Extending non-targeted exposure discovery of environmental chemical exposures during pregnancy and their association with pregnancy complications—a cross-sectional study

Jessica Trowbridge¹, Dimitri Abrahamsson¹, Ting Jiang², Miaomiao Wang², June-Soo Park², Rachel Morello-Frosch³, Marina Sirota⁴, Dana E. Goin¹, Marya Zlatnick¹ and Tracey J. Woodruff¹

¹Department of Obstetrics, Gynecology and Reproductive Sciences, Program on Reproductive Health and the Environment, University of California San Francisco, San Francisco 94143 California

²Department of Toxic Substances Control, Environmental Chemistry Laboratory, California Environmental Protection Agency, Berkeley 94710, California

³Department of Environmental Science and Policy Management and School of Public Health, University of California, Berkeley, Berkeley 94720 California

⁴Bakar Computational Health Sciences Institute and Department of Pediatrics, University of California San Francisco, San Francisco 94158, California

*Corresponding author: Tracey J. Woodruff, tracey.woodruff@ucsf.edu

Abstract

Background: Non-targeted Analysis (NTA) methods identify novel exposures; however, few chemicals have been quantified and interrogated with pregnancy complications.

Objectives: We characterize levels of nine exogenous and endogenous chemicals in maternal and cord blood identified, selected, and confirmed in prior NTA steps including: linear and branched isomers perfluorooctane sulfonate (PFOS); perfluorohexane sulfonate (PFHxS); monoethylhexyl phthalate; 4-nitrophenol; tetraethylene glycol; tridecanedioic acid, octadecanedioic acid; and deoxycholic acid. We evaluate relationships between maternal and cord levels and the relationship gestational diabetes mellitus (GDM) and hypertensive disorders of pregnancy in a diverse pregnancy cohort in San Francisco.

Methods: We collected matched maternal and cord serum samples from 302 pregnant people at delivery from the Chemicals in Our Bodies cohort in San Francisco. Chemicals were identified via NTA and quantified using targeted approaches. We calculate distributions and Spearman correlation coefficients testing the relationship of chemicals within and between the maternal and cord blood matrices. We used logistic regression to calculate the odds of GDM and hypertensive disorders of pregnancy associated with an interquartile range increase in maternal chemical exposures.

Results: We detected linear PFOS, PFHxS, octadecanedioic acid, and deoxycholic acid in at least 97% of maternal samples. Correlations ranged between -0.1 and 0.9. We observed strong correlations between cord and maternal levels of PFHxS (coefficient = 0.9), linear PFOS (0.8), and branched PFOS (0.8). An IQR increase in linear PFOS, branched PFOS, and octadecanedioic acid is associated with increased odds of GDM [OR (95%CI): 1.43 (0.96, 2.14), 1.56 (1.00, 2.44), and 1.26 (0.83, 1.92) respectively] and tridecanedioic acid positively associated with hypertensive disorders of pregnancy [1.28 (0.90, 1.86)].

Discussion: We identified both exogenous and endogenous chemicals, two of which (octadecanedioic acid and tridecanedioic acid) have both endogenous and exogenous sources, and which have seldom been quantified in pregnant people or related to pregnancy complications.

Keywords: Non-targeted analysis, linear and branched PFOS, tridecanedioic acid, fatty acids, gestational diabetes mellitus, pregnancy hypertension

1 1. Introduction

2 Prenatal exposures to environmental chemicals are ubiquitous in the United States (U.S.)¹ and
3 studies have demonstrated they can have lifelong consequences for maternal and child health outcomes,²
4 such as neurodevelopment, cardiovascular disease, and reproduction. However, fewer than 1% of over
5 80,000 chemicals produced and used in the U.S. are regularly biomonitored³ and fewer still have been
6 assessed for adverse health outcomes during pregnancy. Non-targeted analysis (NTA) coupled with a
7 prioritization step and quantification using targeted methods can facilitate the identification and
8 quantitation of previously unmeasured chemical exposures.⁴ Additionally, by identifying and measuring
9 novel chemical exposures we can investigate their relationship with pregnancy complications,
10 facilitating the identification of opportunities for prevention.

11 Increasingly, studies are applying NTA to better characterize exposure to toxic environmental
12 chemicals seldom measured in pregnant people.^{5,6} NTA studies in pregnant and non-pregnant
13 populations have identified both known and novel chemical exposures, including PFAS congeners,
14 pesticides, flame retardants, and other industrial chemicals, which have been linked with pregnancy
15 complications.^{6,7} For example, PFAS has been linked with increased risk of miscarriage and
16 preeclampsia,⁸ and chemicals with fatty acid-like structures, including PFAS and other long-chain
17 hydrocarbons, have been shown to interact with endogenous fatty acid metabolism impacting placental
18 development and may contribute to pregnancy complications and adverse birth outcomes.^{6,9-11} Similarly,
19 NTA studies have been able to link novel chemical exposures and metabolism biomarkers.^{6,12} However,
20 the vast majority of novel chemicals identified through NTA have not been evaluated for associations
21 with pregnancy related complications.

22 This study builds on previous NTA work done on the Chemicals in Our Bodies-2 (CIOB2)
23 cohort, in which we first identified chemical features using high resolution mass spectrometry (HRMS)

24 and confirmed chemical identities via comparison to a standard.^{5,6,13} The aim of the present study is to
25 describe the quantification of a subset of chemicals identified and prioritized through this prior NTA and
26 to evaluate relationships between chemical levels and pregnancy complications. Supplementing the non-
27 targeted discovery of chemicals with targeted quantification described in this analysis addresses some of
28 the limitations of NTA; namely, targeted approaches increase the sensitivity to measure and compare
29 chemical concentrations and facilitate assessment of relationships between chemicals and pregnancy
30 complications.

31 2. Methods

32 2.1 Data and Sample Collection

33 We enrolled pregnant participants in the CIOB2 cohort between 2014 and 2018 from the prenatal
34 clinics at the Zuckerberg San Francisco General Hospital (SFGH) and Mission Bay or Moffit Long
35 (MB/ML) hospitals. Participant recruitment and sample collection is described elsewhere.^{4,5,14} Briefly,
36 participants were recruited during the second trimester of pregnancy, with eligibility criteria including
37 18-40 years of age, English or Spanish speaking, with singleton pregnancies, and no diagnosed
38 pregnancy complications at recruitment. Paired maternal and umbilical cord blood were collected from
39 participants who agreed to bank their samples and be contacted for participation in future studies.
40 Maternal samples were collected at delivery and cord blood was collected after delivery but before
41 clamping whenever possible. Blood was collected in BD Vacutainer Plus serum tubes, spun at 3000 rpm
42 to collect the serum, and stored at -80°C until analysis. Study protocols were approved by the
43 Institutional Review Boards at the University of California, San Francisco and Berkeley (# 13-12160).

44 Demographic information was collected via interviews conducted during the second trimester of
45 pregnancy and included maternal age, race and ethnicity (white, Latinx, and Non-Hispanic Asian/Pacific
46 Islander, other race—due to low numbers of Black, Native American, and multiracial participants, these

47 groups were collapsed into “other” for model specification), maternal education (some college or less,
48 completed a college degree, >college degree), household income (<40k per year, 40 to 100k, >100k),
49 nativity (born in the U.S. yes or no). Pre-pregnancy body mass index (BMI kg/m²; <18.5, 18.5 to <25,
50 25 to <35, >=35)¹⁵ and hospital of delivery [Zuckerberg San Francisco General Hospital (SFGH) or
51 Mission Bay/ Moffit Long (MB/ML)] were abstracted from the medical record.

52 Pregnancy complications were identified by clinical diagnoses and abstracted from the medical
53 record. These included the binary outcomes of: gestational diabetes mellitus (GDM), preeclampsia, and
54 gestational hypertension. Preeclampsia and pregnancy related hypertension were combined into one
55 outcome called hypertensive disorder of pregnancy because they make up part of a larger spectrum of
56 pregnancy complications related to hypertension.^{16,17}

57

58 2.2 Chemical selection

59 Nine environmental chemicals were measured in maternal and cord samples using targeted
60 approaches described below. Linear and branched isomers of perfluorooctane sulfonate (PFOS);
61 perfluorohexane sulfonate (PFHxS); monoethylhexyl phthalate (MEHP), a metabolite of diethylhexyl
62 phthalate, a common plasticizer¹⁸; 4-nitrophenol, a known endocrine disruptor used in pesticides, dyes,
63 and pharmaceuticals¹⁹; tetraethylene glycol, a plasticizer and solvent used in metal lubricants, printing
64 inks, and cosmetics;²⁰ tridecanedioic acid and octadecanedioic acid which are endogenous fatty acids
65 that also have exogenous sources primarily from plastics synthesis;^{21–23} and deoxycholic acid, a bile acid
66 and endogenous biomarker associated with GDM (**Table 1**).²⁴

67 The selection and confirmation procedures for selection of these chemicals is described
68 elsewhere.^{5,6,13} Briefly, in previous NTA studies, 30 matched maternal and cord samples were scanned
69 using HRMS in positive and negative ionization modes identifying suspect features. Detected features

70 were matched to an in-house database and features were then prioritized for MS/MS fragmentation
71 based on the ubiquity of exposure, feature intensity differences across demographic variables, and
72 correlation between maternal and cord samples. Fifteen chemicals were confirmed at level 1 confidence
73 via comparison to analytical standards.²⁸ Level 1 confirmed features included monoethylhexyl phthalate,
74 4-nitrophenol, tridecanedioic acid, and octadecanedioic acid. Subsequent generalized suspect screening
75 analysis of this cohort was expanded to include 295 matched maternal and cord samples and a more
76 comprehensive database including both endogenous and exogenous compounds. This resulted in
77 nineteen chemicals confirmed with analytical standards to level 1 confidence, including those previously
78 identified, and adding PFHxS, PFOS, and deoxycholic acid which were tentatively identified through
79 MS/MS fragmentation (confidence level 2). We selected chemicals for targeted quantification if the
80 chemical was identified in previous NTA with level 1 or level 2 confidence, is a widely used industrial
81 compound with high potential for exposure, is associated with pregnancy complications, or is not
82 regularly biomonitoring by the Center for Disease Control or other large biomonitoring studies.⁵

83

84 2.3 Targeted chemical analysis

85 Quantitation analysis of the nine selected chemicals was carried out using Agilent MassHunter
86 Quantitative Analysis software package (Agilent Technologies Inc., Version 10.0). Analytical grade
87 standards were purchased from suppliers (Sigma Aldrich and Thermo Fisher) and serial dilutions of the
88 standards were prepared in methanol. Calibration curve of each of the specific standard was acquired
89 under the same condition as serum sample analysis, using the LC/QTOF-MS (Agilent 1290 UPLC
90 interfaced with Agilent iFunnel 6550 QTOF-MS system) as described previously.^{5,6,13} M2-PFOA
91 (Wellington Laboratories, Ontario, Canada) was used as internal standard for negative ionization
92 chemicals, while D15-TPP (Cambridge Isotope Laboratories, Tewksbury, MA) for positive ionization

93 chemicals. Particularly, PFOS in serum samples were separated chromatically, thus both total and linear
94 PFOS quantitation were calculated, and branched PFOS concentrations were obtained by subtracting
95 linear portion from the total PFOS.

96

97 2.4 Statistical analysis

98 To address chemical levels below the limit of detection we used machine reported values when
99 available or, if not available, we imputed levels below the LOD with $LOD/\sqrt{2}$. We assessed the
100 detection frequency (percent above the limit of detection) and limited our statistical analysis to
101 chemicals with a detection frequency of 67% or higher. We described the distribution of chemicals in
102 maternal and cord blood samples, and calculated Spearman correlation coefficients for chemicals within
103 and between maternal and cord samples. We natural log-transformed the data for use in statistical
104 models and applied logistic regression models to assess the relationship between maternal chemical
105 exposure and pregnancy complications. Potential confounders were identified *a priori* (**Supplemental**
106 **Figure 1**), and our final models were adjusted for maternal age, hospital of delivery, and race/ethnicity.
107 We considered hospital of delivery as a cumulative indicator of socioeconomic status (including income,
108 educational attainment, and insurance status). Patients in both hospitals are racially and ethnically
109 diverse. However, SFGH is a safety net hospital that serves a large proportion of patients without
110 private health insurance, while MB/ML serves patients of higher socioeconomic status and with private
111 health insurance. Race and ethnicity are included in models as an indicator of the exposure to structural
112 racism, that is independently associated with chemical exposure and adverse pregnancy complications.
113 We calculated the interquartile range (IQR) of each natural log transformed chemical. Beta coefficients
114 and 95% confidence intervals (CI) were multiplied IQR and exponentiated to identify the odds ratio
115 (OR) of each pregnancy complication for each IQR increase in exposure level.

116 3. Results

117 Participants, demographics, and medical record data

118 The analysis included 302 participants (**Table 2**). The average age at delivery was 33 years of
119 age. Forty-three percent of participants identified as white and 32% as Latinx or Hispanic. Most
120 participants completed college or graduate degrees (>65%), were born in the U.S. (54%), and had a
121 household income of over \$100,000 per year (56%). For the outcomes, 21% of participants had a
122 diagnosis of GDM, 13% gestational hypertension, and 7% preeclampsia.

123

124 3.1 Distribution and correlation of chemical levels in maternal samples

125 Participants had at least four chemicals in maternal samples and three chemicals in cord samples.
126 PFHxS, linear PFOS, deoxycholic acid, and octadecanedioic acid were all detected in at least 97% of
127 maternal samples. Deoxycholic acid, tridecanedioic acid, PFHxS, were detected in at least 87% of cord
128 blood samples (Table 3). Since we estimated branched PFOS levels by subtracting linear PFOS from the
129 total PFOS we did not calculate a detection frequency for this chemical. Chemical detection frequency
130 varied between maternal and cord blood samples. Linear PFOS, and 4-nitrophenol were detected more
131 often maternal samples while monoethylhexyl phthalate, and tridecanedioic acid were detected more
132 frequently in cord blood. Similarly, the average chemical levels varied across the different matrices with
133 branched PFOS, linear PFOS, deoxycholic acid, and octadecanedioic acid found at higher levels in
134 maternal samples compared to cord samples, while tridecanedioic acid was higher in cord than in
135 maternal samples.

136 Monoethylhexyl phthalate, tetraethylene glycol, and 4-nitrophenol were each detected in fewer
137 than 50% of maternal samples, however 4-nitrophenol was detected in more maternal serum samples
138 than cord samples, while monoethylhexyl phthalate was found in more cord blood samples. Due to their

139 low detection frequency, monoethylhexyl phthalate, tetraethylene glycol, and 4-nitrophenol, were
140 excluded from further analyses.

141 When assessing the Spearman correlation coefficients (**Figure 2**) of the same chemical across
142 maternal and cord samples, we found that linear PFOS, and branched PFOS and PFHxS were highly
143 correlated in the maternal and cord samples (correlation coefficients of 0.8, 0.8, and 0.9 respectively).
144 When looking at the correlations of different chemicals in the maternal and cord samples we found
145 moderate correlations between cord-linear PFOS and maternal-branched PFOS, as well as between cord-
146 linear PFOS and maternal-PFHxS (coefficients 0.7 and 0.5 respectively). In the cord blood samples,
147 linear PFOS was highly correlated with branched PFOS and moderately with PFHxS (coefficients of 0.8
148 to 0.6 respectively), and octadecanedioic acid was moderately correlated with tridecanedioic acid
149 (coefficient of 0.4). In the maternal samples, the relationship between linear and branched PFOS had a
150 correlation coefficient of 0.8, branched PFOS and PFHxS a correlation of 0.6, and between linear PFOS
151 and PFHxS of 0.6. The PFAS chemicals were weakly correlated with the fatty acids—octadecanedioic
152 acid, tridecanedioic acid—and neither PFAS, nor the fatty acids were correlated with deoxycholic acid,
153 within or between the maternal serum and cord blood samples.

154
155 3.2 Association between chemical exposure with pregnancy complications

156 From our logistic models we observed a trend of increased odds of GDM for each IQR increase in levels
157 of exposure to environmental chemicals and increased odds of pregnancy hypertensive disorders with
158 tridecanedioic acid (Table 4). For the relationship between chemical exposures and GDM, all the
159 chemicals included in this analysis had an OR above 1. Stronger associations were observed for
160 branched PFOS, linear PFOS, and octadecanedioic acid; The unadjusted relationship between linear
161 PFOS and branched PFOS with GDM was statistically significant. The direction and measures of

162 association we observed between chemical levels and hypertensive disorders of pregnancy was less
163 consistent than for GDM. Exposure to tridecanedioic acid was associated with an increase in the odds of
164 hypertensive disorders of pregnancy [OR (95%CI): 1.28(0.90, 1.86)]. Exposure to tridecanedioic acid
165 was also associated in increased odds of the individual outcomes that made up hypertensive disorders of
166 pregnancy—preeclampsia and gestational hypertension (**Supplemental Table 1**). The ORs for PFHxS,
167 octadecanedioic acid, and deoxycholic acid with gestational hypertension approximated 1. Linear and
168 branched PFOS had an OR of <1 but the confidence intervals included the null value.

169 4. Discussion

170 Building on NTA steps from prior studies^{4,6,13}, we confirmed and quantified the presence of nine
171 chemicals, some of which have not been previously measured in pregnant people, such as tridecanedioic
172 acid, octadecanedioic acid, and tetraethylene glycol, or rarely measured in pregnant people like 4-
173 nitrophenol. In addition, a subset of these were moderately to strongly associated with GDM and
174 hypertensive disorders of pregnancy. Linear and Branched PFOS were associated with increased odds of
175 GDM and tridecanedioic acid was associated with hypertensive disorders of pregnancy. This study adds
176 to our understanding of the associations between environmental exposures during pregnancy and
177 pregnancy complications.

178 This study provides further evidence of prenatal exposure to environmental chemicals, some of
179 which have positive associations with pregnancy complications. A 2020 study suggests that PFAS
180 exposure during pregnancy may also affect metabolic pathways that adversely impact placental
181 development⁹ and while evidence remains mixed, PFAS has been shown to be associated with
182 hypertensive disorders of pregnancy, preeclampsia, and GDM.^{9,29-31} The linear and branched forms of
183 PFOS have not previously been well characterized during pregnancy. Furthermore, metabolism,

184 excretion, and placental transfer may vary by the linear and branched forms of this ubiquitous chemical
185 and lead to different health outcomes and pregnancy complications.²⁵

186 Significant positive correlations between branched PFOS, linear PFOS and PFHxS in the
187 maternal and cord samples suggest that they may share a common source of exposure. Indeed,
188 traditional PFOS production yielded a mixture of branched and linear isomers with approximately 30%
189 of isomers in the branched form.²⁵ From linear models we found that the linear and branched forms of
190 PFOS may have a relationship with pregnancy complications. Branched and Linear PFOS were
191 significantly associated with GDM in unadjusted models and although the relationship was attenuated in
192 adjusted models, the direction and magnitude of the association were maintained. Scientific evidence
193 indicates transplacental transfer of PFAS chemicals^{6,32,33} and gestational diabetes may facilitate
194 transplacental transfer of PFOS.³⁴ Linear PFOS makes up a larger proportion of the measured PFOS,²⁵ is
195 hypothesized to remain longer in maternal serum and is therefore more available in maternal samples
196 than branched PFOS, which may contribute to the differences in association we observed with
197 GDM.^{35,36}

198 Previous studies have found that increased exposure to fatty acids may be associated with an
199 increased risk of hypertensive disorders of pregnancy due to disruption of fatty acid regulation and
200 metabolism.³⁷ We saw an association between tridecanedioic acid and hypertensive disorders of
201 pregnancy, but not for octadecanedioic acid. Similarly, although the bile acid, deoxycholic acid, is
202 associated GDM in other studies,²⁴ we did not see elevated risk of these pregnancy complications
203 associated with deoxycholic acid in our study. The endogenous chemicals in our study have seldom been
204 studied in pregnant women and the fatty acids are also known to have exogenous sources and uses.

205 Tetraethylene glycol, 4-nitrophenol and monoethylhexyl phthalate were present in a small subset
206 of our participants but were not evaluated for an association with pregnancy complications because their

207 detection frequency in maternal samples was below 65%. In addition, these chemicals have short half-
208 lives and are quickly metabolized and excreted in urine contributing to their low detection frequency in
209 serum samples.^{19,26,38} These are high production volume chemicals that may interact with endocrine and
210 metabolic pathways; however, their relationship with adverse reproductive or pregnancy outcomes has
211 not been previously assessed, warranting further study. Despite their low detection frequency and that
212 they were measured in serum rather than in urine our study demonstrates that novel exposures to these
213 environmental chemicals can still be identified through non-targeted methods and confirmed through
214 targeted approaches.

215 There are several limitations that may influence the associations we observed in our study. First
216 is the timing of the collection of samples and assessment of health outcomes. This is a cross-sectional
217 study with samples collected at delivery and pregnancy complications were identified from clinical
218 diagnosis abstracted from medical record. Because of this we cannot assess temporality of our exposures
219 and outcome. Nevertheless, the chemicals included in this study are industrial chemicals, many of which
220 are high production volume chemicals that have a high potential for exposure. For example, PFAS
221 chemicals are ubiquitous in biological matrices and the environment, and tridecanedioic acid and
222 octadecanedioic acid are used in the synthesis of plastics and are considered high production volume
223 chemicals.²¹ While the chemical measurements reflect a snapshot of an individual's exposure history
224 during pregnancy the sources and uses of these environmental chemicals imply that there is a potential
225 for prolonged exposure throughout the pregnancy. Studies show that PFAS levels decrease in later
226 trimesters of pregnancy,^{39,40} thus the PFAS levels we measured at delivery may be lower than in other
227 trimesters of pregnancy biasing our results toward to the null. Another potential limitation is that levels
228 of branched PFOS were estimated and not quantified with a standard and thus the levels of branched
229 PFOS may reduce precision of calculated measures of association. Likewise, the low detection

230 frequency of 4-nitrophenol, tetraethylene glycol, or monoethylhexyl phthalate precluded our ability to
231 assess the chemical-outcome relationship. However, these chemicals remain understudied in pregnant
232 people and are of high concern due to their widespread use as industrial chemicals and limited
233 understanding of their health effects. Finally, because our recruitment criteria indicated that participants
234 with pregnancy complications by the second trimester we may have missed earlier manifestations of
235 pregnancy complications. This could undercount the number of GDM and hypertensive disorders of
236 pregnancy biasing our results toward the null. However, diagnosis of hypertensive disorders of
237 pregnancy and testing for GDM usually occurs in the third trimester of pregnancy while our participants
238 were recruited in the second.^{41,42}

239 In this analysis we identified and quantified exogenous and endogenous environmental
240 chemicals, some of which have not been previously measured during pregnancy nor evaluated for
241 associations with pregnancy complications. Our study supports prior reports of associations between the
242 branched and linear forms of PFOS and GDM and adds to the limited research on exposure to
243 tridecanedioic acid and octadecanedioic acid, endogenous fatty acids that also have industrial sources.
244 Finally, this study further illustrates the benefit of pairing non-targeted and targeted approaches to
245 identify, confirm, and quantify novel chemical exposures. Combining NTA methods with targeted
246 chemicals measurement can be further applied to identifying, quantifying, and evaluating metabolism
247 biomarkers that are on the pathway between exposure and downstream adverse health effects.

5. References

- 1 Woodruff TJ, Zota AR, Schwartz JM. Environmental chemicals in pregnant women in the United States: NHANES 2003-2004. *Environ Health Perspect* 2011; **119**: 878–885.
- 2 American College of Obstetricians and Gynecologists' Committee on Obstetric Practice. Reducing Prenatal Exposure to Toxic Environmental Agents: ACOG Committee Opinion, Number 832. *Obstet Gynecol* 2021; **138**: e40–e54.
- 3 CDC. National Report on Human Exposure to Environmental Chemicals | CDC. Center for Disease Control and Prevention. 2019. <https://www.cdc.gov/exposurereport/index.html> (accessed 18 Nov2018).
- 4 Wang A, Gerona RR, Schwartz JM, Lin T, Sirota M, Morello-Frosch R *et al.* A Suspect Screening Method for Characterizing Multiple Chemical Exposures among a Demographically Diverse Population of Pregnant Women in San Francisco. *Environ Health Perspect* 2018; **126**: 077009.
- 5 Wang A, Abrahamsson DP, Jiang T, Wang M, Morello-Frosch R, Park J-S *et al.* Suspect Screening, Prioritization, and Confirmation of Environmental Chemicals in Maternal-Newborn Pairs from San Francisco. *Environ Sci Technol* 2021; **55**: 5037–5049.
- 6 Panagopoulos Abrahamsson D, Wang A, Jiang T, Wang M, Siddharth A, Morello-Frosch R *et al.* A Comprehensive Non-targeted Analysis Study of the Prenatal Exposome. *Environ Sci Technol* 2021; **55**: 10542–10557.
- 7 Grashow R, Bessonneau V, Gerona RR, Wang A, Trowbridge J, Lin T *et al.* Integrating Exposure Knowledge and Serum Suspect Screening as a New Approach to Biomonitoring: An Application in Firefighters and Office Workers. *Environ Sci Technol* 2020; **54**: 4344–4355.
- 8 Gao X, Ni W, Zhu S, Wu Y, Cui Y, Ma J *et al.* Per- and polyfluoroalkyl substances exposure during pregnancy and adverse pregnancy and birth outcomes: A systematic review and meta-analysis. *Environ Res* 2021; **201**: 111632.
- 9 Szilagyi JT, Avula V, Fry RC. Perfluoroalkyl Substances (PFAS) and Their Effects on the Placenta, Pregnancy, and Child Development: a Potential Mechanistic Role for Placental Peroxisome Proliferator-Activated Receptors (PPARs). *Curr Environ Health Rep* 2020; **7**: 222–230.
- 10 Ledda C, Fiore M, Santarelli L, Bracci M, Mascali G, D'Agati MG *et al.* Gestational Hypertension and Organophosphorus Pesticide Exposure: A Cross-Sectional Study. *Biomed Res Int* 2015; **2015**: 280891.
- 11 Kahn LG, Trasande L. Environmental Toxicant Exposure and Hypertensive Disorders of Pregnancy: Recent Findings. *Curr Hypertens Rep* 2018; **20**: 87.
- 12 Bessonneau V, Gerona RR, Trowbridge J, Grashow R, Lin T, Buren H *et al.* Gaussian graphical modeling of the serum exposome and metabolome reveals interactions between environmental chemicals and endogenous metabolites. *Sci Rep* 2021; **11**: 7607.

- 13 Jiang T, Wang M, Wang A, Abrahamsson D, Kuang W, Morello-Frosch R *et al.* Large-Scale Implementation and Flaw Investigation of Human Serum Suspect Screening Analysis for Industrial Chemicals. *J Am Soc Mass Spectrom* 2021; **32**: 2425–2435.
- 14 Eick SM, Hom Thepaksorn EK, Izano MA, Cushing LJ, Wang Y, Smith SC *et al.* Associations between prenatal maternal exposure to per- and polyfluoroalkyl substances (PFAS) and polybrominated diphenyl ethers (PBDEs) and birth outcomes among pregnant women in San Francisco. *Environ Health* 2020; **19**: 100.
- 15 CDC. Defining Adult Overweight and Obesity. Centers for Disease Control and Prevention. 2021. <https://www.cdc.gov/obesity/adult/defining.html> (accessed 12 Feb2022).
- 16 Pinheiro TV, Brunetto S, Ramos JGL, Bernardi JR, Goldani MZ. Hypertensive disorders during pregnancy and health outcomes in the offspring: a systematic review. *J Dev Orig Health Dis* 2016; **7**: 391–407.
- 17 Folk DM. Hypertensive Disorders of Pregnancy: Overview and Current Recommendations. *J Midwifery Womens Health* 2018; **63**: 289–300.
- 18 Martínez-Razo LD, Martínez-Ibarra A, Vázquez-Martínez ER, Cerbón M. The impact of Di-(2-ethylhexyl) Phthalate and Mono(2-ethylhexyl) Phthalate in placental development, function, and pathophysiology. *Environ Int* 2021; **146**: 106228.
- 19 Li R, Song M, Li Z, Li Y, Watanabe G, Nagaoka K *et al.* 4-Nitrophenol exposure alters the AhR signaling pathway and related gene expression in the rat liver. *Journal of Applied Toxicology* 2017; **37**: 150–158.
- 20 Final report on the safety assessment of Triethylene Glycol and PEG-4. *Int J Toxicol* 2006; **25 Suppl 2**: 121–138.
- 21 Tridecanedioic acid - Chemical Details. <https://comptox.epa.gov/dashboard/chemical/details/DTXSID9021683> (accessed 24 Dec2021).
- 22 Cui X, Yan D, Xiao D. Synthesis and characterization of novel polyamides based on tridecanedioic acid: Nylons 3 13, 5 13, 6 13, 7 13, 9 13, 10 13, 11 13. *e-Polymers* 2004; **4**. doi:10.1515/epoly.2004.4.1.784.
- 23 PubChem. Octadecanedioic acid. <https://pubchem.ncbi.nlm.nih.gov/compound/70095> (accessed 1 Mar2022).
- 24 Liu J, Li J, Li S, Leng J, Li W, Yang W *et al.* Circulating Lysophosphatidylcholines in Early Pregnancy and Risk of Gestational Diabetes in Chinese Women. *J Clin Endocrinol Metab* 2020; **105**: dga058.
- 25 Schulz K, Silva MR, Klaper R. Distribution and effects of branched versus linear isomers of PFOA, PFOS, and PFHxS: A review of recent literature. *Science of The Total Environment* 2020; **733**: 139186.

- 26 Koch HM, Bolt HM, Preuss R, Angerer J. New metabolites of di(2-ethylhexyl)phthalate (DEHP) in human urine and serum after single oral doses of deuterium-labelled DEHP. *Arch Toxicol* 2005; **79**: 367–376.
- 27 Jaacks LM, Diao N, Calafat AM, Ospina M, Mazumdar M, Ibne Hasan MOS *et al.* Association of prenatal pesticide exposures with adverse pregnancy outcomes and stunting in rural Bangladesh. *Environ Int* 2019; **133**: 105243.
- 28 Schymanski EL, Jeon J, Gulde R, Fenner K, Ruff M, Singer HP *et al.* Identifying small molecules via high resolution mass spectrometry: communicating confidence. *Environ Sci Technol* 2014; **48**: 2097–2098.
- 29 C8 Science Panel. C8 Science Panel Website. 2019.<http://www.c8sciencepanel.org/> (accessed 30 Jul2019).
- 30 Liu X, Zhang L, Chen L, Li J, Wang Y, Wang J *et al.* Structure-based investigation on the association between perfluoroalkyl acids exposure and both gestational diabetes mellitus and glucose homeostasis in pregnant women. *Environment International* 2019; **127**: 85–93.
- 31 Gao Y, Chen L, Wang C, Zhou Y, Wang Y, Zhang Y *et al.* Exposure to polybrominated diphenyl ethers and female reproductive function: A study in the production area of Shandong, China. *Sci Total Environ* 2016; **572**: 9–15.
- 32 Li A, Zhuang T, Shi W, Liang Y, Liao C, Song M *et al.* Serum concentration of bisphenol analogues in pregnant women in China. *Sci Total Environ* 2020; **707**: 136100.
- 33 Blake BE, Fenton SE. Early life exposure to per- and polyfluoroalkyl substances (PFAS) and latent health outcomes: A review including the placenta as a target tissue and possible driver of peri- and postnatal effects. *Toxicology* 2020; **443**: 152565.
- 34 Eryasa B, Grandjean P, Nielsen F, Valvi D, Zmirou-Navier D, Sunderland E *et al.* Physico-chemical properties and gestational diabetes predict transplacental transfer and partitioning of perfluoroalkyl substances. *Environment International* 2019; **130**: 104874.
- 35 Beeson S, Martin JW. Isomer-Specific Binding Affinity of Perfluorooctanesulfonate (PFOS) and Perfluorooctanoate (PFOA) to Serum Proteins. *Environ Sci Technol* 2015; **49**: 5722–5731.
- 36 Watkins DJ, Vélez-Vega CM, Rosario Z, Cordero JF, Alshawabkeh AN, Meeker JD. Preliminary assessment of exposure to persistent organic pollutants among pregnant women in Puerto Rico. *International Journal of Hygiene and Environmental Health* 2019; **222**: 327–331.
- 37 Varga T, Czimmerer Z, Nagy L. PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 2011; **1812**: 1007–1022.
- 38 Fowles J, Banton M, Klapacz J, Shen H. A toxicological review of the ethylene glycol series: Commonalities and differences in toxicity and modes of action. *Toxicol Lett* 2017; **278**: 66–83.

- 39 Jain RB. Effect of Pregnancy on The Levels of Selected Perfluoroalkyl Compounds for Females Aged 17–39 Years: Data From National Health and Nutrition Examination Survey 2003–2008. *Journal of Toxicology and Environmental Health, Part A* 2013; **76**: 409–421.
- 40 Fisher M, Arbuckle TE, Liang CL, LeBlanc A, Gaudreau E, Foster WG *et al.* Concentrations of persistent organic pollutants in maternal and cord blood from the maternal-infant research on environmental chemicals (MIREC) cohort study. *Environmental Health* 2016; **15**: 59.
- 41 Ohkuchi A, Hirashima C, Takahashi K, Suzuki H, Matsubara S. Prediction and prevention of hypertensive disorders of pregnancy. *Hypertens Res* 2017; **40**: 5–14.
- 42 Rodriguez BSQ, Mahdy H. *Gestational Diabetes*. StatPearls Publishing, 2021 <https://www.ncbi.nlm.nih.gov/books/NBK545196/> (accessed 2 Mar2022).

6. Tables

Table 1. Chemical uses and sources of the nine chemicals identified from non-targeted analyses (NTA) and selected for confirmation and quantification

Name	Sources and Uses
Branched perfluorooctane sulfonate (PFOS)	Industrial Chemicals used for stain, water, and fire resistance. In fabrics and food packaging. ²⁵
Linear PFOS	
Perfluorohexane sulfonate (PFHxS)	
Deoxycholic acid	A naturally occurring bile acid and endogenous biomarkers.
Tridecanedioic acid	Endogenous fatty acid, also used in plastic and resin manufacturing. ^{21,22}
Octadecanedioic acid	Endogenous fatty acid also used in plastic and resin manufacturing. ²³ CDR database and estimates use for 3 companies at <1,000,000 lbs/year each.
Monoethylhexyl phthalate (MEHP)	Secondary diethylhexyl phthalate (DEHP) metabolite. DEHP is a common plasticizer. Studies suggest that secondary oxidized metabolites, not DEHP metabolites, are the developmental toxicants ^{18,26}
Tetraethylene glycol (TEG)	Industrial solvent used in cosmetics, fragrances, manufacturing, food contact, industrial fluids, lubricants. ²⁰ . CDR from 11 companies each reporting 10,000 -50,000 pounds per year
4-nitrophenol	Xenobiotic metabolite of the pesticides parathion and methyl parathion, 4-nitroanisole. ^{19,27}

Chemical Data Reporting (CDR) a USEPA program where companies/manufacturers report quantities of chemicals being used and imported. Tetraethylene Glycol and octadecanedioic acid were the only ones found on the CDR database. [Basic Information about Chemical Data Reporting | US EPA](#)

Table 2. Demographic variables and outcomes in Chemicals in Our Bodies 2 participants (n= 302) recruited between 2014 and 2018 in San Francisco

	Mean(SD) or N(%)
COVARIATES	
Maternal age at delivery (years)	33 (±5)
Race/ethnicity	
White	130 (43%)
Latinx	96 (32%)
Asian	47 (16%)
Other or Unknown ^a	29 (10%)
Education	
Some college or less	94 (31%)
College degree	73 (24%)
Graduate degree	125 (41%)
Missing	10 (3%)
Household income	
<40k	62 (21%)
40k-100k	36 (12%)
>100k	170 (56%)
Missing	34 (11%)
Born in U.S.	
No	110 (36%)
Yes	147 (49%)
Missing	45 (15%)
Pre-pregnancy BMI^b	
<18.5	6 (2%)
18.5 to <25	147 (49%)
25 to <30	73 (24%)
30 or higher	41 (14%)
Missing	35 (12%)
Hospital of Delivery	
San Francisco General Hospital	88 (29%)
Mission Bay	213 (71%)
Missing	1 (<1%)
OUTCOMES	
Gestational diabetes mellitus	
No	237 (78%)
Yes	63 (21%)
Missing	2 (1%)
Preeclampsia	
No	268 (89%)
Yes	20 (7%)
Missing	14 (5%)
Gestational hypertension	
No	248 (82%)
Yes	39 (13%)
Missing	15 (5%)
Hypertensive disorders of pregnancy^c	
None	238 (79%)
Any	50 (17%)
Missing	14 (5%)

^aRace and ethnicity categories were collapsed for use in linear models due to small numbers in some categories. Other/unknown includes Black, Native Hawaiian or Pacific Islander, Native American, multiracial and individuals of unknown race; ^b BMI is calculated as kg/m² and cut-offs are based on the CDC guidelines;¹⁵

^c Combines preeclampsia and gestational hypertension.

Table 3. Distribution of chemical levels in maternal serum (n= 302) and cord blood^a (n= 299) (ng/ml)

Chemical	LOD	DF%	GM	GSD	min	max	Percentile			
							25	50	75	95
Branched PFOS^b (Ratio GM cord/maternal: 0.7)										
cord	NR	NR	0.2	2.4	0	1.3	0.1	0.2	0.3	0.6
mat	NR	NR	0.3	2	0	3	0.2	0.3	0.5	1
Linear PFOS (Ratio: 0.5)										
cord	0.43	67	0.6	1.6	<LOD	3.8	<LOD	0.5	0.8	1.4
mat	0.43	97	1.1	1.8	<LOD	6.7	0.7	1	1.6	3.1
PFHxS (Ratio: 0.8)										
cord	0.11	98	0.3	1.6	<LOD	2.6	0.2	0.3	0.4	0.7
mat	0.11	100	0.4	1.8	<LOD	4.2	0.3	0.4	0.5	1.1
Tridecanedioic acid (Ratio: 1.5)										
cord	6.79	86	11.2	1.6	<LOD	47.7	8.1	11.2	14.9	23.7
mat	6.79	67	7.7	1.5	<LOD	26.2	6.3	8.1	9.9	14
Octadecanedioic acid (Ratio: 0.2)										
cord	0.48	100	3.1	1.8	0.5	35.7	2.2	3	4.3	9.2
mat	0.48	100	13.6	2	1.9	102.8	8.6	13.6	21.8	40.3
Deoxycholic acid (Ratio: 0.2)										
cord	0.81	99	3.7	2.2	<LOD	54.4	2.2	3.2	5.7	15.3
mat	0.81	99	16.1	3.4	<LOD	432.6	7.5	16.5	37.1	104.8
Monoethylhexyl phthalate (Ratio: 1.3)										
cord	5.19	58	6.7	2.7	<LOD	136.1	<LOD	6	10.7	50.4
mat	5.19	44	<LOD	<LOD	<LOD	452.6	<LOD	<LOD	16	60.2
Tetraethylene glycol (Ratio: 1.0)										
cord	2.1	43	3	3.1	<LOD	358	<LOD	<LOD	4.7	35.3
mat	2.1	43	2.9	2.9	<LOD	226.2	<LOD	<LOD	5	24.3
4-nitrophenol (Ratio: 0.8)										
cord	0.67	17	<LOD	1.5	<LOD	1.6	<LOD	<LOD	<LOD	0.9
mat	0.67	27	<LOD	1.6	<LOD	2.5	<LOD	<LOD	0.7	1


Abbreviations: GM geometric mean, GSD geometric standard deviation, LOD limit of detection, DF detection frequency, NR not reportable. ^aN is smaller for cord blood samples because we were not able to collect cord blood from a small number of participants. ^bThe LOD and DF for branched PFOS were not reportable because they were estimated by subtracting the levels of linear PFOS from the total PFOS. Ratio illustrates transfer efficiency between maternal and cord samples and was calculated by GM(Cord)/GM(Mat)

Table 4. Adjusted and unadjusted odds ratios and 95% confidence intervals from linear models of the relationship between an interquartile range (IQR) chemical exposure at delivery and pregnancy complications (n = 302)

Chemical	OR (95% CI)	
	gestational diabetes	hypertensive disorders ^b
Number missing	2	14
N(%) with outcome	63 (21%)	50 (17%)
Branched PFOS		
Model0	1.65 (1.10 , 2.52)	1.05 (0.68 , 1.62)
Model1	1.56 (1.00 , 2.46)	1.06 (0.65 , 1.71)
Linear PFOS		
Model0	1.50 (1.05 , 2.17)	0.85 (0.56 , 1.27)
Model1	1.43 (0.96 , 2.14)	0.89 (0.56 , 1.39)
PFHxS		
Model0	1.15 (0.86 , 1.51)	1.00 (0.72 , 1.35)
Model1	1.09 (0.80 , 1.48)	0.98 (0.67 , 1.38)
Octadecanedioic acid		
Model0	1.20 (0.82 , 1.77)	1.05 (0.69 , 1.63)
Model1	1.26 (0.83 , 1.92)	1.00 (0.65 , 1.54)
Tridecanedioic acid		
Model0	1.08 (0.79 , 1.49)	1.25 (0.88 , 1.80)
Model1	1.03 (0.74 , 1.45)	1.28 (0.90 , 1.86)
Deoxycholic acid		
Model0	1.16 (0.81 , 1.69)	0.96 (0.65 , 1.43)
Model1	1.17 (0.80 , 1.72)	0.97 (0.65 , 1.45)

^a CI = Confidence Interval; Model0 is the unadjusted model; Model1 is adjusted for maternal age, hospital of delivery, and race/ethnicity; ^b includes both preeclampsia and gestational hypertension

7. Figures



Study	HRMS	Database match	Feature selection from <i>a priori</i> criteria	Confirmed with analytical standard	Selected for quantification
Wang <i>et al</i> 2020	17,512 ESI+ 17219 ESI-	557 unique compounds	73 from MS/MS spectra	15	<u>Level 1</u> PFHxS, PFOS, Octadecanedioic acid, Tridecanedioic acid, 4-nitrophenol, Monoethylhexyl phthalate, Tetraethylene glycol
Abrahamsson <i>et al</i> 2021	712 features ESI+ and ESI-	543 exogenous 142 endogenous compounds	73 from MS/MS spectra 98 via annotation algorithm	19	<u>Level 2</u> Deoxychoic Acid

Figure 1. Summary of NTA procedures leading to the selection of nine chemical for quantification.




Figure 2: Spearman correlation coefficients for chemicals detected in maternal and cord samples, chemical have at least 65% DF in maternal and cord samples (n = 302).