Analysis of immunization time, amplitude, and adverse events of seven different vaccines against SARS-CoV-2 across four different countries.

Running Title: Immunization of 7 vaccines vs SARS.CoV-2

Maria Elena Romero-Ibarguengoitia*[1,2], Arnulfo González- Cantú [1,2], Chiara Pozzi[1], Riccardo Levi[3,4], Maximiliano Mollura[5], Riccardo Sarti[3,4], Miguel Angel Sanz-Sánchez[1,2], Diego Rivera-Salinas [1,2], Yodira Guadalupe Hernández-Ruíz[1,2], Ana Gabriela Armendariz-Vázquez[1,2], Gerardo Francisco Del Rio-Parra[1,2], Irene Antonieta Barco-Flores[1], Rosalinda González-Facio[1], Elena Azzolini[3,4], Riccardo Barbieri[5], Alessandro Rodrigo de Azevedo Dias[6], Milton Henriques-Guimaraes Júnior[7], Alessandra Bastos-Borges[7], Cecilia Acciardi[8], Graciela Paez-Bo[9]; Mauro Martins Teixeira[10] , Maria Rescigno*[3, 4].

[1] Research Department, Hospital Clínica Nova de Monterrey, San Nicolás de los Garza, Nuevo León, México.

[2] Vicerrectoría de Ciencias de la Salud, Escuela de Medicina, Universidad de Monterrey, San Pedro Garza García, Nuevo León, México.

[3] IRCCS Humanitas Clinical and Research Center, Rozzano, MI, Italy.

[4] Humanitas University, Department of Biomedical Sciences, Pieve Emanuele, MI, Italy.

[5] Politecnico di Milano, Department of Electronic, Information and Bioengineering, MI, Italy.

[6] Health Department, Ternium Brazil, Rio de Janeiro, Brazil

[7] Research Department Fundação São Francisco Xavier, Minas Gerais, Brazil.

[8] Health Secretary, Unidad Hospitalaria San José, Campana, Buenos Aires, Argentina

[9] Laboratory Department, Hospital Interzonal General de Agudos San Felipe, San Nicolás de los Arroyos, Buenos Aires, Argentina

[10] Biochemistry and Immunology Department, ICB, Universidad Federal de Minas Gerais, Minas Gerais, Brazil

29   Correspondance:

30   Maria Elena Romero

31   MROMEROI@novaservicios.com.mx

33   Maria Rescigno

34   maria.rescigno@hunimed.eu

Abstract

Background: Scarce information exists in relation to the comparison of seroconversion and adverse events following immunization (AEFI) with different SARS-CoV-2 vaccines. Our aim was to correlate the magnitude of the antibody response to vaccination with previous clinical conditions and AEFI.

Methods: A multicentric comparative study where SARS-CoV-2 spike 1-2 IgG antibodies IgG titers were measured at baseline, 21-28 days after the first and second dose (when applicable) of the following vaccines: BNT162b2 mRNA, mRNA-1273, Gam-COVID-Vac, Coronavac, ChAdOx1-S, Ad5-nCoV and Ad26.COV2. Mixed model and Poisson generalized linear models were performed.

Results: We recruited 1867 subjects [52 (SD 16.8) years old, 52% men]. All vaccines enhanced anti-S1 and anti-S2 IgG antibodies over time (p<0.01). The highest increase after the first and second dose was observed in mRNA-1273 (p<0.001). There was an effect of previous SARS-CoV-2 infection; and an interaction of age with SARS-CoV-2, Gam-COVID-Vac and ChAdOx1-S (p<0.01). There was a negative correlation of Severe or Systemic AEFI (AEs) of naïve SARS-CoV-2 subjects with age and sex (p<0.001); a positive interaction between the delta of antibodies with Gam-COVID-Vac (p=0.002). Coronavac, Gam-COVID-Vac and ChAdOx1-S had less AEs compared to BNT162b (p<0.01). mRNA-1273 had a higher number of AEFIs. The delta of the antibodies showed an association with AEFIs in previously infected individuals (p<0.001).

Conclusions: The magnitude of seroconversion is predicted by age, vaccine type and SARS-CoV-2 exposure. AEs are correlated with age, sex, and vaccine type. The delta of the antibody response is positively correlated with AEs in patients previously exposed to SARS-CoV-2.

Registration number: NCT05228912

Keywords: COVID-19, SARS-CoV-2, immunization, vaccines, seroconversion

Introduction

Covid-19 is a pneumonia-like disease caused by a coronavirus, named SARS-CoV-2. It is highly

contagious, and the World Health Organization declared it in 2020 as a pandemic (1). SARS-CoV-2

causes COVID-19 that has a wide range of clinical presentations, from asymptomatic disease to severe

acute respiratory distress syndrome and death(2). SARS-CoV-2 utilizes its surface spike glycoprotein

to enter host cells. Each unit of the spike trimer contains an S1 and S2 subunit, with the N-terminal S1

subunit binding to the cellular angiotensin-converting enzyme 2 (ACE2) receptor through an internal

receptor-binding domain (RBD)(3).

To prevent SARS-CoV-2 infection, 113 vaccines are being tested in human clinical trials, and 44 have

reached the final step of testing(4). Based on the mechanism of action, vaccines can be clustered in

four groups: 1) mRNA vaccines, for example, BNT162b2 mRNA and mRNA-1273. They use

genetically engineered modified RNA to produce the spike protein that prompts an immune response

safely. 2) Viral vector (adenovirus) vaccines, for example, ChAdOx1-S, Ad26.COV2, Ad5-nCoV, and

Gam-COVID-Vac. They use a virus that has been genetically engineered so that it cannot cause

disease but produces coronavirus proteins to safely generate an immune response. 3) Inactivated virus

vaccines, for example, Coronavac, use a form of the virus that has been inactivated or weakened, so it

does not cause disease but still generates an immune response. 4) Protein subunits vaccines, for

example, NVX-CoV2373. It is a protein-based vaccine that uses harmless fragments of proteins or

protein shells that mimic the SARS-CoV-2 to safely generate an immune response(5).

Currently BNT162b2 mRNA, mRNA-1273, ChAdOx1-S , Ad26.COV2 , Ad5-nCoV, Gam-COVID-

Vac, Coronavac, have been administered to people across 587 countries(4). However, no studies have

compared using the same assay and time frame the effectiveness of seroconversion and incidence of

adverse event events in response to vaccines in different countries in the same study.

This study aimed to correlate the magnitude of the antibody response after the first and second dose (if

applicable) between BNT162b2 mRNA, mRNA-1273, ChAdOx1-S, Ad26.COV2, Ad5-nCoV, Gam-

116   COVID-Vac, and Coronavac by measuring SARS-CoV-2 spike 1-2 IgG antibodies, using the same

117   standardized assay, with previous clinical conditions and systemic and severe adverse events (AEs).

Materials and Methods.

This was a multicentric observational study of volunteers who received an approved complete scheme of BNT162b2 mRNA, mRNA-1273, ChAdOx1-S, Ad26.COV2, Ad5-nCoV, Gam-COVID-Vac, or Coronavac COVID-19 vaccine during 2021 in five hospital centers (Hospital Clinica Nova, Humanitas Clinical and Research Center, Fundacion San Francisco Xavier, Ternium Health Center in Rio, Hospital Municipal San Jose, Hospital Interzonal de Agudos San Felipe) from four different countries: Mexico, Italy, Brazil and Argentina. This study followed STROBE guidelines (6). The study was approved by each local Institutional Review Board and conducted per the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments that involve humans.

The inclusion criteria were volunteers of both genders, any age, who consented to participate, planned to conclude the immunization regimen of any vaccine, and agreed to be followed up for the duration of the study. The following vaccines were available: BNT162b2 mRNA, mRNA-1273, ChAdOx1-S, Ad26.COV2, Ad5-nCoV, Gam-COVID-Vac, or Coronavac. The exclusion criteria were to have received any SARS-CoV-2 vaccine prior to study entry.

Each country's Health System defined the vaccines available, the schedule, and dose assignation. On the vaccination day, the research team invited any subject who planned to receive any vaccine scheme, explained the project and asked to sign the informed consent. Inclusion-exclusion criteria were applied, and a plasma sample was collected. The baseline sample was taken before receiving the first dose of any SARS-CoV-2 vaccine (T0). The second (T1) and third samples (T2) were taken after 21 days (+/- 7 days) of the first and second dose, respectively.

At each visit, participants had to answer a questionnaire. The basal questionnaire aimed at obtaining patients' medical history and previous SARS-CoV-2 infections. The questionnaires applied after the first and second dose of vaccines aimed at recognizing adverse events following immunization (AEFI)(7) and identifying a potential SARS-CoV-2 infection after receiving any vaccine dose. SARS-

CoV-2 infection was also monitored by the epidemiology team through PCR testing, which informed

the research team of any new infection.

Primary Outcome, IgG determination

Our primary outcome was to correlate the magnitude of the antibody response to vaccination with

previous clinical conditions and AEs . To determine the amount of specific anti-S1 and anti-S2 IgG

antibodies against SARS-CoV-2 in plasma samples, the laboratory personnel used a

chemiluminescence immunoassay (CLIA) developed by DiaSorin, which had a sensitivity of 97.4%

(95% CI, 86.8-99.5) and a specificity of 98.5% (95% CI, 97.5-99.2). The results were reported as

follows: <12.0 AU/ml was considered negative, 12.0 to 15.0 AU/ml was indeterminate, and > 15

AU/ml was positive. This kit is comparable with other commercial kits and has been used elsewhere

(8–11).

The variables we analyzed were age, sex, personal medical history (for example, the presence of type 2

diabetes, hypertension, obstructive pulmonary disease, any heart condition, obesity, cancer, liver

steatosis, any autoimmune disease), and confirmed SARS-CoV-2 (through nasal swab or serologic

tests). The following AEs were considered of particular interest: fever (>37.5°C), adenopathy, diffuse

rash, edema, facial paralysis, orthostatic hypotension, headache, arthralgia, myalgia, nausea, vomit,

and diarrhea. Anti-S1 and anti-S2 IgG antibodies against SARS-CoV-2 were measured at baseline, 21-

28 days post-first dose (S1), and 21-28 days post-second dose if applicable.

Statistical Methods

The researchers reviewed the quality control and the anonymization of the database. Normality

assumption was evaluated with the Shapiro-Wilk test or Kolmogorov. Descriptive statistics such as

mean, standard deviation, median, interquartile range, frequencies, and percentages were computed.

Kruskal-Wallis test with Dunn's multiple comparisons test was performed for IgG comparisons. We

performed a mixed model where the dependent variable was the delta of the antibodies. The personal

variability and time were constructed as a random effect, whereas each type of vaccine, age, history of

SARS-CoV-2, and the interaction of each vaccine with age were fixed effects. BNT162b mRNA was

the reference vaccine group. For the analysis of AEs, the study population was divided into two

separate cohorts by stratifying according to the history of SARS-CoV-2 infection (1050 Naïve SARS-

CoV-2 subjects and 471 SARS-CoV-2 history subjects). Poisson generalized linear models (PGLM)

were performed on both cohorts with the counts of AEs events after the second dose as the outcome

variables. Included regressors were age, sex, body mass index, 1,000 AU/mL variation of IgG levels

after the second dose compared to the baseline IgG levels, type of vaccine and the interactions between

vaccines and IgG level variation. BNT162b mRNA was the reference vaccine group Missing

completely at random values were analyzed through complete case analysis since missing antibody

levels were less than 5% and we considered it incorrect to impute any AEFI. A sample size of 1870

patients was calculated, according to the primary aim, by using a mixed model formula with an alpha

of 0.05, power of 90%, the effect size of 0.15, and k=7. The statistical programs used were R v.4.0.3

and Python v. 3.8.3. The analyses were two-tailed. A p-value less than 0.05 was statistically

significant.

Results.

A total of 1867 patients were recruited from all countries: 1352 from Mexico, 42 from Italy, 260 from

Brazil, and 213 from Argentina. The most frequent vaccine used was ChAdOx1-S in 666 subjects,

Coronavac in 582, BNT162b2 mRNA in 289, Gam-COVID-Vac in 213, mRNA-1273 in 65,

Ad26.COV2 in 31, and Ad5-nCoV in 19. The mean (SD) age was 52 (16.8) years, being statistically

different across vaccine groups ($p<0.05$) as some vaccines were proposed to a particular age group.

Fifty-two % of subjects were men, 559 (30%) had obesity, and 501 (26.8%) had hypertension. Table 1

shows the medical history of patients divided by vaccine type.

History of SARS-CoV-2

Positive history of SARS-CoV-2 infection was considered if the volunteer had a confirmatory swab or

serological test. Additionally, it was considered if specific anti-S1 and anti-S2 IgG antibodies against

SARS-CoV-2 in plasma samples were > 15 AU/ml at baseline. We had a total of 627 positive cases

before vaccination, of which 165 (24.8%) received ChAdOx1-S, 246 (42.2%) Coronavac, 107 (37.1%)

BNT162b2 mRNA, 55 (25.8%) Gam-COVID-Vac, 31 (47.6%) mRNA-1273, 18 (58%) Ad26.COV2,

and 5 (26%) Ad5-nCoV.

Antibody titers

All vaccines showed significant anti-S1 and anti-S2 IgG antibody changes with significant differences

across vaccines and depending on SARS-CoV-2 history. In naïve patients, the highest increase

[Median (IQR) AU/ml] after the first dose was observed in mRNA-1273 [175.5 (76.7)], then

BNT162b mRNA [78.1(55)], and Ad5-nCoV [43.4(55.9)]. In subjects previously exposed to SARS-

CoV-2, the highest increase after the first dose was observed in mRNA-1273 [5920(4705)], then

BNT162b mRNA [2500(3080)], and ChAdOx1-S [1025(2489)].

In naïve patients, the highest increase [Median (IQR) AU/ml] after the second dose (T2) was observed

in mRNA-1273 [1875 (1190)], then BNT162b mRNA [998 (1241)], and Gam-COVID-Vac [501

(55.9)], whereas in SARS-CoV-2 previously exposed subjects the highest increase after the second

dose was observed in mRNA-1273 [4950(3060)], then Gam-COVID-Vac [3620 (5356)], and

BNT162b mRNA [2630 (2672.5)]. Figure 1 and Figure S1 show the change in antibody levels through

time, classified by vaccine type. Table 2 shows the antibody levels classified by vaccine type.

The mixed model that showed a positive effect of SARS-CoV-2 previous infection (B=767.468,

p<0.001), a positive interaction of age with previous SARS-CoV-2 infection (B=13.03, p=0.003), and

an interaction of Gam-COVID-Vac and ChAdOx1-S with age (B=46.5, p<0.001; B=20.5; p=0.006).

With respect to BNT162b mRNA, mRNA-1273 showed a higher change in antibody titers, while all

the other vaccines had less change (p<0.01). We computed sex in the model, but there was no

significant correlation, so we eliminated it from the final model. These results are shown in Table 3

and Figure 2.

Adverse Events Following Immunization (AEFI)

At least one AEFI was observed after the first dose in 71% of respondents receiving BNT162b2

mRNA, in 93% with mRNA-1273, 38% with Gam-COVID-Vac, 42% with Coronavac, 51% with

ChAdOx1-S, 74% with Ad5-nCoV, and 81% with Ad26.COV2. After the second dose, 65%, 88%,

23%, 33%, and 23% of the respondents experienced at least one AEFI after receiving BNT162b2

mRNA, mRNA-1273, Gam-COVID-Vac, Coronavac, and ChAdOx1-S, respectively. For each

vaccine, the majority of the adverse events occurred during the first 24 hrs after injection, either after

the first or the second dose. Patients receiving BNT162b2 mRNA, mRNA-1273, Gam-COVID-Vac,

Coronavac, ChAdOx1-S, Ad5-nCoV, and Ad26.COV2 subjectively qualified the AEFI after the first

dose as "very mild" or "mild" in 85%, 80%, 95%, 84%, 67%, 93%, and 57% of cases, respectively.

The AEFI after the second dose were qualified as "very mild" or "mild" by 82%, 49%, 98%, 89%, and

76% of patients receiving BNT162b2 mRNA, mRNA-1273, Gam-COVID-Vac, Coronavac, and

ChAdOx1-S respectively. Finally, 49% of patients receiving mRNA-1273 qualified for their adverse

events after the second dose as "moderate". Figure 3 shows the percentages of AEs per vaccine.

AEs were evaluated by using PGLM. The model for naïve SARS-CoV-2 subjects showed a negative

association of the count of adverse events with age (B=-0.0327, p<0.001) and sex (B=-0.8145,

p<0.001). A negative effect with respect to BNT162b mRNA was observed for Gam-COVID-Vac

(B=-1.8582, p<0.001), Coronavac (B=-0.5736, p=0.047) and ChAdOx1-S (B=-0.7871, p=0.006).

Finally, a positive interaction was found between the delta of antibodies due to vaccination and Gam-

COVID-Vac (B=0.5604, p=0.002).

The model for subjects previously exposed to SARS-CoV-2 showed a negative association of the

count of adverse events with age (B=-0.0190, p=0.013) and sex (B=-1.0042, p<0.001). The delta of the

antibodies showed a positive association with the count of adverse events (B=0.1018, p<0.001). A

positive effect with respect to BNT162b mRNA was observed for Gam-COVID-Vac (B=1.2737,

p=0.038), whereas a negative effect was observed for Coronavac (B=-1.1304, p<0.001). Negative

interactions were also found between the delta of antibodies due to vaccination and the vaccines Gam-

COVID-Vac (B=-8.5758, p<0.001) and ChAdOx1-S (B=-1.2386, p=0.013). See Tables 4 and 5 and

Figure 4 for details.

SARS-CoV-2 infection

Subjects were followed up to 28 days after the second dose administration. Between the first and

second dose, one (0.34%) patient with BNT162b2 mRNA, 0 with mRNA-1273, 1 (0.46%) with Gam-

COVID-Vac, 3 (0.51%) with Coronavac, 4 (0.6%) with ChAdOx1-S, 2 (10%) with Ad5-nCoV and 1

(3%) with Ad26.COV2 got the SARS-CoV-2 infection. After the second dose, 4 (1.3%) with

BNT162b2 mRNA, 1 (1.5%) with mRNA-1273, 0 with Gam-COVID-Vac, 18 (3%) with Coronavac,

and 11 (1.6%) with ChAdOx1-S became infected with SARS-CoV-2 (p<0.01).

Discussion

This multicentric study compared the 21-28 day seroconversion, the AEFI after first and second doses, and the associated predictors of AEs of seven of the most common vaccines used worldwide: BNT162b2 mRNA, mRNA-1273, Gam-COVID-Vac, Coronavac, ChAdOx1-S, Ad5-nCoV, and Ad26.COV2(4).

All vaccines showed immunogenicity and seroconversion after the complete vaccination scheme. The greatest change in antibody levels occurred with mRNA-1273, followed by BNT162b2 mRNA and Gam-COVID-Vac. Previous studies have shown a greater change in antibody titers for mRNA vector vaccines(12–14).

Our study found that the previous history of SARS-CoV-2 was related to an increase in change in antibody levels, as confirmed elsewhere (12). Also, we found higher levels in antibodies in older patients when they had previous SARS-CoV-2 infection and received Gam-COVID Vac or ChAdOx1-S. The previous study from Brazil did not find any association of antibody levels with age. We also found that except for the Gam-COVID Vac, in the other vaccines requiring two doses, the first dose was sufficient to induce maximal antibody levels in subjects with a history of SARS-CoV-2, as previously shown for the BNT162b2 mRNA vaccine(9,15–19)

The study of Logunov et al. that tested the efficacy of Gam-COVID Vac reported a higher increase of anti-SARS-CoV-2 IgG levels in the group of 18-30 years and no particular change in other age groups(20); however they excluded patients with previous SARS-CoV-2 infection, so these results are not comparable.

In the study conducted by Ramasamy et al., similar antibody titers were seen 28 days after the boost vaccination of ChAdOx1 across all groups, regardless of age or vaccine dose 21. It is possible that older patients in our group of the study had a more severe disease during SARS-CoV-2 infection and that they developed higher antibody titers, as it has been reported that patients aged higher than 60 can develop this condition (21).

For most patients, at least one AEFI occurred with all types of vaccines during the first 24 hrs. after

injection, and most of them qualified the symptoms as "very mild" or "mild" both after the first and

second dose, except for mRNA-1273 for which nearly half of the volunteers reported moderate

symptoms after the second dose as reported previously(18,20,22–27).

Our models showed a higher risk for women and young people to experience AEs. A real-world study

that reported AEFI of patients receiving BNT162b2 mRNA, mRNA-1273, or Ad26COV2 in one or

two doses (28) , an independent studies conducted with Gam-COVID also confirmed this results

(26,29). The study of Ramasamy et al. on ChAdOx1 showed more adverse events in the group of 18-

55 years old(22). The study conducted by Scott A et al. showed that severe adverse events (grade 5)

were more frequent in women with Ad5-nCoV (25).

On one hand, for the Naïve SARS-CoV-2 cohort, the vaccines Gam-COVID-Vac, Coronavac, and

ChAdOx1-S were related to fewer AEs after the second dose than BNT162b mRNA, and to the best of

our knowledge, there are no studies that compare AEs across these types of vaccines. Also, the delta of

antibodies was related to AEs after the second dose in Gam-COVID-Vac. As previously published,

mRNA-1273 had higher AEFIs when compared to BNT162b2 mRN(30).

On the other hand, for subjects previously exposed to SARS-CoV-2, those receiving Gam-COVID-

Vac showed a greater number of AEs than BNT162b mRNA, while subjects receiving Coronavac

showed significantly fewer events when compared to BNT162b mRNA. In general, an increase in the

antibody levels was related to an increase in the number of AEFIs, but for people receiving Gam-

COVID-Vac or ChAdOx1-S, greater changes in antibody levels corresponded to fewer AEs. This last

result should be taken with caution due to the limited number of patients falling in the category. To the

best of our knowledge, this has not been previously shown. The study conducted by Sung Hee-Lim et

al. that compared the antibody response with adverse events in patients vaccinated with BNT162b

mRNA or ChAdOx1-S reported no association with antibody change and adverse events; however,

patients previously infected with SARS-CoV-2 were excluded (31).

This study shows a low rate of infection in all vaccine groups. However, the highest rate of SARS-

CoV-2 infection in this follow-up period was for the one-dose vaccines Ad5-nCoV and Ad26.COV2.

Also, Coronavac had a high rate of infection. It is important to consider that the patients were followed

for a short period.

As a limitation of our study, we recognize that we had a small sample size in some vaccine groups

such as Ad5-nCoV; however, to the best of our knowledge, there is no other study where this type of

vaccine has been compared to other types of vaccines in a real-world setting. It will be of interest to

follow the IgG titers for a longer period and to evaluate the effect of heterologous combinations of

these vaccines.

This comparative study of vaccine types shows positive immunogenicity and seroconversion of

BNT162b2 mRNA, mRNA-1273, Gam-COVID-Vac, Coronavac, ChAdOx1-S, Ad5-nCoV, and

Ad26.COV2. The highest IgG response was for the mRNA vector vaccines and the lower for the

inactivated vaccine. Women and young individuals developed more AEs. In the naïve SARS-CoV-2

cohort, the vaccines Gam-COVID-Vac, Coronavac, and ChAdOx1-S were related to fewer AEs after

the second dose of BNT162b mRNA. For patients who received Gam-COVID-Vac, higher antibody

levels after the second dose were related to a greater number of AEs. In people previously exposed to

SARS-CoV-2, those receiving Gam-COVID-Vac showed a greater number of AEs than BNT162b

mRNA, while subjects receiving Coronavac showed significantly fewer events when compared to

BNT162b mRNA. In general, an increase in antibody levels was related to AEs.

Abbreviations

Adverse events following immunization (AEFI)

Severe adverse events (AEs).

Data Availability Statement: Data are available upon reasonable request to the authors.

Ethics statement: Ethics committee/local Institutional Review Board from Hospital Clinica Nova,

Humanitas Clinical and Research Center, Fundacion San Francisco Xavier, Ternium Health Center in

Rio, Hospital Municipal San Jose, Hospital Interzonal de Agudos San Felipe gave ethical approval.

Author Approval: all authors all authors read an approved the final version of the manuscript.

Author Contributions:

Conceptualization: MERI, MR, MT, MASS, and EA. Formal analysis: MERI, MR, MT, MASS, AGC,

CP, RL, S, MM, and RB. Investigation: MERI, AGC, MR, MT, MASS, CP, RL, RS, RB, MM, DRS,

YGHR, AGAV, GFDRP, IABF, RGF, EZ, ARAD, MHG, ABB, CA, GPB, MT, and MR. Resources:

MR, MT, and MASS Writing – original draft: MERI, AGC, CP, MM, RS, RL, RB, MASS, EA, DRS,

YGHR, AGAV, GFDRP, IABF, RGF, EZ, ARAD, MHG, ABB, CA, GPB, MT, MR. Writing – review

and editing: MERI, AGC, CP, MM, RS, RL, RB, MASS, EA, DRS, YGHR, AGAV, GFDRP, IABF,

RGF, EZ, ARAD, MHG, ABB, CA, GPB, MT, MR. Project administration: MR, MT, MASS.

Supervision: MERI, MT, MR, and MASS. Funding acquisition: MT, MR, and MASS. All authors

contributed to the article and approved the submitted version.

Funding: This research was conducted using private funding from Techint Group of Companies.

Conflict of Interest: The authors have declared no competing interest.

Acknowledgments:

This project has been supported by the Techint Group of Companies.

We thank Erika Bienek (Techint Group Community Relations Director), Walter Bruno (Humanitas

Communications Director), Cesar Bueno (Fundación São Francisco Xavier President), Mario Galli

(Techint Group Communications Director), Pablo Sturiale (Apsot Director) for their support of the

work and discussions.

References

1. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020 [Accessed April 19, 2020]

2. Kannan S, Shaik Syed Ali P, Sheeza A, Hemalatha K. COVID-19 (Novel Coronavirus 2019) – recent trends. *European Review for Medical and Pharmacological Sciences* (2020) **24**:2006–2011. doi: 10.26355/eurrev_202002_20378

3. Kumar S, Nyodu R, Maurya VK, Saxena SK. "Morphology, Genome Organization, Replication, and Pathogenesis of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2).," In: Saxena SK, editor. *Coronavirus Disease 2019 (COVID-19)*. Medical Virology: From Pathogenesis to Disease Control. Singapore: Springer Singapore (2020). p. 23–31 doi: 10.1007/978-981-15-4814-7_3

4. Covid-19 Vaccine Tracker: Latest Updates - The New York Times. https://www.nytimes.com/interactive/2020/science/coronavirus-vaccine-tracker.html [Accessed January 11, 2022]

5. Mascellino MT, Di Timoteo F, De Angelis M, Oliva A. Overview of the Main Anti-SARS-CoV-2 Vaccines: Mechanism of Action, Efficacy and Safety. *IDR* (2021) **Volume 14**:3459–3476. doi: 10.2147/IDR.S315727

6. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med* (2007) **147**:573–577. doi: 10.7326/0003-4819-147-8-200710160-00010

7. Chen M, Yuan Y, Zhou Y, Deng Z, Zhao J, Feng F, Zou H, Sun C. Safety of SARS-CoV-2 vaccines: a systematic review and meta-analysis of randomized controlled trials. *Infect Dis Poverty* (2021) **10**:94. doi: 10.1186/s40249-021-00878-5

8. Bonelli F, Sarasini A, Zierold C, Calleri M, Bonetti A, Vismara C, Blocki FA, Pallavicini L, Chinali A, Campisi D, et al. Clinical and Analytical Performance of an Automated Serological Test That Identifies S1/S2-Neutralizing IgG in COVID-19 Patients Semiquantitatively. *J Clin Microbiol* (2020) **58**: doi: 10.1128/JCM.01224-20

9. Levi R, Azzolini E, Pozzi C, Ubaldi L, Lagioia M, Mantovani A, Rescigno M. One dose of SARS-CoV-2 vaccine exponentially increases antibodies in individuals who have recovered from symptomatic COVID-19. *Journal of Clinical Investigation* (2021) **131**:e149154. doi: 10.1172/JCI149154

10. Levi R, Ubaldi L, Pozzi C, Angelotti G, Sandri MT, Azzolini E, Salvatici M, Savevski V, Mantovani A, Rescigno M. The antibody response to SARS-CoV-2 infection persists over at least 8 months in symptomatic patients. *Commun Med* (2021) **1**:32. doi: 10.1038/s43856-021-00032-0

11. Sandri MT, Azzolini E, Torri V, Carloni S, Pozzi C, Salvatici M, Tedeschi M, Castoldi M, Mantovani A, Rescigno M. SARS-CoV-2 serology in 4000 health care and administrative staff across seven sites in Lombardy, Italy. *Sci Rep* (2021) **11**:12312. doi: 10.1038/s41598-021-91773-4

12. Szczepanek J, Skorupa M, Goroncy A, Jarkiewicz-Tretyn J, Wypych A, Sandomierz D, Jarkiewicz-Tretyn A, Dejewska J, Ciechanowska K, Pałgan K, et al. Anti-SARS-CoV-2 IgG

against the S Protein: A Comparison of BNT162b2, mRNA-1273, ChAdOx1 nCoV-2019 and
Ad26.COV2.S Vaccines. *Vaccines* (2022) **10**:99. doi: 10.3390/vaccines10010099

13. Lau O, Vadlamudi NK. Immunogenicity and Safety of the COVID-19 Vaccines Compared With
Control in Healthy Adults: A Qualitative and Systematic Review. *Value in Health*
(2021)S1098301521017344. doi: 10.1016/j.jval.2021.09.003

14. Rotshild V, Hirsh-Raccah B, Miskin I, Muszkat M, Matok I. Comparing the clinical efficacy of
COVID-19 vaccines: a systematic review and network meta-analysis. *Sci Rep* (2021) **11**:22777.
doi: 10.1038/s41598-021-02321-z

15. Krammer F, Srivastava K, the PARIS team, Simon V. Robust spike antibody responses and
increased reactogenicity in seropositive individuals after a single dose of SARS-CoV-2 mRNA
vaccine. [preprint]. Allergy and Immunology (2021). doi: 10.1101/2021.01.29.21250653

16. Krammer F, Srivastava K, Alshammary H, Amoako AA, Awawda MH, Beach KF, Bermúdez-
González MC, Bielak DA, Carreño JM, Chernet RL, et al. Antibody Responses in Seropositive
Persons after a Single Dose of SARS-CoV-2 mRNA Vaccine. *N Engl J Med* (2021) **384**:1372–
1374. doi: 10.1056/NEJMc2101667

17. Saadat S, Rikhtegaran Tehrani Z, Logue J, Newman M, Frieman MB, Harris AD, Sajadi MM.
Binding and Neutralization Antibody Titers After a Single Vaccine Dose in Health Care Workers
Previously Infected With SARS-CoV-2. *JAMA* (2021) **325**:1467–1469. doi:
10.1001/jama.2021.3341

18. Sadoff J, Gray G, Vandebosch A, Cárdenas V, Shukarev G, Grinsztejn B, Goepfert PA, Truyers
C, Fennema H, Spiessens B, et al. Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine
against Covid-19. *N Engl J Med* (2021) **384**:2187–2201. doi: 10.1056/NEJMoa2101544

19. Samanovic MI, Cornelius AR, Gray-Gaillard SL, Allen JR, Karmacharya T, Wilson JP, Hyman
SW, Tuen M, Koralov SB, Mulligan MJ, et al. Robust immune responses after one dose of
BNT162b2 mRNA vaccine dose in SARS-CoV-2 experienced individuals. [preprint]. Infectious
Diseases (except HIV/AIDS) (2021). doi: 10.1101/2021.02.07.21251311

20. Logunov DY, Dolzhikova IV, Shcheblyakov DV, Tukhvatulin AI, Zubkova OV, Dzharullaeva
AS, Kovyrshina AV, Lubenets NL, Grousova DM, Erokhova AS, et al. Safety and efficacy of an
rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis
of a randomised controlled phase 3 trial in Russia. *The Lancet* (2021) **397**:671–681. doi:
10.1016/S0140-6736(21)00234-8

21. Cannistraci CV, Valsecchi MG, Capua I. Age-sex population adjusted analysis of disease severity
in epidemics as a tool to devise public health policies for COVID-19. *Sci Rep* (2021) **11**:11787.
doi: 10.1038/s41598-021-89615-4

22. Ramasamy MN, Minassian AM, Ewer KJ, Flaxman AL, Folegatti PM, Owens DR, Voysey M,
Aley PK, Angus B, Babbage G, et al. Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine
administered in a prime-boost regimen in young and old adults (COV002): a single-blind,
randomised, controlled, phase 2/3 trial. *The Lancet* (2020) **396**:1979–1993. doi: 10.1016/S0140-
6736(20)32466-1

23. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, Perez JL, Pérez Marc G,
Moreira ED, Zerbini C, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N
Engl J Med* (2020) **383**:2603–2615. doi: 10.1056/NEJMoa2034577

24. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, Diemert D, Spector SA,
Rouphael N, Creech CB, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N*
Engl J Med (2021) **384**:403–416. doi: 10.1056/NEJMoa2035389

25. Halperin SA, Ye L, MacKinnon-Cameron D, Smith B, Cahn PE, Ruiz-Palacios GM, Ikram A,
Lanas F, Lourdes Guerrero M, Muñoz Navarro SR, et al. Final efficacy analysis, interim safety
analysis, and immunogenicity of a single dose of recombinant novel coronavirus vaccine
(adenovirus type 5 vector) in adults 18 years and older: an international, multicentre, randomised,
double-blinded, placebo-controlled phase 3 trial. *The Lancet* (2022) **399**:237–248. doi:
10.1016/S0140-6736(21)02753-7

26. Jarynowski A, Semenov A, Kamiński M, Belik V. Mild Adverse Events of Sputnik V Vaccine in
Russia: Social Media Content Analysis of Telegram via Deep Learning. *J Med Internet Res*
(2021) **23**:e30529. doi: 10.2196/30529

27. Halder A, Imamura H, Condon S, Boroughs K, Nilsson SC, Anderson T, Caterson ID.
Pfizer/BioNtech BNT162b2: adverse events and insights from an Australian mass vaccination
clinic for COVID-19. *Intern Med J* (2022) **52**:121–124. doi: 10.1111/imj.15623

28. Beatty AL, Peyser ND, Butcher XE, Cocohoba JM, Lin F, Olgin JE, Pletcher MJ, Marcus GM.
Analysis of COVID-19 Vaccine Type and Adverse Effects Following Vaccination. *JAMA Netw*
Open (2021) **4**:e2140364. doi: 10.1001/jamanetworkopen.2021.40364

29. Babamahmoodi F, Saeedi M, Alizadeh-Navaei R, Hedayatizadeh-Omran A, Mousavi SA, Ovaise
G, Kordi S, Akbari Z, Azordeh M, Ahangarkani F, et al. Side effects and Immunogenicity
following administration of the Sputnik V COVID-19 vaccine in health care workers in Iran. *Sci*
Rep (2021) **11**:21464. doi: 10.1038/s41598-021-00963-7

30. Meo SA, Bukhari IA, Akram J, Meo AS, Klonoff DC. COVID-19 vaccines: comparison of
biological, pharmacological characteristics and adverse effects of Pfizer/BioNTech and Moderna
Vaccines. *Eur Rev Med Pharmacol Sci* (2021) **25**:1663–1669. doi:
10.26355/eurrev_202102_24877

31. Lim S-H, Choi S-H, Kim B, Kim J-Y, Ji Y-S, Kim S-H, Kim C-K, Kim T, Choo E-J, Moon J-E,
et al. Serum Antibody Response Comparison and Adverse Reaction Analysis in Healthcare
Workers Vaccinated with the BNT162b2 or ChAdOx1 COVID-19 Vaccine. *Vaccines* (2021)
9:1379. doi: 10.3390/vaccines9121379

Figure 1. IgG antibody levels by vaccine.

IgG antibody response was measured in serum of naïve and SARS-CoV-2 previously exposed (SARS-CoV-2
Exp) subjects at different time points (T0, T1 and T2) and vaccinated with different vaccine types. Samples ≥ 15
AU/mL were considered positive. Log scale on y axis. The box plots show the interquartile range, the horizontal
lines show the median values, and the whiskers indicate the minimum-to-maximum range. Each dot corresponds
to an individual subject. P- values were determined using 2-tailed Kruskal-Wallis test with Dunn's multiple
comparisons test. P values refer to baseline (T0) when there are no connecting lines.

Figure 2. Antibody titers classified by age type of vaccine and Time

Figure shows change in antibody concentration through age, divided SARS-CoV-2 history, and
vaccine type. Antibodies are expressed in logarithm and age has a Z distribution.

Figure 3. Percentage of systemic or severe adverse events after second dose, stratified by vaccine

Graph shows the percentage of systemic or severe adverse events grouped by frequency of appearance

Figure 4. Forest plot for the GLM Poisson model coefficients stratified by SARS-CoV-2 history

Women and young individuals developed more AEs. There is no effect on Body Mass Index. In the
naïve SARS-CoV-2 cohort, the vaccines Gam-COVID-Vac, Coronavac and ChAdOx1-S are related to
fewer AEFIs after the second dose than BNT162b mRNA. Also, for naïve SARS-CoV-2 patients who
received Gam-COVID-Vac, higher antibody levels after the second dose were related to a greater
number of AEs. In people previously exposed to SARS-CoV-2 , those receiving Gam-COVID-Vac
showed a greater number of AEs compared to BNT162b mRNA, while subjects receiving Coronavac
showed significantly fewer events when compared to BNT162b mRNA.

576   Table 1: General characteristics and medical history

	BNT162b2 mRNA (%)	mRNA-1273(%)	Gam-COVID-Vac (%)	Coronavac (%)	ChAdOx1-S (%)	Ad5-nCoV (%)	Ad26.COV 2(%)	p-value
Total	289	65	200	582	666	19	31	
Obesity	54(18.6)	11(16.9)	44(22)	188(32.3)	248(37.2)	8(42.1)	6(19.3)	<0.001
Smoker	37(12.8)	10(15.3)	27(13.5)	60(10.3)	38(5.7)	2(10.5)	2(6.4)	0.001
Hypertension	36(12.4)	2(3)	77(38.5)	116 (19.9)	265(39.7)	3(15.7)	2(6.4)	<0.001
Dyslipidemia	27(9.3)	3(4.6)	55(27.5)	76 (13.5)	150 (22.5)	1(5.2)	1(3.2)	<0.001
Type2 Diabetes	13(4.4)	1(1.5)	32(16)	78(13.4)	164(24.6)	0	0	<0.001
Asthma	9 (3.1)	3(4.6)	6(3)	13 (2.3)	17 (2.5)	0	0	0.79
Other autoimmune disease such as thyroiditis or psoriasis	9(3.1)	0	4(2)	29 (4.9)	60 (9)	1(5.2)	1(3.2)	<0.001
Rheumatoid arthritis	7(2.4)	1(1.5)	16(8)	11(1.8)	56 (8.4)	0	1(3.2)	<0.001
Surgery in the last year	6(2)	0	6(3)	26(4.4)	45(6.7)	3 (15)	1(3.2)	0.002
Previous Cancer	5(1.7)	1(1.5)	1(0.5)	5(0.8)	29(4.3)	0	0	0.001
NAFLD	4 (1.3)	0	7(3.5)	27(4.6)	21(3.1)	2(10.5)	0	0.042
Immunosuppressive treatment	4(1.3)	0	3(1.5)	6(1)	11(1.6)	0	0	0.86
Other liver disease	3 (1)	0	10(5)	10(1.7)	10(1.5)	0	0	0.02

End stage renal disease	2 (0.7)	0	2(1)	3(0.5)	11(1.6)	0	0	0.44
Pregnancy	2(0.7)	1(1.5)	0	3(0.5)	0	1(5.2)	0	0.005
Cirrhosis	2 (0.7)	0	0	2(0.3)	4 (0.6)	0	0	0.88
Pulmonar Obstructive Chronic Disease	1(0.3)	0	3(1.5)	4(0.6)	8 (1.2)	0	0	0.68
Coronary heart disease	1(0.3)	0	15(7.5)	3(0.5)	23(3.4)	0	0	<0.001
Gout	1(0.3)	0	10(5)	19(3.2)	27 (4.05)	0	0	0.02
Active Cancer	0	0	1(0.5)	1(0.17)	10(1.5)	0	0	0.054
Atrial Fibrilation	0	0	13(6.5)	6(1)	36 (5.4)	0	0	0.001
Congestive heart failure	0	0	18(9)	1(0.2)	12(1.8)	0	0	0.001
Stroke	0	0	2(1)	1(0.2)	13(1.9)	0	0	0.01
Trasplant	0	0	0	3(0.5)	5 (0.7)	0	0	0.63
Pregnancy	2	1	0	3(0.5)	0	1(5.2)	0	0.05

578 Table 1 shows the medical history and comparison between vaccine groups, $p<0.05$ was considered
579 statistically significant.

Table 2 Antibody differences between vaccine types

	SARS CoV-2 Naïve							p-value
Vaccine Median (IQR)	BNT162b2 mRNA (n=289)	mRNA-1273 (n=65)	Gam-COVID-Vac(n=213)	Coronavac ((n=582)	ChAdOx1-S (n=666)	Ad5-nCoV Cansino (n=19)	Ad26.COV2(n=31)	
Basal	3.8(0)	3.8 (26.6)	3.8 (0)	3.8(0.9)	3.8 (0)	3.8 (0)	3.8 (13.5)	p<0.005
After First Dose	78.1 (55.0)	175.5 (76.7)	22.1(44.5)	5.59 (8.0)	19.9 (39.6)	43.45 (55.9)	42.6 (57.5)	p<0.001
After Second Dose	998 (1241.0)	1875 (1190.0)	501 (1453.7)	109(97.6)	140.5 (199.5)	NA	NA	p<0.001
	SARS CoV-2 Positive							
Basal	66.9(130.8)	59.3 (67.9)	59.8 (157.3)	72.4 (119.0)	61.9 (134.8)	141.0 (175.2)	24 (66.7)	p<0.001
After First dose	2500.0(3080.0)	5920.0 (4705.0)	400.0 (3252.0)	264.5 (422.7)	1025.0 (2489.0)	182.0 (3734.6)	606.5 (23.69.0)	p<0.001
After Second Dose	2630.0 (2672.5)	4950.0 (3060)	3620 (5356)	279(337)	1020(1538)	NA	NA	p<0.001

*Kruskal-Wallis test was performed for comparison, p<0.05 was considered statistically significant

601   Table 3: Mixed model of antibody changes

	Estimate	Std. Error	95% CI	t value	p-value
(Intercept)	1909.02	345.73	1271.81 – 2546.23	5.52	<0.001
Age	-19.40	6.79	-32.6 7– -6.11	-2.855	0.004
SARS CoV2 previous infection	767.46	231.84	314.45 – 1220.31	3.31	0.001
mRNA-1273	1855.79	501.63	876.08 – 2836.04	3.69	<0.001
Gam-COVID-Vac	-2768.04	482.86	-3711.06 – -1824.4	-5.73	<0.001
Coronavac	-2184.03	430.93	-3025.82- -1342.09	-5.06	<0.001
ChAdOx1-S	-1808.26	361.77	-2514.69 - -1101.2	-4.99	<0.001
Ad5-nCoV	-2710.10	1542.89	-5727.76 – 304.30	-1.75	0.079
Ad26.COV2	-791.46	1291.27	-3317.21 – 1731.09	-0.613	0.54
Age*mRNA-1273	-20.82	15.95	-52 – 10.34	-1.3	0.192
Age*Gam-COVID-Vac	46.57	9.15	28.69 – 64.44	5.09	<0.001
Age*Coronavac	16.78	9.44	-1.67 – 35.22	1.77	0.076

Age*ChAdOx1-S	20.51	7.46	5.92 – 35 .08	2.749	0.006
Age*Ad5-nCoV	44.44	35.86	-25.68 – 114.54	1.23	0.215
Age*Ad26.COV2	-4.24	32.52	-67.82 – 59.33	-0.13	0.896
Age*COVID-19 infection	13.03	4.44	4.35 – 21.71	2.93	0.003

602 Mixed model. The dependent variable is the delta of antibodies. Time and personal variability are
603 random effects. Reference group BNT162b2 mRNA.

625 Table 4: Number of systemic or severe adverse events following vaccination in SARS-CoV-2   Naïve
626 subjects.

	β	e	95% CI	p-value
Intercept	0.73	2.08	0.95-4.57	0.066
Age	-0.03	0.96	0.95-0.97	<0.001
Sex (M)	-0.81	0.44	0.33-0.58	<0.001
Body Mass Index	0.015	1.01	0.99-1.03	0.175
Delta IgG	-0.21	0.80	0.60-1.06	0.128
mRNA-1273	0.46	1.59	0.71-3.54	0.255
Gam-COVID-Vac	-1.85	0.15	0.06-0.35	<0.001
Coronavac	-0.57	0.56	0.31-0.99	0.047
ChAdOx1-S	-0.78	0.45	0.25-0.79	0.006
Delta IgG*mRNA-1273	0.13	1.14	0.78-1.68	0.479
Delta IgG*Gam-COVID-Vac	0.56	1.75	1.22-2.49	0.002
Delta IgG*Coronavac	-1.63	0.19	0.01-2.39	0.201
Delta IgG*ChAdOx1-S	0.97	2.64	0.93-7.44	0.066

627   Poisson Generalized Linear Model regression. The reference group is BNT162b2 mRNA

641 Table 5. Number of systemic or severe adverse events following vaccination in SARS-CoV-2
642 previously exposed subjects.

	β	e	95% CI	p-value
Intercept	0.45	1.57	0.66-3.71	0.297
Age	-0.01	0.98	0.96-0.99	0.013
Sex (M)	-1.00	0.36	0.25-0.51	<0.001
Body Mass Index	0.004	1.00	0.97-1.03	0.776
Delta IgG	0.10	1.10	1.05-1.16	<0.001
mRNA-1273	0.23	1.26	0.48-3.31	0.628
Gam-COVID-Vac	1.27	3.57	1.07-11.89	0.038
Coronavac	-1.13	0.32	0.18-0.57	<0.001
ChAdOx1-S	-0.18	0.82	0.40-1.67	0.597
Delta IgG*mRNA-1273	-0.001	0.99	0.86-1.15	0.982
Delta IgG*Gam-COVID-Vac	-8.57	0.0001	0.00016-0.00022	<0.001
Delta IgG*Coronavac	0.19	1.22	0.52-2.84	0.644
Delta IgG*ChAdOx1-S	-1.23	0.28	0.10-0.76	0.013

643 Poisson Generalized Linear Model regression. Reference group is BNT162b2 mRNA

Percentage of severe or systemic adverse reactions per vaccine, after 2nd dose