Title

Modifiable lifestyle activities affect cognition in cognitively healthy middle-aged individuals

at risk for late-life Alzheimer's Disease

Running title: Midlife lifestyle, AD risk and cognition

Authors

Amy **Heneghan** [ab1], Feng **Deng** [ab1], Katie **Wells** [c], Karen **Ritchie** [cd], Graciela **Muniz-Terrera**

[c], Craig W **Ritchie** [c], Brian **Lawlor** [ab], Lorina **Naci** [ab*]

[a] Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,

Dublin, Ireland.

[b] Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

[c] Edinburgh Dementia Prevention, University of Edinburgh, Edinburgh, UK

[d] INSERM and University of Montpellier, Montpellier, France.

[1] These authors contributed equally to this work

[*] **Corresponding author:**

Lorina Naci

School of Psychology

Trinity College Institute of Neuroscience

Global Brain Health Institute

Trinity College Dublin

Dublin, Ireland

Telephone: +353 (0)87 688 5642

Email: nacil@tcd.ie

Abstract

It is now acknowledged that Alzheimer's Disease (AD) processes are present decades before

the onset of clinical symptoms, but it remains unknown whether lifestyle factors can protect

against these early AD processes in mid-life. We asked whether modifiable lifestyle activities

impact cognition in middle-aged individuals who are cognitively healthy, but at risk for late

life AD. Participants (40–59 years) completed cognitive and clinical assessments at baseline

(N = 206) and two years follow-up (N = 174). Mid-life activities were measured with the

Lifetime of Experiences Questionnaire. We assessed the impact of lifestyle activities, known

risk factors for sporadic late-onset AD (Apolipoprotein E ε4 allele status, family history of

dementia, and the Cardiovascular Risk Factors Aging and Dementia score), and their

interactions on cognition. More frequent engagement in physically, socially and intellectually

stimulating activities was associated with better cognition (verbal, spatial and relational

memory), at baseline and follow-up. Critically, more frequent engagement in these activities

was associated with stronger cognition (verbal and visuospatial functions, and conjunctive

short-term memory binding) in individuals with family history of dementia. Impaired

visuospatial function is one of the earliest cognitive deficits in AD and has previously

associated with increased AD risk in this cohort. Additionally, conjunctive memory functions

have been found impaired in the pre-symptomatic stages of AD. These findings suggest that

modifiable lifestyle activities offset cognitive decrements due to AD risk in mid-life and

support the targeting of modifiable lifestyle activities for the prevention of Alzheimer's

Disease.

Keywords: Cognitively Healthy; Mid-life; Cognition; Lifestyle Factors; Risk Factors;

Alzheimer's Disease

Introduction

Dementia is a growing pandemic that presents profound challenges to health care systems, families, and societies throughout the world. Alzheimer's Disease (AD), the most common etiology of dementia, is characterized by relentless neurodegeneration and accelerated cognitive decline in the years following presentation of clinical symptoms. It is now accepted that AD pathological processes are present decades before clinically relevant symptoms appear [1-2]. In the absence of effective pharmacological treatments, there is an immediate need for early identification, intervention and risk burden modification [3-4] for AD.

Growing evidence suggests that up to 40% of all dementia cases are associated with known lifestyle-related modifiable risk factors, such as alcohol consumption, obesity and hypertension among others [4]. As exposure to most of these risk factors begins decades before dementia onset, interventions must be implemented in mid-life [5-7]. Mid-life, thus, presents a critical and unique window for disease-altering interventions, before the manifestation of substantial brain damage. However, the indicators of AD in mid-life, and the impact of modifiable lifestyle factors on the incipient disease process remain poorly understood.

AD is neuropathologically characterized by the accumulation of beta-amyloid (Aβ) and hyperphosphorylated TAU (pTau) [8-9]. The etiology of the sporadic form of AD remains poorly understood. However, recent studies suggest that tau deposition, is a key etiological factor that presages sporadic AD [10-12]. Analyses of thousands human brains across the lifespan show that tau pathology begins about a decade before formation of Aβ plaques [13]. Additionally, tau pathology, but not Aβ, correlates with progressive grey matter loss [14] and cognitive impairment [15].Neuropathological findings show that tau pathology starts in late young adulthood and early midlife [30–40years], in subcortical nucleus locus coeruleus (LC)

[9] —the key brain site for the production of noradrenaline— then spreads initially to the

transentorhinal and entorhinal cortex [8], to the hippocampal and neocortical association

cortex, and finally throughout neocortex. By contrast, Aβ plaques are usually first deposited

in the association neocortex of the temporal lobe, and then extend throughout the cortex,

including subcortical structures, with disease progression [13]. Therefore, brain changes

related to sporadic AD in early midlife (i.e. 40-50 years) are likely associated with

neurodegeneration from to pTau deposition and, in later age groups, Aβ plaques additionally

contribute strongly to underlying neurodegeneration.

Studies on preclinical AD have used risk stratification approaches to investigate early,

preclinical changes. Key risk factors include the Apolipoprotein E (APOE) ε4 genotype [16-

17], the main genetic risk factor for sporadic late onset AD in the Indo-European population

[18], and family history (FH) of dementia [16, 19-20]. Several dementia risk scores

incorporating lifestyle risk factors have been devised [21-23]. Amongst them, the

Cardiovascular Risk Factors Aging and Dementia (CAIDE) score has been optimized for

middle-aged populations [23] and has been validated in a large US population followed

longitudinally over 40 years [24].

The PREVENT Dementia Program [25] initiated in 2013, is a prospective study of the

cognitively healthy middle-aged children of persons with dementia, designed to seek out

clinical and biological changes, which may subsequently be used as short-term outcome

measures for midlife secondary preventions. Studies of this and other similar cohorts of

cognitively healthy midlife individuals have related these three risk factors to a range of

structural and functional brain changes, including APOE ε4 genotype to loss of volume in the

hippocampal molecular layer [26], cerebral hyperperfusion [27-28], reduced grey matter

volume in the right hippocampus, precentral gyrus, and cerebellar cortex [29], and to

decreased cortical thickness in the frontal cortex [30]; FH to volumetric alterations in

hippocampal subfields and to disrupted white matter integrity [27, 31]; and, CAIDE to whole

brain atrophy [32-34] and to hippocampal volume loss [31, 34]. All three risk groups have

also been found to impact cognition in mid-life. APOE ε4 genotype has been significantly

associated with improvements in verbal, spatial and relational memory [35], immediate recall

[36] and form perception [34], FH with poorer verbal processing and memory performance in

participants under 65 years [37], and poorer executive function [38], and higher CAIDE

scores with impaired verbal and visuospatial functions, and short-term (conjunctive) memory

[34-35], and impaired executive function [39]. Taken together, these studies have established

that risk for late-life AD, including risk that incorporates lifestyle factors, has a significant

impact on the brain health and cognition of middle-aged individuals who are presently

cognitively healthy.

As a multidimensional construct, lifestyle has a multipronged impact on cognition and the

brain. By contrast to aforementioned cardiovascular risk factors captured by the CAIDE

score, several lifestyle activities have been found to protect brain health and cognition in later

life. Factors, such as education and occupational complexity have been associated with

preservation of cognitive function in order adults [40] and reduced symptom severity in

Alzheimer's disease [4, 41-43], a phenomenon known as "cognitive resilience". These factors

are thought to explain why, in late-life Alzheimer's disease, the level of cognitive impairment

shows substantial variability even when accounting for key pathologies including Aβ and

pathological tau [1, 44]. One prominent account of the biological mechanisms mediating the

relationship between stimulating lifestyle activities and cognitive resilience in AD [45-46]

suggests that environmental enrichment upregulates the noradrenergic system which

originated in the locus coeruleus, that otherwise depletes with age [47] and AD pathology

[10], leading to compensatory brain mechanisms for cognitive function.

While epidemiological evidence strongly suggests that education and occupation contribute

to cognitive resilience [48], there is a renewed interest in the additional contribution of other

activities undertaken in mid-life, given their potential modifiability. For example, Gow et al.

(2017) [49] and Chan et al. (2018) [40] used the Lifetime of Experiences Questionnaire

(LEQ) and found that physically, socially and intellectually stimulating lifestyle activities

undertaken in mid-life, independently of education, help to maintain late-life cognitive

performance in older adults, after adjusting for childhood cognitive ability [49]. However, it

is not known whether the protection conferred by lifestyle activities, other than education,

offsets the impact of AD risk in mid-life, or whether it builds up gradually over time with its

benefits viable only in late life or relative to established AD pathology. The answer to this

question is critical for identifying interventions that target modifiable factors for the

prevention of Alzheimer's disease from the earliest life stages.

To address this gap, we investigated the impact of interactions between three risk factors for

late-life AD (APOE ε4 genotype, FH and CAIDE) with lifestyle activities on cognition,

independently of sex, age and years of education. These relationships were investigated in a

large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206),

and at two years follow-up (N=174). To relate our findings to previous studies, we used the

LEQ [50], the same instrument as in aforementioned studies, to evaluate lifestyle activities

specific to mid-life, yielding two composite factors, (a) occupation and managerial

responsibility, and (b) physical, social and intellectual activities. Our hypothesis was that

there would be a significant association between mid-life lifestyle factors and cognitive

performance in domains already shown in this cohort to be affected by risk for late-life AD.

Methods

Participants

PREVENT is an ongoing longitudinal multi-site research programme based across the UK

and Ireland, seeking to identify early biomarkers of AD and elaborate on risk-mechanism

interactions for neurodegenerative diseases decades before the cardinal symptoms of

dementia emerge. Its protocol has been described in detail elsewhere [25]. In the first

PREVENT programme phase, participants were recruited at a single site, via the dementia

register database held at the West London National Health Service (NHS) Trust, of the UK

National Health Service, the Join Dementia Research website

(https://www.joindementiaresearch.nihr.ac.uk/), through public presentations, social media

and word of mouth. Procedures involving experiments on human subjects were done in

accord with the ethical standards of the Institutional Review Board of Imperial College

London and in accord with the Helsinki Declaration of 1975. Approval for the study was

granted by the NHS Research Ethics Committee London Camberwell St Giles. Consented

participants were seen at the West London NHS Trust, where they underwent a range of

clinical and cognitive assessments [25]. The cohort comprised cognitively healthy volunteers

aged 40-59 years. The West London dataset was used here to avail of both baseline and

follow-up testing data acquired for this cohort. 210 individuals (62 male; 148 female) were

tested at baseline, with 188 (89.5%) (55 male; 133 female) retained at 2 years follow-up

(Table 1).

Assessments

Risk factors

APOE□4 Genotyping

The process of APOE □4 allele identification is outlined in detail in Ritchie et al. (2017) [34].

In brief, genomic DNA was isolated from blood samples and APOE genotyping was

performed. All members of the research and clinical teams were blind to the result of APOE

genotyping. In this study, APOE □4 risk is determined by ≥1 APOE □4 allele. 75/210 carried

≥1 APOE □4 allele (See Table 1).

Family History

FH was determined by a 'yes'/'no' question during clinical visits, which asked participants

whether a parent had a diagnosis of dementia. Participants were asked to include the

dementia subtype if known, but answering 'yes' alone categorized a participant as FH+. The

answer 'no' likely captured both participants with no family history of dementia, and

participants for whom FH was unknown. In summary, participants were defined as FH+ if at

least one parent was diagnosed with dementia. Cases where the FH was unknown or partially

known were not recorded outside of the binary yes/no scoring. 103/210 were FH+.

Cardiovascular Risk Factors, Aging, and Incidence of Dementia (CAIDE) score

CAIDE is a composite scale of estimated future dementia risk based on mid-life

cardiovascular measures [51-52]. It takes into consideration the individual's age, sex,

educational attainment, APOE □4 genotype, activity level, BMI, cholesterol and systolic

blood pressure [23] and is scored on a range of 0 – 18. A higher score indicates greater risk.

The CAIDE dementia risk score was calculated for each participant at baseline and follow-

up.

Cognitive testing

The Addenbrooke's Cognitive Assessment (ACE) III [53] was recorded at the follow-up

session, but not at the baseline. Subsequent analyses that focused on the subset of

assessments performed at both testing session did not include the ACE) III.

Cognitive function was assessed at baseline and follow up with the COGNITO

neuropsychological battery [54], designed to examine information processing across a wide

range of cognitive functions in adults of all ages and not restricted to those functions usually

implicated in dementia detection in the elderly. Tests are administered using a tactile screen

to capture information processing time as well as response accuracy and require about 40

minutes to complete. The tests, by order of presentation, are: reaction time; reading;

comprehension of phonemes, phrases, and syntax; focused and divided attention in both

visual and auditory modalities; visual working memory (visual tracking with auditory

interference); the Stroop test; immediate, delayed, and recognition trials for verbal recall

(name list); delayed recognition of spatial stimuli (faces); visuospatial associative learning;

visuospatial span; form perception; denomination of common objects; spatial reasoning;

copying of meaningful and meaningless figures; verbal fluency with se-mantic and phonetic

prompts; immediate recall of a narrative; immediate recall of a description of the relative

position of objects; vocabulary; implicit memory (recognition of new and previously learned

material).

The COGNITO tests are designed to test several aspects of cognition, including attention

(task: visual attention), memory (tasks: narrative recall, description recall, implicit memory,

name-face association, working memory), language (tasks: phoneme comprehension, verbal

fluency) and visuospatial abilities (task: geometric figure recognition) [54-55]. Based on

previous studies [54-55], 11 summary variables from the COGNITO battery capturing the

above functions were used here [for a list see Supporting Information (SI)].

Additionally, we used the Visual Short-Term Memory Binding task (VSTMBT) [56], a

computer-based task that assesses visual short-term memory binding of single features, e.g.,

complex shape or color combinations, or feature conjunctions, e.g., shape and color

combinations. In the single feature condition, participants must identify whether the test

stimuli (three random 6-sided polygons) are the "same" as or "different" to the studied

stimuli in terms of shape (shape only) or color (color only). In the binding condition,

participants are required to correctly identify if both the shape and color of the test stimuli

match studied stimuli. Two summary variables from the VSTMBT were the percentage of

correctly recognized items from the two conditions.

Measurement of lifestyle activities

The Lifetime of Experiences Questionnaire (LEQ) [50], designed to take a lifespan approach

to the measurement of cognitive resilience [57-59] and mental activity, measures engagement

in a broad range of lifestyle activities across three stages of life: young adulthood (13-29

years), mid-life (30-64 years) and late life (65 years onwards). Therefore, the LEQ is

preferable for looking at a mid-life cohort compared to other scales that capture dementia-

specific risk related to modifiable lifestyle factors (e.g., LIBRA [60]). The LEQ comprises

sub-scores capturing "specific" activities, reflecting the primary activity undertaken in each

life stage and "non-specific" activities, reflecting engagement in physical, social and

intellectual activities in any stage. For the purpose of this paper, we define mid-life 'lifestyle'

as all the activities captured by the LEQ (below).

Mid-life specific score

The mid-life specific component score centers on occupation and comprises two sub-scores.

For the first occupational sub-score, participants were asked to record their primary

occupation in each 5-year interval from age 30 to age at assessment. Each reported

occupation was scored according to the International Standard Classification of Occupations

(ISCO 08) guidelines (https://www.ilo.org/public/english/bureau/stat/isco/isco08/), and

scores were inverted and summed. The second sub-score is a measure of the managerial

responsibility associated with reported occupations. If participants indicated that they were

employed in a managerial capacity, the number of people that they oversaw in four of their

reported occupations was documented. Managerial responsibility was scored as follows; 0

people = 8, 1-5 people = 16, 5-10 people = 24 and 10+ people = 32. The highest score is

recorded as the managerial responsibility sub-score. The occupational history and managerial

sub-scores were summed and multiplied by a normalization factor of 0.25. Normalization

ensures that the mid-life specific and non-specific scores have comparable mean values [50].

Mid-life non-specific score

The non-specific score assesses frequency of engagement in 7 activities, capturing those of a

physically, socially and intellectually stimulating nature, scored on a 5-point Likert scale of

frequency (never, less than monthly, monthly, fortnightly, weekly, daily). Scores range from

0 – 35, with higher scores reflecting more frequent engagement in such activities. The items

included in the scale are socializing with family or friends, practicing a musical instrument,

practicing an artistic pastime, engagement in physical activity that is mildly, moderately, or

vigorously energetic, reading, practicing a second language and travel. The travel item asks

participants if they have visited any of a list of continents between the ages of 30-54.

Responses were scored on a 5-point scale as follows; none, 1-2 regions, 3-4 regions, 5

regions, 6-7 regions, all regions.

Statistical analyses

To reduce the number of multiple comparisons between the cognitive tests, given the

multitude of cognitive assessments obtained for this cohort (13 summary variables, see SI and

Table S1), in an independent study of this dataset [35], three composite cognitive components

were extracted from the above cognitive assessments, by using rotated principal component

analysis (rPCA) (Figure S1). In subsequent analyses, we used these cognitive components,

rather than the individual tests to measure the impact of risk and protective lifestyle factors

on cognition.

At baseline, the three components were: 1) C1: verbal, spatial and relational memory; 2) C2:

working and short-term (single-feature) memory; 3) C3: verbal and visuospatial functions,

and short-term (conjunctive) memory (Figure S1 A). Higher scores in C1 and C2 reflected

better performance, whereas higher scores in C3 reflected poorer performance. For ease of

visualization and comparability to C1 and C2, values of C3 were recoded with reversed sign.

The highest loading tasks and their weights for each component differed slightly for the

cross-sectional and longitudinal data. There was high similarity between baseline and follow-

up for C1 ($\phi = 0.90$), and C3 ($\phi = 0.76$), with relatively low similarly for C2 ($\phi = 0.40$)

(Figure S1 B). Although C1 and C3 had relatively high similarity across time, the

longitudinal changes in these components could not be quantified by direct comparison,

because they captured different cognitive functions at each time-point and any longitudinal

change would be hard to interpret.

We used the Statistical Package for Social Sciences (SPSS V.26) and R software for all

statistical analyses. The normality of the data was assessed by combining the visualization of

a quantile-quantile plot and the Shapiro–Wilk test. Demographic and clinical information of

the study cohort was analyzed across risk groups using chi-square (χ^2 tests) for categorical

variables and Mann Whitney U tests for continuous (discrete) variables, given that they were

not normally distributed in this cohort. Subsequently, we used hierarchical regression models

to look at the contribution of mid-life lifestyle factors (LEQ specific and non-specific scores)

risk factors (APOE $_\square$4 genotype and FH) and of their interactions on cognitive performance,

at baseline and follow-up. The effect of APOE $_\square$4 and CAIDE risk factors was modelled

independently, in order to avoid modelling the variance associated with APOE

genotype in the same model twice. In each case, dependent variables were the

aforementioned composite cognitive components, each assessed in a separate model.

The variance of some parts of physical activity is accounted for twice in the model, in the

CAIDE score and in the non-specific LEQ factor, resulting in weakening of its statistical

contribution that can by captured by the lifestyle factor. Therefore, any effect of physical

activity that can be captured by the lifestyle factor (independent variable) in the same model

as CAIDE (independent variable), can only provide a conservative estimate of the

contribution of physical activity to the dependent variable of cognitive performance.

As aforementioned, the baseline and follow-up datasets were analyzed independently, due to

the slightly differing cognitive domains for each visit. Although the two datasets were not

independent, we considered each of them in its own right, because during 2 years, a

proportion of our participants may have had substantial brain health changes that are yet

subthreshold to clinical manifestations. Therefore, the follow-up dataset has the potential to

reveal the impact of age on variables of interest. Corrections for multiple comparisons were

performed across all analyses.

To avoid multicollinearity, we mean centered continuous variables (specific and non-specific

scores). Age, sex and years of education were included as covariates in hierarchical models.

A significant interaction effect between a risk and a lifestyle factor would indicate that the

effect of the lifestyle factor on cognitive performance differs across levels/values of the risk

factor. For any observed interactions, we plotted the regression of the lifestyle factor on

cognitive performance for each level/value of the risk factor [61], to interpret the effect. We

then tested the significance of the slopes of the simple regression lines, to investigate in

which level/value of the risk factor we found an effect of lifestyle on cognitive performance.

Results

Demographic characteristics

In the first PREVENT study phase, 210 participants were recruited from a single site. All 210 completed several cognitive tests and the LEQ questionnaire. 188 were also assessed at the follow up, two years later. At baseline, 2 participants were missing information relating to their APOE ϵ4 genotype, with a further 2 missing cognitive data. The final baseline cohort that could be used for the analyses was N = 206. At follow-up, 12 participants were missing cognitive data, alongside the 2 participants who were missing APOE ϵ4 genotype data. Therefore, the final follow-up cohort was N = 174. Mild cognitive impairment and dementia were ruled out based on detailed clinical assessment at baseline and follow-up. Please see the Supplementary Information and Table S1 for a list of all the outcome variables collected at baseline and follow-up.

Demographic specifications of the cohort at baseline and follow-up, stratified by APOE ϵ4 genotype and family history of dementia, are shown in Table 1. There were no significant differences in age, sex or years of education between the groups. As expected, APOE ϵ4 allele genotype was more frequently found in the FH+ than FH- group at baseline (p = 0.01) and follow up (p = 0.02). CAIDE scores (including APOE status) were significantly higher for the FH+ than FH- group at baseline (p = 0.03) and follow up (p = 0.003). Naturally, CAIDE scores including APOE status were significantly higher for the APOE ϵ4+ than APOE ϵ4- group at baseline (p = 0.0003) and follow up (p = 0.0002), but when APOE status was excluded the CAIDE scores did not differ between the two groups (Table 1).

Table 1 about here please

Verbal, Spatial and Relational memory

At baseline, a hierarchical regression model with lifestyle factors, i.e., the specific and non-

specific LEQ scores, risk factors (APOE ε4 and FH), and age, sex and years of education as

covariates, and verbal, spatial and relational memory as the dependent variable was

significant overall [F (7, 198) = 5.84, R^2 = 0.17, p < 0.0001] (Supplementary Table 2a), and

showed a significant positive association between verbal, spatial and relational memory and

education [β (SE) = 0.09 (0.02), p < 0.0001] (Figure 1a). Similarly, at follow up the model's

performance was significant [F (7, 166) = 5.52, R^2 = 0.19, p < 0.0001], and there was a

significant positive association between verbal, spatial and relational memory and education

[β (SE) = 0.10 (0.02), p < 0.0001] (Figure 1b). Higher education values were significantly

associated with higher performance at both timepoints. The inclusion of risk by lifestyle

interaction terms (FH × specific score, FH × non-specific score, APOE ε4 × specific score,

APOE ε4 × non-specific score) into the hierarchical regression model (Supplementary Table

2b) did not show any significant associations between lifestyle x risk interaction terms and

cognitive performance, at baseline or follow-up.

355   *Figure 1 about here please*

The hierarchical regression models with lifestyle factors and CAIDE as independent

variables, and verbal, spatial and relational memory as the dependent was significant at

baseline [F (3, 202) = 3.11, R^2 = 0.04, p = 0.03] and at follow-up [F (3, 166) = 3.52, R^2 =

0.04, p = 0.02] (Supplementary Table 2c). At both timepoints, we found a significant positive

association between the non-specific LEQ factor and verbal, spatial and relational memory

(baseline: [β (SE) = 0.04 (0.02), p = 0.02], figure 2a) and (follow up: [β (SE) = 0.07 (0.02), p

= 0.002], figure 2b). More frequent engagement in physically, socially and intellectually

stimulating activities was associated with better verbal, spatial and relational memory. This

association was independent of CAIDE and therefore of age, sex and years of education, as

included in this score. The inclusion of CAIDE by lifestyle interaction terms (CAIDE $\times$

specific score, CAIDE $\times$ non-specific score) into the hierarchical regression model

(Supplementary Table 2d) did not show any significant associations between lifestyle x risk

interaction terms and cognitive performance, at baseline or follow-up.

Figure 2 about here please

Working and Short-Term (Single-Feature) Memory

Hierarchical regression analyses showed no significant associations of any of the independent

variables and performance on the working and short-term (single-feature) memory, at

baseline or follow up (Supplementary Table 3a, 3c) Furthermore, no significant associations

with interaction terms were observed at either timepoint (Supplementary Table 3b, 3d).

Verbal and Visuospatial Functions, and Short-Term (Conjunctive) Memory

At baseline, the hierarchical regression model was significant [F (7, 198) = 2.39, R^2 = 0.08, p

= 0.02], and showed a significant negative association between verbal and visuospatial

function and short-term (conjunctive) memory with age [β (SE) = -0.04 (0.01), p = 0.006]

(Supplementary Table 4a). Higher age was associated with poorer performance. At follow-

up, there were no significant associations between the main independent variables with verbal

and visuospatial function and short-term (conjunctive) memory. The inclusion of risk by

lifestyle interaction terms in the hierarchical model revealed a significant association between

the non-specific LEQ score x FH interaction term and cognitive performance [β (SE) = 0.11

(0.05), p = 0.01] (Supplementary Table 4b). To interpret this interaction, we investigated the

relationship between cognitive performance and the non-specific LEQ score for the FH+ and

FH- groups independently (Figure 3). We found a significant negative relationship between

non-specific lifestyle activities and cognition in the FH+ group [β (SE)= 0.08 (0.03), p =

0.02), that was independent of age, sex, or years of education. We did not observe a

relationship between non-specific lifestyle activities and cognition in the FH- group [β (SE) =

-0.03 (0.03), p = 0.34]. These results suggested that for individuals with positive family

history, more frequent engagement in the non-specific LEQ factor — namely physically,

socially and intellectually engaging activities — was associated with better performance in

verbal and visuospatial function and short-term (conjunctive) memory. Finally, we also

observed trend associations between cognitive performance and the specific LEQ factor score

x FH interaction term [β (SE) = -0.07 (0.04), p = 0.06]. This association, while suggestive of

a role for occupational complexity on cognition, stratified by family history risk group, is

weak and needs to be investigated further in future studies. Additionally, as previously

reported [35] we observed a significant negative association between CAIDE and verbal and

visuospatial functions and short-term (conjunctive) memory at baseline (Supplementary

Table 4c). There were no associations between interactions of CAIDE x lifestyle and

cognition at either baseline or follow up (Supplementary Table 4d).

407   *Figure 3 about here please*

Conclusion

It is now acknowledged that Alzheimer's Disease processes are present decades before the

onset of clinical symptoms [2, 62, but, to date, it has remained unknown whether lifestyle

factors can already protect against these early AD processes in mid-life. We asked whether

these modifiable lifestyle activities impact cognition in middle-aged individuals who are

cognitively healthy but at risk for late life AD. Lifestyle activities significantly impacted

cognition in mid-life. Individuals with greater educational attainment showed stronger

cognition in a composite dimension capturing verbal, spatial and relational memory. This

result is consistent with a previous study [34] of this cohort showing that visuospatial abilities

were positively associated with education. It is also consistent with epidemiological studies

on older adults showing that education contributes to cognitive resilience in older life [48].

Our cohort is highly educated, with approximately 30% reaching a post-graduate

qualification [34]. As the LEQ education measure captures total years of education, the effect

of education we observed, reflects long-term effects set in motion from early life and young

adulthood.

The key question in this study, however, was to investigate any additional contribution of

activities undertaken in mid-life, independently of education. The first novel finding of this

study was that more frequent engagement in physically, socially and intellectually stimulating

activities in mid-life was associated with stronger cognition in a composite dimension

capturing verbal, spatial and relational memory, both at baseline and at follow-up. This effect

was independent of sex, age, years of education and cardiovascular factors captured by the

CAIDE score. The second novel finding was that physically, socially and intellectually

stimulating activities undertaken in mid-life had a significant effect in the cognition of

middle-aged individuals at risk for late-life AD. Specifically, higher engagement in these

activities was associated with significantly stronger cognition in another composite

dimension, capturing verbal and visuospatial functions and short-term (conjunctive) memory

in cognitively healthy individuals, who were at risk for late life AD though family history of

dementia, at follow up. Importantly, this effect was independent of age, sex, and years of

education. The presence of this effect in the follow-up but not the baseline dataset is likely

due to the older age of the cohort at follow-up. Short-term (conjunctive) memory functions

have been found impaired in the pre-symptomatic stages of AD [56]. Additionally, impaired

visuospatial function is one of the earliest cognitive deficits observed in AD [63-64] and has

previously been linked to increased AD risk in this cohort [34-35]. In similarly cognitively

healthy, but older (> 65 years) North American cohorts [65-66], changes in cerebrospinal

fluid biomarkers related to inflammation have been associated with changes in visuospatial

cognitive performance, thereby suggesting a biochemically-mediated effect of early

pathology in presymptomatic individuals.

Taken together, our results suggest that stimulating lifestyle activities may boost cognitive

functions that are very vulnerable to AD risk and early AD neuropathology [56, 63-64].

However, we caution that the interpretability of the effect of lifestyle activities on each

individual function is limited by their composite assessment in this study and requires

individuation in future studies with a longer longitudinal follow-up window. Physically,

socially and intellectually stimulating mid-life activities have been shown to influence late-

life cognition, suggesting that they contribute to the cognitive resilience that mitigates the

effect of age-related cognitive decline and AD neuropathology [40, 49]. Our finding advances

understanding by showing that engagement with these activities contributes to cognitive

resilience to risk of AD, or even incipient AD neuropathology, from mid-life, in individuals

who are presently cognitively healthy.

Family history of dementia is a well-established risk factor [16, 19-20], independent of the

genetic risk bestowed by APOE ▯4 genotype, that captures both genetic and environmental

risk influences. For example, an individual who has one or more parent with dementia, may,

independently to the presence of the APOE ▯4 genotype, be exposed to negative

environmental influences that contribute to neuropathology, including increased stress [67],

caregiver burden [68-69] and reduced participation in enriching environments due to

caregiving duties [70]. Therefore, alongside contributing to cognitive resilience, enhanced

engagement with physically, socially and intellectually stimulating activities may positively

impact individuals with a family history of dementia by counteracting these negative

environmental influences.

What might be the mechanism by which stimulating activities impact cognition in midlife?

Models of AD disease delineated by neuropathological staging [9] place the locus coeruleus,

a small nucleus in the pontine tegmentum region of the brainstem, at the pathogenesis of AD

[10-11, 71]. The LC is responsible for the production of neurotransmitter noradrenaline (NA)

[72], a major driver of the brain's arousal system, which strongly modulates high-order

cognition. Novelty provides a key trigger for arousal, and, thus, the LC–NA activity occurs

strongly in response to novel stimuli. Therefore, stimulating activities upregulate the LC and

may protect it from early neurodegeneration due to pTau deposition. Furthermore, as the

brain site of noradrenaline, the LC is key to the protective effect of lifestyle factors on the

whole brain. Studies suggests that environmental enrichment, through activities such as those

measured in this study, upregulates the noradrenergic system, which otherwise depletes with

age [47] and AD pathology [10], leading to compensatory brain mechanisms for cognitive

function, such as the strengthening of the fronto-parietal brain and other large-scale brain

networks [45-46].

In summary, our findings suggest that modifiable lifestyle activities may offset AD risk-

related cognitive decrements in mid-life and support the targeting of certain modifiable

lifestyle activities for the prevention of Alzheimer's Disease, especially in those with a family

history. These activities include socializing with family or friends, practicing a musical

instrument, practicing an artistic pastime, engagement in energetic physical activities,

reading, practicing a second language and travel. Given the apparent lifestyle contributions

both as risk and protective factors of dementia [4], a healthy lifestyle may be the individual's

current best defense against sporadic late onset AD. The modifiability of the lifestyle

activities identified here renders them promising cost-effective candidates for intervention

and prevention strategies from early life. These activities may be particularly significant for

non-pharmacological interventions for AD in low and middle-income countries (LMIC),

where barriers to education are more prevalent than in high income countries [4].

Methodological considerations

As education is strongly positively linked to IQ [73] and our cohort was highly educated, the

question of reverse causation arises [40, 49, 74]. This points to the possibility that cognitive

abilities may determine engagement in stimulating activities, rather than the inverse.

However, the effect on mid-life verbal and visuospatial functions, and short-term

(conjunctive) memory is independent of the total years of education, which shows that

education does not directly drive this effect. Furthermore, we found that mid-life lifestyle

activities were associated with improved cognitive performance only in the family history

positive group, which is not, at least not prima facie, more educated than the family history

negative group. Thus, this effect is likely independent of any indirect effects of education.

The use of composite cognitive domains that capture slightly different functions in the two

assessment points [35] prevents investigation of the impact of lifestyle on cognitive changes

over time in this study. Nevertheless, previous studies from this cohort have shown only

subtle changes over the two-year period [75-76], possibly due to the relatively young age

range of the sample and the short follow up window [34, 55]. Therefore, future studies that

follow this cohort over a longer period and test hypotheses informed by the previous study

waves are needed to determine the longitudinal impact of lifestyle activities in cognitively

healthy middle-aged individuals at risk for late life AD.

It is worth noting that we found no evidence that lifestyle factors can mitigate the impact of

the CAIDE risk, or for interactions between CAIDE, lifestyle and cognition, as reported in

some previous studies (see Kivipelto et al. 2018 [77] for a review). Unlike the CAIDE score

(including systolic blood pressure, BMI, total cholesterol, physical activity), the lifestyle

factors captured by our instrument (the LEQ) are not primarily related to cardiovascular

wellbeing (with the exception of exercise, 1/9 items), and therefore not likely to impact

greatly on cardiovascular health in this relatively young cohort (i.e., 40–59 years). By

contrast, the studies reviewed by Kivipelto et al. (2018) include older populations (> 60–70

years), where cardiovascular factors are more prevalent drivers of health vulnerabilities, and

furthermore those studies, unlike the present one, had a strong focus on cardiovascular health,

including nutritional guidance, exercise and monitoring and management of metabolic and

vascular risk factors.

Acknowledgement

F.D. was funded by the Provost PhD Award Scheme from Trinity College Dublin, to L.N.

L.N. was also funded by a L'Oréal-UNESCO for Women In Science International Rising

Talent Award, the Welcome Trust Institutional Strategic Support grant, and the Global Brain

Health Institute Project Grant.

The PREVENT-Dementia study is supported by the UK Alzheimer's Society (grant numbers

178 and 264), the US Alzheimer's Association (grant number TriBEKa-17-519007) and

philanthropic donations.

We thank all PREVENT-Dementia participants for their enthusiastic participation in this

study. We also thank the Research Delivery service at West London NHS Trust and the

Wolfson Clinical Imaging Facility at Imperial College London for their support in running

the study.

Conflict of interest

The authors declare no conflict of interest.

References

[1] Jack, C. R., Jr, Knopman, D. S., Jagust, W. J., Petersen, R. C., Weiner, M. W., Aisen, P.

S., Shaw, L. M., Vemuri, P., Wiste, H. J., Weigand, S. D., Lesnick, T. G., Pankratz, V. S.,

Donohue, M. C., & Trojanowski, J. Q. (2013). Tracking pathophysiological processes in

Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *The Lancet.*

Neurology, 12(2), 207–216. https://doi.org/10.1016/S1474-4422(12)70291-0

[2] Ritchie, K., Ritchie, C. W., Yaffe, K., Skoog, I., & Scarmeas, N. (2015). Is late-onset

Alzheimer's disease really a disease of midlife?. *Alzheimer's & dementia (New York, N.*

Y.), 1(2), 122–130. https://doi.org/10.1016/j.trci.2015.06.004

[3] Ritchie, C.W., Waymont, J.M.J., Pennington, C. *et al.* The Scottish Brain Health Service

Model: Rationale and Scientific Basis for a National Care Pathway of Brain Health Services

in Scotland. *J Prev Alzheimers Dis* (2021). https://doi.org/10.14283/jpad.2021.63

[4] Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne,

C., Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S. G., Dias, A., Fox, N., Gitlin,

L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Ogunniyi, A., Orgeta, V., …

Mukadam, N. (2020). Dementia prevention, intervention, and care: 2020 report of the Lancet

. *Lancet (London, England), 396*(10248), 413–446. https://doi.org/10.1016/S0140-

6736(20)30367-6

[5] Irwin, K., Sexton, C., Daniel, T., Lawlor, B., & Naci, L. (2018). Healthy Aging and

Dementia: Two Roads Diverging in Midlife?. *Frontiers in aging neuroscience*, *10*, 275.

https://doi.org/10.3389/fnagi.2018.00275

[6] Lachman, M. E., Teshale, S., & Agrigoroaei, S. (2015). Midlife as a Pivotal Period in the

Life Course: Balancing Growth and Decline at the Crossroads of Youth and Old

Age. *International journal of behavioral development*, 39(1), 20–31.

https://doi.org/10.1177/0165025414533223

[7] Gottesman, R. F., Schneider, A. L., Zhou, Y., Coresh, J., Green, E., Gupta, N., Knopman,

D. S., Mintz, A., Rahmim, A., Sharrett, A. R., Wagenknecht, L. E., Wong, D. F., & Mosley,

T. H. (2017). Association Between Midlife Vascular Risk Factors and Estimated Brain

Amyloid Deposition. *JAMA*, *317*(14), 1443–1450. https://doi.org/10.1001/jama.2017.3090

[8] Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the

pathologic process in Alzheimer disease: age categories from 1 to 100 years. *Journal of

neuropathology and experimental neurology*, *70*(11), 960–969.

https://doi.org/10.1097/NEN.0b013e318232a379

[9] Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes.

Acta neuropathologica, *82*(4), 239–259. https://doi.org/10.1007/BF00308809

[10] Jacobs, H., Becker, J. A., Kwong, K., Engels-Domínguez, N., Prokopiou, P. C., Papp, K.

V., Properzi, M., Hampton, O. L., d'Oleire Uquillas, F., Sanchez, J. S., Rentz, D. M., El

Fakhri, G., Normandin, M. D., Price, J. C., Bennett, D. A., Sperling, R. A., & Johnson, K. A.

(2021). In vivo and neuropathology data support locus coeruleus integrity as indicator of

Alzheimer's disease pathology and cognitive decline. *Science translational medicine*,

13(612), eabj2511. https://doi.org/10.1126/scitranslmed.abj2511

[11] Arnsten, A., Datta, D., Del Tredici, K., & Braak, H. (2021). Hypothesis: Tau pathology

is an initiating factor in sporadic Alzheimer's disease. *Alzheimer's & dementia : the journal of

the Alzheimer's Association*, *17*(1), 115–124. https://doi.org/10.1002/alz.12192

[12] Kametani, F., & Hasegawa, M. (2018). Reconsideration of Amyloid Hypothesis and Tau

Hypothesis in Alzheimer's Disease. *Frontiers in neuroscience*, *12*, 25.

https://doi.org/10.3389/fnins.2018.00025

[13] Braak, H., & Del Tredici, K. (2015). Neuroanatomy and pathology of sporadic

Alzheimer's disease. *Advances in anatomy, embryology, and cell biology*, *215*, 1–162.

[14] La Joie, R., Ayakta, N., Seeley, W. W., Borys, E., Boxer, A. L., DeCarli, C., Doré, V.,

Grinberg, L. T., Huang, E., Hwang, J. H., Ikonomovic, M. D., Jack, C., Jr, Jagust, W. J., Jin,

L. W., Klunk, W. E., Kofler, J., Lesman-Segev, O. H., Lockhart, S. N., Lowe, V. J., Masters,

C. L., … Rabinovici, G. D. (2019). Multisite study of the relationships between antemortem

[11C]PIB-PET Centiloid values and postmortem measures of Alzheimer's disease

neuropathology. *Alzheimer's & dementia : the journal of the Alzheimer's Association*, *15*(2),

205–216. https://doi.org/10.1016/j.jalz.2018.09.001

[15] Giannakopoulos, P., Herrmann, F. R., Bussière, T., Bouras, C., Kövari, E., Perl, D. P.,

Morrison, J. H., Gold, G., & Hof, P. R. (2003). Tangle and neuron numbers, but not amyloid

load, predict cognitive status in Alzheimer's disease. *Neurology*, *60*(9), 1495–1500.

https://doi.org/10.1212/01.wnl.0000063311.58879.01

[16] Donix, M., Small, G. W., & Bookheimer, S. Y. (2012). Family history and APOE-4

genetic risk in Alzheimer's disease. *Neuropsychology review*, *22*(3), 298–309.

https://doi.org/10.1007/s11065-012-9193-2

[17] Liu, C. C., Liu, C. C., Kanekiyo, T., Xu, H., & Bu, G. (2013). Apolipoprotein E and

Alzheimer disease: risk, mechanisms and therapy. *Nature reviews. Neurology, 9*(2), 106–118.

https://doi.org/10.1038/nrneurol.2012.263

[18] Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C.,

DeStafano, A. L., Bis, J. C., Beecham, G. W., Grenier-Boley, B., Russo, G., Thorton-Wells,

T. A., Jones, N., Smith, A. V., Chouraki, V., Thomas, C., Ikram, M. A., Zelenika, D.,

Vardarajan, B. N., Kamatani, Y., … Amouyel, P. (2013). Meta-analysis of 74,046 individuals

identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics*, *45*(12), 1452–

1458. https://doi.org/10.1038/ng.2802

[19] Berti, V., Mosconi, L., Glodzik, L., Li, Y., Murray, J., De Santi, S., Pupi, A., Tsui, W.,

& De Leon, M. J. (2011). Structural brain changes in normal individuals with a maternal

history of Alzheimer's. *Neurobiology of aging*, *32*(12), 2325.e17–2325.e2.325E26.

https://doi.org/10.1016/j.neurobiolaging.2011.01.001

[20] Scarabino, D., Gambina, G., Broggio, E., Pelliccia, F., & Corbo, R. M. (2016). Influence

of family history of dementia in the development and progression of late-onset Alzheimer's

disease. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, *171*(2),

250-256. https://doi.org/10.1002/ajmg.b.32399

[21] Barnes, D. E., Beiser, A. S., Lee, A., Langa, K. M., Koyama, A., Preis, S. R., Neuhaus,

J., McCammon, R. J., Yaffe, K., Seshadri, S., Haan, M. N., & Weir, D. R. (2014).

Development and validation of a brief dementia screening indicator for primary

care. *Alzheimer's & dementia : the journal of the Alzheimer's Association*, *10*(6), 656–665.e1.

https://doi.org/10.1016/j.jalz.2013.11.006

[22] Deckers, K., van Boxtel, M. P., Schiepers, O. J., de Vugt, M., Muñoz Sánchez, J. L.,

Anstey, K. J., Brayne, C., Dartigues, J. F., Engedal, K., Kivipelto, M., Ritchie, K., Starr, J.

M., Yaffe, K., Irving, K., Verhey, F. R., & Köhler, S. (2015). Target risk factors for dementia

prevention: a systematic review and Delphi consensus study on the evidence from

observational studies. *International journal of geriatric psychiatry*, *30*(3), 234–246.

https://doi.org/10.1002/gps.4245

[23] Kivipelto, M., Ngandu, T., Laatikainen, T., Winblad, B., Soininen, H., & Tuomilehto, J.

(2006). Risk score for the prediction of dementia risk in 20 years among middle aged people:

a longitudinal, population-based study. *The Lancet. Neurology, 5*(9), 735–741.

https://doi.org/10.1016/S1474-4422(06)70537-3

[24] Exalto, L. G., Quesenberry, C. P., Barnes, D., Kivipelto, M., Biessels, G. J., & Whitmer,

R. A. (2014). Midlife risk score for the prediction of dementia four decades later. *Alzheimer's*

& dementia : the journal of the Alzheimer's Association, *10*(5), 562–570.

https://doi.org/10.1016/j.jalz.2013.05.1772

[25] Ritchie, C. W., & Ritchie, K. (2012). The PREVENT study: a prospective cohort study

to identify mid-life biomarkers of late-onset Alzheimer's disease. *BMJ open*, *2*(6), e001893.

https://doi.org/10.1136/bmjopen-2012-001893

[26] Dounavi, M. E., Mak, E., Wells, K., Ritchie, K., Ritchie, C. W., Su, L., & O'Brien, J. T.

(2020). Volumetric alterations in the hippocampal subfields of subjects at increased risk of

dementia. *Neurobiology of aging*, *91*, 36-44.

https://doi.org/10.1016/j.neurobiolaging.2020.03.006

[27] Mak, E., Dounavi, M. E., Low, A., Carter, S. F., McKiernan, E., Williams, G. B., ... & T

O'Brien, J. (2021). Proximity to dementia onset and multi-modal neuroimaging changes: The

prevent-dementia study. *NeuroImage*, *229*, 117749.

https://doi.org/10.1016/j.neuroimage.2021.117749

[28] McKiernan, E. F., Mak, E., Dounavi, M. E., Wells, K., Ritchie, C., Williams, G., ... &

O'brien, J. (2020). Regional hyperperfusion in cognitively normal APOE ε4 allele carriers in

mid-life: analysis of ASL pilot data from the PREVENT-Dementia cohort. *Journal of*

Neurology, Neurosurgery & Psychiatry, *91*(8), 861-866. https://doi.org/10.1136/jnnp-2020-

322924

[29] Cacciaglia, R., Molinuevo, J. L., Falcón, C., Brugulat-Serrat, A., Sánchez-Benavides, G.,

Gramunt, N., Esteller, M., Morán, S., Minguillón, C., Fauria, K., Gispert, J. D., & ALFA

study (2018). Effects of APOE-ε4 allele load on brain morphology in a cohort of middle-aged

healthy individuals with enriched genetic risk for Alzheimer's disease. *Alzheimer's &*

dementia : the journal of the Alzheimer's Association, *14*(7), 902–912.

https://doi.org/10.1016/j.jalz.2018.01.016

[30] Fennema-Notestine, C., Panizzon, M. S., Thompson, W. R., Chen, C. H., Eyler, L. T.,

Fischl, B., Franz, C. E., Grant, M. D., Jak, A. J., Jernigan, T. L., Lyons, M. J., Neale, M. C.,

Seidman, L. J., Tsuang, M. T., Xian, H., Dale, A. M., & Kremen, W. S. (2011). Presence of

ApoE ε4 allele associated with thinner frontal cortex in middle age. *Journal of Alzheimer's*

disease : JAD, *26 Suppl 3*(Suppl 3), 49–60. https://doi.org/10.3233/JAD-2011-0002

[31] McKeever, A., Paris, A. F., Cullen, J., Hayes, L., Ritchie, C. W., Ritchie, K., ... & Su, L.

(2020). Hippocampal Subfield volumes in middle-aged adults at risk of dementia. *Journal of*

Alzheimer's Disease, *75*(4), 1211-1218. https://doi.org/10.3233/JAD-200238

[32] Liu, X., Dounavi, M. E., Ritchie, K., Wells, K., Ritchie, C. W., Su, L., Muniz-Terrera,

G., & O'Brien, J. T. (2021). Higher midlife CAIDE score is associated with increased brain

atrophy in a cohort of cognitively healthy middle-aged individuals. *Journal of*

neurology, *268*(5), 1962–1971. https://doi.org/10.1007/s00415-020-10383-8

[33] O'Brien, J. T., Firbank, M. J., Ritchie, K., Wells, K., Williams, G. B., Ritchie, C. W., &

Su, L. (2020). Association between midlife dementia risk factors and longitudinal brain

atrophy: the PREVENT-Dementia study. *Journal of neurology, neurosurgery, and*

psychiatry, *91*(2), 158–161. https://doi.org/10.1136/jnnp-2019-321652

[34] Ritchie, K., Carrière, I., Su, L., O'Brien, J. T., Lovestone, S., Wells, K., & Ritchie, C. W.

(2017). The midlife cognitive profiles of adults at high risk of late-onset Alzheimer's disease:

The PREVENT study. *Alzheimer's & dementia : the journal of the Alzheimer's*

Association, *13*(10), 1089–1097. https://doi.org/10.1016/j.jalz.2017.02.008

[35] Deng, F., Wells, K., Ritchie, K., Muniz-Terrera, G., Ritchie, C. W., O'Brien, J.T.,

Lawlor, B., Naci, L. (2022) Functional connectivity of the locus coeruleus with the

hippocampus and the cognitive profiles of asymptomatic mid-life individuals at risk for

Alzheimer's disease. *bioRxiv.*

[36] Jochemsen, H. M., Muller, M., van der Graaf, Y., & Geerlings, M. I. (2012). APOE ε4

differentially influences change in memory performance depending on age. The SMART-MR

study. *Neurobiology of Aging*, *33*(4), 832.e15–832.e8.32E22.

https://doi.org/10.1016/j.neurobiolaging.2011.07.016

[37] Talboom, J. S., Håberg, A., De Both, M. D., Naymik, M. A., Schrauwen, I., Lewis, C.

R., Bertinelli, S. F., Hammersland, C., Fritz, M. A., Myers, A. J., Hay, M., Barnes, C. A.,

Glisky, E., Ryan, L., & Huentelman, M. J. (2019). Family history of Alzheimer's disease

alters cognition and is modified by medical and genetic factors. *eLife*, *8*, e46179.

https://doi.org/10.7554/eLife.46179

[38] Debette, S., Wolf, P. A., Beiser, A., Au, R., Himali, J. J., Pikula, A., Auerbach, S.,

Decarli, C., & Seshadri, S. (2009). Association of parental dementia with cognitive and brain

MRI measures in middle-aged adults. *Neurology*, *73*(24), 2071–2078.

https://doi.org/10.1212/WNL.0b013e3181c67833

[39] Stephen, R., Liu, Y., Ngandu, T., Rinne, J. O., Kemppainen, N., Parkkola, R.,

Laatikainen, T., Paajanen, T., Hänninen, T., Strandberg, T., Antikainen, R., Tuomilehto, J.,

Keinänen Kiukaanniemi, S., Vanninen, R., Helisalmi, S., Levälahti, E., Kivipelto, M.,

Soininen, H., & Solomon, A. (2017). Associations of CAIDE Dementia Risk Score with

MRI, PIB-PET measures, and cognition. *Journal of Alzheimer's disease : JAD*, *59*(2), 695–

705. https://doi.org/10.3233/JAD-170092

[40] Chan, D., Shafto, M., Kievit, R., Matthews, F., Spink, M., Valenzuela, M., Cam-CAN,

& Henson, R. N. (2018). Lifestyle activities in mid-life contribute to cognitive reserve in late-

life, independent of education, occupation, and late-life activities. *Neurobiology of aging*, *70*,

180–183. https://doi.org/10.1016/j.neurobiolaging.2018.06.012

[41] Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D.,

Ballard, C., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L. N.,

Howard, R., Kales, H. C., Larson, E. B., Ritchie, K., Rockwood, K., Sampson, E. L., Samus,

Q., … Mukadam, N. (2017). Dementia prevention, intervention, and care. *Lancet (London,*

England), *390*(10113), 2673–2734. https://doi.org/10.1016/S0140-6736(17)31363-6

[42] Wang, H. X., MacDonald, S. W., Dekhtyar, S., & Fratiglioni, L. (2017). Association of

lifelong exposure to cognitive reserve-enhancing factors with dementia risk: A community-

based cohort study. *PLoS medicine*, *14*(3), e1002251.

https://doi.org/10.1371/journal.pmed.1002251

[43] Dekhtyar, S., Marseglia, A., Xu, W., Darin-Mattsson, A., Wang, H. X., & Fratiglioni, L.

(2019). Genetic risk of dementia mitigated by cognitive reserve: A cohort study. *Annals of*

neurology, *86*(1), 68–78. https://doi.org/10.1002/ana.25501

[44] Franzmeier, N., Koutsouleris, N., Benzinger, T., Goate, A., Karch, C. M., Fagan, A. M.,

McDade, E., Duering, M., Dichgans, M., Levin, J., Gordon, B. A., Lim, Y. Y., Masters, C. L.,

Rossor, M., Fox, N. C., O'Connor, A., Chhatwal, J., Salloway, S., Danek, A., Hassenstab, J.,

… Ewers, M. (2020). Predicting sporadic Alzheimer's disease progression via inherited

Alzheimer's disease-informed machine-learning. *Alzheimer's & dementia : the journal of the*

Alzheimer's Association, *16*(3), 501–511. https://doi.org/10.1002/alz.12032

[45] Robertson I. H. (2014). Right hemisphere role in cognitive reserve. *Neurobiology of*

aging, *35*(6), 1375–1385. https://doi.org/10.1016/j.neurobiolaging.2013.11.028

[46] Robertson I. H. (2013). A noradrenergic theory of cognitive reserve: implications for

Alzheimer's disease. *Neurobiology of aging*, *34*(1), 298–308.

https://doi.org/10.1016/j.neurobiolaging.2012.05.019

[47] Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Cam-

CAN, Howard, R., & Hämmerer, D. (2020). Noradrenergic-dependent functions are

associated with age-related locus coeruleus signal intensity differences. *Nature*

communications, *11*(1), 1712.

[48] Richards, M., & Deary, I. J. (2005). A life course approach to cognitive reserve: a model

for cognitive aging and development?. *Annals of neurology*, *58*(4), 617–622.

https://doi.org/10.1002/ana.20637

[49] Gow, A. J., Pattie, A., & Deary, I. J. (2017). Lifecourse Activity Participation From

Early, Mid, and Later Adulthood as Determinants of Cognitive Aging: The Lothian Birth

Cohort 1921. The journals of gerontology. Series B, *Psychological sciences and social*

sciences, *72*(1), 25–37. https://doi.org/10.1093/geronb/gbw124

[50] Valenzuela, M. J., & Sachdev, P. (2007). Assessment of complex mental activity across

the lifespan: development of the Lifetime of Experiences Questionnaire

(LEQ). *Psychological medicine, 37*(7), 1015–1025. https://doi.org/10.1017/S003329170600938X

[51] Fayosse, A., Nguyen, D. P., Dugravot, A., Dumurgier, J., Tabak, A. G., Kivimäki, M.,

Sabia, S., & Singh-Manoux, A. (2020). Risk prediction models for dementia: role of age and

cardiometabolic risk factors. *BMC medicine*, *18*(1), 107. https://doi.org/10.1186/s12916-020-

01578-x

[52] Sindi, S., Calov, E., Fokkens, J., Ngandu, T., Soininen, H., Tuomilehto, J., & Kivipelto,

M. (2015). The CAIDE Dementia Risk Score App: The development of an evidence-based

mobile application to predict the risk of dementia. *Alzheimer's & dementia (Amsterdam,*

Netherlands), *1*(3), 328–333. https://doi.org/10.1016/j.dadm.2015.06.005

[53] Hsieh, S., Schubert, S., Hoon, C., Mioshi, E., & Hodges, J. R. (2013). Validation of the

Addenbrooke's Cognitive Examination III in frontotemporal dementia and Alzheimer's

disease. *Dementia and geriatric cognitive disorders*, *36*(3-4), 242–250.

https://doi.org/10.1159/000351671

[54] Ritchie, K., de Roquefeuil, G., Ritchie, C., Besset, A., Poulain, V., Artero, S., &

Ancelin, M. L. (2014). COGNITO: computerized assessment of information

processing. *Journal of Psychology and Psychotherapy*, *4*(2). https://doi.org/10.4172/2161-

0487.1000136

[55] Ritchie, K., Carrière, I., Howett, D., Su, L., Hornberger, M., O'Brien, J. T., Ritchie, C.

W., & Chan, D. (2018). Allocentric and Egocentric Spatial Processing in Middle-Aged

Adults at High Risk of Late-Onset Alzheimer's Disease: The PREVENT Dementia

Study. *Journal of Alzheimer's disease : JAD*, *65*(3), 885–896. https://doi.org/10.3233/JAD-

180432

[56] Parra, M. A., Abrahams, S., Logie, R. H., Méndez, L. G., Lopera, F., & Della Sala, S.

(2010). Visual short-term memory binding deficits in familial Alzheimer's disease. *Brain : a*

journal of neurology, *133*(9), 2702–2713. https://doi.org/10.1093/brain/awq148

[57] Stern Y. (2009). Cognitive reserve. *Neuropsychologia*, 47(10), 2015–2028.

https://doi.org/10.1016/j.neuropsychologia.2009.03.004

[58] Stern Y. (2012). Cognitive reserve in ageing and Alzheimer's disease. *The Lancet.*

Neurology, *11*(11), 1006–1012. https://doi.org/10.1016/S1474-4422(12)70191-6

[59] Stern, Y., Barnes, C. A., Grady, C., Jones, R. N., & Raz, N. (2019). Brain reserve,

cognitive reserve, compensation, and maintenance: operationalization, validity, and

mechanisms of cognitive resilience. *Neurobiology of aging*, *83*, 124–129.

https://doi.org/10.1016/j.neurobiolaging.2019.03.022

[60] Schiepers, O., Köhler, S., Deckers, K., Irving, K., O'Donnell, C. A., van den Akker, M.,

Verhey, F., Vos, S., de Vugt, M. E., & van Boxtel, M. (2018). Lifestyle for Brain Health

(LIBRA): a new model for dementia prevention. *International journal of geriatric*

psychiatry, *33*(1), 167–175. https://doi.org/10.1002/gps.4700

[61] Aiken, L. S., & West, S. G. (1991). Multiple regression: Testing and interpreting

interactions. Newbury Park, CA: Sage.

[62] Jack, C. R., Jr, Knopman, D. S., Jagust, W. J., Shaw, L. M., Aisen, P. S., Weiner, M. W.,

Petersen, R. C., & Trojanowski, J. Q. (2010). Hypothetical model of dynamic biomarkers of

the Alzheimer's pathological cascade. *The Lancet. Neurology*, *9*(1), 119–128.

https://doi.org/10.1016/S1474-4422(09)70299-6

[63] Williams, O. A., An, Y., Armstrong, N. M., Kitner-Triolo, M., Ferrucci, L., & Resnick,

S. M. (2020). Profiles of Cognitive Change in Preclinical and Prodromal Alzheimer's Disease

Using Change-Point Analysis. *Journal of Alzheimer's disease : JAD, 75*(4), 1169–1180.

https://doi.org/10.3233/JAD-191268

[64] Laukka, E. J., Macdonald, S. W., Fratiglioni, L., & Bäckman, L. (2012). Preclinical

cognitive trajectories differ for Alzheimer's disease and vascular dementia. *Journal of the*

International Neuropsychological Society : JINS, *18*(2), 191–199.

https://doi.org/10.1017/S1355617711001718

[65] Miron, J., Picard, C., Lafaille-Magnan, M. É., Savard, M., Labonté, A., Breitner, J.,

Rosa-Neto, P., Auld, D., Poirier, J., & PREVENT-AD research group (2019). Association of

TLR4 with Alzheimer's disease risk and presymptomatic biomarkers of inflammation.

Alzheimer's & dementia : the journal of the Alzheimer's Association, *15*(7), 951–960.

https://doi.org/10.1016/j.jalz.2019.03.012

[66] Noble, J. M., Manly, J. J., Schupf, N., Tang, M. X., Mayeux, R., & Luchsinger, J. A.

(2010). Association of C-reactive protein with cognitive impairment. *Archives of neurology*,

67(1), 87–92. https://doi.org/10.1001/archneurol.2009.308

[67] Franks, K. H., Bransby, L., Saling, M. M., & Pase, M. P. (2021). Association of Stress

with Risk of Dementia and Mild Cognitive Impairment: A Systematic Review and Meta-

Analysis. *Journal of Alzheimer's disease : JAD*, *82*(4), 1573–1590.

https://doi.org/10.3233/JAD-210094

[68] Chiao, C. Y., Wu, H. S., & Hsiao, C. Y. (2015). Caregiver burden for informal

caregivers of patients with dementia: A systematic review. *International nursing*

review, *62*(3), 340–350. https://doi.org/10.1111/inr.12194

[69] Yu, H., Wang, X., He, R., Liang, R., & Zhou, L. (2015). Measuring the Caregiver

Burden of Caring for Community-Residing People with Alzheimer's Disease. *PloS*

one, 10(7), e0132168. https://doi.org/10.1371/journal.pone.0132168

[70] Ross, K. M., & Carroll, J. E. (2017). Alzheimer's Disease Caregivers' Health and

Important Resources: Next Steps in Intervention Research. *Psychosomatic medicine*, *79*(7),

732–734. https://doi.org/10.1097/PSY.0000000000000518

[71] Márquez, F., & Yassa, M. A. (2019). Neuroimaging Biomarkers for Alzheimer's

Disease. *Molecular neurodegeneration*, *14*(1), 21. https://doi.org/10.1186/s13024-019-0325-

5

[72] Amaral, D. G., & Sinnamon, H. M. (1977). The locus coeruleus: neurobiology of a

central noradrenergic nucleus. *Progress in neurobiology*, *9*(3), 147–196.

https://doi.org/10.1016/0301-0082(77)90016-8

[73] Ritchie, S. J., & Tucker-Drob, E. M. (2018). How Much Does Education Improve

Intelligence? A Meta-Analysis. *Psychological science*, *29*(8), 1358–1369.

https://doi.org/10.1177/0956797618774253

[74] Borgeest, G. S., Henson, R. N., Shafto, M., Samu, D., Cam-CAN, & Kievit, R. A.

(2020). Greater lifestyle engagement is associated with better age-adjusted cognitive

abilities. *PloS one*, *15*(5), e0230077. https://doi.org/10.1371/journal.pone.0230077

[75] Dounavi, M. E., Low, A., McKiernan, E. F., Mak, E., Muniz-Terrera, G., Ritchie, K.,

Ritchie, C. W., Su, L., & O'Brien, J. T. (2021). Evidence of cerebral hemodynamic

dysregulation in middle-aged APOE ε4 carriers: The PREVENT-Dementia study. *Journal of*

cerebral blood flow and metabolism : official journal of the International Society of Cerebral

Blood Flow and Metabolism, *41*(11), 2844–2855.

https://doi.org/10.1177/0271678X211020863

[76] Low, A., Su, L., Stefaniak, J. D., Mak, E., Dounavi, M. E., Muniz-Terrera, G., Ritchie,

K., Ritchie, C. W., Markus, H. S., & O'Brien, J. T. (2021). Inherited risk of dementia and the

progression of cerebral small vessel disease and inflammatory markers in cognitively healthy

midlife adults: the PREVENT-Dementia study. *Neurobiology of aging*, *98*, 124–133.

https://doi.org/10.1016/j.neurobiolaging.2020.10.029

[77] Kivipelto, M., Mangialasche, F., & Ngandu, T. (2018). Lifestyle interventions to prevent

cognitive impairment, dementia and Alzheimer disease. *Nature reviews. Neurology*, *14*(11),

653–666. https://doi.org/10.1038/s41582-018-0070-3

Figures and Legends

Table 1. Demographic specifications of the cohort at baseline and follow-up based on the dementia family history and on APOE genotype

	Baseline (n=210)			Follow-up (n=188)		
	FH- (n=107)	FH+ (n=103)	p (Mann-Whitney U)	FH- (n=89)	FH+ (n=99)	p (Mann-Whitney U)
Age (y)	52.0 ± 11.0	53.0 ± 7.0	0.54	54.0 ± 11.0	55.0 ± 6.0	0.22
Years of Education	16.0 ± 6.0	16.0 ± 5.0	0.34	17.0 ± 5.0	16.0 ± 5.0	0.21
CAIDE	5.0 ± 4.0	6.0 ± 2.0	0.03*	6.0 ± 4.0	7.0 ± 3.0	0.003**
			p (Chi-Square)			p (Chi-Square)
Sex (%f)	69.2%	71.8%	0.67	69.7%	71.7%	0.76
APOE e4 (% Carriers) [§]	28.0%	44.6%	0.01*	28.1%	44.3%	0.02*
	APOEe4- (n=133)	APOEe4+ (n=75)	p (Mann-Whitney U)	APOEe4- (n=118)	APOEe4+ (n=68)	p (Mann-Whitney U)
Age (y)	53.0 ± 8.0	52.0 ± 8.0	0.06	55.0 ± 8.0	54.5 ± 8.0	0.08
Years of Education	16.0 ± 5.0	17.0 ± 5.0	0.50	16.0 ± 5.0	17.0 ± 5.0	0.47
CAIDE (incl. APOE status)	5.0 ± 3.0	7.0 ± 3.0	0.0003***	6.0 ± 3.0	7.0 ± 4.0	0.0002***
CAIDE (excl. APOE status)	5.0 ± 2.0	4.0 ± 3.0	0.12	5.0 ± 2.0	4.5 ± 3.0	0.25
			p (Chi-Square)			p (Chi-Square)
Sex (%f)	69.9%	70.7%	0.91	71.2%	69.1%	0.77

Note: median ± interquartile range (IQR) was reported for continuous variables. Abbreviations: FH-, Negative family history of dementia; FH+, positive family history of dementia; APOE ε4 +, Apolipoprotein ε4 genotype positive: APOE ε4 -, Apolipoprotein ε4 genotype negative. [§] Information for the APOE ε4 genotype was not collected for 2 subjects.

Figure 1. Association of years of education with verbal, spatial and relational memory performance at baseline (a) and follow-up (b). The x axis displays total reported years of education attained. On the y axis, higher scores represent better verbal, spatial and relational memory performance.

Figure 2. Association of physical, social and intellectual activities with verbal, spatial and relational memory performance at baseline (a) and follow-up (b). Physical, social and intellectual activities are mean-centred. On the x axis, higher scores represent more frequent engagement in physical, social and intellectual lifestyle activities, and on the y axis, higher scores represent better verbal, spatial and relational memory performance.

Figure 3. Interaction of physical, social and intellectual activities and family history of dementia on verbal and visuospatial functions, and short-term (conjunctive) memory at follow-up. On the x axis, higher scores represent more frequent engagement in physical, social and intellectual lifestyle activities, and on the y axis, higher scores represent poorer verbal and visuospatial functions, and short-term (conjunctive) memory. Physical, social and intellectual activities are mean-centred. The positive family history of dementia group showed a significant positive association between engagement in these activities and improved verbal and visuospatial functions, and short-term (conjunctive) memory. No significant association was seen for the negative family history group. Abbreviations: FH, family history; FH+, family history positive; FH-, family history negative.