

TITLE: Comorbidities Diminish the Likelihood of Seropositivity After SARS-CoV-2 Vaccination

SHORT TITLE: COMORBIDITY & SARS-COV-2 SEROSTATUS

ARTICLE TYPE: Brief Report

AUTHORS:

Alok R. Amraotkar, MD, MPH¹ (Twitter Handle: @AlokAmraotkar)

Adrienne M. Bushau-Sprinkle, PhD²

Rachel J. Keith, PhD¹

Krystal T. Hamorsky, PhD²

Kenneth R. Palmer, PhD²

Hong Gao, PhD¹

Shesh N. Rai, PhD^{1,4-8}

Aruni Bhatnagar, PhD^{1,3}

AFFILIATIONS:

¹Division of Environmental Medicine, Christina Lee Brown Envirome Institute, University of Louisville School of Medicine, Louisville, KY

² Center for Predictive Medicine for Biodefense and Emerging Infectious Diseases, University of Louisville School of Medicine, University of Louisville, Louisville, KY, USA

³Center for Integrative Environmental Health Sciences, University of Louisville, Louisville, Kentucky

⁴Biostatistics and Bioinformatics Facility, Brown Cancer Centre, University of Louisville, Louisville, KY 40202, USA

⁵Biostatistics and Informatics Facility, Centre for Integrative Environmental Research Sciences, University of Louisville, Louisville, KY, 40202, USA

⁶Hepatobiology and Toxicology Center, University of Louisville, Louisville, KY 40202, USA

⁷School of Interdisciplinary and Graduate Studies, University of Louisville, Louisville, KY 40292, USA

⁸Department of Bioinformatics and Biostatistics, University of Louisville, Louisville, KY 40202, USA

COUNTS:

Abstract: (Unstructured) 100 words

Main Text: 1,372

of Tables: 2

of Figures: 1

AUTHOR CONTRIBUTIONS:

Conception and Design: A.R. Amraotkar, A. Bhatnagar, R. Keith

Collection, Analysis, or Interpretation of Data: A.R. Amraotkar, R. J. Keith, A. M. Bushau-Sprinkle, K. T. Hamorsky.

Statistical Analysis: Hong Gao and Shesh N. Rai

Drafting of Article: All Authors

Critical Revision for Important Intellectual Content: All authors

CORRESPONDING AUTHOR:

Aruni Bhatnagar, PhD, FAHA

Director, Christina Lee Brown Envirome Institute

Chief, Division of Environmental Medicine

University of Louisville School of Medicine

302 E. Muhammad Ali Blvd, Suite 107, Louisville KY 40202

Email: Aruni@Louisville.edu

ABSTRACT

Background: The impact of chronic health conditions (CHC) on serostatus post-severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) vaccination is unknown.

Methods: We assessed serostatus post-SARS-CoV-2 vaccination among fully vaccinated participants recruited between April 2021 through August 2021 in 18 years and older residents of Jefferson County, Kentucky, USA. Serostatus was determined by measuring SARS-CoV-2 Spike protein specific immunoglobulin (Ig) G (Spike IgG) antibodies via enzyme-linked immunoassay (ELISA) in peripheral blood samples.

Results: Of the 5,178 fully vaccinated participants, 51 were seronegative and 5,127 were seropositive. Chronic kidney disease (CKD) (OR=13.49; 95% CI: 4.88–37.3; P<0.0001) and autoimmune disease (OR=11.34; 95% CI: 5.21–24.69; P<0.0001) showed highest association with negative serostatus in fully vaccinated participants. The absence of any CHC was strongly associated with positive serostatus (OR=0.37; 95% CI: 0.19–0.73; P=0.003). The risk of negative serostatus increased in the presence of two CHCs (OR=2.82; 95% CI: 1.14–7) to three or more CHCs (OR=4.52; 95% CI: 1.68–12.14). Similarly, use of 2 or more CHC related medications was significantly associated with seronegative status (OR=6.08; 95%: 2.01–18.35).

Conclusions: Presence of any CHC, especially CKD or autoimmune disease, increased the likelihood of seronegative status among individuals who were fully vaccinated to SAR-CoV-2. This risk increased with a concurrent increase in number of comorbidities, especially with multiple medications. Absence of any CHC was protective and increased the likelihood of a positive serological response post-vaccination. These results will help develop appropriate guidelines for booster doses and targeted vaccination programs.

Keywords: antibody response; serology; serostatus; SARS-CoV-2; comorbidity

BACKGROUND

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and its mutated variants continue to increase morbidity and mortality worldwide.(1) Congruent with infection prevention and control measures, wide-spread vaccination is an effective means of containing and ending the Coronavirus Disease 2019 (COVID-19) pandemic.(2) However, in addition to the continued expansion of vaccination efforts, it is important to identify factors that impact vaccine efficacy. Effectiveness of the vaccines among non-immunocompromised individuals in the US is documented.(3) But, there are sparse data comparing vaccine efficacy between individuals with and without pre-existing comorbidities, and their relative effect on vaccination response. Gaining insights into factors which affect the immunological response to vaccines will guide the deployment of boosters worldwide and help identify individuals at greater risk of breakthrough infections. Therefore, we sought to study the effect of comorbidities on serostatus post-SARS-CoV-2 vaccination.

METHODS

Design and Study Population

Data for the study were collected under the Co-Immunity Project, which is a federally funded ongoing population-based study for the surveillance of SARS-CoV-2 in Jefferson County, Kentucky. The study was initiated in June 2020 and has completed 8 rounds of testing through October 2021. The SARS-CoV-2 vaccines were made widely available to the public in Kentucky starting April 2021. Data reported here are from April 2021 through August 2021. Participants were 18 years and older residents of Jefferson County who provided signed consent. This study and all the protocols were approved by the Institutional Review Board of the University of Louisville (IRB # 20.0393).

Vaccination Status

For analysis, only fully vaccinated participants were included. As per the Centers for Disease Control and Prevention (CDC), participants were considered “Fully Vaccinated” only if their final dose (2nd dose for Moderna or Pfizer-BioNTech and 1st dose for Janssen) was ≥ 14 days prior to the study appointment.(4) All participants in this study were within 9 months of their final vaccination dose.

Health History

Health history regarding pre-existing chronic health conditions and medications was self-reported by the study participants. History of lupus, rheumatoid arthritis, celiac disease, spondylitis, graves disease, etc. was combined into a single variable “Autoimmune Diseases”. Regular use of medications involved with immune-suppression like steroids, Secukinumab, Baricitinib, Tofacitinib, Methotrexate, etc. was combined into a single variable “Immunosuppressants”. History of all modalities of cancer treatments including chemotherapy, radiation therapy, surgery, immunotherapy, etc. were combined into a single variable “Cancer Treatment”. Use of angiotensin converting enzyme inhibitors (ACEi) or angiotensin receptor blocker (ARB) was combined into a single variable “ACEi or ARB”.

Human Samples and Serology

Trained staff collected nasopharyngeal swab and blood finger prick samples. Samples were analyzed for infection by reverse transcription polymerase chain reaction (RT-PCR).(5) Serostatus was determined by measuring levels of SARS-CoV-2 Spike protein specific immunoglobulin (Ig) G (Spike IgG) antibodies in peripheral blood samples as reported previously.(6) Serostatus was a qualitative assessment (positive or negative) based on the enzyme-linked immunoassay (ELISA) results from above.

Statistical Analysis

Two study groups were defined as: seronegative and seropositive. The primary objective was to compare the univariate relationship of chronic health conditions (CHC) between seronegative versus seropositive statuses. The secondary objective was to examine the adjusted

associations between clinical characteristics and seronegative status. Participant characteristics are expressed as mean \pm standard deviation for continuous variables, and frequency (%) for categorical variables. Relative magnitude of negative serostatus was estimated by odds ratio (OR) and 95% confidence intervals (CI).

For the secondary objective, multivariable logistic regression models were built to calculate the adjusted OR for negative serostatus in participants with CHCs or taking medications. Models were adjusted for age, sex, CHCs, and medications. P-value was considered significant at <0.05 . All statistical analyses were performed using SAS, version 9.4 (SAS Institute, Inc., Cary, North Carolina).

RESULTS

Of the 7,046 participants enrolled from April 2021 through August 2021, a total of 1,868 were excluded from the analysis because 802 were unvaccinated, 487 were missing vaccination dates or other vaccination related critical data, 247 were missing critical demographic information or medical history, and 332 were not “Fully Vaccinated” as defined in Methods (Figure 1). The final study dataset included 5,178 participants, of which, 51 were seronegative and 5,127 were seropositive. Univariate comparison of clinical characteristics between the two study groups is presented in Table 1.

Serostatus

Among those with no-CHC, 22% were seronegative, whereas 42% were seropositive (OR=0.37, 95% CI: 0.19-.073; P=0.003) (Table 1). Among chronic kidney disease (CKD), 12% were seronegative, and 2% were seropositive (OR=13.49, 95% CI: 4.88-37.3; P<0.0001). For autoimmune diseases, 31% were seronegative, whereas 5% were seropositive (OR=11.34, 95% CI: 5.21-24.69; P<0.0001). Similarly, diabetes (OR=3.27, 95% CI: 1.35-7.94; P=0.01),

hypertension (OR=2.9, 95% CI: 1.38-6.11; P=0.003), and cardiovascular disease (CVD) (OR=3.56, 95% CI: 1.37-9.23; P=0.013) had significantly higher rates of seronegative statuses.

The presence of any CHC was 78% in the serogroup, but 58% in the seropositive group (OR=2.68, 95% CI: 1.37-5.24; P=0.003). Similarly, the presence of CVD was 49% in seronegative and 36% in the seropositive group (OR=2.93, 95% CI: 1.4-6.11; P=0.003). History of immunosuppressants (OR=23, 95% CI: 12.1-43.25; P<0.0001) and cancer treatments (OR=12, 95% CI: 4.17-36.88; P=0.001) were also higher in the seronegative group. History of ACEi or ARB was not associated with serostatus (OR=0.82; 95% CI: 0.34-1.99; P=0.66).

Adjusted Associations Between Clinical Characteristics and Seronegative Status

Age \geq 65 years, sex, or reported history of tobacco product use were not significantly associated with seronegative status (Table 2). However, the presence of even one reported CHC was significantly associated with seronegative status (OR=2.69; 95%: 1.25-5.79). Similarly, the association between pre-existing CHCs and seronegative status strengthened with an increase in the number of CHCs from two (OR=2.82; 95%: 1.14-7.0) to three or more (OR=4.52; 95%: 1.68-12.14).

DISCUSSION

In this study of 5,178 fully vaccinated participants enrolled during the beginning of the Delta variant wave, we found that those reporting no CHCs were more likely to develop detectable antibody levels (a.k.a., seropositivity) after completing vaccination regimen. Conversely, those reporting a pre-existing comorbid condition were less likely to develop a seropositive status after immunization. The highest risk of not developing a detectable antibody response was seen with CKD, although those reporting CVD, diabetes, or hypertension were also less likely to develop seropositive status after vaccination. The odds of a seronegative status increased with an increase in the number of comorbidities.

More than 10.6 billion COVID-19 vaccine doses have been administered worldwide.(1) Although it is widely suspected that pre-existing comorbidities play a critical role in determining responses to immunization,(7) few data are available to determine the role of comorbidities on serological response or vaccine efficacy in the general public. We found that <1% of our study population failed to develop an appreciable response to the vaccine despite the high rates of diabetes (10.7%), hypertension (33.7%), and cancer (6.8%) in our cohort, which are comparable to national rates of these comorbidities. Even though serological tests do not represent the competence of an immune response, there is a significant relationship between a positive serology and SARS-CoV-2 neutralization potential,(6) suggesting that those who fail to establish a measurable seropositive status are more likely to remain vulnerable to SARS-CoV-2 infection. Furthermore, even though a seropositive status does not guarantee successful vaccination effect (i.e., SARS-CoV-2 neutralization), a seronegative status excludes any possibility of virus neutralization.(8)

About 52% adults in the United States have at least 1 CHC and about 27% have multiple CHCs.(9) According to the data presented in our study, almost half of the adult population in the United States has a two-fold chance of being vulnerable to SARS-CoV-2 (or re-infection), even if they are fully vaccinated. Although diabetes, hypertension, and CVD are well-known risk factors for severe COVID-19 outcomes,(10) our study shows that CKD is the biggest threat, even among fully vaccinated individuals as has been suspected previously.(11, 12) Our findings suggest that presence of these comorbidities significantly increases the likelihood of vaccine failure.

The Global Burden of Disease Study (GBD) has estimated one in five individuals worldwide to be at risk for severe COVID-19, primarily due to their burden of pre-existing CHCs.(11) There is a significant overlap in diseases reported by the GBD and those found to increase the likelihood of a seronegative status in our study.(13) Cancer, a major CHC in GBD, showed a very modest association with seronegative status in our study. However, current

treatment for cancer was strongly associated with seronegative status, indicating that treatment status might be a more accurate clinical indicator of impaired immune response rather than a clinical diagnosis of cancer. Some studies have shown a significant association between ACEi or ARB use and risk of SARS-CoV-2 infection and severity of health outcomes.(14, 15) In contrast to that, our study did not find any association, positive or negative, between ACEi or ARB use and serostatus. This suggests that use of ACEi or ARB may not affect vaccine efficacy, even if their use has the potential to influence risk of disease or its severity. Interestingly, our study

Additional work is required to elucidate whether smoking and other comorbidities influence the effectiveness of the virus neutralization capabilities post-vaccination. The results of such studies will help better shape the guidelines for booster doses and future vaccination programs.

Limitations

Limitations of the study include a lack of SARS-CoV-2 titer or neutralization data, which was not the primary objective of the study, but can evaluate a continuous relationship between comorbid conditions and antibody titer and neutralization efficacy. Serological assessment is not a definitive method to measure antibody efficacy, but seropositive status has a strong correlation with virus neutralization potential. The main goal of this study was to identify differences in serostatus among fully vaccinated individuals and was not powered to study the difference between individual vaccines. Finally, this study is cross-sectional in design, which does not allow us to report change over-time.

CONCLUSIONS

Among fully vaccinated individuals, presence of any CHC, especially CKD or autoimmune disease, increased the likelihood of seronegative status by multi-fold. This risk increased with a concurrent increase in number of comorbidities. CHCs requiring long-term treatment, especially

immunosuppressants or cancer treatments, increased the likelihood of a seronegative status. Absence of any CHC was protective and increased the likelihood of a positive serological response post-vaccination. These results will help develop appropriate guidelines for booster doses and targeted vaccination programs.

LEGENDS

Figure 1. CONSORT Flow Diagram of Dataset Identification

ACKNOWLEDEMENTS

We would like to thank all study participants who generously consented to participate in this study. We would also like to thank the City of Louisville Metro, Kentucky and various other collaborators and partners who helped us complete this important study.

INFORMED CONSENT STATEMENT

All the study participants provided signed consent. This study and all the protocols were approved by the Institutional Review Board of the University of Louisville (IRB # 20.0393), which conforms to the standards currently applied in the United States.

FUNDING

Coronavirus Aid, Relief, and Economic Security (CARES) Act – Recipient: Aruni Bhatnagar

Centers for Disease Control and Prevention (# 75D30121C10273) – Recipient: Aruni Bhatnagar

James Graham Brown Foundation – Recipient: Aruni Bhatnagar

Owsley Brown II Family Foundation – Recipient: Aruni Bhatnagar

DATA AVAILABILITY STATEMENT

Primary de-identified data supporting the findings of this study are available through the corresponding author (Aruni Bhatnagar: Aruni.bhatnagar@louisville.edu) upon request.

CONFLICT OF INTEREST

All authors have no financial, personal, or other conflicts of interests to disclose.

REFERENCES

1. World Health Organization (W.H.O). Who Coronavirus (Covid-19) Dashboard. (2021).
2. Centers for Disease Control and Prevention (C.D.C). Covid-19. , (2021).
3. Self WH, Tenforde MW, Rhoads JP, Gaglani M, Ginde AA, Douin DJ, et al. Comparative Effectiveness of Moderna, Pfizer-Biontech, and Janssen (Johnson & Johnson) Vaccines in Preventing Covid-19 Hospitalizations among Adults without Immunocompromising Conditions - United States, March-August 2021. *MMWR Morb Mortal Wkly Rep* (2021) 70(38):1337-43. Epub 2021/09/24. doi: 10.15585/mmwr.mm7038e1.
4. Centers for Disease Control and Prevention (C.D.C). Fully Vaccinated. (2021).
5. Centers for Disease Control and Prevention (C.D.C). Cdc Diagnostic Tests for Covid-19. (2020).
6. Hamorsky KT, Bushau-Sprinkle AM, Kitterman K, Corman JM, DeMarco J, Keith RJ, et al. Serological Assessment of Sars-Cov-2 Infection During the First Wave of the Pandemic in Louisville Kentucky. *Sci Rep* (2021) 11(1):18285. Epub 2021/09/16. doi: 10.1038/s41598-021-97423-z.
7. Kompaniyets L, Pennington AF, Goodman AB, Rosenblum HG, Belay B, Ko JY, et al. Underlying Medical Conditions and Severe Illness among 540,667 Adults Hospitalized with Covid-19, March 2020-March 2021. *Prev Chronic Dis* (2021) 18:E66. Epub 2021/07/02. doi: 10.5888/pcd18.210123.
8. Aziz NA, Corman VM, Echterhoff AKC, Muller MA, Richter A, Schmandke A, et al. Seroprevalence and Correlates of Sars-Cov-2 Neutralizing Antibodies from a Population-Based

Study in Bonn, Germany. *Nat Commun* (2021) 12(1):2117. Epub 2021/04/11. doi:

10.1038/s41467-021-22351-5.

9. Boersma P, Black LI, Ward BW. Prevalence of Multiple Chronic Conditions among Us Adults, 2018. *Prev Chronic Dis* (2020) 17:E106. Epub 2020/09/19. doi: 10.5888/pcd17.200130.

10. Li J, He X, Yuan Y, Zhang W, Li X, Zhang Y, et al. Meta-Analysis Investigating the Relationship between Clinical Features, Outcomes, and Severity of Severe Acute Respiratory Syndrome Coronavirus 2 (Sars-Cov-2) Pneumonia. *Am J Infect Control* (2021) 49(1):82-9. Epub 2020/06/17. doi: 10.1016/j.ajic.2020.06.008.

11. Clark A, Jit M, Warren-Gash C, Guthrie B, Wang HHX, Mercer SW, et al. Global, Regional, and National Estimates of the Population at Increased Risk of Severe Covid-19 Due to Underlying Health Conditions in 2020: A Modelling Study. *Lancet Glob Health* (2020) 8(8):e1003-e17. Epub 2020/06/20. doi: 10.1016/S2214-109X(20)30264-3.

12. Council E-E, Group EW. Chronic Kidney Disease Is a Key Risk Factor for Severe Covid-19: A Call to Action by the Era-Edta. *Nephrol Dial Transplant* (2021) 36(1):87-94. Epub 2020/12/20. doi: 10.1093/ndt/gfaa314.

13. Diseases GBD, Injuries C. Global Burden of 369 Diseases and Injuries in 204 Countries and Territories, 1990-2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *Lancet* (2020) 396(10258):1204-22. Epub 2020/10/19. doi: 10.1016/S0140-6736(20)30925-9.

14. Armstrong K, Soltoff A, Rieu-Werden M, Metlay J, Haas J. Use of Angiotensin Converting Enzyme Inhibitors and Angiotensin Receptor Blockers Associated with Lower Risk of Covid-19 in Household Contacts. *PLoS One* (2021) 16(3):e0247548. Epub 2021/03/03. doi: 10.1371/journal.pone.0247548.

15. Pan M, Vasbinder A, Anderson E, Catalan T, Shadid HR, Berlin H, et al. Angiotensin-Converting Enzyme Inhibitors, Angiotensin II Receptor Blockers, and Outcomes in Patients Hospitalized for Covid-19. *J Am Heart Assoc* (2021) 10(24):e023535. Epub 2021/12/11. doi: 10.1161/JAHA.121.023535.

TABLES

Table 1. Univariate Comparison of Clinical Characteristics Between Study Groups

Characteristics	Antibody Negative* (n = 51)	Antibody Positive* (n = 5,127)	Odds Ratio (95%CI)	P-Value
Age, Median ± IQR	69 ± 25	62 ± 23		0.024
Sex, Female N (%)	37 (72.6)	3,393 (66.2)		0.34
Race, White N (%)	48 (94.1)	4,457 (86.9)	2.41 (0.75 - 7.74)	0.13
Any Tobacco Use	2 (3.9)	201 (3.9)	1 (0.24 - 4.14)	1.00
Chronic Health Conditions (CHC)				
None, N (%)	11 (21.6)	2,176 (42.4)	0.37 (0.19 - 0.73)	0.003
Diabetes, N (%)	9 (17.7)	544 (10.6)	3.27 (1.35 - 7.94)	0.010
Hypertension, N (%)	23 (45.1)	1723 (33.6)	2.9 (1.38 - 6.11)	0.003
Heart Disease, N (%)	7 (13.7)	389 (7.6)	3.56 (1.37 - 9.23)	0.013
Autoimmune Disease, N (%)	16 (31.4)	279 (5.4)	11.34 (5.21 - 24.69)	<.0001
Cancer, N (%)	5 (9.8)	347 (6.8)	2.85 (0.98 - 8.25)	0.06
Thyroid Disease, N (%)	6 (11.8)	547 (10.7)	2.17 (0.8 - 5.89)	0.13
Chronic Kidney Disease, N (%)	6 (11.8)	88 (1.7)	13.49 (4.88 - 37.3)	<.0001
Composite CHC				
Cardiovascular Disease, N (%)	25 (49.0)	1,856 (36.2)	2.93 (1.4 - 6.11)	0.003
Any CHC, N (%)	40 (78.4)	2,951 (57.6)	2.68 (1.37 - 5.24)	0.003
Medications				
None, N (%)	27 (52.94)	3,852 (75.1)	0.37 (0.21 - 0.65)	0.0003
All Medications, N (%)	24 (47.06)	1,275 (24.9)	2.69 (1.54 - 4.67)	0.0003
ACEI or ARB, N (%)	6 (11.76)	1,041 (20.3)	0.82 (0.34 - 1.99)	0.66
Immunosuppressants, N (%)	17 (33.33)	106 (2.1)	22.88 (12.1 - 43.25)	<.0001
Cancer Treatments, N (%)	4 (7.84)	46 (0.9)	12.41 (4.17 - 36.88)	0.001

Abbreviations: ACEI – Angiotensin Converting Enzyme Inhibitor; ARB – Angiotensin Receptor Blocker; CHC – Chronic Health Conditions; IQR – Interquartile Range

*Serostatus Determined by Presence/Absence of SARS-CoV-2 Spike Protein (IgG) Antibodies in peripheral blood via ELISA

Table 2. Multivariable Association Between Clinical Characteristics and SARS-CoV-2 Spike IgG Antibody Negative Status

Characteristics	Referent Group	Odds Ratio	95%CI
Age \geq 65 years	< 65 Years	1.13	0.64 – 1.98
Sex, Female (%)	Male	1.4	0.77 – 2.54
Composite Chronic Health Conditions (CHC)			
1 CHC	No CHC	2.69	1.25 – 5.79
2 CHCs	No CHC	2.82	1.14 – 7.0
\geq 3 CHCs	No CHC	4.52	1.68 – 12.14
Composite Medications			
1 medication	No Medications	1.43	0.76 – 2.71
\geq 2 medications	No Medications	6.08	2.01 – 18.35

Abbreviations: CHC – Chronic Health Conditions; CI – Confidence Interval

Figure 1. CONSORT Flow Diagram of Dataset Identification



