Modelling the impact of non-pharmaceutical interventions on workplace transmission of SARS-CoV-2 in the home-delivery sector

Carl A. Whitfield[a],[1,2,3] Martie van Tongeren,[4,3] Yang Han,[1] Hua Wei,[4,3] Sarah Daniels,[4,3] Martyn Regan,[5,4,3] David W. Denning,[2,3] Arpana Verma,[4,3] Lorenzo Pellis,[1] University of Manchester COVID-19 Modelling Group[b],[1] and Ian Hall[1,6,3]

[1]*Department of Mathematics, University of Manchester, Manchester, England*
[2]*Division of Infection, Immunity & Respiratory Medicine,*
School of Biological Sciences, University of Manchester, Manchester, England
[3]*Manchester Academic Health Science Centre,*
University of Manchester, Manchester, England
[4]*Division of Population Health, Health Services Research & Primary Care,*
School of Health Sciences, University of Manchester, Manchester, England
[5]*National COVID-19 Response Centre,*
UK Health Security Agency, London, England
[6]*Public Health Advice, Guidance and Expertise,*
UK Health Security Agency, London, England

---

[a] Corresponding author: carl.whitfield@manchester.ac.uk

Abstract

Objective: We aimed to use mathematical models of SARS-COV-2 to assess the potential efficacy of non-pharmaceutical interventions on transmission in the parcel delivery and logistics sector.

Methods: We developed a network-based model of workplace contacts based on data and consultations from companies in the parcel delivery and logistics sectors. We used these in stochastic simulations of disease transmission to predict the probability of workplace outbreaks in this settings. Individuals in the model have different viral load trajectories based on SARS-CoV-2 in-host dynamics, which couple to their infectiousness and test positive probability over time, in order to determine the impact of testing and isolation measures.

Results: The baseline model (without any interventions) showed different workplace infection rates for staff in different job roles. Based on our assumptions of contact patterns in the parcel delivery work setting we found that when a delivery driver was the index case, on average they infect only 0.14 other employees , while for warehouse and office workers this went up to 0.65 and 2.24 respectively. In the large-item delivery setting this was predicted to be 1.40, 0.98, and 1.34 respectively. Nonetheless, the vast majority of simulations resulted in 0 secondary cases among customers (even without contact-free delivery). Our results showed that a combination of social distancing, office staff working from home, and fixed driver pairings (all interventions carried out by the companies we consulted) reduce the risk of workplace outbreaks by 3-4 times.

Conclusion: This work suggests that, without interventions, significant transmission could have occured in these workplaces, but that these posed minimal risk to customers. We found that identifying and isolating regular close-contacts of infectious individuals (i.e. house-share, carpools, or delivery pairs) is an efficient measure for stopping workplace outbreaks. Regular testing can make these isolation measures even more effective but also increases the number of staff isolating at one time. It is therefore more efficient to use these isolation measures in addition to social distancing and contact reduction interventions, rather than instead of, as these reduce both transmission and the number of people needing to isolate at one time.

IMPORTANCE

During the COVID-19 pandemic the home-delivery sector was vital to maintaining people's access to certain goods, and sustaining levels of economic activity for a variety of businesses. However, this important work necessarily involved contact with a large number of customers as well as colleagues. This means that questions have often been raised about

whether enough was being done to keep customers and staff safe. Estimating the potential risk to customers and staff is complex, but here we tackle this problem by building a model of workplace and customer contacts, from which we simulate SARS-CoV-2 transmission. By involving industry representatives in the development of this model, we have simulated interventions that have either been applied or considered, and so the findings of this study are relevant to decisions made in that sector. Furthermore, we can learn generic lessons from this specific case study which apply to many types of shared workplace as well as highlighting implications of the highly stochastic nature of disease transmission in small populations.

I. INTRODUCTION

Demand for home-delivery services spiked globally during the COVID-19 pandemic, as people stayed at home to reduce transmission [1]. In the UK, non-essential retail shops were closed for much of 2020 and 2021, increasing the demand for online retail and home delivery. Additionally, stay-at-home orders brought new demand for large items such as furniture and white goods as many people adjusted to spending more time at home [2]. This new and displaced demand has, on the whole, been successfully absorbed and managed by the delivery and logistics sector, due in no small part to the efforts of the key workers in those sectors to keep business moving, while adapting to a changing work environment. Meanwhile, key workers in all sectors were disproportionately exposed to transmission of SARS-CoV-2 [3]. In the delivery sector, drivers and warehouse workers were also at risk, given their exposure to a large number of contacts, the likelihood of asymptomatic transmission in SARS-CoV-2, and the potential economic impact of absence due to the prevalence of flexible or zero-hours contracts in this sector. Furthermore, studies from other countries indicate that delivery drivers there could be at much greater risk [4] than the general population, and so is a sector that requires greater attention.

Mathematical models have been central to understanding transmission of SARS-CoV-2 and in predicting the impact of various interventions. As more data has become available, models have been developed for a number of specific settings, including schools, hospitals, prisons and workplaces [5–8], to take into account the nuances and unique features of each setting. In this paper we present a model of delivery sector that has been uses to assess the impact of various measures that some companies have taken, as well as measures that were under consideration. One unique feature of these settings is the high number of brief contacts that delivery drivers have with members of the public, who themselves may otherwise have

very limited contacts. Another feature in the delivery of heavy or large items is the safety requirement for employees to handle and deliver goods in pairs, often requiring prolonged close contact and entry into customers' properties. Finally, there is still the poorly understood route of fomite transmission that has the potential to be important in this setting, due to the large volume of packages being handled. The model we present considers all of these aspects, and where data is unavailable or uncertain (e.g. for risk of fomite transmission), we consider a wide range of possible scenarios.

We have developed an agent-based network model with stochastic transmission. Therefore, each realisation of the simulation represents a possible chain of transmission within a workplace, and so conclusions can only be drawn from the aggregated results of many simulations. There are commonalities with several models in the literature, including the network models for COVID-19 transmission in workplaces [8]. The stochastic infection and isolation model is similar to other agent-based and branching process models [7, 9]. The model was developed based on a combination of epidemiological data and data and qualitative information gained from consultations with companies in the logistics sector in the UK.

With global rollout of SARS-CoV-2 vaccines, the most severe impacts of COVID-19 on public health to be curtailed and so have most of the restrictions and measures in place to reduce transmission. However, containing the spread of new variants is likely to require good surveillance testing. There has been considerable debate around the usefulness of Lateral Flow Device (LFD) tests that can be self-administered and give rapid results [10–12]. Primarily, this centres around the lower sensitivity of LFD against Polymerase Chain Reaction testing (PCR), particularly at low viral loads [13], and the potential impact of false positives. However, recent data suggests that LFD specificity may be at least 99.9% [14], suggesting that false positives will have a negligible impact. Furthermore, culturable SARS-CoV-2 virus is only found, at most, in the first 8-10 days following symptom onset [15–17], when viral load is higher. This suggests that lower sensitivity tests may still be useful at detecting people when they are most infectious. However, the way tests are performed (e.g. self-administered vs. trained tester) can have an impact on sensitivity [18], plus the method of rollout (e.g. supervised vs. unsupervised testing) can affect the adherence to the testing policy. The model we present accounts for these various factors.

The aim of this paper is to estimate the efficacy of different workplace interventions with a model particularly tailored to the home-delivery sector. We considered several interventions and scenarios based on formal consultations with company representative from this sector.

A secondary aim is to estimate the potential impact of presenteeism (working while sick) with COVID-19 symptoms. Flexible or 'gig' contracts are common in the home delivery sector, as well as the use of self-employed couriers, all of which are factors associated with increased presenteeism [19], so this is an important factor to consider.

II. METHODS

A. Data Collection and Company Consultations

We carried out recorded consultations via teleconference with representatives from six companies between July and August 2020 (Round 1), and May and June 2021 (Round 2), three of these companies were interviewed in both rounds. Each semi-structured interview lasted 60-90 mins and was based around a set of open-ended questions regarding how the pandemic had impacted on the operations of the business and what measures had been put in place to protect staff and customers. As part of these consultations we asked questions regarding the number of staff working at typical sites and the frequency of contacts between employees and the public. Additionally, two companies provided data on staff numbers and deliveries, which are detailed in Appendix A. A summary report was sent to each company for comments and corrections to verify that we had interpreted their answers accurately, and the data correctly. Further details on the consultations are published in [2].

We also used data from an online contact survey aimed at delivery drivers in the UK [20], which received 170 responses (104 of which were from the workers involved in the delivery of small packages and/or large items). This survey was elective so was not statistically representative. The results of this survey are to be published elsewhere but a few results are utilised in this paper. Namely, only 5.3% reported working while having symptoms of COVID-19 or with a member of their household having a suspected or confirmed case of COVID-19. Conversely, 17.2% reported having isolated with symptoms of COVID-19 or due to a member of their household having a suspected or confirmed case of COVID-19. This suggests approximately 1 in 4 failing to isolate for one of these reasons. For this reason we consider two p_{isol} values (0.5 and 0.9) as 'low' and 'high' isolation rates, noting the likely caveat of reporting biases. Staff reported large numbers of daily contacts (mean 15.0) at their place of work, which, tallying with the results of consultations, we interpreted as a result of repeated interaction within a work cohort (with only rare random interactions on top). Hence our assumed cohort size for drivers of ≈ 13.

B. Workplace Network Model

In this section we present an overview of the model details, with further details supplied in appendix B. The model we use is a stochastic agent-based network model of disease transmission. The parameters and symbols used in the following section are all described in table I.

TABLE I: Model parameters for workplace contacts and transmission. The values given are the values used unless otherwise stated for a given figure or section. The "perceived uncertainty" is simply to indicate the level of confidence we have in the parameter values – Low: based on primary data or peer-reviewed sources; Moderate: based on literature reviews, surveys, or specific consultation questions; High: assumed or extrapolated from consultation answers.

Parameter	Description	Value	Source	Perceived uncertainty
$N_D, N_L,$ N_O	Total number of drivers, pickers, and office staff employed in the workplace respectively.	Parcel: $\{50, 25, 15\}$ Large-item: $\{20, 10, 5\}$	Company data and consultation	Low
T_D, T_L, T_O	Total number of driver, picker, and office staff cohorts/teams.	Parcel: $\{3, 2, 1\}$ Large-item: $\{2, 2, 1\}$	Consulations and survey	Moderate
$n_D(t),$ $n_L(t),$ $n_O(t)$	Number of drivers, pickers, and office staff working on day t respectively.	Variable	Company data and consultation	Low
$D_P(t)$	Total number of packages delivered on day t.	Variable	Company data	Low
p_c	Probability of two individuals at work having a random F2F contact in a given day.	$2/(N_D+N_O+N_L)$	Consultation and survey	High

ρ_D	For contacts including drivers, p_c is scaled by this factor.	0.05	Consultation	Moderate
f_c	Cohort flux rate. The probability each day of a worker switching to a different cohort.	0.01 per day	Consultation	High
β_{F2F}	Infection rate for F2F contact at 1m distance while speaking (with a person with unit infectiousness)	0.15 h^{-1}	A plausible range of 0.03–0.24 was inferred from [21–23]	Moderate
c_i	Modifier for exposure due to type of contact i	1 (inside, talking) ×0.2 (outside) ×0.2 (not talking)	[21]	Moderate
β_{SS}	Infection rate via room-sharing (with a person with unit infectiousness)	0.002 h^{-1}	See appendix A.4	High
x_{ss}	Effective distance for room-sharing interaction	4.3m (shared spaces), 3.6m (office)	see appendix A.4	Moderate
β_{FOM}	Package-mediated fomite infection rate (from a person with unit infectiousness) if time between handling is 0.	0.001 per contact	Assumed	Very high
λ	Half-life of virus deposited on packages.	3 h^{-1}	[24]	Low
τ_{office}	Time office staff spend in shared office each day.	6 h	Consultation	Low

τ_{break}	Time office and picker staff spend in shared break rooms	1 h	Consultation	Moderate
H	Average number of employees per employee household minus 1 ($H = 0$ means no employees live together, $H = 1$ means the average household has two employees)	0.05, 0.5	Assumed	High
C	Number of cars/shared commutes per households minus 1.	0.05, 0.5	Assumed	High
$J_k[V_k(t - t_k)]$	Relative infectiousness of person k with viral load V_k infected at time t_k.	See appendix A.3 and [25]	[26]	Moderate
$S_k(t - t_k)$	Relative susceptibility of person k infected at time t_k.	$S_k(t - t_k < 0) = 1$ $S_k(t - t_k \geq 0) = 0$	Basic SIR model assumption	Low
p_{symp}	Probability that an individual develops symptoms relevant for self-isolation guidance.	0.5	[3]	Moderate (is age, variant, and guidance dependent).
p_{isol}	Probability that an individual adheres to self-isolation guidance.	0.5, 0.9	[19] and Survey	Moderate.
p_{miss}	Probability that an 'adherent' person misses a test.	0.4	Based on adherence rates in the UK public sector.	Moderate.

The model considers contacts between all employees working in a home delivery depot (i.e. engaged in business-to-consumer delivery or B2C) that has a warehouse and onsite offices. The workplace is populated by 3 groups: drivers, who deliver packages from the warehouse to customers; pickers, who transport and load packages within the warehouse; and office/admin staff, who work in the same building but in shared offices. There exists a pool of N_D drivers , N_L pickers and N_O office staff available for work each day. Workforce turnover is ignored, as it is assumed negligible over the time scales considered, however it may play a role over long time periods.

1. Employee work schedules

The model network consists of all within-workplace contacts between employees, as well as contacts between employees due to house-sharing or carpooling, in order to simulate workplace outbreaks in detail. Thus we assume, unless they share a household, employees only make contact with other employees if they are both at work on that day. We use an idealised model for the work schedule, whereby the number of employees in work depends on the day of the week, this pattern was calculated from data from two UK logistics firms (see Appendix A for details). For all pickers and drivers, we randomly assign consignments (i.e. deliveries/packages) for loading and delivery (as detailed in Appendix A). We assume that each consignment is first handled by pickers, then subsequently by drivers, and finally by the customer. Drivers are the only group of employees that have direct contact with members of the public while on shift. For simulation efficiency, repeat interactions with customers are not considered (as contacts via this route have a very low probability of infection, so double counting of infections is very unlikely), but these contacts are simulated and infection ingress/egress through this route is included in the model.

We also consider the case where drivers and pickers work in pairs (i.e. large goods delivery), we round the number of staff required in these roles to the nearest even number, and then assign pairings randomly each day. One intervention simulated is fixed pairings; in this case, these are assigned a priori and we pick the pairs working on a given day at random from those available. A pair is unavailable if either worker in that pair is isolating, therefore this intervention is always used alongside "pair isolation", where one member of the fixed pair isolates for the same period as their partner (whether or not they develop symptoms).

2. *Workplace contacts and infections*

Infections are modelled to occur via four routes; face-to-face (F2F) contact with infectious individuals, indirect contact via sharing a space with infectious individuals, household sharing, and fomite transmission via goods handling.

The model generates direct F2F contacts between employees through three different mechanisms, summarised in table II. Table II also lists the parameters for the different contact routes simulated. Contacts made via these routes are assumed to be dominated by face-to-face transmission.

Indirect aerosol-mediated transmission is taken to occur on a one-to-all basis. Given the well-ventilated nature of warehouses, we assume that this kind of transmission only occurs in offices, or in lunch/break rooms. Finally, fomite transmission via package handling is simulated as a decaying random process, such that the probability of onward transmission depends on the time between package handling events by infectious and susceptible individuals. See appendix A.4 for the justification of the various transmission parameters used.

Contact-type	Description	Transmission modifier (c_i)	Duration (τ_i)	Distance
Cohort	F2F contacts that occur within a team or cohort (see appendix A.5 for more information).	0.4 (either indoor or "loud-talking" outdoors, 25% of time)	Drivers: $\tau_{\mathrm{coh}} = 15$min Pickers/Office: $\tau_{\mathrm{coh}} = 1$h	1m
Random	F2F contacts that occur randomly in the workplace with weighted probability towards contacts between same job roles (see appendix A.5 for more information).	0.4 (25% of time talking)	$\tau_{\mathrm{rand}} = 15$ min	1m
Large-item handling	Time spent lifting and moving packages in pairs (see section appendix A.5 for more information).	0.08 (outdoor, 25% of time talking)	$\tau_{\mathrm{han}} = 5$ min per delivery	1m

Pair delivery	Contact via sharing a cabin while delivering large-items (see appendix A.5 for more information).	0.4 (window closed) 0.08 (window open)	$\tau_{cab} = 10$ min per delivery	1m
Pair dropoff	Contact between driver pairs during dropping large-item off at customer's property (see appendix A.5 for more information).	0.4 (25% of time talking)	$\tau_{drop} = 5$ min per delivery	1m
Customer	Contact between driver(s) and customer during item delivery	Parcel: 0.08 (outside, talking 25% of time) Large-item: 0.24 (Inside 50% of time, talking 25% of time)	Parcel: 30s per delivery Large-item: 5min per delivery	1m
Room-share	Aerosol-mediated contact in poorly ventilated rooms	0.2 (no talking)	$\tau_{off} = 6$h $\tau_{lun} = 1$h	$\sim$ 6m (see Appendix A.4 based on [27–29])
Car-share	Contact via carpooling to and from work.	0.4 (25% of time talking)	0.5h	1m
House-share	Contact via shared accommodation.	1	0.5h	1m (see appendix A.4 based on [21, 30–32]))

TABLE II: Summary of the direct contact routes simulated and the associated transmission rate modifier, duration of contact, and contact distance. These are the values used in all simulations in the main text unless explicitly stated otherwise. Note that this table does not include fomite transmission routes, which are simulated and are described in detail appendix A.4.

The transmission routes between different groups are illustrated in fig 1.

FIG. 1. Sketch of workplace staff groups and the potential transmission routes between them. Blue lines indicate face-to-face contacts, with dashed lines indicating transmission routes with either a lower contact rate or less contact time. Orange arrows are fomite transmission routes (via packages) and green indicates aerosol transmission in shared rooms. Arrows indicate direction of transmission.

C. Individual characteristics: viral load, infectiousness, and test positivity

Viral load trajectories are generated from the individual level data in [26]. The algorithm to generate individual viral load and infectiousness profiles is described in further detail in appendix B and in [33] is available at [34]. The method is detailed in [25] and summarised in appendix A.3.

D. Simulation algorithm

We simulate two types of scenario:

- **Point-source outbreak:** A single index case is chosen and we assume that there are no other introductions during the simulation. All other employees are susceptible at the simulation start (i.e. zero prevalence). The simulation terminates when there are no infectious cases remaining. This type of scenario is modelled in section III A and in appendix C.

- **Continuous-source outbreak:** No index cases are chosen initially and introductions occur randomly (Poisson process) based on the community incidence and prevalence in March-June 2020. The simulation runs for a fixed time window, and the number of customer contacts and packages delivered follow the pattern of demand experienced during that period of time (see appendix A). This scenario is modelled in section III B 1.

In the point-source outbreak scenarios, in order to define a 'successful' outbreak we arbitrarily set a threshold of a final attack rate of 5%. Note that we choose this as it is a low-threshold, like the epidemiological definition of an outbreak as a single linked secondary case. However, by defining it as a percentage of workplace size this makes the results from the two different settings more comparable. Therefore, if $R - 1 > 0.05(N_D + N_L + N_O)$, where R is the number of recovered individuals at the end of the simulation, then we record this simulation as a successful outbreak. The fraction of simulations where a successful outbreak occurs is then used as an estimate of the probability of an index case resulting in an outbreak.

For continuous-source outbreaks, there is random ingress of new cases, so instead we compare the number of workplace infections (ignoring introductions) as well as the number of isolation days to measure impacts on productivity. Introductions can occur in these simulations through two routes:

- **Community ingress:** Each susceptible individual in the workplace has probability $I(t)$ of being infected outside of work, where $I(t)$ is the community incidence at time t.

- **Customer ingress:** For each delivery a driver makes, there is probability $P(t)$ that the customer is currently infectious, where $P(t)$ is the community prevalence at time t. When a susceptible driver interacts with an infectious customer, there is probability $p_{\text{cust}} = 1 - \exp(-c_i \beta_{F2F} \tau_{\text{doorstep}})$ of an infection.

This is a very simple model of case ingress and does not account for household structure, the geographical/individual variability in the wider population, or repeat deliveries to customers.

The testing strategy we model here is non-directed mass testing, i.e. all employees are tested regularly every τ_p days. A random day in the period $[1,\tau_p]$ is drawn as the first test day, and all subsequent test days follow sequentially τ_p days after the previous. Following a positive test, an individual cannot be tested again for τ_{pause} days after their positive test. Other testing strategies may be beneficial, particularly if looking to reduce the burden on employees or because of affordability, and we address some of these in the discussion.

The simulation follows an SIR-type structure, such that individuals who have previously been infected cannot be re-infected. This is a reasonable assumption over the timescales of up to 3 months that we consider here. An example visualisation of a single realisation of the simulation is shown in figure 2. The source code for the simulations can be found at [35].

III.   RESULTS

Appendix C contains results pertaining to the baseline transmission dynamics in the two work settings that we focus on. This includes an investigation of changing parameters that affect the mixing rates in the settings, including changing work cohort size and close-contact work in pairs. This shows that, in the parcel workplace, we predict that office size and occupancy is a more important potential factor in workplace outbreaks than transmission between drivers at the workplace, even though office workers are in the minority (appendix C.1). In the large-item delivery workplace, close-contact working pairs (primarily delivery pairs, who share a vehicle for much of the day) were predicted to be very important, and keeping these pairs fixed had a significant impact on reducing workplace spread (appendix C.2). Finally, we also present the effect of presenteeism, which in this model we define as workers with symptomatic COVID-19 attending work, which we find can have a notable effect

FIG. 2. Example outbreak in a parcel delivery workplace, where the simulation terminates when no infectious cases remain. (a) The evolution of the number of recovered, infectious and quarantined (isolated) people in the model on each day (dashed and dotted lines indicate the same quantities for each subgroup as labelled). (b) Example network of the "cohort" contacts, each cohort has edges between all member nodes, additionally each driver cohort (blue D nodes) is supervised by a member of staff from the warehouse (red P nodes). Office staff are disconnected (green O nodes), but make contact through random interactions, break rooms, and house/car sharing arrangements.

on transmission, particularly when coupled with other measures to isolate close-contacts of symptomatic individuals (appendix C.3).

Furthermore, appendix D contains a sensitivity analysis of some of the pertinent baseline model parameters, to show how these affect predicted transmission rates.

In this following section, we focus on the impacts of testing and the combination of different workplace interventions to analyse their potential effectiveness.

A. Impact of mass testing on point-source outbreaks

Given the long incubation period of COVID-19 (compared to flu) and the significant proportion of asymptomatic cases, regular mass testing has been proposed and deployed in various settings to screen asymptomatic and pre-symptomatic cases. Figures 3 and 4 show the reduction in outbreak probability resulting from testing at different frequencies with different test types in the parcel and large-item delivery work settings respectively. Overall,

the results show that LFD tests have a similar effect to PCR with a 2-day turnaround (given that lab turnaround targets were 24h in the UK, a 2-day turnaround was typical for much of the pandemic). Therefore, considering LFDs relative low cost, this suggests that they are a better option for mass testing [36]. Note this estimate for the sensitivity of LFD tests is based on estimates sensitivity in phase 3b testing in [18] adjusted for the relative error induced by self vs. trained swabbing (see [25] for further details).

In each figure, two cases, representing idealised behaviours, are shown. In the first case testing is voluntary meaning 90% of people do 60% of the required tests on average (missing tests at random), while the other 10% do no tests (3(a) and 4(a)). This therefore reduces the potential benefits of testing. In the second case testing is enforced(3(b) and 4(b)) meaning that all workers test and report their results. This is the theoretical maximum effect that we could expect testing to have.

FIG. 3. Probability of an outbreak in the model parcel delivery workplace from a single introduction (selected at random), where the black line is the mean baseline case (no testing, estimated from 10,000 simulations). Each coloured marker shows the mean result of 10,000 simulations with the labelled testing intervention. In (a) testing is not enforced so $p_{\mathrm{miss}} = 0.4$ and and in (b) it is so $p_{\mathrm{miss}} = 0.0$ and all people isolate with a positive test. In both cases we use $p_{\mathrm{isol}} = 0.9$ for symptomatic isolation.

Comparing figures 3 and 4 shows that testing has a similar proportional impact in the

FIG. 4. The same as figure 3 but for the model large-item delivery workplace with fixed pairings and pair isolation in place.

large-item delivery delivery setting. However, in the large-item delivery case testing has a more noticeable effect even when performed as infrequently as 14 days. With total compliance to testing, the probability of outbreaks in both workplaces is reduced by approximately 75%, by twice weekly LFD tests (which have been deployed in other sectors). Note that this intervention is not as effective in the large-item delivery workplace when fixed-pairing and pair isolation policies are not in place (a reduction of approx. xx% under the same circumstances, data not shown). Therefore, targeted isolation policies can improve the efficacy of testing.

To conclude, we have found that regular testing, particularly in combination with close-contact isolation, can have a very significant effect on workplace transmission. Any testing intervention needs to be weighed against potential costs, at low community prevalence the vast majority of tests are likely to be negative, and those that are positive are more likely to be false positives and so the intervention may not represent good value for money. Alternatively, at high community prevalence, testing and close-contact isolation could result in many isolations, some of which are only precautionary, which can have a huge impact on business. The latter case is not well described by the point-source outbreak considered in this section, as introductions into the workplace are more likely to occur in quick succession. Therefore in the following section we look at the impacts of a range of interventions in the

case of a continuous-source outbreak.

B.   Impact of testing in the presence of household transmission

There are a number of confounding factors in reality that mean testing interventions may not be as effective as outlined in the previous section. One of these is the potential for household transmission between co-workers who share accommodation. In the previous sections we have considered 5% of worker households in the model to be shared, which is a significant fraction but not enough to have a large effect on transmission dynamics. It was suggested in consultations that it is likely that this will vary widely by workplace location and recruitment. Therefore, in this section we test what effects changing this fraction has on these predictions.

FIG. 5.   Mean number of secondary cases resulting from a single random introduction plotted against the house share factor H. Different colour lines show the different intervention scenarios as labelled where "Testing" means LFD testing every 3 days (with default adherence rates) and "HS isol." means that the household isolation policy was implemented. (a) and (b) show the results for the two workplace types, as labelled. Each point plotted shows the mean of 10,000 simulations, with shaded error region estimated using a bootstrapping process [37]

Figure 5 shows that increasing the household sharing factor H from 0.05 to 2.0 increases transmission but the relative effect of testing (regular LFD testing every 3 days) remains approximately unchanged. Interestingly, a household isolation policy (i.e. the whole household

isolates if one member isolates due to symptoms or a positive test) only has a minor effect for $H < 0.5$ and this is because we assume that a household transmission event between two cohabiting employees can still occur even if both are isolating (and this still contributes to the total number of infections). In reality it is likely that this risk of household transmission may reduce during the shared isolation period if the cohabiting employees are able to remain physically separate, however it is also possible the risk will increase as they would spend more time in the shared accommodation during isolation. At very high rates of house sharing ($H > 1$ i.e. 2 or more employees in the average household), household isolation has a larger impact as this mode of transmission is prominent enough to dominate the workplace chains of transmission and household isolation can break a significant fraction of those transmission chains.

To conclude, we see that in cases where there is high rates of household-sharing (or, more generally, any contacts between employees outside of work during isolation) then this can continue to drive transmission between employees and is difficult to distinguish from workplace transmission. Nonetheless, for all values simulated, regular mass asymptomatic testing has a sizeable effect on transmission rates.

1. *Impact of interventions in a real-world context*

In this section we model each workplace in the context of realistic community SARS-CoV-2 incidence rates. We used incidence rates inferred from deaths and hospitalisations in the UK during the period 1st March 2020 until 31st May 2020 (see appendix A.2). We then applied ran simulations with different interventions in place, for each scenario we added an extra intervention to the ones applied before, the interventions are:

1. Symptom isolation only: People who develop symptoms self-isolate with probability $p_{\mathrm{isol}} = 0.5$.

2. Improved isolation: To mimic the impact of pandemic messaging, isolation probability is increased to $p_{\mathrm{isol}} = 0.9$.

3. Distancing: All F2F interactions, except those involved in pair work, have interaction distance $x = 2$m.

4. Cohort Size Reduction: In the parcel delivery setting, the number of cohorts for all job types is doubled.

5. House share isolation: All employees who share a household isolate when one self-isolates.

6. Fixed-pairings: In the large-items delivery setting, driver and picker pairs are fixed and both self-isolate if one self-isolates.

7. Office WFH: Office staff do not enter work, they only make contact other employees if they share a household.

8. Testing: Twice weekly lateral flow testing is introduced for all employees.

9. Enforced testing: Testing becomes mandatory so no tests are missed.

10. Car share isolation: If a person travels to work with someone who self-isolates, they self-isolate.

11. Cohort isolation: If one member of the cohort isolates, all people in the cohort isolate.

In the model, introductions due to customer interactions only had a small but noticeable effect meaning that drivers were slightly more exposed than other employees (mean 0.11 introductions per driver for both work settings, averaged over all scenarios vs. 0.09 for other staff respectively). Nonetheless, over the period, around 10% of the workforce is infected purely due to the imposed prevalence and incidence.

Figure 6 shows the cumulative impact of interventions on secondary cases and isolations in the parcel delivery workplace. The interventions are applied in approximately the sequence that was reported by companies we consulted. The intervention "Distancing" increases all "cohort" and "random" contacts to 2m interactions and has a large effect. Reducing cohort size and office staff working from home ("Office WFH") have a big impact on reducing transmission since this model predicts that outbreaks are most likely to start in this group. Interventions beyond "enforced testing" are predicted to increase isolation levels without much greater impact on transmission, particularly "cohort isolation" which likely causes a great deal of disruption despite these groups being unlikely to be infected. Note this becomes a much more viable option though if cohorts are much smaller, which is one major benefit of reducing cohort size if possible. Comparing the two graphs in 6 we see that is a slightly more efficient to have contact reduction measures in place before adding isolation-based measures, as these reduce the number of workers who will need to isolate. When isolation measures are implemented alone, we see an increase in the predicted number of isolations even though the relative reduction in transmission is similar.

FIG. 6. Boxen plots [37] of the number of secondary cases divided by the number of introductions in a parcel delivery workplace over a 3 month period. Each distribution shows all the simulations (from 10000) with more than one introduction. The labels on the x-axis indicate the addition of an intervention (in-tandem with all the interventions to the left). In (a) the measures restricting contacts are introduced first and in (b) the isolation-based measure.

As shown in section III A, moving from voluntary to mandatory testing has a sizeable impact on transmission risk and this is reproduced here (compare "testing" to "enforced testing" in figures 6 and 7). Interestingly, we also see it has only a small impact on the number of isolations. This is because the reduction in transmission means fewer cases, which acts to counteract the increased rate of people entering isolation. This effect is even more stark if testing is enforced in the absence of other measures (see figure 8). In that case, the imapct of testing is significant enough to mean that the number of isolations actually reduces by switching from voluntary to mandatory testing.

FIG. 7. The same as figure 6 but for the large-item delivery workplace.

The impact of interventions for the large-item delivery workplace is very similar (see figure 7). We see that the "fixed pairings" intervention (which includes pair isolation) has a marked effect on transmission. The extra benefit gained from testing is clearly visible too, but again isolation measures beyond this appear have little further effect.

To conclude, figures 6 and 7 demonstrate some of the trade-offs for different intervention measures in terms of their impact on transmission and their impact on the number of isolating employees. Certain interventions act to reduce both (social distancing, Office staff WFH) but potentially have other costs for business/feasibility issues that need to be considered. When there are employees that still need to be in close-contact (e.g. driver and picker pairs in this model) the combination of fixed pairings, pair isolation, and regular testing is highly effective for reducing transmission.

FIG. 8. The same plots as figures 6 and 7 except with interventions applied separately (rather than cumulatively). (a) The number of secondary cases and isolations in simulations of the parcel delivery workplace. (b) The same quantities for the large-item delivery workplace.

IV. DISCUSSION

In this paper we have developed a stochastic model of SARS-CoV-2 spread in small/medium size workplaces. The contact patterns simulated were designed to represent warehouses/depots in the home-delivery sector, particularly those focusing on B2C delivery. To our knowledge this is the first model to consider SARS-CoV-2 transmission in this sector specifically. While the parameterisation of these models has significant uncertainty, we have been able to test the relative impact of various interventions that companies in this sector deployed to reduce

SARS-CoV-2 transmission over a range of scenarios and parameter regimes.

The results predict that workplace transmission in this sector is modest, due to the bulk of the staff, drivers, working alone most of the day. Without any interventions there is predicted to be a small risk to customers for an individual delivery, but in workplaces undergoing an outbreak, home-installation of items can pose a risk to customers without other interventions. The companies we consulted discontinued home-installation during in the spring of 2020, but later re-introduced it with social distancing measures. Parcel delivery companies switched to "contactless" delivery, meaning that signatures are no longer required, essentially eliminating the only route of transmission to customers. Overall, this suggests that this sector played a key role in reducing community transmission of SARS-CoV-2, as it allowed people to stay at home during periods of high-prevalence. Quantifying this impact is more difficult though as the counterfactual situation (i.e. how people would have behaved if this sector failed to keep up with increased demand or shops had remained open) is unknown.

Safeguarding the key workers in this sector was a broader challenge and companies reported implementing multiple measures based on government guidelines and their own judgement. A key result of this paper is that identifying high-risk contacts (due to e.g. shared accommodation or work tasks requiring prolonged close-contact) is very important and forms the basis of contact-tracing interventions. Workplaces have an extra advantage over contact-tracers in that they have control and knowledge over some of the contacts that employees are required to make in their line of work. In this case, high-risk contacts can be limited by using fixed pairs for large-item delivery, reducing then number of people sharing an office, and reducing the occupancy in shared spaces. This then allows efficient isolation policies to be implemented based on knowledge of the limited number of high-risk contacts people have made (e.g. workers are given paid isolation leave if they share accommodation or work in a delivery pair with an employee who has tested positive or reported COVID-like symptoms). Supplementary figure S7 and figure 4 show that this combination can be very effective in small workplaces.

Rates of presenteeism, which in this case we define as those who do not self-isolate when they develop symptoms, has been shown to be much less likely if fully-paid sick leave is offered [38, 39]. Therefore, in order to be effective, such isolation policies could incur considerable costs to a business as well as reducing productivity. Similarly, company-backed testing and isolation interventions will incur further costs and mean that asymptomatic cases are detected, potentially resulting in even more isolations. Therefore, companies may be apprehensive about deploying such strategies. Here, we showed that these strategies are more

efficient when close-contacts of index cases can be identified and isolated too, particularly when such contacts are necessary for the job. This can mean that chains of transmission are quickly shut down, and outbreaks are much less likely to occur. Furthermore, combining these measures with social distancing, WFH, and similar interventions that reduce transmission (e.g. masking) can reduce the number of isolations since workplace outbreaks become less likely. Combining testing measures with these can reduce risk of infection in the workplace and therefore reduce the costs of employee isolation (as fewer people will need to isolate).

The model developed has several limitations which are important for the interpretation of the results presented. First, the contact model has been developed based on a mix of quantitative (survey data, staff numbers and demand levels) and qualitative data (consultations). Novel insight was gained by speaking directly to representatives from the sector, but their position was not objective and so there may have been implicit biases in the descriptions of the nature of workplace contacts and some potential routes of infection contacts could have been missed. Furthermore, simplifying assumptions, such as all contact durations being identical for the same mode of contact, mean that this model is idealised compared to reality. Third, the transmission rate and the modifiers used for different types of contact are uncertain, and is based on a combination of peer-reviewed [22, 23, 29] and non-peer-reviewed literature [21]. Improvements to this transmission model from the adaption of more mechanistic modelling approaches that predict explicitly the infectious dose associated with different modes of contact [23, 29, 40–46], as well as updating with data on new variants and vaccines, will mean that this model could be applied to numerous future workplace scenarios to test the impact of different non-pharmaceutical interventions.

There are also some complicating factors we choose to ignore in this model, that may be important to consider when interpreting these results. First, we do not model severe illness, which can impact results by increasing the time away from work of individuals with COVID-19. Second, we do not model the complex relationship between interventions and behaviour. It is possible that as more interventions are introduced, adherence with other interventions wanes so the expected impact of combined interventions may not be as high as predicted. This behavioural change is difficult to predict, and so would need to be monitored by companies to gauge whether interventions are working as expected. Furthermore, even with high adherence there is no guarantee that people will use the test as intended. For example, people may be inclined to test more regularly when feeling run down or 'paucisymptomatic', i.e. exhibiting very mild COVID-19 symptoms, whereas in the absence of testing they may

have simply isolated from work. In this case, much of the benefit of testing can be lost [47] because asymptomatic carriers will be less likely to be detected while symptomatic carriers who would have otherwise isolated may be given a false negative and choose not to. For this reason, in some sectors, mandatory regular testing (i.e. carried out by trained swabbers at the workplace) may be the preferred option, because with the adherence rates assumed in this paper, one mandatory test per week has a similar impact to two voluntary ones (see figures 3 and 4).

One major benefit of the model presented here is that it incorporates the dynamics and variability in individual viral load, and simulates its impact on test sensitivity and infectiousness. This, means that the correlation between test-positivity and infectiousness is incorporated, meaning that impacts of these interventions can be more accurately estimated. Thus we were able to estimate not only the effect on average transmission rates, but also the frequency of rare superspreading events. This has highlighted the importance of stacking interventions that reduce transmission through different mechanisms. The source code is open access [35] and the underlying network transmission model is malleable enough to be applied to any small closed populations.

In conclusion, this paper has shown that the multiple interventions put in place by the logistics and home delivery sector during the early stages of the pandemic are likely to have reduced the risk of community and workplace transmission by safeguarding customers and staff. The availability of lateral flow tests is another valuable layer of protection that could have been added, and that this would have been most effective when combined with isolation measures that target the most high-risk contacts.

ETHICS STATEMENT

The project was reviewed and approved by the University Research Ethics Committee at University of Manchester, Ref: 2020-9787-15953. Consent to participation was verbally obtained before the commencement of the interviews. Written informed consent for participation was not required for this study in accordance with the national legislation and the institutional requirements.

have simply isolated from work. In this case, much of the benefit of testing can be lost [47] because asymptomatic carriers will be less likely to be detected while symptomatic carriers who would have otherwise isolated may be given a false negative and choose not to. For this reason, in some sectors, mandatory regular testing (i.e. carried out by trained swabbers at the workplace) may be the preferred option, because with the adherence rates assumed in this paper, one mandatory test per week has a similar impact to two voluntary ones (see figures 3 and 4).

One major benefit of the model presented here is that it incorporates the dynamics and variability in individual viral load, and simulates its impact on test sensitivity and infectiousness. This, means that the correlation between test-positivity and infectiousness is incorporated, meaning that impacts of these interventions can be more accurately estimated. Thus we were able to estimate not only the effect on average transmission rates, but also the frequency of rare superspreading events. This has highlighted the importance of stacking interventions that reduce transmission through different mechanisms. The source code is open access [35] and the underlying network transmission model is malleable enough to be applied to any small closed populations.

In conclusion, this paper has shown that the multiple interventions put in place by the logistics and home delivery sector during the early stages of the pandemic are likely to have reduced the risk of community and workplace transmission by safeguarding customers and staff. The availability of lateral flow tests is another valuable layer of protection that could have been added, and that this would have been most effective when combined with isolation measures that target the most high-risk contacts.

ETHICS STATEMENT

The project was reviewed and approved by the University Research Ethics Committee at University of Manchester, Ref: 2020-9787-15953. Consent to participation was verbally obtained before the commencement of the interviews. Written informed consent for participation was not required for this study in accordance with the national legislation and the institutional requirements.

FUNDING

This project was funded by the UK Research and Innovation (UKRI) and National Institute for Health Research (NIHR) COVID-19 Rapid Response call, Grant Ref: MC_PC_19083. MvT is the Principal Investigator of the project. CAW and SD held post-doctoral posts funded by this grant.

LP is supported by the Wellcome Trust and the Royal Society (grant no. 202562/Z/16/Z). IH is supported by the National Institute for Health Research Policy Research Programme in Operational Research (OPERA, PR-R17-0916-21001). IH and LP are supported by The Alan Turing Institute for Data Science and Artificial Intelligence, EPSRC (EP/V027468/1). CAW, IH, and LP were also supported by UKRI through the JUNIPER modelling consortium (grant no. MR/V038613/1).

ACKNOWLEDGEMENTS

The authors would like to thank the project's advisory group that consist of Catherine Noakes, Chris Armitage, Sheena Johnson, Jeanette Edwards, Barbara Hockey, Nina Day, Nick Gent and Thomas House, for their advice that helped refine the aims and objectives of this article. We would also like to thank Helen Beers and Peter Baldwin from HSE for their advice on the business engagement and feedback to companies.

[1] A. Insight, UK Parcels Market Insight Report, Tech. rep. (2020).

[2] H. Wei, S. Daniels, C. A. Whitfield, Y. Han, D. W. Denning, I. Hall, M. Regan, A. Verma, M. van Tongeren, Agility and Sustainability: A Qualitative Evaluation of COVID-19 Non-pharmaceutical Interventions in the UK Logistics Sector, Frontiers in Public Health 10 (2022).

[3] K. B. Pouwels, T. House, E. Pritchard, J. V. Robotham, P. J. Birrell, A. Gelman, K.-D. Vihta, N. Bowers, I. Boreham, H. Thomas, J. Lewis, I. Bell, J. I. Bell, J. N. Newton, J. Farrar, I. Diamond, P. Benton, A. S. Walker, K. B. Pouwels, A. S. Walker, D. Crook, P. C. Matthews, T. Peto, E. Pritchard, N. Stoesser, K.-D. Vihta, A. Howarth, G. Doherty, J. Kavanagh, K. K. Chau, S. B. Hatch, D. Ebner, L. M. Ferreira, T. Christott, B. D. Marsden, W. Dejnirattisai, J. Mongkolsapaya, S. Hoosdally, R. Cornall, D. I. Stuart, G. Screaton, D. Eyre, J. Bell, S. Cox, K. Paddon, T. James, T. House, J. N. Newton, J. V. Robotham, P. Birrell, H. Jordan,

T. Sheppard, G. Athey, D. Moody, L. Curry, P. Brereton, J. Hay, H. Vansteenhouse, I. Bell, I. Diamond, A. Lambert, P. Benton, E. Rourke, S. Hawkes, S. Henry, J. Scruton, P. Stokes, T. Thomas, J. Allen, R. Black, H. Bovill, D. Braunholtz, D. Brown, S. Collyer, M. Crees, C. Daglish, B. Davies, H. Donnarumma, J. Douglas-Mann, A. Felton, H. Finselbach, E. Fordham, A. Ipser, J. Jenkins, J. Jones, K. Kent, G. Kerai, L. Lloyd, V. Masding, E. Osborn, A. Patel, E. Pereira, T. Pett, M. Randall, D. Reeve, P. Shah, R. Snook, R. Studley, E. Sutherland, E. Swinn, H. Thomas, A. Tudor, J. Weston, S. Leib, J. Tierney, G. Farkas, R. Cobb, F. V. Galen, L. Compton, J. Irving, J. Clarke, R. Mullis, L. Ireland, D. Airimitoaie, C. Nash, D. Cox, S. Fisher, Z. Moore, J. McLean, M. Kerby, Community prevalence of SARS-CoV-2 in England from April to November, 2020: Results from the ONS Coronavirus Infection Survey, The Lancet Public Health 6 (1) (2021) e30–e38. `doi:10.1016/S2468-2667(20)30282-6`.

[4] E. Ortiz-Prado, A. R. Henriquez-Trujillo, I. A. Rivera-Olivero, T. Lozada, M. A. Garcia-Bereguiain, High prevalence of SARS-CoV-2 infection among food delivery riders. A case study from Quito, Ecuador, Sci. Total Environ. 770 (2021). `doi:10.1016/j.scitotenv.2021.145225`.

[5] T. Leng, E. M. Hill, R. N. Thompson, M. J. Tildesley, M. J. Keeling, L. Dyson, Assessing the impact of secondary school reopening strategies on within-school COVID-19 transmission and absences: A modelling study, medRxiv (2021) 2021.02.11.21251587`doi:10.1101/2021.02.11.21251587`.

[6] G. S. Malloy, L. Puglisi, M. L. Brandeau, T. D. Harvey, E. A. Wang, Effectiveness of interventions to reduce COVID-19 transmission in a large urban jail: A model-based analysis, BMJ Open 11 (2) (2021). `doi:10.1136/bmjopen-2020-042898`.

[7] C. C. Kerr, R. M. Stuart, D. Mistry, R. G. Abeysuriya, K. Rosenfeld, G. R. Hart, R. C. Núñez, J. A. Cohen, P. Selvaraj, B. Hagedorn, L. George, M. Jastrzębski, A. S. Izzo, G. Fowler, A. Palmer, D. Delport, N. Scott, S. L. Kelly, C. S. Bennette, B. G. Wagner, S. T. Chang, A. P. Oron, E. A. Wenger, J. Panovska-Griffiths, M. Famulare, D. J. Klein, Covasim: An agent-based model of COVID-19 dynamics and interventions, PLOS Computational Biology 17 (7) (2021) e1009149. `doi:10.1371/journal.pcbi.1009149`.

[8] E. M. Hill, B. D. Atkins, M. J. Keeling, L. Dyson, M. J. Tildesley, A network modelling approach to assess non-pharmaceutical disease controls in a worker population: An application to SARS-CoV-2, PLOS Computational Biology 17 (6) (2021) e1009058. `doi:10.1371/journal.pcbi.1009058`.

[9] M. Fyles, E. Fearon, C. Overton, n. null, T. Wingfield, G. F. Medley, I. Hall, L. Pellis, T. House, Using a household-structured branching process to analyse contact tracing in the SARS-CoV-2 pandemic, Philosophical Transactions of the Royal Society B: Biological Sciences 376 (1829) (2021) 20200267. `doi:10.1098/rstb.2020.0267`.

[10] J. J. Deeks, A. E. Raffle, Lateral flow tests cannot rule out SARS-CoV-2 infection, BMJ 371 (December) (2020) 1–2. `doi:10.1136/bmj.m4787`.

[11] J. Wise, Covid-19: Lateral flow tests miss over half of cases, Liverpool pilot data show, BMJ 371 (2020) m4848. `doi:10.1136/bmj.m4848`.

[12] H. Houston, A. Gupta-Wright, E. Toke-Bjolgerud, J. Biggin-Lamming, L. John, Diagnostic accuracy and utility of SARS-CoV-2 antigen lateral flow assays in medical admissions with possible COVID-19, J. Hosp. Infect. (xxxx) (2021) 18–20. `doi:10.1016/j.jhin.2021.01.018`.

[13] J. Ferguson, S. Dunn, A. Best, J. Mirza, B. Percival, M. Mayhew, O. Megram, F. Ashford, T. White, E. Moles-Garcia, L. Crawford, T. Plant, A. Bosworth, M. Kidd, A. Richter, J. Deeks, A. McNally, Validation testing to determine the sensitivity of lateral flow testing for asymptomatic SARS-CoV-2 detection in low prevalence settings: Testing frequency and public health messaging is key, PLOS Biology 19 (4) (2021) e3001216. `doi:10.1371/journal.pbio.3001216`.

[14] A. Wolf, J. Hulmes, S. Hopkins, Lateral flow device specificity in phase 4 (post-marketing) surveillance, Tech. rep. (2021).

[15] K. A. Walsh, K. Jordan, B. Clyne, D. Rohde, L. Drummond, P. Byrne, S. Ahern, P. G. Carty, K. K. O'Brien, E. O'Murchu, M. O'Neill, S. M. Smith, M. Ryan, P. Harrington, SARS-CoV-2 detection, viral load and infectivity over the course of an infection, J. Infect. 81 (3) (2020) 357–371. `doi:10.1016/j.jinf.2020.06.067`.

[16] B. La Scola, M. Le Bideau, J. Andreani, V. T. Hoang, C. Grimaldier, P. Colson, P. Gautret, D. Raoult, Viral RNA load as determined by cell culture as a management tool for discharge of SARS-CoV-2 patients from infectious disease wards, Eur. J. Clin. Microbiol. Infect. Dis. 39 (6) (2020) 1059–1061. `doi:10.1007/s10096-020-03913-9`.

[17] J. J. van Kampen, D. A. van de Vijver, P. L. Fraaij, B. L. Haagmans, M. M. Lamers, N. Okba, J. P. van den Akker, H. Endeman, D. A. Gommers, J. J. Cornelissen, R. A. Hoek, M. M. van der Eerden, D. A. Hesselink, H. J. Metselaar, A. Verbon, J. E. de Steenwinkel, G. I. Aron, E. C. van Gorp, S. van Boheemen, J. C. Voermans, C. A. Boucher, R. Molenkamp, M. P. Koopmans, C. Geurtsvankessel, A. A. van der Eijk, Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-19),

Nat. Commun. 12 (1) (2021) 8–13. doi:10.1038/s41467-020-20568-4.

[18] T. Peto, COVID-19: Rapid Antigen detection for SARS-CoV-2 by lateral flow assay: A national systematic evaluation for mass-testing, medRxiv (2021) 2021.01.13.21249563.

[19] S. Daniels, H. Wei, Y. Han, H. Catt, D. W. Denning, I. Hall, M. Regan, A. Verma, C. A. Whitfield, M. van Tongeren, Risk factors associated with respiratory infectious disease-related presenteeism: A rapid review, BMC Public Health 21 (1) (2021) 1955. doi:10.1186/s12889-021-12008-9.

[20] J. R. Bridgen, H. Wei, C. Whitfield, Y. Han, I. Hall, C. P. Jewell, M. J. van Tongeren, J. M. Read, Contact patterns of UK home delivery drivers and their use of protective measures during the COVID-19 pandemic: A cross-sectional study, medRxiv (2022) 2022.09.09.22279754doi:10.1101/2022.09.09.22279754.

[21] The microCOVID Project, White paper (2020).

[22] L. Y. W. Lee, S. Rozmanowski, M. Pang, A. Charlett, C. Anderson, G. J. Hughes, M. Barnard, L. Peto, R. Vipond, A. Sienkiewicz, S. Hopkins, J. Bell, D. W. Crook, N. Gent, A. S. Walker, T. E. A. Peto, D. W. Eyre, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infectivity by Viral Load, S Gene Variants and Demographic Factors, and the Utility of Lateral Flow Devices to Prevent Transmission, Clinical Infectious Diseases 74 (3) (2022) 407–415. doi:10.1093/cid/ciab421.

[23] G. Cortellessa, L. Stabile, F. Arpino, D. E. Faleiros, W. van den Bos, L. Morawska, G. Buonanno, Close proximity risk assessment for SARS-CoV-2 infection, Science of The Total Environment 794 (2021) 148749. doi:10.1016/j.scitotenv.2021.148749.

[24] N. van Doremalen, T. Bushmaker, D. H. Morris, M. G. Holbrook, A. Gamble, B. N. Williamson, A. Tamin, J. L. Harcourt, N. J. Thornburg, S. I. Gerber, J. O. Lloyd-Smith, E. de Wit, V. J. Munster, Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1, New England Journal of Medicine 382 (16) (2020) 1564–1567. doi:10.1056/NEJMc2004973.

[25] C. A. Whitfield, U. o. M. C.-. M. Group, I. Hall, Modelling the impact of repeat asymptomatic testing policies for staff on SARS-CoV-2 transmission potential, arXiv (Oct. 2022). doi:10.48550/arXiv.2210.08888.

[26] R. Ke, P. P. Martinez, R. L. Smith, L. L. Gibson, A. Mirza, M. Conte, N. Gallagher, C. H. Luo, J. Jarrett, R. Zhou, A. Conte, T. Liu, M. Farjo, K. K. O. Walden, G. Rendon, C. J. Fields, L. Wang, R. Fredrickson, D. C. Edmonson, M. E. Baughman, K. K. Chiu, H. Choi, K. R. Scardina, S. Bradley, S. L. Gloss, C. Reinhart, J. Yedetore, J. Quicksall, A. N. Owens,

J. Broach, B. Barton, P. Lazar, W. J. Heetderks, M. L. Robinson, H. H. Mostafa, Y. C. Manabe, A. Pekosz, D. D. McManus, C. B. Brooke, Daily longitudinal sampling of SARS-CoV-2 infection reveals substantial heterogeneity in infectiousness, Nat Microbiol 7 (5) (2022) 640–652. `doi:10.1038/s41564-022-01105-z`.

[27] T. Watanabe, T. A. Bartrand, M. H. Weir, T. Omura, C. N. Haas, Development of a dose-response model for SARS coronavirus, Risk Anal. 30 (7) (2010) 1129–1138. `doi:10.1111/j.1539-6924.2010.01427.x`.

[28] X. Zhang, J. Wang, Dose-response Relation Deduced for Coronaviruses from COVID-19, SARS and MERS Meta-analysis Results and its Application for Infection Risk Assessment of Aerosol Transmission, Clin. Infect. Dis. (2020). `doi:10.1093/cid/ciaa1675`.

[29] F. Poydenot, I. Abdourahamane, E. Caplain, S. Der, J. Haiech, A. Jallon, I. Khoutami, A. Loucif, E. Marinov, B. Andreotti, Risk assessment for long and short range airborne transmission of SARS-CoV-2, indoors and outdoors, PNAS Nexus (2022) pgac223`doi:10.1093/pnasnexus/pgac223`.

[30] Z. J. Madewell, Y. Yang, I. M. Longini, Jr, M. E. Halloran, N. E. Dean, Household Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis, JAMA Network Open 3 (12) (2020) e2031756. `doi:10.1001/jamanetworkopen.2020.31756`.

[31] K. Kuwelker, F. Zhou, B. Blomberg, S. Lartey, K. A. Brokstad, M. C. Trieu, A. Madsen, F. Krammer, K. G. Mohn, C. Tøndel, D. W. Linchausen, R. J. Cox, N. Langeland, High attack rates of SARS-CoV-2 infection through household-transmission: A prospective study, medRxiv (2020). `doi:10.1101/2020.11.02.20224485`.

[32] Q. L. Jing, M. J. Liu, Z. B. Zhang, L. Q. Fang, J. Yuan, A. R. Zhang, N. E. Dean, L. Luo, M. M. Ma, I. Longini, E. Kenah, Y. Lu, Y. Ma, N. Jalali, Z. C. Yang, Y. Yang, Household secondary attack rate of COVID-19 and associated determinants in Guangzhou, China: A retrospective cohort study, Lancet Infect. Dis. 20 (10) (2020) 1141–1150. `doi:10.1016/S1473-3099(20)30471-0`.

[33] S. C. W. Group, SCWG Chairs: Summary of role of shielding, 20 December 2021, https://www.gov.uk/government/publications/scwg-chairs-summary-of-role-of-shielding-20-december-2021 (2021).

[34] C. A. Whitfield, Model of SARS-CoV-2 transmission in delivery workplaces (2022).

[35] C. A. Whitfield, Model of SARS-CoV-2 viral load dynamics, infectivity profile, and test-positivity. (2022).

[36] D. B. Larremore, B. Wilder, E. Lester, S. Shehata, J. M. Burke, J. A. Hay, M. Tambe, M. J. Mina, R. Parker, Test sensitivity is secondary to frequency and turnaround time for COVID-19 screening, Science Advances 7 (1) (2021) eabd5393. `doi:10.1126/sciadv.abd5393`.

[37] M. L. Waskom, Seaborn: Statistical data visualization, J. Open Source Softw. 6 (60) (2021) 3021. `doi:10.21105/joss.03021`.

[38] K. Piper, A. Youk, A. E. J. Iii, S. Kumar, Paid sick days and stay-at-home behavior for influenza, PLOS ONE 12 (2) (2017) e0170698. `doi:10.1371/journal.pone.0170698`.

[39] F. Ahmed, S. Kim, M. P. Nowalk, J. P. King, J. J. VanWormer, M. Gaglani, R. K. Zimmerman, T. Bear, M. L. Jackson, L. A. Jackson, E. Martin, C. Cheng, B. Flannery, J. R. Chung, A. Uzicanin, Paid Leave and Access to Telework as Work Attendance Determinants during Acute Respiratory Illness, United States, 2017–2018, Emerg Infect Dis 26 (1) (2020) 26–33. `doi:10.3201/eid2601.190743`.

[40] A. L. Freeman, S. Parker, C. Noakes, S. Fitzgerald, A. Smyth, R. Macbeth, D. Spiegelhalter, H. Rutter, Expert elicitation on the relative importance of possible SARS-CoV-2 transmission routes and the effectiveness of mitigations, BMJ Open 11 (12) (2021) e050869. `doi:10.1136/bmjopen-2021-050869`.

[41] A. K. Pitol, T. R. Julian, Community Transmission of SARS-CoV-2 by Surfaces: Risks and Risk Reduction Strategies, Environ. Sci. Technol. Lett. 8 (3) (2021) 263–269. `doi:10.1021/acs.estlett.0c00966`.

[42] H. C. Burridge, S. Fan, R. L. Jones, C. J. Noakes, P. F. Linden, Predictive and retrospective modelling of airborne infection risk using monitored carbon dioxide, Indoor and Built Environment 31 (5) (2022) 1363–1380. `doi:10.1177/1420326X211043564`.

[43] G. Buonanno, A. Robotto, E. Brizio, L. Morawska, A. Civra, F. Corino, D. Lembo, G. Ficco, L. Stabile, Link between SARS-CoV-2 emissions and airborne concentrations: Closing the gap in understanding, Journal of Hazardous Materials 428 (2022) 128279. `doi:10.1016/j.jhazmat.2022.128279`.

[44] S. L. Miller, W. W. Nazaroff, J. L. Jimenez, A. Boerstra, G. Buonanno, S. J. Dancer, J. Kurnitski, L. C. Marr, L. Morawska, C. Noakes, Transmission of SARS-CoV-2 by inhalation of respiratory aerosol in the Skagit Valley Chorale superspreading event, Indoor Air 31 (2) (2021) 314–323. `doi:10.1111/ina.12751`.

[45] P. Azimi, Z. Keshavarz, J. G. C. Laurent, B. Stephens, J. G. Allen, Mechanistic transmission modeling of COVID-19 on the Diamond Princess cruise ship demonstrates the importance of aerosol transmission, PNAS 118 (8) (Feb. 2021). `doi:10.1073/pnas.2015482118`.

[46] L. Morawska, J. W. Tang, W. Bahnfleth, P. M. Bluyssen, A. Boerstra, G. Buonanno, J. Cao, S. Dancer, A. Floto, F. Franchimon, C. Haworth, J. Hogeling, C. Isaxon, J. L. Jimenez, J. Kurnitski, Y. Li, M. Loomans, G. Marks, L. C. Marr, L. Mazzarella, A. K. Melikov, S. Miller, D. K. Milton, W. Nazaroff, P. V. Nielsen, C. Noakes, J. Peccia, X. Querol, C. Sekhar, O. Seppänen, S.-i. Tanabe, R. Tellier, K. W. Tham, P. Wargocki, A. Wierzbicka, M. Yao, How can airborne transmission of COVID-19 indoors be minimised?, Environment International 142 (2020) 105832. `doi:10.1016/j.envint.2020.105832`.

[47] J. P. Skittrall, SARS-CoV-2 screening: Effectiveness and risk of increasing transmission, Journal of The Royal Society Interface 18 (180) (2021) 20210164. `doi:10.1098/rsif.2021.0164`.