Updated population-level estimates of child restraint practices among children aged 0-12 years in Australia, ten years after introduction of age-appropriate restraint use legislation

Julie Brown[a,b,c*], Bianca Albanese[a,b,c], Catherine Ho [b], Jane Elkington [b], Sjaan Koppel[d], Judith L. Charlton[d], Jake Olivier[e], Lisa Keay[b,f], Lynne E Bilston[a,c]

a. Neuroscience Research Australia, Barker St, Randwick 2031, Australia
b. The George Institute for Global Health, University of New South Wales, 1 King St Newtown 2042, Australia
c. Faculty of Medicine, University of New South Wales, Sydney 2052, Australia
d. Monash University Accident Research Centre, Monash University, Clayton 3800, Australia
e. School of Mathematics and Statistics, University of New South Wales, Sydney 2052, Australia
f. School of Optometry and Vision Science, University of New South Wales, Sydney 2033, Australia

*Corresponding author

Abstract

Objective: To determine child restraint practices approximately 10 years after introduction of legislation requiring correct use of age-appropriate restraints for all children aged up to 7 years.

Methods: A stratified cluster sample was constructed to collect observational data from children aged 0-12 years across the Greater Sydney region of NSW. Methods replicated those used in a similar 2008 study. Population weighted estimates for restraint practices were generated, and logistic regression used to examine associations between restraint type, and child age with correct use accounting for the complex sample.

Results: Almost all children were appropriately restrained (99.3%, 95% CI 98.4-100). However, less than half were correctly restrained (No error = 27.3%, 95% CI 10.8-43.8, No serious error = 43.8%, 95% CI 35.0-52.7). For *any* error, the odds of error decreased by 39% per year of age (OR 0.61, 95% 0.46-0.81), and for *serious error* by 25% per year (OR 0.75 95% CI 0.60-0.93).

Conclusion: The findings demonstrate a substantial increase in appropriate child restraint, but no real change in correct use.

Implications for Public Health: Given the negative impact incorrect use has on crash protection, continuing high rates of incorrect use may reduce effectiveness of legislative change on injury reduction.

Key Words

Child passenger, child restraints, car seats, seat belts, appropriate use, correct use. legislation

1. Introduction

Seat belts and child restraints are effective countermeasures to serious injury and death in car crashes.[1,2] However, optimal crash protection for child passengers requires the correct use of a restraint that is most appropriate for the child's size.[1-3] New legislation was introduced in Australia in

2009/2010 requiring all child passengers aged up to 7 years to correctly use an age-appropriate

restraint. Similar child restraint laws exist in many other jurisdictions.[4,5]

In Australia, like many developed countries with long histories of seat belt and child restraint use,

the rate of restraint use by children is high. In Australia, restraint use rates have been >90% since the

late 1990s,[6] yet children continue to be seriously and fatally injured in crashes. Population

observation studies prior to the introduction of the 2009/2010 legislation indicated around 20% of

child passengers were optimally restrained, with approximately one third using restraints

inappropriately, another third using their restraint incorrectly, and the remaining one in five

incorrectly using an inappropriate type of restraint. [7] As the intention of the legislative change was

to ensure the proportion of children correctly using age-appropriate restraints increased, we might

postulate that child restraint practices in Australia have improved over the last decade.

Pre- and post-observations of children aged 2-5 years in low socioeconomic areas of Sydney

immediately before and after introduction of the 2009/2010 age-appropriate restraint use laws

found the odds of age-appropriate restraint was almost three times higher, and the odds of correct

use was 1.6 times higher post legislation.[8] However, Koppel et al[9] found no significant difference in

appropriate restraint or correct use in samples of child restraint observations made in the two years

before and two years after the introduction of the legislation. While there were substantial

differences in the ages of children studied, sampling methods and definitions used for inappropriate

and incorrect use between these Australian studies, the contradictions in results reflect similar

contradictions in evidence for the impact of child restraint legislation in the literature more broadly.[5]

From a recent systematic review[5], substantial heterogeneity between different studies makes it

difficult to interpret the evidence for the impact of legislation on restraint use, particularly in terms

of appropriate and correct restraint use. In addition to heterogeneity issues, there is also a paucity of

studies that incorporate gold-standard approaches such as: using observed restraint practices rather

than self-reported practices, randomly selected samples rather than self-selected samples, and

samples that are population representative rather than convenience samples or focused on one

segment of the population. There have been no population-level examinations of observed restraint

practices in Australia since our 2008 study or after the introduction of the legislation.

In this study, we aimed to provide an update on child restraint practices in NSW, approximately 10

years after the introduction of age-appropriate legislation. While we do not intend to test the impact

of the legislation, we used the same methodological approaches as those in our 2008 study[7] to allow

comparison of the situation before and after the legislation using homogenous and gold-standard

observational approaches with stratified random sampling to reduce selection bias. Our original

intention was to provide updated restraint practice estimates for the state of NSW; however, our

study was impacted by COVID-19 restrictions in 2020 and data collection was limited to the Greater

Sydney region. We therefore aim to provide updated estimates for this region only.

76   2. Material and methods

We used a cross-sectional design with a stratified cluster sample constructed to observe restraint

practices across a population-representative sample of children aged 0-12 years in the greater

Sydney region. Data collection occurred during 2019-2020. Six strata based on Local Government

Areas (LGAs), four from inner metropolitan areas and two from outer areas, were randomly selected

from all LGAs in the region. The 4:2 ratio was used to select inner and outer areas as corresponding

to population distribution across Greater Sydney. Sites where children of different age groups visit

regularly were then randomly selected from each LGA: preschool/long daycare centres to capture

infants and pre-school aged children, and primary schools to capture school aged children, resulting

in 12 data collection clusters.

Participants were children aged 0-12 years and their carer who attended a data collection site during

a data collection period. Child-carer dyads were randomly selected as they arrived at the site and

invited to participate by on-site researchers before they left their vehicles. Refusals were recorded,

and reasons for non-participation noted. If more than one eligible child was in the vehicle, the

participant was the one who had last had a birthday. An inability to converse in English, and previous

participation in the study were the only exclusion criteria.

The study was approved in August 2018 by the South Eastern Sydney Local Health District Human

Research Ethics Committee. Approval to collect data at education sites was also obtained via the

NSW School Education Research Applications Process.

2.1 Data collection

Data collection methods were the same as that used in the 2008 observational study.[7] In brief,

trained researchers attended the sites over a one-two hour period corresponding with child

care/school morning drop-off times. If the driver/carer agreed to participate, the child's initial

observations were recorded as they remained in their restraint within the vehicle. Once the child left

the vehicle, a structured interview was conducted with the carer, the height

and weight of the child were measured using scales and a portable potentiometer, and a detailed

inspection of the restraint installation was conducted. The latter included detail on restraint type,

make and model and each possible form of correct/incorrect use of the restraint.

The only difference in data collection method from earlier studies was the addition of a standardised

series of photographs used for data quality control.

2.2 Variables

Appropriate restraint use was defined using the current legal definitions of appropriate restraint in

NSW, Australia (see Table 1). Children were categorised as 'appropriately' or 'inappropriately'

restrained using their age and the restraint type they were using.

Child age	Restraint type
0 – 6 months	Rearward-facing restraint
6 months – <4 years	Rearward- or forward-facing restraint
4 years – <7 years	Forward-facing restraint or booster seat
7+ years	Forward-facing restraint, booster seat or adult seat belt.

Table 1: Appropriate restraint definitions based on current restraint use laws in NSW, Australia

Correct use was defined as use exactly as the manufacturer intended. Each type of error was

categorised as an 'installation error' or a 'securing error', and as 'minor' or 'serious' depending on

the likely degradation in protection or increased risk of injury in a crash introduced by that error.[7]

(see Table 2). This resulted in six correct use outcome variables: (i) any error (whether minor or

serious), (ii) any serious error, (iii) any installation error (whether minor or serious) (iv) any serious

installation error, (v) any securing error (whether minor or serious), and (vi) any serious securing

errors.

Restraints were categorised by type (i.e., rearward-facing, forward-facing, booster seat or adult seat

belt) and whether they were convertible or non-convertible restraint types, i.e. designed to be used

in more than one mode (e.g., rearward- and forward-facing restraint or forward-facing and a booster

seat).

In Australia, rearward and forward facing child restraints can be installed using vehicle seat belt, or

ISOFIX compatible connectors. Therefore restraints were also categorised as 'ISOFIX compatible' or

'seatbelt'.

Child age was calculated from date of birth and rounded down to their last birthday.

Restraint types	Minor	Serious
Rearward facing and forward facing restraints	Harness height error	Harness buckle not used/engaged correctly
	Harness slack <25mm	Arms not within harness
	Harness webbing twists	Harness slack >25mm
		Harness off one or both shoulders
	Seat belt and ISOFIX compatible system used	No lower anchorage system used
	ISOFIX compatible anchorage webbing or seat belt webbing twists	ISOFIX compatible anchorage or seat belt buckle not engaged correctly
	ISOFIX compatible anchorage webbing or seat belt <25mm slack	ISOFIX compatible anchorage webbing or seat belt >25mm slack
		Incorrect seat belt routing
		Infant carrier not correctly engaged with base
		Infant restraint stabiliser bar required but not used
Rearward facing, forward facing restraints & booster seats	Top tether slack <25mm	Top tether required but not used
	Top tether webbing twists	Top tether slack >25mm
		Incorrect Top tether routing
		Top Tether anchored to incorrect point
Booster seats		sash guide not used (inc. anti-submarine clips)
Booster seats and adult seat belts	Seat belt webbing twists	Sash belt used incorrectly e.g. placed behind back, under arm
	Seat belt slack <25mm	Seat belt buckle not engaged
		Seat belt slack >25mm

Table 2: Minor and Serious Errors

2.3 Data Analysis

All analysis was conducted using SAS Version 9.2 (SAS Institute, 2008). Sample weights were

constructed using standard weighting procedures.[10,11] Post-stratification weighting for age

distribution, and under-and over sampling at different sites were used to generate population-level

estimates for the Greater Sydney region. The SURVEYFREQ procedure was used to generate

population weighted estimates and 95% confidence intervals (CI) for the proportion of children

appropriately and correctly restrained, and those using different restraint types. The

SURVEYLOGISTIC procedure was used to examine associations between different restraint types,

anchorage methods and child age with correct use. The relationship between restraint type and

correct use was also examined while controlling for child age, and significant interactions explored

through sub-group analysis. Odds ratios (OR) and 95% CI were calculated, and associations assessed

as significant when the confidence interval did not pass through 1.

3. Results

Data were collected from 213 children aged 0-12 years, sample characteristics are summarised in

Table 3. The refusal rate was 40%, with 'lack of time' given as the primary reason for refusal. No

children were unrestrained, and almost all (99.3%, 95% CI 98.4-100) were using an appropriate

restraint as defined by the current legislation. However, fewer children were correctly restrained

(i.e., no error = 27.3%, 95% CI 10.8-43.8; no serious error= 43.8%, 95% CI 35.0-52.7). Both

installation (any installation error = 53.9%, 95% CI 33.7-74.0, any serious installation error = 36%, 95%

CI 24.6-47.4) and securing errors (any securing error = 65%, 95% CI 52.6-77.2, any serious securing

error = 40.7%, 95% CI 34.1-47.2) were common.

Characteristic		Unweighted N (%)	Weighted/Age adjusted estimate (95% CI)
Age	0-3 years	66 (31.0)	33.81 (12.19-55.42)
	4-8 years	108 (50.7)	41.22 (29.97-52.48)
	9-12 years	39 (18.3)	24.97 (3.02-46.92)
Gender	Male	108 (51.2)	49.05 (40.36-57.54)
	Female	103 (48.8)	50.95 (42.25-59.64)
Restraint Type	Rear-facing restraint	12 (5.7)	5.09 (0.01-11.08)
	Forward-facing restraint	124 (58.5)	54.84 (26.65-83.02)
	Booster	35 (16.5)	14.01 (3.93-24.09)
	Adult Seat Belt	41 (19.3)	26.06 (2.85-49.27)
Appropriate *	Yes	211 (99.1)	99.32 (98.39-100.00)
	No	2 (0.9)	0.68 (0.00-1.61)
Any Error	No	51 (24.1)	27.31 (10.82-43.79)
	Yes	161 (75.9)	72.69 (56.21-89.18)
Installation error	No	90 (42.4)	46.14 (26.02-66.26)
	Yes	122 (57.6)	53.86 (33.74-73.98)
Securing error	No	70 (33.0)	35.07 (22.79-47.34)
	Yes	142 (67.0)	64.93 (52.65-77.21)
Serious error	No	92 (43.4)	43.85 (54.99-52.72)
	Yes	120 (56.6)	56.15 (47.28-65.01)
Installation error (Serious)	No	130 (61.3)	64.01 (52.64-75.36)
	Yes	82 (38.7)	35.99 (24.63-47.35)
Securing (Serious)	No	126 (59.4)	59.33 (52.79-65.88)
	Yes	86 (40.6)	40.67 (34.12-47.21)
Characteristic		M (SE)	95% CI for mean
Weigh (kg)	Overall	23.4(3.4)	15.4-31.4

	Rear-facing restraint	8.7(0.3)	7.9-9.4
	Forward-facing restraint	15.5(0.7)	13.9-17.0
	Booster	22.4(1.4)	19.2-25.6
	Adult Seat Belt	43.7(3.4)	35.5-51.8
Height(mm)	Overall	111.8(6.2)	97.1-126.5
	Rear-facing restraint	70.6(3.4)	62.6-78.5
	Forward-facing restraint	98.1(2.0)	93.4-102.8
	Booster	116.6(1.9)	112.0-121.2
	Adult Seat Belt	143.7(1.4)	140.4-147.0

*Table 3: Sample characteristics *NSW legal definition*

Across the sample, 5.1% (95% CI 0-11.1) used a rearward-facing restraint, 54.8% (95% CI 26.6-83.0)

used a forward-facing restraint, 14.0% (95% CI 26.6-83.0) used a booster seat, and 9.8% (95% CI 2.9-

49.3) used an adult seat belt. Most of the rearward- and forward-facing restraints were installed

using a seat belt, with 12.5% using an ISOFIX compatible lower anchorage.

Among the children using inappropriate restraint types, all were aged three years and had been

prematurely graduated to a booster seat. In addition, all had serious errors in the way the restraint

was used with 3/3 failing to use available belt guides resulting in incorrectly positioned belts; one

failing to use the top tether, and one failing to engage the seat belt buckle.

Among dedicated restraints (i.e. rear or forward-facing restraints and booster seats), 84% were

convertible restraints. There was no association between whether dedicated restraint were single

mode or convertible restraints and errors in use (for *any error*, p=0.67; for *any serious error* p=0.84).

Figure 1 illustrates the variation in proportion of children with observed errors by restraint type.

Seat belts were used correctly more often than all dedicated child restraint types. Among the

dedicated restraints, there were more errors in forward-facing restraints than in rearward-facing

restraints (for *any error* in forward-facing restraints compared to rearward facing restraints (OR 6.3,

95% CI 1.18-34.03) but this difference was not significant for serious errors (OR 2.1, 95% CI 0.64-

6.88). There were no significant differences in booster seat users compared to rearward-facing

restraints (*any error* OR 4.7, 95% CI 0.37-60.7, for *serious errors* OR 4.5, 95% CI 0.84-23.81), or

compared to forward-facing restraints (for *any error* OR 0.75, 95% CI 0.17-3.22, for *serious errors* OR

2.1 95% CI 0.49-9.24).

Examining the types of errors, there were no significant differences in the likelihood of installation

errors among the different types of dedicated restraint; however there were significant differences

in the odds of securing errors. Forward facing restraint users had over 7 times the odds of securing

errors than rearward facing restraints (*any error* OR 7.7, 95% CI 4.9-66.7) but the difference for

serious errors was not significant (OR 4.8, 95% CI 0.39-58.8). Booster seat users had significantly

greater odds for any securing error and serious securing errors than rearward facing restraint users.

Booster seat users also had more securing errors than forward facing restraint users but this

difference was only significant for serious securing errors (*any error* OR 2.3, 95% CI 0.7-7.8, *serious*

errors OR 3.2 95% CI 1.0-10.4).

Any error	RF	FF	BS	SB
■ Error	70.07	93.68	91.73	18.81
□ No error	29.93	6.32	8.27	81.19

Serious error	RF	FF	BS	SB
■ Error	54.04	71.25	84.05	9.78
□ No Error	45.96	28.75	15.95	90.22

Figure 1: Proportion of errorr, and serious errors by restraint type. RF= Rearward-Facing Restraint, FF=Forward-Facing
Restraint, BS=Booster Seat, SB= Seat belt.

Figures 2 and 3 illustrate the proportion of children of different ages with and without errors in

different types of restraint.

The odds of errors decreased with increasing age of the child. For *any error,* the odds decreased by

39% with every year older, and for *serious error* by 25% with every year older (for *any error* OR 0.61,

95% 0.46-0.81; for *any serious error* OR 0.75 95% CI 0.60-0.93). When age was added to the child

restraint type models, there was no significant associations between age and errors, or differences

between restraint types, for *'any error'* or *'serious errors'*. However, there was a significant

interaction between age and booster seat use in the 'any error' model, and between age and

forward-facing restraints in the 'serious error' model. This indicates age may have a moderating

effect on the likelihood of error by restraint type.

Figure 2: Proportion of children with any error by age for each restraint type.

Figure 3: Proportion of children with serious error by age for each restraint type.

Stratifying by restraint type, age had different relationships with incorrect use. While there was no

significant relationship between age and errors in the forward-facing restraints (for *any error*, OR

0.67 95% CI 0.37-1.24/*any serious error* OR 0.91, 95% CI 0.69-1.2), and adult seat belts (for *any error,*

OR 1.5 95% CI 0.39-6.09/*any serious error* OR 6.4 95% CI 0.87-46.9), however there was a tendency

for the odds of an error to <u>decrease </u>in forward-facing restraints and to <u>increase </u>with increasing age

in adult seat belts. For rearward-facing restraints, the odds of errors <u>increased </u>with age *for any*

serious error (OR 2.7 95% CI 1.1-6.6), and while there was a similar trend for *any error,* this did not

reach significance (OR 2.2, 95% CI 0.86-5.6). For booster seats, the odds of *any error* <u>decreased </u>by 40%

with every year increase in age (OR 0.59, 95% CI 0.39-0.89). This relationship was less clear for *any*

serious error (OR 0.62, 95% CI 0.17-2.2). Therefore, while overall the odds of errors decreased with

increasing age, the relationship between age and errors varied in different restraint types as

indicated by the significant interaction in the logistic regression model.

The proportion of errors in rear and forward facing restraints installed using ISOFIX compatible lower

anchorage systems were not statistically different (for *any error,* OR 0.29 95% CI 0.05-1.69, for *any*

serious error 0.31 95% CI 0.10-1.11, for *any installation error* OR 0.60, 95% CI 0.17-2.05, for *any*

serious installation error OR 0.44, 95%CI 0.07-2.62).

4. Discussion

This study demonstrates high rates of age-appropriate restraint use among children aged 0-12 years in the greater Sydney region, approximately ten years after introduction of legislation requiring age-appropriate restraint use until age 7. This appears to represent a substantial increase compared to prior to the legislation's introduction with 99.3% appropriate use in the current study compared to 48.8% in 2008.[7] However, incorrect restraint use continues to be widespread (51.4% had any error in 2008[7] and 72.7% in the current study; 38.3% had a serious error in 2008[7] and 56.2% in this current study).

The results also indicate the uptake of restraints with ISOFIX compatible anchorages is relatively low (~12% of rearward and forward-facing restraints) compared to North America where the use of these systems is more common than not.[13] Reasons for this are unknown but might relate to the more recent adoption of ISOFIC compatible restraints and generally older vehicle fleet in Australia. While the lack of significant difference in errors by anchorage method suggests the expected benefit of ISOFIX in reducing installation errors has not been realised, it also possible that user characteristics of those choosing to use or not use this method of anchorage might confound this association. Further exploration of these issues in future work is warranted.

While inappropriate restraint was rare, the characteristics were broadly like those reported in 2008[7], commonly involving premature graduation to booster seats, and serious errors.[7] However, there are stark differences between the 2008[7] and current study in premature graduation to adult seat belts with no observed premature graduation to seat belts in the current study. One important intrinsic difference before and after the introduction of the age-appropriate legislation is how appropriate use is defined. In this study we used the legal age-based definition. Before the legislation was enacted there was no legal definition, and we used restraint type designations set by the product Standard AS/NZS 1754 based on child height and weight to guide appropriate use definitions. To examine the effect of this, the same definitions used in the 2008 study were applied to the current

dataset. With those definitions appropriate use in the current sample was slightly lower at 94.2%,

(95% CI 87.5-100.0), compared to 99.3%, (95% CI 98.4-100) using the age-based definition, indicating

a substantial increase regardless of the definition used. Interestingly, the difference was due to a

small number of children aged 7 or 8 years who by the current age-based definitions were

appropriately using seat belts but who weighed ≤26kg which was the old upper boundary for

appropriate use of booster seats.

It is also important to note that the definitions of appropriate use in this study and in the earlier

2008 study reflect minimum acceptable practices which may be different to best practice. For

example, best practice in Australia encourages children be kept in a restraint of a specific type until

they reach shoulder height markers fixed to the restraint. With many rearward facing restraints

accommodating children up to the age of 2 or 3 years, and some forward-facing restraint

accommodating children up to age, best practice would see children remain in these restraint types

longer. However, as this depends on the specific make and model which was not always available, it

was not possible to compute estimates based on best practice. Similarly, while children aged over 7

years are appropriately restrained by adult seat belts according to Australian law, best practice

would be for children to only use an adult seat belt once they can sit with their knees bent over the

seat cushion and they achieve a good fit with the adult belt. In a separate analysis using data from

the current study[14] we found that only 40% of children deemed to be appropriately using adult seat

belts according to the legal definition actually achieved a good seat belt fit.

Restraint designs change over time, and since the legislation was introduced in Australia there have

been two substantial revisions to the Australian Child Restraint Standard (in 2010 and in 2013). It is

possible that the increasing complexity of restraint design may be a factor in the higher rates of

incorrect use in this current study, as there was an increase in the number of possible compared to

the earlier 2008 study.[7] However, an additional explanation is the shift of children from

inappropriate forms of restraint with lower propensity for errors, like seat belts, to more complex

dedicated child restraint systems with more opportunity for errors. While the same basic data

collection and method of identifying errors was used in this and the earlier 2008 study,[7] the new

quality control measures introduced in this current study (i.e. the secondary check of photographs

described in the methods) may also have led to more errors being detected than would have

without the secondary review of all cases.

In the current study, we found saw an overall decrease in likelihood of errors with increasing age,

consistent with observations made in the earlier study. However, this association was lost when

restraint type was controlled for, and we found interactions between age and some restraint types.

Our sub-group analysis suggests a more complex relationship between child age, restraint type and

incorrect restraint use, but the results of this analysis are limited by a loss of power with smaller

sample sizes in the sub-groups. The results from these analyses must therefore be viewed with

caution. However, more targeted investigation of this issue in future studies is warranted.

There are several limitations to keep in mind. As data were collected in metropolitan regions, rates

of appropriate and correct restraint may be slightly higher than in regional areas, as restraint

practices are generally reported to be poorer in rural and regional areas.[5,15] As this analysis sought to

estimate population-level estimates, no attempt was made to estimate differences in restraint use

by socioeconomic level and it is possible within the greater metropolitan area, restraint use

characteristics might vary from location to location. Estimates from this study may also be biased

due to a high refusal rate (~40%). Ethical conduct of research requires consent for participation, and

it is possible that people who are knowingly not using optimal restraint practice may choose not to

participate, leading to over-estimations of optimal restraint. Finally, this study did not attempt to

examine the impact of vehicle design on correctness of restraint use and the influence of any vehicle

design change over time on rates of incorrect restraint use remains unknown.

Strengths of this study include the randomly selected sample and the homogeneity of methods with

studies conducted prior to the introduction of the legislation in NSW in 2009/2010. While direct

estimates of the effectiveness of the legislation are not possible as other potential influences are not

accounted for, use of the same methods allows quantitative comparison of restraint practices before

and after the legislative change.

5. Conclusion

This study indicates a substantial increase in appropriate use of restraints by child passengers since

2008, likely influenced by the introduction of new legislation requiring age-appropriate restraint use.

However, this has not been accompanied by an improvement in correct use of restraints. Given the

negative influence of incorrect restraint on the level of protection in a crash, continuing high rates of

incorrect use may negatively impact effectiveness of legislation change on reducing injury among

child passengers.

6. Author Contributions

Julie Brown Conceptualization, Methodology, Formal Analysis, Writing -Original draft, Supervision,

Funding acquisition; *Bianca Albanese* Validation, Investigation, Data curation, Writing – review and

editing, Project Administration; *Catherine Ho*, Investigation, Data curation, Writing – review and

editing, Project Administration; *Jane Elkington*, Investigation, Data curation, Writing – review and

editing, Project Administration; *Sjaan Koppel*, Conceptualisation, Methodology, Writing – review and

editing, Funding acquisition; *Judith L. Charlton,* Conceptualisation, Methodology, Writing – review

and editing, Funding acquisition: *Jake Olivier*, Methodology, Formal Analysis, Writing – review and

editing, Funding acquisition *Lisa Keay,* Conceptualisation, Methodology, Writing – review and

editing, Funding acquisition; *Lynne E Bilston*' Conceptualisation, Methodology, Writing – review and

editing, Funding acquisition

7. Funding

This study was supported by Australian Research Council Discovery Project DP170103517, and funding from the NSW Centre for Road Safety. LEB is supported by an NHMRC Investigator Leadership grant.

8. Declaration of interest

The authors declare no competing interests.

9. References

1. Elliott MR, Kallan MJ, Durbin DR, Winston FK. Effectiveness of child safety seats vs seat belts in reducing risk for death in children in passenger vehicle crashes. Archives of pediatrics & adolescent medicine. 2006 Jun 1;160(6):617-21.

2. Du W, Finch CF, Hayen A, Bilston L, Brown J, Hatfield J. Relative benefits of population-level interventions targeting restraint-use in child car passengers. Pediatrics. 2010;125(2):304-12

3. Brown J, McCaskill ME, Henderson M, Bilston LE. Serious injury is associated with suboptimal restraint use in child motor vehicle occupants. Journal of paediatrics and child health. 2006 Jun;42(6):345-9.

4. Klinich KD, Benedetti M, Manary MA, Flannagan CA. Rating child passenger safety laws relative to best practice recommendations for occupant protection. Traffic injury prevention. 2017 May 19;18(4):406-11.

5. Sartin EB, Lombardi LR, Mirman JH. Systematic review of child passenger safety laws and their associations with child restraint system use, injuries and deaths. Injury prevention. 2021 Dec 1;27(6):577-81.

327 6. Brown J, Bilston LE. Child occupant protection in Australia. Journal of the Australasian College of

328 Road Safety. 2012 Jun;23(2):37-45.

329 7. Brown J, Hatfield J, Du W, Finch CF, Bilston LE. Population-level estimates of child restraint

330 practices among children aged 0–12 years in NSW, Australia. Accident Analysis & Prevention. 2010

331 Nov 1;42(6):2144-8.

332 8. Brown J, Keay L, Hunter K, Bilston LE, Simpson JM, Ivers R. Increase in best practice child car

333 restraint use for children aged 2–5 years in low socioeconomic areas after introduction of mandatory

334 child restraint laws. Australian and New Zealand journal of public health. 2013 Jun;37(3):272-7.

335 9. Koppel S, Charlton JL, Rudin-Brown CM. The impact of new legislation on child restraint system

336 (CRS) misuse and inappropriate use in Australia. Traffic injury prevention. 2013 Mar 25;14(4):387-96.

337 10.Korn, E.L., Graubard, B.I., 1999. Analysis of Health Surveys. Wiley, New York.

338 11. Lohr, S., 1999. Sampling: Design and Analysis. Duxbury Press, Pacific Grove

339 12. Brown J, Finch CF, Hatfield J, Bilston LE. Child restraint fitting stations reduce incorrect restraint

340 use among child occupants. Accident Analysis & Prevention. 2011 May 1;43(3):1128-33.

341 13. Greenwell, N. K. (2015, May). Results of the national child restraint use special study. (Report No.

342 DOT HS 812 142). Washington, DC: National Highway Traffic Safety Administration.

343 14. Parab A, Whyte T, Albanese B, Bilston L, Koppel S, Charlton JL, Olivier J, Keay L, Brown J. Can age

344 or height define appropriate thresholds for transition to adult seat belts? An analysis of observed

345 seat belt fit in children aged 7–12 years. International journal of environmental research and public

346 health. 2022 Jan 28;19(3):1524.

347 15. Whyte T, Albanese B, Elkington J, Bilston L, Brown J. Restraint factors and child passenger deaths

348 in New South Wales, Australia. International journal of environmental research and public health.

349 2020 Jan;17(4):1147.

352