1 The impact of social and environmental extremes on cholera time varying reproduction

2 number in Nigeria

4 Gina E C Charnley[1,2], Sebastian Yennan[3], Chinwe Ochu[3], Ilan Kelman[4,5,6], Katy A M

5 Gaythorpe[1,2], Kris A Murray[1,2,7]

1. Department of Infectious Disease Epidemiology, School of Public Health, Imperial College

London, London, UK

2. MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College

London, London, UK

3. Surveillance and Epidemiology Department/IM Cholera, Nigeria Centre for Disease Control,

Abuja, Nigeria

4. Institute for Risk and Disaster Reduction, University College London, London, UK

5. Institute for Global Health, University College London, London, UK

6. University of Agder, Kristiansand, Norway

7. MRC Unit The Gambia at London School of Hygiene and Tropical Medicine, Fajara, The

Gamiba

Correspondence to: Gina E C Charnley, g.charnley19@imperial.ac.uk

https://orcid.org/0000-0003-2087-7822

22 Abstract

Nigeria currently reports the second highest number of cholera cases in Africa, with numerous

socioeconomic and environmental risk factors. Less investigated are the role of extreme events,

despite recent work showing their potential importance. To address this gap, we estimated time

varying reproductive number (R) from cholera incidence in Nigeria and used a machine learning

approach to evaluate its association with extreme events (conflict, flood, drought) and pre-existing

vulnerabilities (poverty, sanitation, healthcare). We then created a traffic-light system for cholera

outbreak risk, using three hypothetical traffic-light scenarios (Red, Amber and Green) and used this

to predict R. The system highlighted potential extreme events and socioeconomic thresholds for

outbreaks to occur. We found that reducing poverty and increasing access to sanitation lessened

vulnerability to increased cholera risk caused by extreme events (monthly conflicts and the

Palmers Drought Severity Index). The work presented here shows the need for sustainable

development for disaster prevention and mitigation and to improve health and quality of life.

Introduction

Cholera was reintroduced into Africa in the 1970s during the seventh and continuing cholera

pandemic. It has since caused significant mortality and morbidity, especially amongst the most

vulnerable, such as children under five[1]. Despite this, other disease outbreaks have drawn

attention away from cholera in Africa in recent years, including COVID-19 and Ebola[2,3]. Explosive

cholera outbreaks are not uncommon due to the short incubation period (2 hours to 5 days) and

high numbers of asymptomatic infections, which when contaminating the environment can sustain

transmission[4]. Cholera is considered a disease of inequity and is preventable through wide-spread

access to safe drinking water and sanitation[5]. However, the effect of these pre-existing

vulnerabilities on disease risk can be exacerbated in times of environmental and social extremes,

which can in turn act as a catalyst for, or exacerbate the impacts of, outbreaks.

Previous research has found several links between extreme events and cholera including floods,

drought and conflict[6-8]. Disaster-related risk factors leading to disease outbreaks include an

inability to access routine care such as vaccination, fears over safety, destruction of infrastructure,

disruption of water, sanitation and hygiene (WASH) services and human displacement[9,10]. Previous

research on disaster-related infectious disease outbreaks have examined extreme events in

isolation[7,10], while others do not include multiple pre-existing socio-economic factors into the

methodology[11,12]. Research linking several social and environmental extremes to diseases,

including via risk factor cascades, is a global research gap and is important for predicting cholera

transmission and mitigating outbreaks[13].

Nigeria currently reports the second highest number of estimated cholera cases in Africa[1,14] and

has experienced many large outbreaks[15-18]. This is likely due to the presence of many underlying

social and environmental risk factors, including a favourable climate[19,20], poor access to WASH[21,22]

and a high proportion of the population living in poverty (62% at <$1.25/day)[23-25]. It also has a

relatively robust reporting system which may correlate with more cases, as cholera is an under-

reported disease and cases and deaths are often missed or misattributed. The country has been

frequently challenged by both social and environmental extremes such as drought and floods,

which may alter in intensity and frequency with climate change[13,24], along with ongoing conflict in

the northeastern region due to Boko Haram (Islamic State West Africa Province)[8,13].

Here, we aim to resolve the role of extreme events in causing or contributing to cholera and

increase the attention on cholera in Nigeria. In collaboration with the Nigeria Centre for Disease

Control (NCDC), we evaluated by way of machine learning how a range of environmental and

social covariates influence time-varying reproductive number (R) of cholera. Using the model with

the best predictive power, we predicted a traffic-light system of cholera risk to illustrate how

disasters and pre-existing vulnerabilities alter R and therefore the risk of cholera outbreaks. We

anticipate that this novel and relatively simple framework of cholera outbreak risks could be

employed by a range of professionals working in fragile settings by targeting interventions towards

key disaster-related risk factors.

Results

Incidence and R

In Nigeria, there were 837 and 564 confirmed cholera cases for 2018 and 2019, respectively. The

geographic distribution of confirmed cases is shown in Fig. 1a and are concentrated in the northeast

of the country, with Adamawa, Borno, Katsina and Yobe having the highest burden. The number of

cases declined steeply with age to a minimum in the 35-44 years category, before increasing again

84   over 45 years. Whereas, cases were relatively evenly split by sex overall, with slightly more males

85   affected in 2018 (51.6% male) and more females in 2019 (43.6% male) (Fig. 1b).

Fig. 1: Number of confirmed cholera cases. a, by state, grey indicates states that had no

reported confirmed cases and **b**, by sex and age group, all for 2018 and 2019.

87   Six states for 2018 and two states for 2019 had sufficient cases to be included for R calculations,

88   including Adamawa (2018 & 2019), Bauchi (2018), Borno (2018 & 2019), Gombe (2018), Katsina

89   (2018) and Yobe (2018). Both the R values and the incidence data used to calculate R are shown

90   temporally in Fig. 2 for each state and year. Some states appear to have a peak in transmission

91   around June-July, whereas others appear later during September to October.

Fig. 2: R values (line) calculated from the incidence (bar) of cholera. Data is for the

confirmed cholera cases for 2018 and 2019 of states which met the threshold equal to or

more than 40 cases.

Covariate Selection and Random Forest Models

Twenty-one covariates were included in the clustering and variable importance analyses and were

grouped into nine clusters. The clusters and variable importance (based on reducing node impurity)

of each covariate are shown in Fig. 3. Stepping through different covariate combinations, the best

fit model included number of monthly conflict events, Multidimensional Poverty Index (MPI),

Palmers Drought Severity Index (PDSI) and improved access to sanitation, fitted to R values with a

serial interval of 5 days (standard deviation: 8 days). The fit of the incidence-based vs covariate-

based R values (including error) are shown in Fig. 4 and had a correlation of 0.87, with the model

RMSE at 0.33 and R² of 0.32.

Fig. 3: The variable importance for the 21 tested for inclusion in the best fit model. All three

serial interval values tested are shown (Rt3 - 3 days, Rt5 - 5 days, Rt8 - 8 days) and the numbers

represent the clusters. Variable importance is measured through node impurity (see Methods for

details). SPEI01, 12, 48 - Standardised Precipitation Index calculated on 1, 12 and 48 month

scale. PDSI - Palmers Drought Severity Index. MPI - Multidimensional Poverty Index. IDP –

Internally Displaced Persons.

Fig. 4: Incidence-based vs covariate-based R values for the best fit model fitted to the testing dataset. The error bars show mean absolute error and the line is a linear trend line intercepting at 0.

Nowcasting

Using the best fit model, R was predicted for the remaining 31 states which did not have sufficient

cases to be included in the R calculations and any missing dates for the six states which were

included. This created estimates of R for all 37 states on a monthly temporal scale for 2018 and

2019. The predictions provide further evidence that the model accurately predicts R, as the higher

R values were in areas with known elevated cholera burden (northern and northeastern regions)

and the states which only marginally fell below the threshold for R calculations (Fig. 5).

Fig. 5: Average R values for 2018 and 2019 for all 37 Nigerian states. Incidence-based (green) - the five states which met the equal to or more than 40 case thresholds. Covariate-based (purple) - the 31 states which did not meet the threshold and had R predicted using the best fit model. State label colour shows which states had an average R of R = >1 (black) and R = <1 (orange).

Traffic-Light System for Cholera Outbreak Risk

Fig. 6 shows the predicted R values for the three traffic-light scenarios (Red = R over 1, Amber = R

around 1 and Green = R less than 1) of cholera outbreak risk, based on the four selected covariates.

Sanitation and MPI had a clear relationship with the R threshold, with consistently lower MPI (less

poverty) and a higher proportion of people with access to sanitation seeing lower R values. R

increased above 1 at 50% or lower for improved sanitation access and MPI values of above 0.32.

The historical average sanitation level for R = >1 was 52.8% for the full dataset, whereas for R <1

it was 61.2%, for MPI the mean values were 0.27 and 0.13 for R = >1 and R <1, respectively.

In contrast, monthly conflict events and PDSI shows a less defined relationship, with conflict having

a wide range of values in each of the three traffic-light scenarios. For PDSI and conflict, R values

increased above 1 at around -1.1 for PDSI and monthly conflict events of 1.6. The historical spatial

trends for conflict and PDSI are presented in Supplementary Figure 1 and shows polarity in the

relationships between the selected social and environmental extremes and R values, which differ

between states.

Fig. 6: Traffic-light system of cholera risk. The three traffic-light scenarios (Red = R over 1,

Amber = R around 1 and Green = R less than 1) for each of the four covariates in the best fit

model and the corresponding predicated R value using the best fit model.

Spatial Heterogeneities

Conflict

Borno and Kaduna were selected due to their clear positive relationship between conflict and R

(increased conflict and R = >1). The three traffic-light scenarios created for conflict in these two

states found a consistently high cholera outbreak risk. The Green traffic-light scenario was relatively

small, with only a narrow range of conflict values causing R values less than 1. Both Kaduna and

Borno have high levels of poverty and low access to sanitation (40-41% access). For Borno, raising

monthly conflict events from 1 to 2 increased R above 1, but an increase in access to sanitation

from 41-46% pushed the R value back below one. This relationship continued in a stepwise pattern

and in a similar way for MPI but to a lesser degree. This showed that increasing sanitation and

therefore decreasing vulnerability, allowed the states to adapt to increasing conflict and keep the R

value below 1 (See Supplementary Figure 2).

Drought

Four states were investigated to evaluate the differences between extreme wetness (Lagos and

Ekiti) and extreme dryness (Nasarawa and Kwara) and R values over 1. In contrast to Borno and

Kaduna, all four states predicted consistently low R values (Supplementary Figs. 3 & 4), a potential

explanation for this is the high variable importance of PDSI (Fig. 3) and the high levels of sanitation

and low levels of poverty in all four states, contributing to overall lower predicted levels of cholera.

Therefore, the model was detecting a signal in only small changes in PDSI, that resulted in changing

R values which have not been detected in other states with higher rates of poverty and lower levels

of sanitation access. It also helps to highlight the multi-directionality of the relationship between

PDSI and cholera transmission, with both extreme wetness and extreme dryness causing increases

in R.

Discussion

The results presented here show the importance of social and environmental extremes on cholera

outbreaks in Nigeria, along with the importance of underlying vulnerability and socioeconomic

factors. Of the 1,401 positive cases for Nigeria in 2018 and 2019, the northeast of the country and

children under 5 carried the highest burden of disease, whereas there was minimal differentiation

in cases between sex. Six states were used to calculate the R values, including Adamawa, Bauchi,

Borno, Gombe, Katsina and Yobe. Twenty-one covariates were considered for model inclusion and

the best fit model according to the selected model performance measures (variable importance

based on node impurity, RMSE, R^2 and correlations) included monthly conflict events, percentage

of the population with access to sanitation, MPI and PDSI. Using the best fit model, nowcasting was

used to calculate the R values for the remaining thirty-one states which did not meet the threshold.

The predicted R values from the three traffic-light scenarios helped to shed light on the thresholds

and triggers for raising R values above 1 in Nigeria. MPI and sanitation showed a well-defined

relationship with R, with consistently higher access to sanitation and less poverty (lower MPI value)

when R was less than 1. Thresholds which pushed R above one included decreasing access to

sanitation below 50% and increasing the MPI above 0.32. Whereas the relationship between R and

conflict events and PDSI appeared to vary spatially, with some states showing a negative and some

states a positive association. For these two covariates, the effect on R was largely dependent on

the access to sanitation and poverty within the states, with high levels of sanitation and low poverty

resulting in a decreased effect of PDSI and conflict. This showed that better sustainable

development in the state acted as a buffer to social and environmental extremes and allowed people

to adapt to these events better, due to less pre-existing vulnerability.

According to the World Bank[26], up to 47.3% (98 million people) of Nigeria's population live in

multidimensional poverty. Poverty is a well-known risk factor for cholera, which is considered a

disease of inequity[27]. Poverty can result in several risk factor cascades, which puts people at risk of

not just cholera but several other diseases. Examples of these risks include poor access to WASH[21],

inadequate housing[28], malnutrition[29] and overcrowding[30]. The expansion of sustainable

development helps to reduce these risks and meeting or exceeding the Sustainable Development

Goals would see significant gains in global health[31]. People living in poverty have fewer options and

abilities to adapt to new and extreme situations, becoming trapped in the affected area or displaced

to areas where their needs are not met. This provides further evidence for the need to reduce pre-

existing vulnerabilities and to implement known techniques for reducing disasters[32,33].

Poverty when measured in monetary terms alone can create issues due to its impact on the risk

factors stated and is an advantage of using the MPI as a poverty indicator. Nigeria's cash transfer

scheme has allowed many Nigerians to meet the household income limit for poverty but there is a

case for turning these funds and attention onto structural reform[34]. Nigeria's nationwide average

access to sanitation is around 25%, therefore using these funds to increase access to sanitation

may significantly improve health[35]. Currently, 73% of the enteric disease burden in Nigeria is

associated with inadequate WASH[36] and here we show the need for expansion of sanitation to

reduce cholera risks and the shocks of extremes on its transmission. In a recent review on the

implementation of non-pharmaceutical cholera interventions, there was generally a high acceptance

of several WASH interventions. Despite this, education was key and building community

relationships is needed to achieve this, such as understanding cultural differences and barriers[37].

This is especially important in areas with conflict, where trust between the government and residents

may have been lost[29].

Since 2002, Boko Haram (and Islamic State's West Africa Province) has been gaining a foothold

and territory in northeastern Nigeria which has resulted in ongoing conflict, unrest and oppression

of civilians[38]. Currently 5,860,200 people live in Borno state[39], where the fighting has been most

concentrated. Millions of people comprise conflict-affected populations globally and there is an

increasing proportion of people living in early post conflict areas[40]. This is significant in terms of

health and disease, as conflict has known risk factors for cholera along with several other

diseases[8,10,41] and can worsen several of the social risk factors discussed above. Here, conflict was

included in the best fit model and in some states, highly influential in terms of cholera transmission.

Providing services and protecting health in conflict zones is especially challenging and coordination

across organisations in reporting and operations are needed to streamline resources and prevent

duplication of services[42]. The traffic-light system used here helps highlight what is needed in these

situations to protect health and when outbreaks may occur.

PDSI and several of the other drought indices tested here showed high variable importance but, in

some states, had only marginal influence on R predictions when the PDSI values were manipulated.

When analysing spatial differences between R and PDSI, the relationship appears to be multi-

directional, with both extreme wetness (PDSI = +4) and extreme dryness (PDSI = -4) associated

with R values above 1. Furthermore, access to sanitation and poverty were important in how PDSI

impacted R, similar to the impacts of conflict. There is significant evidence to show that both

droughts[7,11] and floods[12,43] can cause cholera outbreaks and elevated transmission. Mechanisms

through which this can occur includes a lack of water increasing risky drinking water behaviour and

floods allowing for the dispersal of the pathogen. Nigeria has a varied climate across the country

and therefore both extremes are likely to be felt by those living there. Cholera outbreaks have been

seen in both the rainy and the dry season and the work presented here shows potential triggers for

when extra vigilance is needed, especially in certain states. This immediate insight is important,

while continually working to offset cholera risks from extremes through sanitation and hygiene,

which can take significant time and resources[44].

Despite adapting the methodology to account for this, a potential limitation may be lagged effects

of the covariates on cholera[45,46]. Both long-term and short-term changes to the population may take

time before changes in cholera transmission are evident. While some influences may be considered

slow-onset or rapid-onset and therefore defining their beginning is subjective. Despite this, the

incubation period of cholera is short (<2 hours - 5 days) and previous research has suggested that

acute impacts cause increases in cholera cases within the first week of the event[47-49]. Calculating

R on monthly sliding windows and using monthly covariate data helped to reduce potential lagged

effects on the R values, which would be captured if the one-week lag estimate is applicable here.

Although beyond the scope of the research presented here, the impacts of different lagged periods

for several of these covariates and cholera outbreaks is an essential area of future research.

Cholera is considered an under-reported disease, and the lack of symptomatic cases means that

many are likely to be missed. There are also incentives not to report cholera cases, due to travel

restrictions and isolations and implications for trade and tourism[50]. However, the robust reporting

system in Nigeria suggests that the data used here is the best available for analysis. While during

times of crisis, cholera may be over-reported or more accurately represent the cholera burden in

the area. This is due to the presence of cholera treatment centres, increased awareness among the

population and external assistance from non-governmental organization, detecting cases that may

have been missed previously[8].

The Global Task Force on Cholera Control's 2030 target of reducing cholera deaths by 90%[51] will

require acceleration of current efforts and significant commitment. Increasing cholera research and

data are important in achieving this and the traffic-light system for cholera risk presented here sheds

light on ways to reduce cholera outbreaks in fragile settings. The results highlight the importance of

extreme events on cholera transmission and how reducing pre-existing vulnerability could offset the

resultant cholera risk. This research is the first time several disaster types and measures of

population vulnerability have been evaluated together quantitatively in terms of cholera. We hope it

shows the importance of doing so to gain a more accurate understanding of disease outbreaks in

complex emergencies. Nigeria is currently working towards its ambitious goal of lifting 100 million

people out of poverty by 2030[34]. If it is successful, this could significantly improve health, increase

quality of life and decrease the risks of social and environmental extremes.

Methods

Datasets

Cholera data were obtained from NCDC and contained linelist data for 2018 and 2019. The data

were age and sex-disaggregated, on a daily temporal scale and to administrative level 4. The data

also provided information on the outcome of infection and whether the patient was hospitalised.

The data were subset to only include cases that were confirmed either by rapid diagnostic tests or

by laboratory culture and only these confirmed cases were used in the analyses.

A range of covariates were investigated based on previously understood cholera risk factors.

Covariates included factors related to conflict (monthly, daily)[52], drought (Palmers Drought Severity

Index, Standardised Precipitation Index)[53,54], internally displaced persons (IDPs) (households,

individuals)[55], WASH (improved drinking water, piped water, improved sanitation, open defecation,

basic hygiene)[56], healthcare (total facilities, facilities per 100,000 people)[52], population (total)[57] and

poverty (MPI, headcount ratio in poverty, intensity of deprivation among the poor, severe poverty

and population vulnerable to poverty)[52].

Covariate data were on a range of spatial and temporal scales, therefore administrative level one

(state) was set as the spatial granularity (data on a finer spatial scale were attributed to

administrative level 1) and the finest temporal scale possible (daily) was used for covariate

selection (repeating values if data were not available at the daily level). The datasets and methods

used here were approved by Imperial College Research Ethics Committee and a data sharing

agreement between NCDC and the authors.

Incidence and R

The 2018 and 2019 laboratory confirmed linelist data were used to calculate incidence. Incidence

was calculated on a daily scale by taking the sum of the cases reported by state and date of onset

of symptoms. This created a new dataset with a list of dates and corresponding daily incidence for

each state. All analysis was completed in R with R Studio version 4.1.0. (packages "incidence"[58] &

"EpiEstim"[59]).

Rather than using incidence as the outcome variable (which has less implicit assumptions), R was

calculated, as it is more descriptive providing information on epidemic evolution (e.g., R = >1, cases

are increasing), instead of new reported disease cases for a single time point. R was calculated

from incidence using the parametric standard interval method, which uses the mean and the

standard deviation of the standard interval (SI). SI is the time from illness onset in the primary case

to onset in the secondary case and therefore impacts the evolution of the epidemic and speed of

transmission. The SI for cholera is well-documented and there are several estimates in the

literature[60-62]. To account for this reported variation in SI, a sensitivity analysis was conducted with

SI set at 3, 5 and 8 days with a standard deviation of 8 days. The parametric method was used (vs

the non-parametric which uses a discrete distribution), as this can be adequately modelled by

a normal probability distribution and has a fixed set of parameters.

Estimating R too early in an epidemic increases error, as R calculations are less accurate when

there is lower incidence over a time window. A way to understand how much this impacts R values

is to use the coefficient of variation (CV), which is a measure of how spread out the dataset values

are relative to the mean. The lower the value, the lower the degree of variation in the data. A

coefficient of variation threshold was set to 0.3 (or less) as standard, based on previous work[59]. To

reach the CV threshold, calculation start date for each state was altered until the threshold CV was

reached. States with <40 cases were removed, as states with fewer cases did not have high enough

incidence across the time window to reach the CV threshold. Additionally, R values were calculated

over monthly sliding windows, to ensure sufficient cases were available for analysis within the time

window.

Covariate Selection and Random Forest Models

Supervised machine learning algorithms such as decision-tree based algorithms, are now a widely

used method for predicting disease outcomes and risk mapping[63,64]. They work by choosing data

points randomly from a training set and building a decision tree to predict the expected value given

the attributes of these points. Transparency is increased by allowing the number of trees

(estimators), number of features at each node split and resampling method to be specified. Random

Forests (RF) then combines several decision trees into one model, which has been shown to

increase predictive accuracy over single tree approaches, while also dealing well with interactions

and non-linear relationships[65,66].

The covariates listed above (conflict, drought, IDPs, WASH, healthcare, population and poverty)

were first clustered to assist in the selection of covariates for model inclusion and to understand any

multicollinearities. Despite RF automatically reducing correlation through subsetting data and tuning

the number of trees and depth[64,67], the process here lends support that the final model is measuring

somewhat independent processes and not purely overfitting the same patterns[63]. The clustering

was based on the correction between the covariates meeting an absolute pairwise correlation of

above 0.75. A secondary covariate selection process was run during preliminary analysis and acted

as a method of validation. The process is detailed in Supplementary Information 1.

Random forest variable importance was used to rank all 21 clustered covariates. Variable

importance provided an additional method of guiding the fitting of the best fit model, by testing the

covariates which found the highest variable importance first. In this context, variable importance is

a measure of the cumulative decreasing mean standard error each time a variable is used as a

node split in a tree. The remaining error left in predictive accuracy after a node split is known as

node impurity and a variable which reduces this impurity is considered more important.

Training (70% of data) and testing (30%) datasets were created to train the model and test the

model's predictive performance. Random forest regression models (as opposed to classification

models) were used since the outcome variable (R) is continuous. The parameters for training were

set to repeated cross-validation for the resampling method, with ten resampling interactions and

five complete sets of folds to complete. The model was tuned and estimated an optimal number of

predictors at each split of 2, based on the lowest out-of-bag (OOB) error rate with RMSE used as

the evaluation metric (package "caret"[68]).

A stepwise analysis was used to fit the models under each SI condition (3, 5 & 8 days), taking into

consideration the covariate clustering and variable importance. One covariate was selected from

each cluster, and all combinations of covariates were tested until the best-fit model was found.

Models were assessed against each other in terms of predictive accuracy, based upon R^2 and

RMSE. Predictions were then calculated on the testing dataset to compare incidence-based (R

values calculated using the incidence data) vs covariate-based R values (R values calculated

through model predictions). The terms, actual vs predicted was not used here, as all R values

were modelled making the term "actual" misleading in this context. Model performance

evaluations were built on multiple metrics including correlation, R^2 and RMSE.

Despite random forest models being accurate and powerful for prediction, they are easily over-fit

(fitting to the testing dataset too closely or exactly) and therefore calculating error for the

predictions are important. Little to no error in the predictions are an indication of over-fitting which

can occur through predictions based off too small a dataset, more parameters than can be

justified by the data and multicollinearity. Here, error was calculated using mean absolute error

(MAE), where y_i is the prediction and x_i is the true value, with the total number of data points as n.

$$MAE = \frac{\sum_{i=1}^{n}|y_i - x_i|}{n}$$

Nowcasting

The best fit model, in terms of predictive power according to the metrics above, was used to predict

R for the remaining states which did not have sufficient reported cases to calculate R using

incidence or had missing data for certain dates. Data for the best fit model covariates were collected

for the states and missing dates from the sources given above. The data for the selected covariates

are shown spatially in Supplementary Figure 5.

Traffic-Light System for Cholera Outbreak Risk

The best fit model was then used to predict the traffic-light system for cholera outbreak risk, by

manipulating the covariates values and using these to predict R. The traffic light system was defined

by:

• Red - Covariate values which pushed R over 1

• Amber - Covariates values with predicted R around 1

• Green - Covariate values which predicted R below 1.

By using these three traffic-light scenarios, cholera outbreak triggers were identified based on the

conditions of the four selected covariates. No specific R value had to be met for each traffic-light

scenario, to account for the complexity of the relationships and non-linearity (Supplementary Figs.

6 & 7). To illustrate the historical trends between the best fit model covariates and the R thresholds

(R = >1, R <1), the data is split both spatially (by month) and temporally (by state) in Supplementary

Figs. 1 & 8.

Spatial Heterogeneities

To understand spatial differences in the relationship between the selected social and environmental

extremes (conflict and PDSI) and cholera outbreak risk and the role pre-exiting vulnerabilities played

in altering these relationships, six states were selected for additional analysis. These states were

selected because they had either a clear positive or clear negative relationship with conflict or PDSI

and R (PDSI is hypothesised to increase R at either end of the scale, +4/-4) and included Borno,

Kaduna, Nasarawa, Ekiti, Lagos and Kwara (see Supplementary Figure 1). The processes above

for predicting R under the three traffic-light scenarios was repeated for the six states but only PDSI

and conflict values were manipulated, keeping the other three covariates at the mean value for R =

>1 across the full dataset for the state. The spatial analyses identified the thresholds in conflict and

PDSI needed to push R values below 1.

References

1. Ali, M., Nelson, A. R., Lopez, A. L. & Sack, D. A. Updated global burden of cholera in endemic

countries. *PLoS Neglect. Trop. Dis.* **9**, e0003832 (2015).

2. Carter, S. E. et al. What questions we should be asking about COVID-19 in humanitarian

settings: perspectives from the social sciences analysis cell in the Democratic Republic of the

Congo. *BMJ Glob. Health.* **5**, e003607 (2020).

3. Musa, S. S. et al. Dual tension as Nigeria battles cholera during the COVID-19 pandemic. *Clin.*

Epidemiology Glob Health. **12** (2021).

4. King, A. A., Ionides, E. L., Pascual, M. & Bouma, M. J. Inapparent infections and cholera

dynamics. *Nature* **454**, 877-880 (2008).

5. Anbarci, N., Escaleras, M. & Register, C. A. From cholera outbreaks to pandemics: the role of

poverty and inequality. *Working Paper 05003* (Florida Atlantic University, FL, 2006).

6. Elimian, K.O. et al. Descriptive epidemiology of cholera outbreak in Nigeria, January–November,

2018: implications for the global roadmap strategy. *BMC Public Health* **19**, 1-11 (2019).

7. Charnley, G. E. C., Kelman, I., Green, N., Hinsley, W., Gaythorpe, K. A. M. & Murray, K. A.

Exploring relationships between drought and epidemic cholera in Africa using generalised linear

models. *BMC Infect. Dis*. **21,** 1-12 (2021).

8. Charnley, G. E. C., Jean, K., Kelman, I., Gaythorpe, K. A. M. & Murray, K. A. Using self-

controlled case series to understand the relationship between conflict and cholera in Nigeria and

the Democratic Republic of Congo. Preprint at https://doi.org/10.1101/2021.10.19.21265191

(2021).

9. Charnley, G. E. C., Kelman, I., Gaythorpe, K. A. M. & Murray, K. A. Traits and risk factors of

post-disaster infectious disease outbreaks: a systematic review. *Sci. Rep*. **11**, 1-4 (2021).

10. Wells, C. R. et al. The exacerbation of Ebola outbreaks by conflict in the Democratic Republic

of the Congo. *PNAS*. **116**, 24366-72 (2019).

11. Rieckmann, A., Tamason, C. C., Gurley, E. S., Rod, N. H. & Jensen, P. K. Exploring droughts

and floods and their association with cholera o20utbreaks in sub-Saharan Africa: a register-

based ecological study from 1990 to 2010. *Am. J. Trop. Med. Hyg*. **98**, 1269 (2018).

12. Jutla, A. et al. Environmental factors influencing epidemic cholera. *Am. J. Trop. Med. Hyg*. **89**,

597 (2013).

13. Elimian, K. O. et al. What are the drivers of recurrent cholera transmission in Nigeria? Evidence

from a scoping review. *BMC Public Health*. **20**, 1-3 (2020).

14. Lessler, J. et al. Mapping the burden of cholera in sub-Saharan Africa and implications for control: an analysis of data across geographical scales. *Lancet* **391**, 1908-1915 (2018).

15. Dalhat, M. M. et al. Descriptive characterization of the 2010 cholera outbreak in Nigeria. *BMC Public Health*. **14**, 1-7 (2014).

16. Ngwa, M. C. et al. The multi-sectorial emergency response to a cholera outbreak in internally displaced persons camps in Borno state, Nigeria, 2017. *BMJ Glob. Health*. **5**, e002000 (2020).

17. Sule, I. B., Yahaya, M., Aisha, A. A., Zainab, A. D., Ummulkhulthum, B. & Nguku, P. Descriptive epidemiology of a cholera outbreak in Kaduna State, Northwest Nigeria, 2014. *Pan Afr. Med. J.* **27**, (2017).

18. Adeneye, A. K. et al. Risk factors associated with cholera outbreak in Bauchi and Gombe States in North East Nigeria. *J. Public Health Epidemiol*. **8**, 286-296 (2016).

19. De Magny, G. C., Guégan, J. F., Petit, M. & Cazelles, B. Regional-scale climate-variability synchrony of cholera epidemics in West Africa. *BMC Infect. Dis.* **7**, 1-9 (2007).

20. Abdussalam, A. F. Modelling the climatic drivers of cholera dynamics in Northern Nigeria using generalised additive models. *Int. J. Geogr. Environ. Manage*. **2**, 84-97 (2016).

21. Gidado, S. et al. Cholera outbreak in a naïve rural community in Northern Nigeria: the importance of hand washing with soap, September 2010. *Pan Afr. Med. J.* **30** (2018).

22. Hutin, Y., Luby, S., Paquet, C. A large cholera outbreak in Kano City, Nigeria: the importance of hand washing with soap and the danger of street-vended water. *J. Water Health*. **1**, 45-52 (2003).

23. Dan-Nwafor, C. C. et al. A cholera outbreak in a rural north central Nigerian community: an unmatched case-control study. *BMC Public Health* **19**, 1-7 (2019).

24. Leckebusch, G. C. & Abdussalam, A. F. Climate and socioeconomic influences on interannual variability of cholera in Nigeria. *Health Place*. **34**, 107-17 (2015).

25. United Nations Statistical Division. Millennium Development Goal Indicators. https://unstats.un.org/unsd/mdg/SeriesDetail.aspx?srid=580 (2015).

26. World Bank. Tackling poverty in multiple dimensions: A proving ground in Nigeria.

https://blogs.worldbank.org/opendata/tackling-poverty-multiple-dimensions-proving-ground-

nigeria (2021).

27. Talavera, A. & Perez, E. M. Is cholera disease associated with poverty?. *J. Infect. in Dev.*

Countr. **3**, 408-11 (2009).

28. Penrose, K., Castro, M. C., Werema, J. & Ryan, E. T. Informal urban settlements and cholera

risk in Dar es Salaam, Tanzania. *PLoS Neglect. Trop. Dis.* **4**, e631 (2010).

29. Charnley, G. E. C., Kelman, I. & Murray, K. A. Drought-related cholera outbreaks in Africa and

the implications for climate change: a narrative review. *Pathog. Glob. Health.* 1-10 (2021).

30. Ververs. M. & Narra, R. Treating cholera in severely malnourished children in the Horn of Africa

and Yemen. *Lancet.* **390**, 1945-6 (2017).

31. von Schirnding Y. Health and sustainable development: can we rise to the challenge?. *Lancet.*

360, 632-7 (2002).

32. Masozera, M., Bailey, M. & Kerchner, C. Distribution of impacts of natural disasters across

income groups: A case study of New Orleans. *Ecol. Econ.* **63,** 299-306 (2007).

33. Lahsen, M. & Ribot, J. Politics of attributing extreme events and disasters to climate change.

Wiley Interdiscip. Rev. Clim. Change. **13**, e750 (2022).

34. Onyeiwu, S. *Nigeria's poverty profile is grim. It's time to move beyond handouts.*

https://theconversation.com/nigerias-poverty-profile-is-grim-its-time-to-move-beyond-handouts-

163302 (2021).

35. Ajisegiri, B. et al. Geo-spatial modeling of access to water and sanitation in Nigeria. *J. Water*

Sanit. Hyg. Dev. **9**, 258-80 (2019).

36. World Bank Group. A Wake Up Call: Nigeria Water Supply, Sanitation, and Hygiene Poverty

Diagnostic. World Bank; 2017 Aug.

37. Polonsky, J. A. et al. Feasibility, acceptability, and effectiveness of non-pharmaceutical

interventions against infectious diseases among crisis-affected populations: a scoping review.

Infect. Dis. Poverty. **11**, 1-9 (2022).

38. Falode, J. A. The nature of Nigeria's Boko Haram war, 2010-2015: A strategic analysis.

Perspect. Terror. **10**, 41-52 (2016).

39. Borno State Government. Population. https://bornostate.gov.ng/population/ (2016).

40. Garfield, R. M., Polonsky, J. & Burkle, F. M. Changes in size of populations and level of conflict

since World War II: implications for health and health services. *Disaster Med Public Health Prep*.

6, 241-6 (2012).

41. Federspiel F, Ali M. The cholera outbreak in Yemen: lessons learned and way forward. BMC

public health. 2018 Dec;18(1):1-8.

42. Ricau, M., Lacan, L., Ihemezue, E., Lantagne, D. & String, G. Evaluation of monitoring tools

for WASH response in a cholera outbreak in northeast Nigeria. *J. Water Sanit. Hyg. Dev.* **11**,

972-82 (2021).

43. Sidley, P. Floods in southern Africa result in cholera outbreak and displacement. *BMJ* **336**, 471

(2008).

44. Onwe, F. I., Agu, A. P., Umezuruike, D. & Ogbonna, C. Factors responsible for the 2015

Cholera outbreak and spread in Ebonyi state, Nigeria. *J. Epidemiol. Soc. Nigeria*. **2**, 53-58

(2018).

45. Reyburn, R., Kim, D. R., Emch, M., Khatib, A., Von Seidlein, L. & Ali, M. Climate variability and

the outbreaks of cholera in Zanzibar, East Africa: a time series analysis. *Am. J. Trop. Med. Hyg*.

84, 862 (2011).

46. Emch, M., Feldacker, C., Yunus, M., Streatfield, P. K., DinhThiem, V. & Ali, M. Local

environmental predictors of cholera in Bangladesh and Vietnam. *Am. J. Trop Med. Hyg.* **78**, 823-

32 (2008).

47. Fredrick, T. et al. Cholera outbreak linked with lack of safe water supply following a tropical

cyclone in Pondicherry, India, 2012. *J. Health. Popul. Nutr*. **33**, 31 (2015).

48. Bhunia, R. & Ghosh, S. Waterborne cholera outbreak following cyclone Aila in Sundarban area

of West Bengal, India, 2009. *Trans R Soc Trop*. **105**, 214-219 (2011).

49. Jeandron, A. et al. Water supply interruptions and suspected cholera incidence: a time-series

regression in the Democratic Republic of the Congo. *PLoS Med.* **12**, e1001893 (2015).

50. Ganesan, D., Gupta, S. S. & Legros, D. Cholera surveillance and estimation of burden of

cholera. *Vaccine* **38**, A13-7 (2020).

51. Global Task Force on Cholera Control. Roadmap 2030. https://www.gtfcc.org/about-

gtfcc/roadmap-2030/ (2020).

52. HDX. The Humanitarian Data Exchange. https://data.humdata.org (2021).

53. University of East Anglia. Climate Research Unit. https://www.uea.ac.uk/groups-and-

centres/climatic-research-unit (2020).

54. CEDA. High resolution Standardized Precipitation Evapotranspiration Index (SPEI) dataset for

Africa. https://catalogue.ceda.ac.uk/uuid/bbdfd09a04304158b366777eba0d2aeb (2019).

55. IOM. DTM Nigeria. https://displacement.iom.int/nigeria (2021).

56. JMP. Nigeria. https://washdata.org (2020).

57. WorldBank.   Data   Bank   Subnational   Population.

https://databank.worldbank.org/source/subnational-population (2021).

58. Kamvar, Z. N., Cai, J., Pulliam, J. R. C., Schumacher, J. & Jombart, T. Epidemic curves made

easy using the R package incidence https://doi.org/10.12688/f1000research.18002.1 (2019).

59. Cori, A. EpiEstim: Estimate Time Varying Reproduction Numbers from Epidemic Curves. R

package version 2.2-4. https://CRAN.R-project.org/package=EpiEstim (2021).

60. Azman, A. S. et al. Urban cholera transmission hotspots and their implications for reactive

vaccination: evidence from Bissau city, Guinea bissau. *PLoS Neglect Trop. Dis.* **6**, e1901 (2012).

61. Azman, A. S. et al. Population-level effect of cholera vaccine on displaced populations, South

Sudan, 2014. *Emerg. Infect. Dis.* **22**, 1067 (2016).

62. Kahn, R. et al. Incubation periods impact the spatial predictability of cholera and Ebola outbreaks

in Sierra Leone. *PNAS.* **117**, 5067-73 (2020).

63. Hamlet A, Ramos DG, Gaythorpe KA, Romano AP, Garske T, Ferguson NM. Seasonality of

agricultural exposure as an important predictor of seasonal yellow fever spillover in Brazil. *Nat.*

Commun. **12**, 1-1 (2021).

64. Kapwata T, Gebreslasie MT. Random forest variable selection in spatial malaria transmission

modelling in Mpumalanga Province, South Africa. *Geospat. Health*. **11**, 251-262 (2016).

65. Breiman, L. Random forests. *Mach. Learn.* **45**, 5-32 (2001).

66. Biau, G. Analysis of a random forests model. *J. Mach. Learn. Res.* **13**, 1063-95 (2012).

67. Genuer R, Poggi JM, Tuleau-Malot C. Variable selection using random forests. *Pattern*

Recognit. Lett. **31**, 2225-36 (2010).

68. Kuhn, M. caret: Classification and Regression Training. https://CRAN.R-

project.org/package=caret (2021).

Acknowledgements

We would like to thank and acknowledgement the Nigeria Centre for Disease Control for providing

the data used here and those who work for the NCDC who collected the data in the field. We would

also like to thank Anwar Musah (University College London) and Kelly Elimian (Karolinska Institutet)

for their guidance on cholera data for Nigeria and facilitating the partnership with NCDC. This work

was supported by the Natural Environmental Research Council [NE/S007415/1], as part of the

Grantham Institute for Climate Change and the Environment's (Imperial College London) Science

and Solutions for a Changing Planet Doctoral Training Partnership. We also acknowledge joint

Centre funding from the UK Medical Research Council and Department for International

Development [MR/R0156600/1].

Author Contributions

GECC was part of the study design and conceptualisation of ideas, ran the analysis, wrote and

finalised the manuscript and incorporated any feedback. SY & CO provided the cholera datasets,

facilitated the data sharing agreement and provided expertise on cholera in Nigeria. IK offered

expertise on disasters and health, provided expertise in the methodology and revised several

drafts. KAMG was part of the study design and conceptualisation of ideas, provided expertise in

the methodology, provided supervision and revised several drafts. KAM was part of the study

design and conceptualisation of ideas, provided expertise in the methodology, provided

supervision and revised several drafts. All authors have read and approved the manuscript.

Competing interests

The authors declare no competing interests.

Supplementary Material

Supplementary Information 1: Additional covariate selection using linear regression
The same 21 covariates (conflict, drought IDPs, WASH, healthcare, population and poverty) analysed using variable importance were also run through an additional covariate selection process and stepwise analysis as developed by:

1. Garske, T. et al. Yellow fever in Africa: estimating the burden of disease and impact of mass vaccination from outbreak and serological data. *PLoS Med.* **11**, e1001638 (2014).

2. Gaythorpe, K. A. M. et al. The global burden of yellow fever. *Elife* **10**, e64670 (2021).

The selection process removes covariates that are not significantly associated with the outcome variable (Rt3, Rt5, Rt8) at p = <0.1 using linear regression. It then clusters the remaining covariates based on the correction between them at an absolute pairwise correlation of above 0.75.

Ten were removed, either because they were not significantly associated with the outcome variable (R) or because they were too highly correlated with other covariates (healthcare facilities, piped water, open defecation, population, IDPs, severe poverty, vulnerable to poverty, basic hygiene). Eleven covariates remained and were grouped into five clusters, the clusters and variable importance of each covariate are shown below

The variable importance for the eleven remaining covariates after variable selection. All three serial interval values tested are shown (Rt3 - 3 days, Rt5 - 5 days, Rt8 - 8 days) and the numbers represent the clusters. SPEI01, 12, 48 - Standardised Precipitation Index calculated on 1, 12 and 48 month scale. PDSI - Palmers Drought Severity Index. MPI - Multidimensional Poverty Index.

Supplementary Figure 1: Historical spatial trends between the selected social and environmental extremes (conflict and PDSI) and the R thresholds (R = >1, R <1). The mean and standard error for the two covariates for the full dataset split by state and R threshold. The red "x" shows the states which were included in the sub-national analysis: Conflict (Borno and Kaduna), extreme wetness (Lagos and Ekiti), extreme dryness (Nasarawa and Kwara).

Supplementary Figure 2: Three traffic-light scenarios for conflict only and the corresponding predicted R values. The other three (PDSI, Sanitation and MPI) covariate values were retained at the mean value for R = >1 for the full dataset (values shown in the plot) for **a**, Borno and **b**, Kaduna.

Supplementary Figure 3: Three traffic-light scenarios for PDSI (drier conditions) only and the corresponding predicted R values. The other three (Conflict, Sanitation and MPI) covariate values were retained at the mean value for R = >1 for the full dataset (values shown in the plot) for **a**, Kwara and **b**, Nasarawa.

Supplementary Figure 4: Three traffic-light scenarios for PDSI (wetter conditions) only and the corresponding predicted R values. The other three (Conflict, Sanitation and MPI) covariate values were retained at the mean value for R = >1 for the full dataset (values shown in the plot) for **a**, Ekiti and **b**, Lagos.

Supplementary Figure 5: Average values of the four covariates included in the best fit model. By state, covariates included: **a**, monthly conflict events, **b**, Palmers Drought Severity Index (PDSI), **c**, percentage access to sanitation and **d**, Multidimensional Poverty Index (MPI).

Supplementary Figure 6: Single predictor partial dependency plots for the covariates in the best fit model. Showing the relationships between **a**, monthly conflict events, **b**, access to sanitation, **c**, Palmers Drought Severity Index (PDSI) and **d**, Multidimensional poverty Index (MPI) and R.

Supplementary Figure 7: Multi predictor partial dependency plots for the covariates in the best fit model. Showing the relationships between **a**, Palmers Drought Severity Index (PDSI) & Multidimensional poverty Index (MPI), **b**, PDSI & Sanitation, **c**, Monthly conflict & MPI, **d**, Monthly conflict & Sanitation, **e**, Sanitation & MPI and R.

Supplementary Figure 8: Historical temporal trends between the best fit model covariates and the R thresholds (R = >1, R <1). The mean and standard error for the four covariates included in the best fit model for the full dataset split by month and R threshold.