Title Page

Full title: Age-stratified normative cognitive scores in urban and rural Maharashtra — a

community-based study.

Short title: Community-based study of cognitive scores in Maharashtra

Authors:

1. Dr Uma Sundar[1], MD.   Email: umasundar2@gmail.com.

2. Dr Amita Mukhopadhyay[2], MD.   Email: dr.amukho@gmail.com.

3. Dr Suresh Sundar[3], MD.   Email: sundarhcca@hotmail.com.

4. Dr Nilesh Shah[4], MD.   Email: drnilshah@hotmail.com

Author Affiliations (as per superscripted numbers above)**:**

1. Professor, Medicine. Lokmanya Tilak Municipal Medical College and General Hospital

(LTMMC&GH), Sion, Mumbai 400022, Maharashtra, India.

2. Associate Professor, Community Medicine. Dr Chandramma Dayananda Sagar Institute

of Medical Education and Research, Kanakapura 562112, Ramanagara District,

Karnataka, India.

3. Professor, Community Medicine. BKL Walawalkar Rural Medical College, Chiplun

415606, Ratnagiri District, Maharashtra, India.

4. Professor, Psychiatry. LTMMC&GH, Mumbai, India.

Corresponding Author: Dr Amita Mukhopadhyay, dr.amukho@gmail.com

Total number of tables and figures: Tables 8; Figures 0.

Word Count: Abstract 199; Text 2969.

Keywords: Neuro-epidemiology, Cognitive testing, Kolkata Cognitive Battery, Geriatric

Depression Scale, Rural-urban cognition gap

Ethical approval: The study was approved by the Human Research Institutional Ethics

Committee, Lokmanya Tilak Municipal Medical College and General Hospital, Sion, Mumbai.

Financial support: This study was part of a larger study on "Age-stratified normative cognitive

scores in the community and their correlation with MRI volumetric studies", fully funded by

Cognitive science research initiative 2012, Department of Science and Technology, Government

of India.

Conflict of interest: Nil

47   **Age-stratified normative cognitive scores in urban and rural Maharashtra — a**

48   **community-based study.**

Abstract

Background and objectives: Indian ethnic and educational diversities necessitate obtaining

normative cognitive data in different populations. We aimed to evaluate cognitive scores using a

Marathi translation of the Kolkata Cognitive Battery (KCB), and to study the association of KCB

components with depression and socio-demographic variables.

Methodology: We studied 2651 individuals aged ≥40 years, without pre-existing

neuropsychiatric conditions, from urban (Mumbai) and rural districts of Maharashtra. For each

component of KCB, the lowest 10th percentile score was used to define cognitive impairment.

Results: We studied 1435 (54%) rural and 1216 (46%) urban residents equally divided by

gender (1316 women, 1335 men), average age 54 years. KCB scores were significantly lower

with female sex, older age, illiteracy and depression. The largest effect sizes attributable to these

factors were in the domains of calculation (gender), visuo-constructional ability (VCA; rurality)

and verbal fluency (VF; depression). Scores remained significantly lower in rural residents after

controlling for age, sex and education, particularly for VCA, immediate recall and calculation.

Conclusion: This Marathi KCB having been validated on large urban as well as rural samples,

may be used to study cognition in Marathi speaking populations with appropriate cut-offs

tailored to the degree of urbanization of the population.

Keywords

Neuro-epidemiology, Cognitive testing, Kolkata Cognitive Battery, Geriatric Depression Scale,

Rural-urban cognition gap

71
72

Age-stratified normative cognitive scores in urban and rural Maharashtra — a community-based study

Background

Community based studies on cognition in India, especially in rural areas, are few. Ethnic, socio-cultural, educational and linguistic diversities in India make it necessary to obtain normative data in cognition in different populations. Tripathi et al have reported objective cognitive dysfunction in the areas of "orientation" and "concentration" and "functioning/self-care" in normally aging older adults in urban Lucknow.[1] Gambhir et al, in a rural study of elderly, found a Dementia prevalence of 2.7% and suggested that Hindi Mental State Examination (HMSE) score of 17 should be used as cutoff for the illiterate population.[2] Ganguli et al described a battery of cognitive tests that was developed for screening an illiterate elderly rural Hindi-speaking population, based on their research conducted in Ballabhgarh district in Haryana.[3] Further, Banerjee et al[4] and Das et al[5] have obtained normative data for cognitive functions in a Bengali urban population utilizing the Kolkata Cognitive Screening Battery (KCB).

The mini mental state examination (MMSE) and the Hindi mental state examination (HMSE), although popular and easy to administer, suffer from a lack of sensitivity to early dementia, especially of the subcortical variety.[6] We aimed to evaluate cognitive scores using the KCB, in an age-stratified manner, in Maharashtrian rural and urban communities, and evolve cut-off scores based on individual test scores within the KCB, as a step towards refining its sensitivity and specificity as a screening instrument to detect cognitive impairment in the specific population under study. Further, we aimed to compare differences in individual test scores classed according to different socio-demographic variables as well as Geriatric Depression Scale (GDS)[7] scores and finally, compare our sample's scores with those of the Kolkata and Ballabhgarh study populations.

Methodology

Study design

The KCB has been validated in urban populations by Das et al,[5] and the Hindi cognitive battery from which it was derived, has been validated in the rural setting by Ganguli et al.[3] In the first part of this study, the partially modified Hindi cognitive screening battery used by Ganguli et al was translated into Marathi and again back - translated to Hindi by two independent bilingual translators. The GDS was also translated into Marathi and back- translated into Hindi. Field testing on 40 persons was done initially, for both KCB and GDS, to ensure inter and intra investigator reliability, being scored by the chief investigator and a qualified psychologist each time. Intra- rater reliability was 1, and inter-rater reliability was over 0.9 for both KCB and GDS. Qualified psychologists and final year residents in psychiatry training were trained in administration of the translated KCB and GDS test by the chief investigator and a consultant psychiatrist. The translated test battery has been appended as the Marathi Kolkata Cognitive Battery (KCB) screening (Appendix 1).

In the second part, a cross sectional interview-based survey was conducted in urban, semi urban and rural areas of Maharashtra.

Setting

The study was conducted in the villages of Vaitarna (Vadataluka, Thane district, population 671 as per 2011 census), Birwadi (Shahapurtaluka, Thane district, population 3415), and Penn taluka (Raigad district, population 37,852) over a period of 16 months from March 2015 to July 2016. These areas were chosen considering that they are the setting of ongoing primary health care projects under the community health department of the parent hospital, which made it feasible to obtain information on household numbers, size and the number of people over 40 years of age in these areas. The urban areas in Mumbai included Kumbharwada in Dharavi, Kala Chowki and Antop Hill areas.

Data collection

All persons aged 40 and over and residing in the study areas were invited to take part in the

survey. We excluded persons with known mental retardation, previous stroke, any known

psychiatric illness, or those reported by family to have previously diagnosed dementia or

significant memory impairment impeding daily activities. Healthy individuals, who met these

criteria and consented to take part in the study, comprised the final sample.

Trained personnel first recorded all epidemiological and demographic data for the consenting

respondents, and then administered the KCB. We used the KCB version validated by Ganguli et

al, as it had questions on post office, district, village and block, relevant to a rural community.[3]

The Marathi KCB was validated by re-administering 30% of the cognition tasks from each of the

eight domains of the KCB on 40 randomly chosen study participants—by a second field

researcher for inter-rater, and by the same researcher after a gap of 6 weeks for intra-rater

reliability. Test administration and scoring were supervised by the chief investigator.

Our primary outcome measure was the age-stratified KCB score. Data collected on completed

years of schooling was classified into educational categories namely, illiterate, primary (1-5

years), secondary (6-12 years), and graduation level (13 years and above).

Cut-off scores: For screening a large population, operational cutoff points are essential to

differentiate between the cognitively impaired and the unimpaired groups.[8] Following the

precedent set by Das et al,[5] we chose the lowest 10th percentile score as the cut-off point to

identify the significantly cognitively impaired tenth of the population. In GDS, scores range from

0-30. The higher scores indicate undetected depression. We chose the 90th percentile as the

operational cut-off point to identify respondents with masked depression.

Statistical methods

The data was analyzed using appropriate SPSS[9] and JASP.[10] Values were expressed in

frequencies as well as percentages in relevant rows. Spearman rank correlation coefficients were

calculated to determine intra and inter observer reliability. Anova was used to compare means

between groups. Kruskal Wallis H test was used to analyze overall differences between age

groups and educational classes, and Mann Whitney U test was used to compare the two genders,

urban and rural groups, groups with and without masked depression, and to compare each of the

subgroups in age and education with the other subgroups.

The study was cleared by the Institutional Ethical Committee of Lokmanya Tilak Municipal

Medical College and General Hospital, Sion, Mumbai.

Results

Demographic profile: We screened 2752 persons over 40 years of age; we excluded 19 persons

with known intellectual developmental disorder, 31 with possible previous stroke, 36 with

previous psychiatric illness, and 15 persons reported by family to have previously diagnosed

dementia or significant memory impairment impeding daily activities. Our final sample size was

2651 individuals who satisfied the inclusion and exclusion criteria and consented to participate in

an interview. Their socio-demographic characteristics are summarized in Table 1. Among these

participants, 1435 (54%) were from rural localities while 1216 (46%) belonged to urban

localities. The study group was equally divided with respect to gender, comprising 1335 men and

1316 women. However, in the younger age groups (<60 years), men outnumbered women and

this trend was reversed for the older age groups.   The mean age of the study participants was 54

years and women were slightly older (mean 54.28 years) than men (mean 53.87 years), though

this difference was not significant. Most of the participants were poorly educated. Out of 1335

men, 752 (56.3%) had studied for five years or less, and among women the corresponding

proportion was significantly higher (1001 of 1316, 76.1%, p<0.01). Correspondingly, men

significantly outnumbered women in the groups with higher education. We also noted a

significant gender difference in occupation, with most of the women participants reporting as

housewives (1108, 84.2% of 1316) and most of the remainder in unskilled and semiskilled

occupations. Among men, 516/1335, i.e., 38.4%, reported employment in unskilled vocations,

followed by semiskilled and clerical work respectively. We observed a significant downtrend in

mean years of education with increasing age (table 1). We observed a significant difference in

scores between rural and urban subsets of the study sample. The urban residents were

significantly younger and had significantly higher years of education than the rural residents

(table 1). While the study sample comprised roughly equal numbers of men and women with no

significant difference in their mean age, the two gender groups differed notably in education,

with men on average having 6.5 years of schooling compared to 4.7 years for women (table 1).

Test scores and validation: Individual test scores with mean, median, standard deviation,

interquartile range and 10th and 90th percentile scores are recorded in table 2. Spearman rank

correlation coefficients for both inter and intra rater reliability were high for each test, ranging

from 0.92 to 1.0. The score range for each domain spanned from zero to the maximum attainable

score, except in the case of immediate recall, in which the maximum score attained was 28 out of

30.

Gender: The individual test scores refined by gender, shown in table 3, highlight a significant

difference in the mean scores, with women scoring lower than men on each test. The 90[th]

percentile scores were higher for men in tests of verbal fluency, immediate recall, visuo-

constructional ability and delayed recognition, and the 10[th] percentile scores were higher for

MMSE, visuo-constructional ability and delayed recognition. The difference in test scores

attributable to gender, i.e., effect size, ranged from 5% to 20%, with the largest effect size

observed for test 4, calculation.

Rural vs Urban residence: The individual test scores for rural and urban subsets, shown in table

4, demonstrate higher mean scores in each test for the urban subset. Tenth percentile scores were

higher for urban residents in all tests except delayed word list, and 90[th] percentile scores were

also higher in four of the eight tests. Effect size ranged from 10% to 47%, with the largest

difference observed for test 6, visuo-constructional ability followed by test 5, immediate recall.

Presence vs Absence of depression: In our study sample, the 90th percentile score for GDS was

17, which we used as the operational cut off to identify masked depression. We observed

significantly lower mean scores on all tests among the respondents who scored 17 and above on

the GDS (table 5). The largest effect size, up to 18%, was seen in Test 3, Mental State

Examination, and Test 1, Verbal Fluency.

Table 6 shows the comparison of mean scores between different educational groups. Significant

differences were observed in all tests between the illiterate group and every other educational

category, except for the comparison between illiterate and primary education group for object

naming and delayed recognition tests. Similarly, in the comparison between secondary education

and graduate level groups, the difference was not significant for several tests including

immediate recall, visuo-constructional ability and delayed recognition.

The comparison of means scores between age groups is summarized in table 7. Whereas

between adjacent age bins, there was no significant difference in individual test scores (at both

extremes of age groups), significant differences were observed when the age groups were

sufficiently apart, typically between the youngest two bins and the older bins.

Table 8 summarizes the comparison between the four educational classes at each age level. We

observed that scores differed significantly between educational groups in the younger age levels

across all tests, but with increasing age, differences ceased to be significant in the object naming

and delayed word list domains.

Discussion

Discussion

The ethnic and educational diversity of the Indian population makes it necessary to obtain

normative cognitive data for different regions. The most notable among the very few validated

cognitive screening tools are those developed by Ganguli et al[3] for rural Hindi speaking

communities, and the Kolkata Cognitive Battery by Das et al.[5] We now attempt to modify and

tailor the KCB for Maharashtrian populations.

The GDS is designed such that high scores correlate with severe depression. Thus, the 90th

percentile score of the GDS is used as the cut off to screen in individuals who are potentially

clinically depressed. For our study sample, this cut off is 17 which is lower than the accepted

standard score of 22[7] in rural communities (the cut off in Das et al's sample[5] was 21). Our camp-

based methodology was dependent on respondents being sufficiently ambulatory and motivated

to come to the screening site, which may have excluded the most depressed individuals with the

uppermost range of GDS scores. However, even this lower cut off showed a significant

association with low cognitive scores. While the depressed tenth of the study group had

statistically significant lower scores in each of the tests, the gap in mean scores was notably wide

in the fields of immediate recall and verbal fluency. Research indicates that depression affects

cognitive performance in older as well as younger subjects, affecting ability to complete complex

tasks.[11] Sheline et al,[12] studying the effect of late life depression on cognitive performance,

assessed five major cognitive domains in patients with major depression, and reported that

changes in processing speed were found to most fully mediate the influence of predictor

variables on all other cognitive domains.

Our study showed significantly poorer performance by women in each of the tests, compared to

men. While the largest gender attributable score difference was observed in testing calculation,

the difference in visuo-constructional ability scores was striking even in the lowest tenth

percentile. The women were significantly less educated than men. Few among them (208/1316;

15.8%) had jobs outside the home, the majority being homemakers. The proportion of illiteracy

was more than twice as high among women compared to men (f: 244/1316, 18.5%; m: 105/1335,

7.9%). The general awareness, and comfort with writing and drawing among women may have

been adversely affected, as a result. Education, in particular literacy, has been shown to be

associated with high cognition in the elderly.[13] In a study of ethnically diverse elders,

respondents with lower literacy levels had lower cognitive scores, this difference being attributed

to poorer organization of visuospatial information, lack of previous exposure to stimuli, and

difficulties with interpretation of the logical functions of language.[14] Thus, it may be surmised

that the observed differences in cognitive scores between men and women in our study are

attributable to differences in their education levels.

Our study showed that test scores decreased (albeit not uniformly) with increasing age, with

differences more pronounced between the youngest and oldest age groups; no significant

difference being observed comparing the scores of the moderately older or younger age groups

with each other. Das et al, in their Kolkata study, observed a similar pattern of significant

differences in overall test scores mainly when comparing the relatively younger age groups (50-

59) with elderly subjects aged above 70 years.[5]

The most remarkable finding of our study is the difference in test scores between the rural and

urban subsets of the study group, particularly in the domains of visuo-constructional ability and

immediate recall. Compared to rural, our urban respondents were significantly younger and

better educated, with 50% of the urban dwellers having secondary level or higher education

versus only 21% rural dwellers. The rural subset also had proportionately more women

compared to urban (R: 791/1435; 55.1% vs. U: 525/1216; 43.2%). We noted a positive

correlation of test scores with urban residence even after controlling for age, sex and education,

with the correlation coefficients being highest in the domains of visuo-constructional ability

(0.323, p< 0.01), immediate recall (0.313, p< 0.01), and calculation (0.239, p< 0.01). The

modern, fast-paced urban life, involving daily commutes, monetary calculations, workplace

interactions and a variety of cultural exchanges with different occupations and ethnicities, could

have contributed to better general awareness, alertness and mental processing speed, and

subsequently higher test scores. Researchers from other countries have noted a similar rural-

urban gap in cognitive scores, attributing it to educational disadvantage, selection through

migration, and the existence of potentially modifiable risk factors other than lifestyle.[15,16] There

is thus considerable scope for further, larger studies of cognitive differences in urban and rural

India, exploring factors such as education, income, childhood illnesses and substance use habits.

Ceiling effect: Das et al,[5] in their study of cognitive scores in an urban elderly population,

noticed possible ceiling effects in tests of object naming, calculation, mental state examination

and delayed recognition, which they attributed to the large number of literate subjects in an urban

milieu. However, no such effect has been observed in the present study, either overall or in the

urban subset of the participants group. This may be consequent to the difference in the

educational distribution of the two study sets, with two-thirds (501 of 745, 67%) of the Kolkata

sample being educated at the secondary or higher level, compared to only a third of the present

study group (898 of 2651, 34%).

Limitations:

A limitation of our study which may have introduced a bias, (in addition to the camp-based

methodology which may have excluded the severely depressed) is that many of our older

subjects were unable to recall their exact age, and there may have been marginal

misclassification of age bins.

Conclusion

We have presented and discussed the formulation and validation of a Marathi translation of the

Kolkata Cognitive Battery. We found that cognitive scores declined with increasing age and poor

education; however, our chief finding is the striking positive difference that an urban residential

environment made to cognitive scores, when other variables were comparable. This Marathi

KCB having been validated on a large sample of urban as well as rural residents, may be used to

study cognition in Marathi speaking populations with appropriate cut-offs tailored to the degree

of urbanization of the locality.

Our study participants performed poorly in the domains of verbal fluency, object naming and

delayed recognition in comparison to the Ballabhgarh and Kolkata study groups with whom the

original Hindi and Bengali cognitive testing instrument versions were validated. Further research

is warranted to explore the factors leading to these differences. Women had poorer cognitive

scores than men, which we surmise is due to their homebound lives, lesser education and lower

involvement in financial management.

Funding: This study was part of a larger study on "Age-stratified normative cognitive scores in

the community and their correlation with MRI volumetric studies", fully funded by Cognitive

science research initiative 2012, Department of Science and Technology, Government of India.

References:

1.   Tripathi RK, Tiwari SC. Cognitive Dysfunction in Normally Aging Urban Older Adults: A
Community-based Study. *Indian J Psychol Med* 2011; 33: 177–181.

2.   Gambhir IS, Khurana V, Kishore D, et al. A clinico-epidemiological study of cognitive
function status of community-dwelling elderly. *Indian J Psychiatry* 2014; 56: 365–370.

3.   Ganguli M, Chandra V, Gilby JE, et al. Cognitive test performance in a community-based
nondemented elderly sample in rural India: the Indo-U.S. Cross-National Dementia
Epidemiology Study. *Int Psychogeriatr* 1996; 8: 507–524.

4.   Banerjee TK, Mukherjee CS, Dutt A, et al. Cognitive dysfunction in an urban Indian
population--some observations. *Neuroepidemiology* 2008; 31: 109–114.

5.   Das SK, Banerjee TK, Mukherjee CS, et al. An urban community-based study of cognitive
function among non-demented elderly population in India. *Neurol Asia* 2006; 11: 37–48.

6.   Issac TG, Chandra SR, Rajeswaran J, et al. Demographic Features and Neuropsychological
Correlates in a Cohort of 200 Patients with Vascular Cognitive Decline Due to Cerebral
Small Vessel Disease. *Indian J Psychol Med* 2016; 38: 127–132.

7.   Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression
screening scale: a preliminary report. *J Psychiatr Res* 1982; 17: 37–49.

8. Paulsen JS, Hoth KF. Neuropsychology. In: *Bradley WG, Daroff RB, Fenichel GM, Jankovic*
J eds: Neurology in clinical practice – principles of diagnosis and management.
Philadelphia: Lippincott Williams & Wilkins, 2004, pp. 679–80.

9. IBM Corp. Released 2011. *IBM SPSS Statistics for Windows, Version 20.0.* Armonk, NY:
IBM Corp.

10. JASP Team. *JASP (Version 0.13)[Computer software]*, https://jasp-stats.org/ (2020, accessed
15 July 2020).

11. Tarbuck AF, Paykel ES. Effects of major depression on the cognitive function of younger
and older subjects. *Psychol Med* 1995; 25: 285–295.

12. Sheline YI, Barch DM, Garcia K, et al. Cognitive Function in Late Life Depression:
Relationships to Depression Severity, Cerebrovascular Risk Factors and Processing Speed.
Biol Psychiatry 2006; 60: 58–65.

13. Barnes DE, Tager IB, Satariano WA, et al. The Relationship Between Literacy and
Cognition in Well-Educated Elders. *J Gerontol Ser A* 2004; 59: M390–M395.

14. Manly J, Desiree B, Pegah T, et al. Literacy and cognitive change among ethnically diverse
elders. *Int J Psychol*; 39: 47–60.

15. Saenz J, Downer B, Garcia M, et al. Cognition and Context: Rural-Urban Differences in
Cognitive Function in Mexico. *J Aging Health* 2018; 30: 965–986.

16. Nakamura K, Kitamura K, Watanabe Y, et al. Rural-urban differences in the prevalence of
cognitive impairment in independent community-dwelling elderly residents of Ojiya city,
Niigata Prefecture, Japan. *Environ Health Prev Med* 2016; 21: 422–429.

356   Table 1: Demographic characteristics

Variables	Sex				Total	Mean age in years (±SD)	Mean years of education (±SD)
	Male		Female			M 53.9(±10.3)	M 6.5(±4.3)
						F 54.3(±10.5)	F 4.7(±4.2)
						T 54.1(±10.4)	T 5.6(±4.3)
	n	%	n	%	n		
Age group (years)							
40-49	525	51.9	487	48.1	1012	44.2(±2.9)	6.4(±4.5)
50-59	441	52.6	398	47.4	839	53.5(±2.7)	5.8(±4.4)
60-69	232	43.3	304	56.7	536	63.2(±2.7)	4.5(±3.7)
70-79	116	55.2	94	44.8	210	73.1(±3.0)	4.2(±3.7)
80+	21	38.9	33	61.1	54	83.1(±3.7)	4.4(±3.6)
Total	1335	50.4	1316	49.6	2651		
Educational level							
Illiterate	105	30.1	244	69.9	349	56.9(±10.6)	0(0)
Primary	647	46.1	757	53.9	1404	54.5(±10.7)	3.7(±1.5)
Secondary	403	65.4	213	34.6	616	53.2(±9.8)	9.0(±.4)
Graduate	180	63.8	102	36.2	282	50.2(±8.1)	14.6(±.9)
Place of residence							
Rural	644	44.9	791	55.1	1435	55.1(±10.8)	4.4(±3.9)
Urban	691	56.8	525	43.2	1216	52.8(±9.8)	7.1(±4.3)
Occupation							
Clerical	128	85.3	22	14.7	150		
Housewife	0	0	1108	100	1108		
Professional	71	76.3	22	23.7	93		
Retired	75	97.4	2	2.6	77		
Semiskilled	309	84.7	56	15.3	365		
Skilled	170	83.3	34	16.7	204		
Unemployed	66	94.3	4	5.7	70		
Unskilled	516	88.4	68	11.6	584		

357   M = Male; F = Female; T = Total
358   *Significant difference noted in mean years of education across age groups, mean age across educational groups, and mean age*
359   *and years of education between urban and rural subgroups;* p value <0.01 (anova) for all
360

361   Table 2: Overview of test scores

KCB Component Tests	Maximum score	n	Range	Mean	Std. Deviation	Median	Inter-quartile range	10th percentile score	90th percentile score
Test 1 Verbal Fluency	NS	2651	0-32	11.86	5.35	12	7	5	20
Test 2 Object Naming	15	2650	0-15	10.00	3.80	10	8	5	15
Test 3 Mental State Examination	30	2635	0-30	24.56	5.31	26	8	17	30
Test 4 Calculation	5	2567	0-5	3.45	1.69	4	3	1	5
Test 5 Immediate Recall	30	2651	0-28	12.01	5.89	12	8	4	20
Test 6 Visuo-constructional Ability	13	2646	0-13	6.68	3.79	6	6	2	12
Test 7 Delayed Word List	10	2623	0-10	4.77	2.14	5	2	2	7
Test 8 Delayed Recognition	20	2637	0-20	8.57	4.36	9	6	2	14

370   Table 3: Test scores sorted by gender and their comparison by Mann-Whitney U test

Parameters	Gender	n	Mean	Std. Deviation	10th percentile score	90th percentile score	p value	Effect size*	95% CI for effect size	
									Lower	Upper
Test 1 Verbal Fluency	Men	1335	12.36	5.4	5	20	<0.01	0.113	0.070	0.157
	Women	1316	11.35	5.25	5	19				
Test 2 Object Naming	Men	1335	10.16	3.62	5	15	<0.05	0.051	0.007	0.095
	Women	1315	9.85	3.97	5	15				
Test 3 Mental State Examination	Men	1324	25.17	5.21	17	30	<0.01	0.16	0.117	0.203
	Women	1311	23.94	5.35	16	30				
Test 4 Calculation	Men	1282	3.74	1.6	1	5	<0.01	0.203	0.16	0.246
	Women	1285	3.15	1.72	1	5				
Test 5 Immediate Recall	Men	1335	12.75	5.92	4	20	<0.01	0.144	0.101	0.187
	Women	1316	11.27	5.76	4	19				
Test 6 Visuo-constructional Ability	Men	1332	7.13	3.76	2	12	<0.01	0.139	0.095	0.181
	Women	1314	6.23	3.78	0	11				
Test 7 Delayed Word List	Men	1319	4.93	2.07	2	7	<0.01	0.089	0.045	0.132
	Women	1304	4.61	2.19	2	7				
Test 8 Delayed Recognition	Men	1327	8.91	4.32	3	14	<0.01	0.085	0.041	0.129
	Women	1310	8.22	4.35	2	13				

371   * For the Mann-Whitney test, effect size is given by the rank biserial correlation

379   Table 4: Test scores sorted by urban/rural residence, and their comparison by Mann-Whitney U

380   test

Parameters	Residence	n	Mean	Std. Deviation	10th percentile score	90th percentile score	p value	Effect size*	95% CI for effect size	
									Lower	Upper
Test 1 Verbal Fluency	Rural	1435	10.55	5.27	5	18	<0.01	-0.326	-0.365	-0.286
	Urban	1216	13.41	5.02	7	20				
Test 2 Object Naming	Rural	1434	9.68	4.25	4	15	<0.01	-0.109	-0.153	-0.065
	Urban	1216	10.39	3.15	6	15				
Test 3 Mental State Examination	Rural	1427	23.14	5.33	15	30	<0.01	-0.381	-0.419	-0.343
	Urban	1208	26.23	4.78	19	30				
Test 4 Calculation	Rural	1424	2.94	1.73	0	5	<0.01	-0.378	-0.416	-0.339
	Urban	1143	4.07	1.41	2	5				
Test 5 Immediate Recall	Rural	1435	9.87	5.18	4	16	<0.01	-0.466	-0.5	-0.43
	Urban	1216	14.54	5.66	7	22				
Test 6 Visuo-constructional Ability	Rural	1432	5.27	3.43	0	10	<0.01	-0.472	-0.505	-0.437
	Urban	1214	8.34	3.52	3	12				
Test 7 Delayed Word List	Rural	1427	4.5	2.17	2	7	<0.01	-0.168	-0.211	-0.125
	Urban	1196	5.09	2.06	2	7				
Test 8 Delayed Recognition	Rural	1435	7.78	4.46	2	13	<0.01	-0.248	-0.289	-0.206
	Urban	1202	9.51	4.06	4	14				

381   * For the Mann-Whitney test, effect size is given by the rank biserial correlation

389   Table 5: Test scores sorted by presence/absence of masked depression, and their comparison by

390   Mann-Whitney U test

Parameters	Depression[#]	n	Mean	Std. Deviation	10th percentile score	90th percentile score	p value	Effect size*	95% CI for effect size	
									Lower	Upper
Test 1 Verbal Fluency	No	2376	12.03	5.41	5	20	<0.01	0.173	0.102	0.242
	Yes	275	10.39	4.49	5	16				
Test 2 Object Naming	No	2375	10.11	3.76	5	15	<0.01	0.141	0.07	0.211
	Yes	275	9.09	4.01	5	15				
Test 3 Mental State Examination	No	2364	24.71	5.27	17	30	<0.01	0.187	0.116	0.256
	Yes	271	23.25	5.54	16	29				
Test 4 Calculation	No	2299	3.49	1.68	1	5	<0.01	0.152	0.08	0.222
	Yes	268	3.06	1.76	0	5				
Test 5 Immediate Recall	No	2376	12.20	5.89	5	20	<0.01	0.165	0.094	0.234
	Yes	275	10.40	5.60	3	18				
Test 6 Visuo-constructional Ability	No	2371	6.78	3.76	2	12	<0.01	0.135	0.063	0.205
	Yes	275	5.81	4.01	0	11				
Test 7 Delayed Word List	No	2351	4.84	2.12	2	7	<0.01	0.159	0.087	0.228
	Yes	272	4.21	2.25	1	7				
Test 8 Delayed Recognition	No	2362	8.68	4.33	3	14	<0.01	0.136	0.065	0.206
	Yes	275	7.59	4.58	2	12				

392   * For the Mann-Whitney test, effect size is given by the rank biserial correlation

393   [#]Presence of depression operationally defined as GDS score ≥17

399   Table 6: Test scores of each educational level compared with higher and lower levels using

400   Mann Whitney U test

Parameters	Illiterate vs Primary	Illiterate vs Secondary	Illiterate vs Graduate	Primary vs Secondary	Primary vs Graduate	Secondary vs Graduate
Test 1 Verbal Fluency	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01	P<0.05
Test 2 Object Naming	NS	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01
Test 3 Mental State Examination	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01
Test 4 Calculation	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01
Test 5 Immediate Recall	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01	NS
Test 6 Visuo-constructional Ability	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01	NS
Test 7 Delayed Word List	P<0.01	P<0.01	P<0.01	NS	P<0.01	P<0.01
Test 8 Delayed Recognition	NS	P<0.01	P<0.01	P<0.01	P<0.01	NS

413   Table 7: Test scores of each age group compared with older and younger groups using Mann

414   Whitney U test

Parameters	40-49 vs 50-59	40-49 vs 60-69	40-49 vs 70-79	40-49 vs 80+	50-59 vs 60-69	50-59 vs 70-79	50-59 vs 80+	60-69 vs 70-79	60-69 vs 80+	70-79 vs 80+
Test 1 Verbal Fluency	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	NS	NS	NS
Test 2 Object Naming	NS	$P<0.01$	$P<0.01$	NS	$P<0.01$	$P<0.01$	NS	NS	NS	NS
Test 3 Mental State Examination	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.05$	$P<0.05$	NS
Test 4 Calculation	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	NS	NS	NS
Test 5 Immediate Recall	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	NS	$P<0.05$	NS
Test 6 Visuo-constructional Ability	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	NS	NS
Test 7 Delayed Word List	NS	$P<0.01$	$P<0.01$	$P<0.05$	$P<0.01$	$P<0.01$	NS	NS	NS	NS
Test 8 Delayed Recognition	NS	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P<0.01$	$P=0.05$	NS	NS

424   Table 8: Comparison between the four educational classes at each age bin, using Kruskal-Wallis

425   Test

Parameters	Within 40-49	Within 50-59	Within 60-69	Within 70-79	Within 80+
Test 1 Verbal Fluency	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01
Test 2 Object Naming	P<0.01	P<0.01	NS	NS	P<0.05
Test 3 Mental State Examination	P<0.01	P<0.01	P<0.01	P<0.01	P<0.01
Test 4 Calculation	P<0.01	P<0.01	P<0.01	NS	P<0.01
Test 5 Immediate Recall	P<0.01	P<0.01	P<0.01	P<0.01	P<0.05
Test 6 Visuo-constructional Ability	P<0.01	P<0.01	P<0.01	P<0.01	NS
Test 7 Delayed Word List	P<0.01	P<0.01	P<0.01	NS	NS
Test 8 Delayed Recognition	P<0.01	P<0.01	P<0.01	P<0.01	P<0.05

434