

1 Title: Enteric viruses isolates and rotavirus genotypes during the seasonal surge of acute watery
2 childhood diarrhoea in South-Eastern Nigeria

3 Short title: Viruses isolated from childhood diarrhoea in Nigeria
4

5

6 Akinwale Michale Efunshile*^{1,2}, Chukwuemeka Nwangwu.¹, Bethrand Amechi Ngwu.¹

7

8 ¹ Department of Medical Microbiology, Ebonyi State University, Abakaliki, Nigeria

9 ² Department of Medical Microbiology, Federal Teaching Hospital, Abakaliki, Nigeria

10 * Address for correspondence: AME akinwale.dr.efunshile@daad-alumni.de

11

12

13

14

15

16 **Abstract**

17 Diarrhoea remains one of the top three causes of death in Africa. However, data on the
18 molecular epidemiology of enteric viruses in Nigeria is rare.

19 Two hundred and forty three infants and children below the age of 5 years with acute watery
20 diarrhoea were evaluated for the presence of enteric viruses in stool by real-time PCR
21 (rtPCR) during the dry months of December to April 2017, which correspond to diarrhoea
22 season in South-East Nigeria. At least one viral pathogen was detected in 95.1% (231/243) of
23 the study population. Rotavirus, 231(95.1%) was the most common followed by adenovirus,
24 103(42.4%) and enterovirus 32(13.2%). Other viruses seen in the stool samples include
25 astrovirus 7.3 % (11/151), parechovirus 3.5 % (4/113), sapovirus, 2.8 % (4/145), bocavirus
26 6.8% (5/73) and human coronavirus 13.9% (10/73). Norovirus and hepatitis A and E viruses
27 were not detected. Children that consumed factory packaged water had the lowest prevalence
28 of rotavirus infection ($p=0.044$). A significant association between the viral pathogen and
29 home treatment of drinking water or exclusive breastfeeding was not observed. Randomly
30 selected 46 samples were genotyped for rotavirus, which showed that G3P[6] (39.1%) was
31 the most common, followed by G1P[8] (15.2%), G9P[8] (13.0%), G12P[8] (6.5%),
32 G9P[6]P[8] (2.2%), and G12P[6] (2.2%).

33 This was the highest rotavirus prevalence in any published African study, and may be a
34 reflection of inadequate water sanitation/hygiene in the study area, a peculiar epidemiological
35 situation and/or the sensitivity of the diagnostic method used.

36 The present study suggests that the burden of rotavirus is bigger than previously reported; and
37 that morbidity can be greatly reduced if rotavirus vaccine is included in Nigerian national
38 immunization policy.

39

40 **Keywords:** Acute diarrhoea, Enteric viruses, Rotavirus, Rotavirus genotypes.

41 **Introduction**

42

43 Diarrhoea caused an estimate of 1.3 million deaths globally in 2015 and it was reported to be
44 the 4th leading cause of death among children less than 5 years old [1].

45 Diarrhoea was reported to have killed more than 500,000 children under 5 years in 2015
46 alone, with Asia and Sub-Saharan Africa recording the highest number of incidence and
47 deaths [2]. Although, deaths from diarrhoea and other diseases declined globally between
48 2010 and 2015, the reduction rate was not remarkable in Nigeria and many other Sub-Sahara
49 Africa countries as a result of inadequate water, poor sanitation and hygiene; as well as
50 predisposition from malnutrition [2]. Rotavirus, cryptosporidium and shigella *spp* were the
51 major agents which accounted for more than 50% of diarrhoea deaths among children below
52 the age of 5 years in 2015 [1]. In its comprehensive assessment, the Global Burden of
53 Diseases in 2015 also listed enteric adenovirus and noroviruses as other viral agents that
54 significantly contributed to diarrhoea deaths [1]. Though, the burden of specific viral agent
55 depends on the study region and the season. Because rotavirus is the commonest diarrhoea
56 agent worldwide, the incidence and mortality of its associated diarrhoea were shown to have
57 significantly declined in countries where rotavirus vaccines were introduced [3]. The WHO
58 therefore recommends that rotavirus vaccine is included in routine immunization policy,
59 especially in developing countries [4]. Rotavirus gastroenteritis is more seasonal in temperate
60 regions compared to the tropical areas where it occurs more or less all the year round. And
61 this difference was found to be associated with socio-economic factors rather than being an
62 effect of climate [5]. A systematic review of rotavirus diarrhoea in several African countries
63 concluded that the disease usually peak during the dry period of the year in most countries
64 [6]. In Asia, where diarrhoea burden is similar to what obtains in Africa, findings also
65 showed that diarrhoea occurs all the year round, and that 45% of diarrhoea in children below

66 the age of 5 years was attributed to rotavirus [7]. Co-infection between rotavirus and other
67 viral diarrhoea agents have also been reported. For instance, rotavirus (26.6%) was found to
68 be more common among Moroccan diarrhoea children, while mixed infection with norovirus
69 was found in 2.7% of the cases [8]. Another diarrhoea study reported rotavirus and norovirus
70 in a cohort of 50 diarrheic South-West Nigerian children with a prevalence of 34.5% and
71 25.5% respectively, while co-infection with both viruses was seen in 9.1% of cases [9]. Three
72 enteric viruses, rotavirus (28.3%), Adenovirus (19.3%), and norovirus (3.6%) were also
73 reported among diarrhoea children in South-South region of Nigeria with co-infections by 2
74 or 3 viruses in 7.6% cases [10]. A recent report evaluating diarrhoea agents over a 12 months
75 period in Nigeria showed that at least one was identified in 58.3% of cases with co-infections
76 involving norovirus or astrovirus or aichivirus in some cases [11].

77 Countries with the highest rotavirus burden including Nigeria are yet to implement the WHO
78 vaccine policy despite its proven efficacy [4,12]. Changes in the aetiology and epidemiology
79 of viral agents of diarrhoea were reported following the introduction of rotavirus vaccines
80 [13, 14]. This underlines the importance of continuous surveillance of enteric viral agents to
81 enable effective planning for future control efforts. Such surveillance will also help to bring
82 the public health impact and effectiveness of the vaccines to the forefront, thereby
83 encouraging policy implementation in countries that are yet to comply. The present study
84 aimed at evaluating enteric viral agents associated with acute diarrhoea in the study area and
85 to provide baseline epidemiologic data ahead of the national rotavirus vaccine
86 implementation. Additionally, the relationship between feeding practices and sources of
87 drinking water and enteric viruses was evaluated.

88
89
90

91 **Methods**

92 **Study design**

93 This is a cross sectional study primarily designed to identify the viral agents of acute
94 diarrhoea in infants and children below 5 years of age. Inclusion criteria were passage of
95 watery stool three or more times within the last 24 hours, age of 5 years or less and
96 willingness to participate in the study. The Ethical Committee of the Federal Teaching
97 Hospital Abakaliki, Nigeria, approved the study. Written informed consent were obtained
98 from the parents/guardian of the children after the purpose of the study were translated to
99 them in local dialects.

100 **Study Area**

101 The study was carried out at University Teaching Hospitals (Ebonyi and Enugu) in the South-
102 Eastern Region of Nigeria between latitudes 04⁰30'N and 07⁰30'N and longitudes 06⁰45'E
103 and 08⁰45'E [15], during the peak diarrhoea period, in the months of December 2016 to April
104 2017.

105 **Data collection and management**

106 Questionnaires were administered to the parents/guardian of the participants to collect data
107 regarding age, frequency and duration of diarrhoea, risk factors and other relevant
108 information. This information were subsequently transferred into excel sheets for analysis.

109 **Specimen management and preliminary examination**

110 About 5 ml of watery stool were collected from the participants into wide mouth bottles.
111 Inactivated aliquots were shipped to the Institute of Virology, Leipzig University, Germany.

112

113 **Virus PCR**

114 Total nucleic acids was extracted from 10 % stool suspensions in PBS by MagNA Pure 96
115 system (Roche Applied Science, Mannheim, Germany), aliquoted, and stored at -80°C.

116 Aliquots were subsequently screened by real-time (RT)-PCRs for presence of viral genome of
117 species A rotavirus, norovirus (genogroup I and II), sapovirus (genogroup I, II, IV and V),
118 adenovirus (group A to F), parechovirus, astrovirus (HastV 1–8), and hepatitis E virus and
119 coronavirus [16-24]. Primers and probes used for detection of enterovirus (EV) species A to
120 D (including coxsackieviruses, echoviruses, enterovirus and poliovirus), human bocavirus and
121 hepatitis A virus (HAV) targeted the 5’NTR-regions of the viral genome: EV-1s (5’-
122 GGTGCGAAGAGTCT ATTGAGCTA-3’), EV-1as (5’-ACCCAAAGTAGTCGGTTCC-3’),
123 EV-probe (5’-6FAM-TGAATGCGGCTAATCCTAACTGCG-DB-3’), hBoV_5UTR_Q_s1
124 (5’-CCTATATAARCCGATGCACT), hBoV_5UTR_Q_s2 (5’-
125 CCTATATAAGCTGCTGCACT), hBOV_NS1_Q_as1 (5’-
126 AGGAGCAGAAAAGGCCATA), hBoV_NS1_Q_as2 (5’-AGGRGGATTGAAAGCCATA),
127 hBoV_Q-TaqFAM (5’-6FAM-TCCGGCGAGTGAACATCTCT—BBQ), HAV-5UTR-Qs
128 (5’AATATTAATTCCTGCAG GTTCAG-3’), HAV-5UTR-Qas
129 (5’AGCCAAGTTAACTGCAAG-3’), and HAV-5UTR QFAM (5’6FAM-
130 CAATTTAGACTCCTACAGCTCCATGCTA-BBQ-3’).

131 **Rotavirus G and P typing**

132 Randomly selected 46 samples tested by quantitative real time PCR for rotavirus were
133 sequenced as previously described using published methodology [25,26]. The G and P type-
134 specific amplification of VP7 and VP4 fragments was performed using QIAgen One-step RT-
135 PCR kit (QIAgen, Hilden, Germany). For the first run, primers used were VP7-F and VP7-R
136 for the VP7 gene while VP4-F and VP4-R were used for the VP4 gene. A second semi-nested
137 multiplex PCR was done using G1-, G2-, G3-, G4-, G8-,G9-, G10- and G12-specific sense
138 primers for G types, while P-typing included P[4]-, P[6]-, P[8]-, P[9]-, P[10]-, and P[11]-
139 specific antisense primers together with VP7-R and VP4-F, respectively.

140 Gel purified amplicons (Wizard SV Gel and PCR Clean-Up System; Promega, Mannheim,
141 Germany) were partially sequenced (approximately 600 bp each) using BigDye Terminator
142 v1.1 Cycle Sequencing kit (PE Applied Biosystems, Foster City, CA) on an ABI Prism 310
143 Genetic Analyzer. Nucleotide sequences read from the chromatograms were aligned.
144 Sequencing of the r amplicons was done for the 46 randomly selected rotavirus isolates.

145 **Statistical analysis**

146 Data were analysed with the freely available GNU PSPP Statistical Analysis Software
147 version 1.0.1-g818227 [27]. Descriptive statistics was used to compare the various variables,
148 while Chi-square tests were used to derive significant associations which was set at p-value
149 <0.05.

150

151 **Results**

152 For the present investigations, stool samples obtained from 243 children were available. Most
153 of the children, 71.2% (173/243) were below the age of 2 years, and 65% (158/243) were
154 males (table 1).

155

156 **Table 1. Age and sex distribution of participants**

	Gender (N (%))		Total
	Male	Female	
Age group			
<2	111 (64.2)	62 (35.8)	173 (100)
2-3	40 (64.5)	22 (35.5)	62 (100)
>3	7 (87.5)	1 (12.5)	8 (100)
Total	158 (65.2)	85 (34.8)	243 (100)

157

158

159 Eleven enteric viruses were tested. Not all samples were tested for each of the viruses (table
160 2). Ninety-four (38.7%) of the adenovirus positive cases were also co-infected with *Rotavirus*
161 while triple infection between rotavirus, adenovirus and enterovirus was observed in 7.0% of
162 the cases (table 2).

163

164 **Table 2. Enteric virus distribution in the study population**

Enteric viruses	Number of samples tested	Prevalence, N (%)
Rotavirus	243	231 (95.1)
Adenovirus	243	103 (42.4)
Enterovirus	243	32(13.2)
Astrovirus	151	11(7.3)
Sapovirus	150	3(2.0)
Norovirus	145	0(0.0)
Parechovirus	113	4 (3.5)
Bocavirus	73	5(6.8)
Human Coronavirus	73	10(13.7)
Hepatitis A virus	51	0(0)
Hepatitis E virus	51	0(0)
Rotavirus +Adenovirus	243	94(38.7)
Rotavirus +Enterovirus	243	30(12.3)
Rotavirus +Astrovirus [#]	151	7(4.6)
Adenovirus + Enterovirus	243	24(9.9)
Rotavirus +Adenovirus + Enterovirus	243	17(7.0)
Enterovirus +Sapovirus [#]	150	1(0.7)
Parechovirus [#] +Adenovirus	113	1(0.9)

165 N; number of positives [#] See discussion

166

167 There was no significant association between ages or sex and any of the 3 most commonly
168 encountered enteric viruses (table 3).

169

170 **Table 3. Age and sex distribution of Rotavirus, Adenovirus and Enterovirus among**
171 **diarrheic children below the age of 5 years**

	Rotavirus , N (%)		Adenovirus, N (%)		Enterovirus, N (%)	
	Positive	Negative	Positive	Negative	Positive	Negative
Age (years)						
<1	140 (99.2)	1 (0.8)	76 (53.1)	65 (46.9)	25 (19.4)	116 (17.7)
1-2	50 (84.7)	9 (15.3)	22 (37.3)	37 (62.7)	4 (6.8)	55 (93.2)
2-5	41 (95.3)	2 (4.7)	5 (11.6)	38 (88.4)	3 (7.0)	40 (93.0)
Total	231 (95.1)	12(4.9)	103(42.4)	140 (57.6)	32 (13.2)	211 (86.8)
P-value	0.676		0.164		0.824	
Sex						
Male	150 (94.9)	8 (5.1)	71 (44.9)	87 (55.1)	23 (14.6)	135 (85.4)
Female	81 (95.3)	4 (4.7)	32 (37.6)	53 (62.4)	9 (10.6)	76 (89.4)
Total	231 (95.1)	12 (4.9)	103(42.4)	140 (57.6)	32 (13.2)	211 (86.8)
P-value	0.378		0.243		0.210	

172

173 The present study did not establish a significant association between the three common
174 enteric viruses (i.e. rotavirus, enterovirus and adenovirus) and past history of exclusive
175 breastfeeding or the current feeding practice of the children (table 4).

176 **Table 4. Rotavirus, adenovirus and enterovirus prevalence in relation to feeding**
177 **practices in diarrheic children below the age of 5 years**

	Rotavirus		Adenovirus		Enterovirus	
	Pos, N (%)	Neg , N (%)	Pos, N (%)	Neg, N (%)	Pos, N (%)	Neg, N (%)
EBF history						
Yes	102 (95.3)	5 (4.7)	47 (43.9)	60 (56.1)	37 (34.6)	70 (65.4)
No	129 (95.6)	6 (4.4)	61 (45.2)	74 (54.8)	24 (17.8)	111 (82.2)
Total	231(95.1)	12(4.9)	103(42.4)	140 (57.6)	32 (13.2)	211 (86.8)
P-value	0.378		0.055		0.496	
Current Feeding						
EBF	16 (100)	0(0)	7 (43.8)	9 (65.2)	2 (12.5)	14 (87.5)
Complimentary	195 (95.1)	10 (4.9)	87 (44.5)	118 (57.6)	28 (13.7)	177 (86.3)
PBF	19 (95)	1 (5)	8 (40.0)	12 (60.0)	1 (5.0)	19 (95.0)
Supplementary	1 (50)	1 (50)	1 (50.0)	1 (50.0)	1 (50.0)	1 (50.0)

Total	231(95.1)	12(4.9)	103(42.4)	140 (57.6)	32 (13.2)	211 (86.8)
P-value	0.85		0.981		0.153	

178

179 Key-Pos= Positive, Neg= Negative, EBF= Exclusive breastfeeding, PBF = Predominant
180 breastfeeding

181

182

183

184 But there was a significant association between the sources of drinking water and prevalence
185 of rotavirus infection (P=0.044), as children that consumed factory packaged water had the
186 lowest prevalence of rotavirus infection. But this association was not significant for
187 adenovirus and enterovirus (table 5). Approximately half of the children consumed factory
188 packaged water from plastic bottles. An association between home treatments of drinking
189 water with any of the viruses was not observed, though the majority of children consumed
190 water without any form of home treatment (table 5).

191

192 **Table 5. Rotavirus, adenovirus and enterovirus relationship with sources of drinking**
193 **water and treatment practices among children below the age of 5 years**

	Rotavirus		Adenovirus		Enterovirus	
	Pos, N (%)	Neg, N (%)	Pos, N (%)	Neg, N (%)	Pos, N (%)	Neg, N (%)
Source of drinking water						
Cart vendor	66 (100)	0 (0)	30 (45.5)	36 (55.5)	8 (14.0)	49 (86.0)
Borehole	13 (92.9)	1 (7.1)	8 (57.1)	6 (42.9)	6 (42.9)	8 (57.1)
Dug well	4 (100)	0 (0)	1 (15.0)	3 (75.0)	2 (50.0)	2 (50.0)
Bottled water	116 (96.7)	4 (3.3)	57 (43.2)	75 (56.8)	16 (13.5)	102 (86.5)
Total#	199 (97.5)	5 (2.5)	96(44.4)	120 (55.6)	32(16.6)	161(83.4)

P-value	0.044*		0.691		0.097	
Home water treatment						
None	165 (97.1)	5 (2.9)	75 (41.2)	107 (58.8)	30 (18.6)	130 (81.4)
Boiling	30 (100)	0 (0)	17 (56.7)	13 (43.3)	2 (6.7)	28 (93.3)
Filtration	1 (100)	0 (0)	1 (100.0)	0 (0)	0 (0)	1 (100)
Chemical (Chlorination)	3 (100)	0 (0)	3 (100.0)	0 (0)	0 (0)	2 (100)
Total	199 (97.5)	5 (2.5)	96(44.4)	120 (55.6)	32 (16.6)	161 (83.4)
P-value	0.338		0.051		0.155	

194 Key-Pos= Positive, Neg= Negative, *= Significant value, # Not all children drink water
195 because of exclusive breastfeeding
196

197 There was no significant difference between rotavirus infections in the children across the
198 two cities that were studied. Adenovirus was significantly more common among the children
199 from Enugu (P=0.001), (table 6).

200

201 **Table 6. Rotavirus, adenovirus and enterovirus distribution across the study areas**

Study sites	Rotavirus, N (%)		Adenovirus, N (%)		Enterovirus, N (%)	
	Pos	Neg	Pos	Neg	Pos	Neg
Enugu	161 (95.3)	8 (4.7)	86 (50.8)	83 (49.2)	31 (18.3)	138 (81.7)
Abakaliki	70 (94.5)	4 (5.5)	17 (22.9)	57 (77.1)	1 (1.4)	73 (98.6)
Total	231(95.1)	12(4.9)	103(42.4)	140(57.6)	32 (13.2)	211 (86.8)
P-value	0.418		0.001*		0.048*	

202 Key-Pos= Positive, Neg= Negative, *= Significant value

203

204 The genotypes of the 46 randomly selected rotavirus isolates were described in the table 7.

205 Genotype G3P[6] was the predominant genotype, 18(39.1) followed by G1P[8], 7(15.2).

206

207

208

209 **Table 7 : Distribution of G and P genotype combination of RVA strains detected among**
210 **gastroenteritis children below 5years old**

Rotavirus genotype					
	No. (%) of strains				
	P[6]	P[8]	P[6]/P[8]	P n/t	Total
G1	0	7(15.2)	0	0	7(15.2)
G3	18(39.1)	0	0	0	18(39.1)
G9	0	6(13.0)	1(2.2)	0	7(15.2)
G12	1(2.2)	3(6.5)	2(4.3)	0	6(13.0)
G1/G9 (mixed)	0	0	1(2.2)	0	1(2.2)
G3/G12 (mixed)	1(2.2)	0	2(4.3)	0	3(6.5)
G9/G12 (mixed)	0	1(2.2)	1(2.2)	0	2(4.3)
G n/t	2(4.3)	0	0	0	2(4.3)
Total	22(47.8)	17(14.0)	7(15.2)	0	46(100)

211 n/t= not typeable

212

213

214 Discussion

215 The prevalence of rotavirus observed in our study (95.1%) was far higher than the average
216 report among African children, which is in the range of 25 to 40% [6]. It was also higher than
217 the WHO rotavirus surveillance data which reported a mean prevalence of 32% among 37
218 countries including Nigeria [28]. The High prevalence might be due to differences in water
219 supply and hygiene practices between our study area and the other regions with lower
220 prevalence. For instance, a recent review showed that packaged water in developing countries
221 was 4.6-13.6 times more likely to contain bacteria indicator of faecal contamination
222 compared to developed countries [29]. Sizable numbers of the children also drank from cart
223 water vendors, while most of the water consumed was not subjected to any form of home
224 treatment. Differences in the prevalence of rotavirus may also depend on the diagnostic
225 method used and the season of the year when the study was conducted. It is likely that the
226 present study would have reported a lower prevalence if ELISA had been used, because a

227 study by the research group of Adah reported a rotavirus prevalence of 11.8% and 88.2%
228 from the same stool samples, when ELISA and RT-PCR methods were used respectively
229 [30]. Similar studies from our study area reported a lower rotavirus prevalence of 46% and
230 56% in 2 separate studies carried out over a period of 5 and 2 years respectively using ELISA
231 test methods [31, 33]. But another Nigerian study that used RT-PCR also reported lower
232 rotavirus prevalence of 48%. This might be related to the differences in the study regions and
233 study period [33].

234 Lack of significant association between enteric viruses and exclusive breastfeeding in the
235 present study seems unpredictable. This is possibly because we did not evaluate non-
236 diarrhoeic children as negative controls, which of course is beyond the scope of our study.
237 But there are studies which seem to agree with our findings, for instance, the research group
238 of Clemens showed that breastfeeding does not reduce incidence of diarrhoea, it only reduces
239 the severity [34]. Golding *et al* also concluded in their research that breastfeeding only
240 protect against diarrhoea of non-viral aetiologies [35].

241 The 42.4% prevalence of Adenovirus in our study was higher than the result from a study
242 from North-Western Nigeria [36] which was 22.3%. The difference may also be because we
243 used RT-PCR technique in our study which is more sensitive than the rapid kits used by the
244 other study. Unlike our findings, another study from Nigeria reported higher prevalence of
245 astrovirus (40.4%) [37], and this might be due to regional differences in viral diarrhoea
246 agents. Absence of norovirus in our study was contrary to findings from other African
247 studies. Norovirus prevalence of 25.5% was once reported in South-western region of Nigeria
248 [9] while a review of 19 studies from Sub-Sahara Africa showed that the prevalence ranged
249 from 4.6% to 32.4% [38]. The season of our study was probably responsible. There is a need
250 to further investigate the epidemiology of this virus in South-East Nigeria in a larger cohort
251 of diarrheic children, with specimen collection throughout the year. Studies relating to

252 sapovirus are very rare in Nigeria. A similar study in 2 areas in Kenya also reported low
253 prevalence of 4% and 6% at both areas [39]. This supports the impression that this enteric
254 virus played less significant role in our current study. Like sapovirus, this is the first report of
255 parechovirus study in Nigeria. Two of the three samples that were positive for this virus were
256 negative for all other enteric viruses tested, and they also had the cycle threshold (C_T) of the
257 RT-PCR below 30 (C_t data not shown), suggesting significantly high viral load and
258 probability of being responsible for the diarrhoea cases. But it had been reported that the
259 types, rather than viral load of parechovirus is more important in clinical settings. Infections
260 with type 3 HPeV is said to be clinically relevant when detected, regardless of the viral load
261 [40]. Co-infection of 2 or 3 enteric viruses observed in our study was likely due to the fact
262 that they share the same faecal-oral transmission pathway, as a result of inadequate water,
263 sanitation and hygiene. The present result of co-infection with astrovirus, sapovirus and
264 parechovirus also called for interpretation with caution since only a subset of the stool
265 samples were tested for these particular viruses. Further studies will need to simultaneously
266 test larger sample size to get a more accurate co-infection rate. The relatively lower C_T of
267 *rotavirus* compared to other viruses is a reflection of its higher viral load [41], which may
268 suggest its relative importance in the cases with co-infections.

269 From this study, genotype G3P[6] was the commonest genotype observed. This supports the
270 earlier suggestion that the high prevalence of the RTV reported in this study could be largely
271 due to ingestion of contaminated water since type G3 is known to be conserved in human
272 without zoonotic transmission from animals.⁴² Some authors have observed a significant
273 correlation between G3 genotype and poor sanitation and infection control, and low rotavirus
274 vaccine coverage [42-44]. Although most recent studies in Nigeria, reported other rotavirus G
275 types as the most common. For instance in studies done by Japhet (2017) and Motayo (2018),
276 the prevalent genotypes were G12 and G1 respectively [43, 44]. The two studies were done

277 throughout the year in the Western part of Nigeria. Apart from the differences in the study
278 period, the study areas were different from ours with respect to social, cultural and
279 economical determinants. These factors were known to determine the spread of rotavirus
280 genotype [45], hence there is need for national surveillance and characterization of rotavirus
281 in Nigeria for accurate data to better influence policy, monitoring and evaluation of
282 interventions. In a six year (2011-2016) rotavirus surveillance study done at Enugu by Tagbo
283 (2019), G3P [6] was the most common genotype reported in 2016, which is the part of the
284 period our study was conducted [46]. This also affirms the yearly and seasonal variation of
285 rotavirus strains as wildly reported [45]. In the same study by Tagbo, the cumulative
286 prevalence of rotavirus genotypes throughout the six year study period was G12, G1 and G3
287 accounting for 27.6%, 21.0% and 16.3% respectively [46]. Therefore rotavirus strains
288 distribution and prevalence is determined by a lot of factors and study design.

289

290 **Conclusion**

291 The present study suggests that the burden of rotavirus in Nigeria was likely underestimated
292 by previous studies that used non-molecular epidemiologic tool. It also underscores the need
293 to properly investigate the accurate epidemiology of enteropathogenic viruses at a national
294 level so that attention of policy makers and other healthcare stakeholders can urgently be
295 drawn to the need for prompt implementation of rotavirus vaccine as part of the national
296 immunization policy.

297

298 **Limitation of the study**

299

300 The study did not rule out all other possible cause of diarrhoea including bacteria and
301 parasites among the study participants, so it was difficult to establish a causal relationship
302 between the viruses and the diarrhoea.

303 **Competing interest**

304 The authors declared no conflict of interest

305 **Author's contributions**

306 EAM, NC conceived the project idea and developed the proposal. EAM, NC carried out
307 sample and data collection, NC went to Germany to analyze the samples, EAM, NC, NBAF
308 wrote the manuscript, EAM, BAF carried out the statistics analysis. All the authors edited
309 and approved final version of the manuscript

310 **Acknowledgment**

311 We sincerely thank the Institute of Virology, Leipzig University, Germany for the technical
312 assistance for the molecular diagnostics

313

314 **References**

- 315 1. Troeger C, Forouzanfar M, Rao PC, Khalil I, Brown A, Reiner Jr RC, Fullman N,
316 Thompson RL, Abajobir A, Ahmed M, Alemayohu MA. Estimates of global,
317 regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a
318 systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis.*
319 2017;17(9):909-948.
- 320 2. Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, Lawn JE, Cousens S, Mathers C,
321 Black RE. Global, regional, and national causes of under-5 mortality in 2000–15: an
322 updated systematic analysis with implications for the Sustainable Development Goals.
323 *Lancet.* 2016;388(10063):3027-3035.
- 324 3. Troeger C, Khalil IA, Rao PC, Cao S, Blacker BF, Ahmed T, Armah G, Bines JE,
325 Brewer TG, Colombara DV, Kang G. Rotavirus vaccination and the global burden of
326 rotavirus diarrhea among children younger than 5 years. *JAMA Pediatr*
327 2018;172(10):958-965.
- 328 4. World Health Organization Rotavirus vaccines. WHO position paper – January 2013.
329 *Wkly Epidemiol Rec.* 2013;88:49–64.
- 330 5. Patel MM, Pitzer V, Alonso WJ, Vera D, Lopman B, Tate J, Viboud C, Parashar UD.
331 Global seasonality of rotavirus disease. *Pediatr Infect Dis J.* 2013; 32(4): e134.
- 332 6. Cunliffe NA, Kilgore PE, Bresee JS, Hart CA, Glass RI Epidemiology of rotavirus
333 diarrhoea in Africa: a review to assess the need for rotavirus immunization. *Bull*
334 *World Health Organ.* 1998; 76:525–537.
- 335 7. Bresee J, Fang ZY, Wang B, Nelson EA, Tam J, Soenarto Y, Wilopo SA, Kilgore P,
336 Kim JS, Kang JO, Lan WS. First report from the Asian Rotavirus Surveillance
337 Network. *Emerg Infect Dis.* 2004;10(6):988-995.

- 338 8. El Qazoui M, Oumzil H, Baassi L, El Omari N, Sadki K, Amzazi S, Benhafid M, El
339 Aouad R. *Rotavirus* and *Norovirus* infections among acute gastroenteritis children in
340 Morocco. *BMC Infect Dis*. 2014;14(1):1-9.
- 341 9. Japhet MO, Adesina OA, Famurewa O, Svensson L, Nordgren J. Molecular
342 epidemiology of rotavirus and norovirus in Ile-Ife, Nigeria: High prevalence of
343 G12P[8] rotavirus strains and detection of a rare *norovirus* genotype. *J Med Virol*.
344 2012; 84(9):1489-1496.
- 345 10. Imade PE, Eghafona NO. Viral Agents of Diarrhea in Young Children in Two
346 Primary Health Centers in Edo State, Nigeria. *Int J Microbiol*. 2015:685821.
347 doi:10.1155/2015/685821.
- 348 11. Japhet MO, Famurewa O, Adesina OA, Opaleye OO, Wang B, Höhne M, Bock CT,
349 Marques AM, Niendorf S. Viral gastroenteritis among children of 0-5 years in
350 Nigeria: Characterization of the first Nigerian aichivirus, recombinant noroviruses and
351 detection of a zoonotic astrovirus. *J Clin Virol*. 2019;111:4-11.
- 352 12. Yen C, Tate JE, Hyde TB, Cortese MM, Lopman BA, Jiang B, Parashar UD.
353 Rotavirus vaccines: Current status and future considerations. *Hum Vaccin*
354 *Immunother*. 2014;10(6):1436-1448.
- 355 13. Becker-Dreps S, Bucardo F, Vilchez S, Zambrana LE, Liu L, Weber DJ, Peña R,
356 Barclay L, Vinjé J, Hudgens MG, Nordgren J, Lennart Svensson, Douglas R. Morgan,
357 Félix Espinoza, Margarita Paniagua. Etiology of childhood diarrhea following
358 rotavirus vaccine introduction: a prospective, population-based study in Nicaragua.
359 *Pediatr Infect Dis J*. 2014;33(11):1156.
- 360 14. Chen CJ, Wu FT, Huang YC, Chang WC, Wu HS, Wu CY Lin JS, Huang FC, Hsiung
361 CA. Clinical and Epidemiologic Features of Severe Viral Gastroenteritis in Children:

- 362 A 3-Year Surveillance, Multicentered Study in Taiwan with Partial Rotavirus
363 Immunization. *Med*. 2015;*94*(33): e1372.
- 364 15. Igbokwe JI, Akinyede JO, Dang B, Alaga T, Ono MN, Nnodu VC, Anike LO.
365 Mapping and monitoring of the impact of gully erosion in South eastern Nigeria with
366 satellite remote sensing and Geographic Information System. *Intl Arch*
367 *Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2008;*37*:
368 (B8):875-872
- 369 16. Freeman MM, Kerin T, Hull J, McCaustl and K, Gentsch J. Enhancement of detection
370 and quantification of rotavirus in stool using a modified real-time RT-PCR assay. *J*
371 *Med Virol*. 2008;*80*:1489-1496.
- 372 17. Hoehne M, Schreier E. Detection of Norovirus genogroup I and II by multiplex real-
373 time RT-PCR using a 3'-minor groove binder-DNA probe. *BMC Infect Dis*.
374 2006;*6*(1):1-6.
- 375 18. Kageyama T, Kojima S, Shinohara M, Uchida K, Fukushi S, Hoshino FB, Takeda N,
376 Katayama K. Broadly reactive and highly sensitive assay for Norwalk-like viruses
377 based on real-time quantitative reverse transcription-PCR. *J Clin Microbiol*. 2003;
378 41:1548-57.
- 379 19. Oka T, Katayama K, Hansman GS, Kageyama T, Ogawa S, Wu FT, White PA,
380 Takeda N. Detection of human sapovirus by real-time reverse transcription-
381 polymerase chain reaction. *J Med Virol*. 2006;*78*:1347-1353.
- 382 20. Heim A, Ebnet C, Harste G, Pring-Akerblom P. Rapid and quantitative detection of
383 human adenovirus DNA by real-time PCR. *J Med Virol*. 2003;*70*:228-239.
- 384 21. Logan C, O'Leary JJ, O'Sullivan N. Real-time reverse transcription-PCR for detection
385 of rotavirus and adenovirus as causative agents of acute viral gastroenteritis in
386 children. *J Clin Microbiol*. 2006;*44*:3189-3195.

- 387 22. Jothikumar N, Cromeans TL, Robertson BH, Meng XJ, Hill VR. A broadly reactive
388 one-step real-time RT-PCR assay for rapid and sensitive detection of hepatitis E virus.
389 J Virol Methods. 2006;131:65-71.
- 390 23. Pietsch C, Liebert UG. "Genetic diversity of human parechoviruses in stool samples,
391 Germany." Infect Genet Evol. 2019;68:280-285.
- 392 24. Escutenaire S, Mohamed N, Isaksson M, Thoren P, Klingeborn B, Belák S, Berg M,
393 Blomberg J. SYBR Green real-time reverse transcription-polymerase chain reaction
394 assay for the generic detection of coronaviruses. Arch Virol. 2007;152(1):41-58.
- 395 25. Pietsch C, Schuster V, Liebert UG. A hospital based study on inter-and intragenotypic
396 diversity of human rotavirus A VP4 and VP7 gene segments, Germany. J Clin Virol.
397 2011;50(2):136-141.
- 398 26. Marques AM, Diedrich S, Huth C, Schreier E. Group A rotavirus genotypes in
399 Germany during 2005/2006. Arch Virol. 2007;152(9):1743-1749.
- 400 27. PSPP statistical software. Downloaded from
401 <https://www.gnu.org/software/pspp/get.html>
- 402 28. Agócs MM, Serhan F, Yen C, Mwenda JM, de Oliveira LH, Teleb N, Wasley A,
403 Wijesinghe PR, Fox K, Tate JE, Gentsch JR. WHO global rotavirus surveillance
404 network: a strategic review of the first 5 years, 2008–2012. MMWR Morb Mortal
405 Wkly Rep. 2014;63(29):634.
- 406 29. Williams AR, Bain RES, Fisher MB, Cronk R, Kelly ER, Bartram J. A Systematic
407 Review and Meta-Analysis of Fecal Contamination and Inadequate Treatment of
408 Packaged Water. Li M, ed. PLoS One. 2015; 10(10): e0140899.
- 409 30. Adah MI, Wade A, Taniguchi K. Molecular epidemiology of Rotaviruses in Nigeria:
410 detection of unusual strains with G2P [6] and G8P [1] specificities. J Clin Microbiol.
411 2001;39(11):3969-3975.

- 412 31. Tagbo BN, Mwenda JM, Armah G, Obidike EO, Okafor UH, Oguonu T, Ozumba
413 UC, Eke CB, Chukwubuike C, Edelu BO, Ezeonwu BU. Epidemiology of rotavirus
414 diarrhea among children younger than 5 years in Enugu, South East, Nigeria. *Pediatr*
415 *Infect Dis J.* 2014;33:S19-22.
- 416 32. Tagbo BN, Mwenda JM, Eke CB, Edelu BO, Chukwubuike C, Armah G, Seheri ML,
417 Isiaka A, Namadi L, Okafor HU, Ozumba UC. Rotavirus diarrhoea hospitalizations
418 among children under 5 years of age in Nigeria, 2011–2016. *Vaccine.*
419 2018;36(51):7759-7764.
- 420 33. Japhet MO, Famurewa O, Iturriza-Gomara M, Adesina OA, Opaleye OO, Niendorf S,
421 Bock CT, Mas Marques A. Group A rotaviruses circulating prior to a national
422 immunization programme in Nigeria: Clinical manifestations, high G12P [8]
423 frequency, intra-genotypic divergence of VP4 and VP7. *J of Med Virol.*
424 2018;90(2):239-249.
- 425 34. Clemens J, Rao M, Ahmed F, Ward R, Huda S, Chakraborty J, Yunus M, Khan MR,
426 Ali M, Kay B. et al. Breast-feeding and the risk of life-threatening *Rotavirus*diarrhea:
427 prevention or postponement? *Pediatr.* 1993;92(5):680–685
- 428 35. Golding J, Emmett PM, Rogers IS. Gastroenteritis, diarrhoea and breast feeding.
429 *Early Hum Dev.* 1997; 49 (Suppl): S83–103. doi: 10.1016/S0378-3782(97)00055-8.
- 430 36. Aminu M, Ahmad AA, Umoh JU, De Beer MC, Esona MD, Steele AD. Adenovirus
431 infection in children with diarrhea disease in North western Nigeria. *Ann Afr Med.*
432 2007;6(4):168-173.
- 433 37. Ayolabi C, Ojo D, Akpan I. Astrovirus Infection in Children in Lagos, Nigeria. *Afr J*
434 *Infect Dis.* 2012;6(1):1-4
- 435 38. Munjita SM, Current Status of Norovirus Infections in Children in Sub-Saharan
436 Africa. *J Trop Med.* 2015;1-7 doi:10.1155/2015/309648

- 437 39. Shioda K, Cosmas L, Audi A, Gregoricus N, Vinjé J, Parashar UD, Hall AJ.
438 Population-Based Incidence Rates of Diarrheal Disease Associated with Norovirus,
439 Sapovirus, and Astrovirus in Kenya. PLoS One. 2016;11(4):e0145943.
440 <http://doi.org/10.1371/journal.pone.0145943>
- 441 40. Wildenbeest JG, Benschop KS, Minnaar RP, Jongh SB, Wolthers KC, Pajkrt D.
442 Clinical relevance of positive human parechovirus type 1 and 3 PCR in stool samples.
443 Clin Microbiol Infect. 2014;20(10):O640-0647.
- 444 41. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT
445 method. Nat Protoc. 2008;3(6):1101-1108.
- 446 42. Ianiro G, Delogu R, Baba M, Oderinde BS, Dawurung J, Ruggeri FM, Fiore L.
447 Molecular characterization of group A rotavirus strains detected in children with
448 diarrhea admitted to Nigerian hospitals in 2013. Arch Virol. 2015;160(6):1511-1517.
- 449 43. Motayo BO, Faneye AO, Adeniji JA. Epidemiology of rotavirus A in Nigeria:
450 Molecular diversity and current Insights. J Pathog. 2018;2018:1-7
- 451 44. Japhet MO, Famurewa O, Iturriza-Gomara M, Adesina OA, Opaleye OO, Niendorf S,
452 Bock CT, Mas Marques A. Group A rotaviruses circulating prior to a national
453 immunization programme in Nigeria: Clinical manifestations, high G12P [8]
454 frequency, intra-genotypic divergence of VP4 and VP7. J Med Virol. 2018;90(2):239-
455 49.
- 456 45. Jagai JS, Sarkar R, Castronovo D, Kattula D, McEntee J, Ward H, Kang G, Naumova
457 EN. Seasonality of rotavirus in South Asia: a meta-analysis approach assessing
458 associations with temperature, precipitation, and vegetation index. PloS One. 2012
459 May 31;7(5):e38168.
- 460 46. Tagbo BN, Chukwubike C, Mwenda JM, Seheri ML, Armah G, Mphahlele JM,
461 Ozumba UC, Benjamin-Puja C, Azubuike C, Okafor HU, Nnani RO. Molecular

462 characterization of rotavirus strains circulating in Enugu Nigeria: 2011 to 2016.

463 World J Vaccines. 2019;9(01):22.

464

465

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484