Reproductive health needs of Human papillomavirus (HPV) Positive women: A systematic 1 review 2 Running Title: HPV women Reproductive health needs 3 4 Mina Galeshi¹, Hoda Shirafkan², Shahla Yazdani³, Zahra Motaghi^{4*} 5 1. Student Research Committee, School of Nursing and Midwifery, Shahroud University of 6 Medical Sciences, Shahroud, Iran 7 2. Social Determinants of Health Research Center, Health Research Institute, Babol 8 University of Medical Sciences, Babol, Iran. 9 3. Cancer Research Center, Health Research Institute, Babol University of Medical 10 11 Sciences, Babol I.R.Iran 4. Department of Reproductive Health, School of Nursing and Midwifery, Shahroud 12 13 University of Medical Sciences, Shahroud, Iran 14 15 16 17 Corresponding Author: DR .Zahra Motaghi Email: galeshi m@yahoo.com 18 Phone:00989119119331 19 Address: Shahroud University of Medical Sciences and Health Services, Hafte Tir Square, 20 21 Shahroud, Iran 22 Fax: 009823-32391609 23 Postal Code: 3614773955 24 25 26 27 28 29

31 Abstract

30

- 32 **Background:**Human papillomavirus is one of the most important transmitted viruses that causes
- 33 of cervical cancer.
- In women undergoing cervical screening, it is important to be aware of ways to reduce anxiety.
- 35 The present systematic review conducted to determine the reproductive health needs of women
- with HPV.
- 37 **Methods:** In this systematic review, articles without time constraints searched in PubMed,
- 38 Scopus, Web of Science, Google Scholar and Iranian Magiran, SID and Iranmedex.
- 39 Keywords used: HPV, Information, Want, Need, Know, etc. and their Persian equivalents in the
- 40 title and abstract of the articles.
- 41 Papers after identification by two researchers Contradictions evaluated and discussed with the
- 42 third author.
- 43 **Results:** At first 1056articles retrieved which after removing, 13articles published between 2004-
- 44 2021 were entered. The studies were qualitative (N=9), quantitative (N=3), and one was
- unclear.Most qualitative studies collected data using individual interviews(N=7), two qualitative
- studies, narratives of HPV patients from a website of patient experiences and questions. The
- 47 quality evaluation showed that good=8, Average=2, and one was of poor. There is not enough
- 48 information about Friedman AL et al and Garland SM articles to check the quality, but since
- 49 these studies were valuable, they included in the study according to the opinion of the research
- 50 team.

59

- Conclusion: Surveys showed that the majority of women had unanswered questions about their
- 52 HPV test results. The information that women thought was helpful in interpreting their test
- results included having a high-risk type of HPV, the risk of short-term and long-term cancer, and
- 54 cancer survival statistics for the virus. Women also needed information about sexual
- 55 transmission, how HPV tested positive in a long-term relationship, and the potential
- consequences for their partners and the risk of re-infection. Younger women had questions about
- whether HPV could affect fertility.
- 58 **Keywords:** HPV, Reproductive health, need

Introduction

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

Human papillomavirus (HPV) is one of the most important sexually transmitted viruses (1). This virus is one of the causes of cervical cancer (2). Although this cancer has many environmental and genetic causes (such as smoking, early marriage, high parity, etc.), the most important cause is infection with HPV (3). Cervical cancer is the second most common cancer in women, with 500,000 new cases reported each year (4). Various and extensive studies have been conducted to investigate the prevalence of HPV infections in different countries as well as in Iran, which indicates its increasing prevalence in the world (5-7). The prevalence of this virus in the world is between 10.5-55.4% and in Iran, it is 49.5% (6, 7). The prevalence of genital HPV infection in the United States among women aged 18 to 59 years was 40% for all types of HPV and 20% for high-risk HPVs (8). In general, cervical cancer is both preventable and treatable, and its prevalence is 6 times higher in developing countries than in developed countries, mainly due to differences and the lack of appropriate screening programs in these countries (9). More than 95% of cervical cancers are attributed to the papillomavirus (10). More than 100 strains of the virus have been identified, but types 16 and 18 are the most important strains of the virus in developing cervical cancer (1). Types 6 and 11 of this virus cause warts in the vulva (11), vagina, cervix, and penis (12). This disease affects the quality of life of individuals and imposes financial costs on health systems (13). It has been found that 35-65% of people become infected with HPV during sex (14). Understanding cervical cancer, along with being involved with HPV infection, raises questions about needs and preferences. In women undergoing cervical cancer screening, it is important to be aware of ways to minimize anxiety. Previous research has been more limited to the field of HPV DNA testing (15, 16). HPV DNA testing included information on HPV disease, prevention, treatment, and risk of cancer (17). Having an HPV test and getting the result can cause undesirable psychosocial reactions. To prevent anxiety and psychological distress, education and counseling are increasingly important for women diagnosed with HPV. Therefore, more attention should be paid to how to communicate and provide information (18-20).

Information challenges and communication gaps for women with HPV in the stages of cervical 89 cancer screening in primary care, waiting time to refer to a specialist, first consultation, and 90 91 Occurs after consultation in specialized care. Relatively little research has shown that health care providers have knowledge gaps and there is no adequate flow of HPV information between 92 health care providers and women with HPV (21, 22). 93 Counseling with HPV patients poses special challenges for physicians and patients, especially in 94 communities where HPV is rare and associated with severe stigma. It will be invaluable to 95 provide a deep understanding of the challenges, perspectives, experiences, and needs of women 96 97 living with HPV positively about healthcare providers at the primary and specialist care levels. 98 Therefore, the present systematic review was performed to determine the needs of women with HPV. 99 **Material and Methods** 100 101 The present study was approved by the Ethics Committee of Shahroud University of Medical Sciences: IR.SHE.REC.1400.154 and after receiving the code from the Prospero system Code: 102 CRD42021293223 Based on the proposed systematic review and meta-analysis checklist 103 (PRISMA) was done. 104 105 Search strategy for identifying papers Related articles, with electronic search in medical databases such as: PubMed, SCOPUS, Web of 106 Science, Google Scholar search engine and Iranian Magiran, Scientific Information Database and 107 Iranmedex and additional articles with a gray literature search using OpenGrey (www.open 108 109 grey.eu)were identified. Without language restrictions and the date of publication of the article until December 2021 was reviewed. 110 Keywords used included: (HPV Human Papillomavirus), Alphapapillomavirus, (Human 111 Papillomavirus, HPV), Information, Want, Need, Know, Qualitative Research, theoretical 112 framework, Comparative Analysis, and their Persian equivalents and the combination of 113 114 keywords with or, and in the title and abstract of the articles.

Selection process 117 As the reproductive health needs of women with HPV were not fully understood, a 118 comprehensive and systematic study was conducted on the subject. Inclusion criteria include 119 Studies related to the needs of women with HPV, Conference-related abstracts, and Exclusion 120 criteria: Studies related to HPV in men, Complications of HPV, HPV-related risk factors, 121 cervical cancer, Treatment of cervical cancer or colposcopy, commentaries, opinion pieces, and 122 editorials. 123 Data extraction 124 125 All the articles obtained in the introductory search at first reviewed by title, then Abstract, and finally their full text. The articles were reviewed by two people independently. The names of the 126 authors and the journal were not hidden from reviewers. Disagreements between reviewers were 127 decided by negotiating with a third reviewer until a final agreement was reached. The data 128 129 extracted for analysis including the author's name, year of the study, place of the study, study method, and the sample size in each group were entered into the electronic datasheet. 130 Researchers contacted their authors for more information when they could not retrieve articles 131 from authorized databases. All studies were reviewed for duplication. For each study, the data is 132 listed in the table based on the study checklist. 133 Quality assessment 134 A quantitative/qualitative adjusted checklist (CASP) was used to critically evaluate the articles. 135 The Checklist (CASP) is a standard tool for evaluating articles, developed by the JAMA Group 136 in 1994, and is the oldest and most widely used critical evaluation tool for a variety of 137 articles/studies (23). 138 139 The present checklist was 10 items to help understand qualitative research. How to use this tool for evaluating qualitative studies is as follows: When evaluating, three general issues should be 140 considered first. A: Are the study results valid enough?, B: What are the results? A: Will the 141 results help locally? The first two questions are systematically screening questions and can be 142 answered quickly. If the answer to both is "yes", the remaining questions are continued each item 143 144 was assigned a score of 2 (including that item in the article) or zero (not paying attention to the

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

(Flowchart).

item in the article) and one (cannot be said) and the total score of this checklist was between 0-20. To evaluate cross-sectional studies, this tool consisted of 18 items, and each item was given a score of one. Including that item in the article (or zero) was considered as not paying attention to the item in the article. Attitude evaluation (3 items), study design (5 items), and results (5 items) were divided and the total score of this checklist varied between 0-18. After carefully reading the full text of each article, an article quality evaluation checklist was completed by the first researcher and items were scored. Reassessment was performed in the same way by the second researcher. If there was no agreement on scoring the items, the final score would be taken in a joint session. In the next step, these articles were compared in terms of scores obtained in each of the four areas as well as in terms of total scores. Also, the qualitatively reviewed articles were divided into three categories of good, medium, and poor quality articles according to the scores obtained from this checklist. Total score 20 high quality; 16-19 Medium quality; ≤15 was considered low quality, also the articles reviewed cross-sectional in consultation with experts, scores of 75% of the total score and above as good quality (score 13), scores between 25-25% of the total score as average quality (scores 6-12) and scores less than 25% of the total score (score 5 and below) were divided as poor quality (24, 25). Analysis Ouantitative and qualitative findings were analyzed separately. The results of quantitative studies were reported descriptively by first creating an initial combination of findings, existing relationships and data strength were examined. For qualitative studies, a thematic synthesis was performed by examining the text line by line in the results section and discussing and then identifying descriptive topics. **Results** In the search of databases, at first 1056 articles were retrieved, which after removing 963 duplicate and unrelated articles, 61 articles were evaluated based on inclusion criteria

- Studies were conducted in the United States (N=5), Europe (N=4), Asia (N=4), and published
- between 2004-2021. The studies were mostly qualitative (N=9), the study was quantitative
- 175 (N=3), and one type of study was unclear. Most qualitative studies collected data using
- individual interviews (N=7), two qualitative studies, narratives of HPV patients from a website
- of patient experiences and questions. The characteristics of the participant and the study are
- shown in the table (Table 1).
- The results of the overall quality evaluation of these articles showed that 8 articles were of good
- quality, 2 were of average quality and one was of poor quality. There is not enough information
- about Friedman AL et al and Garland SM articles to evaluate the quality, but since these studies
- were valuable, they were included in the study according to the opinion of the research team.
- The results of the article evaluation are divided into four areas of the checklist, the total score of
- each article, and article quality in (Tables 2 and 3).
- Summary of the needs presented by HPV women in qualitative studies including:
- People with the virus cognition about HPV and cervical cancer, but did not have the necessary
- 187 knowledge and awareness about it (26). They were confused about the nature of the disease, its
- severity, transmission, types, treatment, and instructions for the HPV, and needed to have
- complete information (27). Therefore, it is necessary to help patients understand the results of
- their test and their importance. Do not experience unnecessary fear, anxiety, or stigma and seek
- 191 follow-up and recommendations to meet the patient's needs.
- 192 Signs, Symptoms, Transmitted, Cause, Consequences of HPV
- 193 People with the cause of HPV and how it is transmitted, the signs and symptoms of HPV
- infection and cervical cancer, how to get screened for cervical cancer, the consequences of the
- virus and its effects (for example, does the virus always cause cervical cancer?), Timeline (for
- 196 example, how long does HPV cause cervical cancer?), Control and treatment (can it be
- 197 prevented?), Information on how long the infection stays in the body, And does it lead to
- 198 cancer? They needed to have complete information because they had stated that it would reduce
- 199 their anxiety (25, 28-30).
- 200 HPV and sexual partner
- Women with HPV needed counseling on how to communicate and deal with a sexual partner.
- Because some women expressed concern about their husbands' infidelity and a lack of trust was
- 203 expressed between the couple (31).

When I got HPV, How do I tell my partner what to say? What effect does HPV have on sexual 204 life? Because sometimes these issues lead to uncertainty and stress and in severe cases lead to the 205 206 breakdown of relationships (28). The most common question asked by women with HPV was: which sexual partner (active or 207 former) did the infection come from? Did my partner get it from a previous relationship or did 208 he or she betray me? When HPV is diagnosed in a couple, what are the consequences for the 209 spouse and sexual partner? Will her sexual partner get it too? Should her partner test and follow 210 up? When a couple is infected with HPV and is being treated, will they get the virus again from 211 their partner? Will they constantly infect each other? Will they use a condom? Can I have sex? 212 Can a person get HPV orally or by hand? (32, 33). 213 Seeking needed information by women with HPV 214 Because many women with the condition felt that they did not have enough information about 215 HPV, many sought more information, including the Internet (social media, blogs, and websites 216 provided by labs and private professionals) (34). Their health care providers and other women 217 living with HPV). They found finding information reliable and up-to-date challenging. Most 218 219 women prefer the information provided by official websites. Women who were regular Internet users reported finding useful information (31). However, it was often not easy for them to 220 interpret and determine the correctness or incorrectness of the material. Women reported that 221 most cases of finding information about HPV, along with the effect of the virus on genital warts 222 223 and other sexually transmitted diseases, in which risk factors such as having multiple sexual partners were highlighted, some women were very upset and experienced a sense of shame and 224 225 embarrassment (35). Or, in many cases, this information was in addition to information related to uterine cancer, which led to fear, anxiety, and stress in them. It seems that the information that 226 227 women are looking for and the way their doctor informs them can affect their psychological response to HPV infection and is one of the issues that is considered important in follow-up (36). 228 HPV and social consequences 229 Dysfunction or inability to perform daily activities, limitation in relationships with others, 230 231 limitation of social activities, shock, insecurity, and fear of deformity of the limbs following the development of genital warts and its impact on quality of life were among the challenges (37). 232

Women with HPV experienced it during the illness. Therefore, women with HPV need

psychosocial care at different stages of the disease, so that non-fulfillment of psychosocial needs 234 can complicate the course of treatment and recovery (38). 235 236 Reproductive health needs in women with HPV Women with HPV raised some issues related to fertility, and the need for proper counseling on 237 the impact of the virus on reproductive health was felt (34). 238 Many women reported seeking information about fertility and pregnancy from various sources. 239 Many women seek information about the adverse effects of HPV on male fertility, the negative 240 impact of HPV on female fertility, the effect of treatments and vaccinations on female fertility 241 and its relationship to female fertility, maternal health threats during pregnancy, adverse 242 pregnancy outcomes, fetal harm, Fear of having a newborn This fear was so great that some 243 people who decided to become pregnant postponed their pregnancy program until the cytology 244 results improved so that they could receive appropriate counseling during this time. Women also 245 noted the association between genital HPV infection and various maternal-fetal variables and 246 pregnancy complications such as miscarriage and preterm delivery Therefore, these women need 247 to know what to do to protect their reproductive health and reduce its complications against 248 249 infection (30). Since genital warts can multiply during pregnancy, removing warts during pregnancy was one of 250 the most important issues. Women with genital warts have revealed the need for information 251 about the teratogenicity of some wart removal treatments. They were anxious about any threat it 252 253 might pose to the fetus (37). Information on contraception, fear of premature menopause, fear of cervical cancer, fear of 254 255 hereditary cancer caused by HPV were other issues raised in this area (30). To prevent unplanned pregnancies, women with HPV needed more information to choose the 256 257 preferred method of contraception. Because long-term use of birth control pills increases the risk of cervical cancer for women with persistent HPV, they needed guidance in changing their 258 method of contraception. Women sought information on the negative effects of combined oral 259 contraceptive pills (COCs) and levonorgestrel (LNG) pills on their cellular changes (39). 260 261 HPV screening and cervical cancer and vaccinations 262 Another piece of information that infected women needed to know about was screening. They needed to know how helpful these screenings can be in diagnosing uterine cancer (37). These 263

include: How is screening done? Is a Pap smear enough to diagnose? Can HPV be diagnosed

- before cancer develops? In addition to annual checkups, what type of checkups are started at
- what intervals and at what age? What does the [HPV] test include? Can I get an HPV test from
- a doctor? How long does it take to determine the result of an HPV test? What is the difference
- between an HPV test and a Pap smear? (28)
- The women also asked for information about the effect of the vaccine, side effects, long-term
- effects, mechanism of action of the vaccine, number of doses received, and the age range for
- vaccination (29).

Quantitative studies

- SungJong Lee, et al. In their quantitative study showed 10 categories for topics and sample
- 274 questions with different topics. Of the 3,062 people who visited the HPV website, 2,330 asked
- 275 general questions and 732 asked private questions. The type and frequency of public and private
- 276 questions showed a statistical difference between the two groups (p<0.001).
- 277 From 2004 to 2011 the most common questions, respectively: a. Treatment of HPV and cervical
- dysplasia (37.8) 1156, b. How HPV transmission (22.3) 684, c. HPV regression (15.7) 481, d.
- Genital warts (5.7) 174, e. Risk factors for cervical cancer (4.8) 147, f. HPV vaccination 116
- 280 (8.3), g. Prevention of HPV 90 (9.2), h. Vaginal discharge (2.8) 86, i. HPV and pregnancy (2.6)
- 281 81, j. Laryngeal cancer and HPV 47 (1.5) was (37).
- Sandra Millon Underwood et al. showed that sexual behavior and chronic HPV infections are
- 283 risk factors for cervical cancer. Sexual behavior and chronic HPV infections are risk factors for
- cervical cancer. When asked to describe their sexual activity, 76.8% (352) of women reported
- having had sexual activity in the past six months. Among them, 2.62 (219) reported, having a
- sexual partner, 15.6% (55) two sexual partners and 22.2% (78) three or more partners in the past
- six months. The most common questions included: cause of HPV, cause of cervical cancer,
- signs, and symptoms of HPV infection, signs and symptoms of cervical cancer, HPV and
- cervical cancer screening, HPV vaccine, HPV and cervical cancer risk management (28).
- 290 Kathrin Milbury, et, al in their study showed that 66% of patients were aware of their HPV
- status, but only 35 percent identified HPV as a possible cause of their cancer. Most patients
- 292 disclosed their HPV status to their partner, 41% talked about transmitting the virus, and only
- 293 23% felt aware of the potential risks of transmission and precautions. 39% of their oncologists
- want to discuss more HPV issues, and 58% sought it from other sources. More than a third said
- 295 they were interested in more information about HPV.

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

Women are interested in receiving any information. 18% want more information on how HPV causes cancer. 15% wanted information on HPV vaccination (especially whether vaccinating their children to prevent cancer), 10% wanted information on how to prevent transmission to their partner, and 10% wanted to know if there was any cure for Is there HPV or not? (33). **Discussion** Based on the review, this systematic review is one of the few review studies that comprehensively addresses the needs of women with HPV. The potential advantage of this study is that the HPV-related questions and needs raised by patients were collected. Recently, many researchers have used written surveys, interviews, and other data collection methods to assess knowledge about HPV among women. They found that women have relatively little knowledge of HPV and suggested using more training programs (40, 41). In addition, many patients become confused after hearing their doctor's description of HPV. As a result, many patients search the Internet for seeking information about HPV, but there are still many questions for them (42). According to the findings, most people, especially when it comes to frequently asked questions, really want to find out about the whole disease process, from HPV infection to recovery (35). Since most of the studies studied were qualitative studies, this study was able to identify a wide range of needs of affected women. The needs of women with HPV lead to fear, anxiety and worry. Their anxiety is largely due to their poor knowledge of the virus. Therefore, women who are screened for or exposed to the virus also need quality information about HPV and its role in cervical cancer when faced with an abnormal result. Physicians potentially play an important role in modulating the effects of diagnosis through how HPV is diagnosed (43). Therefore, recognizing these needs can help health professionals understand what questions they are expected to answer. KirstenMcCaffery and Les Irwig showed in her study that women needed more information about different types of HPV, transmission, effects of the virus on sexual partners, prevalence, delay and regression of HPV, management options, and infection consequences for cancer and fertility risk. Uncertainty about key aspects of HPV, how the physician announced the outcome, and how the outcome was presented (letter, telephone, or consultation) affected the psychological response of women with HPV (35). Women's experience of searching the Internet for more information on HPV has been reported as problematic, anxious, and stigmatizing infection in the community, as the information provided

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

is often related to other sexually transmitted infections and multiple sexual partners as a risk factor for Infections are prominent (44). Sung-Jong Lee shows in his study that when people go to a website about HPV, they are looking for "HPV treatment and cervical dysplasia", "HPV transmission", "HPV cure", "Genital warts" And "risk of cervical cancer." Also, our results showed that "genital warts" and "HPV and pregnancy" may be considered embarrassing topics. Therefore, these findings can be used to provide informed recommendations for clinical or Internet-based communication with patients and the general public (37). Sandra Millon Underwood In reviewing the questions reported by women with HPV, mentioned five related topics such as: cervical cancer, HPV/cervical cancer prevention, and risk management (28). Numerous questions asked by women focused on the origin of HPV, the different types of HPV, HPV and risk factors for cervical cancer, the risk factors associated with HPV transmission, and the development of cervical cancer. Similar questions were asked about "early signs", "warning signs", symptoms associated with HPV infection and cervical cancer, and if, when and how often women should be tested. However, most of the questions asked by women focused on HPV and cervical cancer prevention and risk management. Among them were specific questions about how HPV was transmitted. Questions about the proper use and effectiveness of male and female inhibitory methods (eg. condoms) to reduce the risk of HPV and cervical cancer; Questions about the safety, availability and cost of the HPV vaccine; and questions about the need for further community education and information (28). **Conclusion** Surveys showed that the majority of women had unanswered questions about their HPV test results. The information that women thought helped interpret their test results included having a high-risk type of HPV, the risk of short-term and long-term cancer, and cancer survival statistics for the virus. Women also needed information about sexual transmission, how HPV tested positive in a long-term relationship, and the potential consequences for their partners and the risk of re-infection. Younger women had questions about whether HPV could affect fertility. HPV information should be scientifically accurate and minimize any possible stigma that may be associated with the infection due to its sexually transmitted nature. It was found that seeking information from alternative sources also increased women's anxiety by providing information

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

Availability of data and materials

about HPV in the context of other highly sexually transmitted diseases and highlighting several sexual partners as a risk factor for infection. The strength of this study is that it was a comprehensive and systematic review using PRISMA guidelines. In addition, an extensive search strategy was performed without any history or language restrictions. It is possible that due to the range of terms that can be used to describe the needs, some eligible studies may not have been identified in our search, however, we did forward and backward citation searching to reduce this possibility. The data were also extracted by one author and reviewed independently by the second author. While qualitative synthesis allows us to identify a range of different factors that help women with HPV. However, since the majority of studies were qualitative, meta-analysis was not possible. It is suggested that future studies should measure the effect of raising awareness about HPV over time to determine whether raising awareness leads to improved health of infected women. Knowing when and how this response to needs is most effective can help determine the most appropriate interventions. **Abbreviations** HPV: Human Papillomavirus COCs: contraceptive pills LNG: levonorgestrel **Declarations Details of Ethics Approval** This study has been performed in accordance with the Declaration of Helsinki and has been reviewed and approved by the Ethics Committee of Shahroud University of Medical Sciences The: (IR.SHE.REC.1400.154) and after receiving the code from the Prospero system Code: CRD42021293223 Based on the proposed systematic review and meta-analysis checklist (PRISMA) was done. **Consent of participate** Not applicable **Consent for publication** Not applicable

The data that support the findings of this study are available from the corresponding author, [MG], upon reasonable request. **Competing Interests** The authors have no conflict of interest. **Funding** Not applicable Acknowledgments Not applicable **Authors' Contributions** MG, ZM,SHZ,HSH designed the studyHSH and ZM planned and undertook the analysis. MG,ZM, and HSH wrote the initial and subsequent drafts of the manuscript. MG,SHZ,ZM and HSH contributed to revising the manuscript. All authors read and approved the final manuscript.

References

- 1. Berek JS. Berek & Novak's Gynecology Essentials: Lippincott Williams & Wilkins;
- 414 2020.

- 415 2. Wu J, Ding C, Liu X, Zhou Y, Tian G, Lan L, et al. Worldwide burden of genital human
- 416 papillomavirus infection in female sex workers: a systematic review and meta-analysis.
- 417 International Journal of Epidemiology. 2021;50(2):527-37.
- 418 3. Nassar ACF, de Oliveira Garcia D, Salvio JRC, Altieri NGGL, Izique LP, Isper C. HPV
- infection as a risk factor of esophagus squamous cell carcinoma and it prognosis. Rev Med (São
- 420 Paulo). 2021;100(2):144-51.
- 421 4. Sethi S, Ali A, Ju X, Antonsson A, Logan R, Canfell K, et al. A systematic review and
- meta-analysis of the prevalence of human papillomavirus infection in Indigenous populations-A
- 423 Global Picture. Journal of Oral Pathology & Medicine. 2021.
- 424 5. Bruce J, Drury N, Poobalan AS, Jeffrey R, Smith WCS, Chambers WJP. The prevalence
- of chronic chest and leg pain following cardiac surgery: a historical cohort study. 2003;104(1-
- 426 2):265-73.
- 427 6. Escobar N, Plugge E. Prevalence of human papillomavirus infection, cervical
- 428 intraepithelial neoplasia and cervical cancer in imprisoned women worldwide: a systematic
- review and meta-analysis. J Epidemiol Community Health. 2020;74(1):95-102.
- 430 7. Kesheh MM, Keyvani H. The prevalence of HPV genotypes in Iranian population: An
- 431 Update. Iranian journal of pathology. 2019;14(3):197.
- 432 8. McQuillan GM, Kruszon-Moran D, Markowitz LE, Unger ER, Paulose-Ram R.
- Prevalence of HPV in adults aged 18-69: United States, 2011-2014: US Department of Health
- and Human Services, Centers for Disease Control and ...; 2017.
- 435 9. Safaei A, Khanlari M, Momtahen M, Monabati A, Robati M, Amooei S, et al. Prevalence
- of high-risk human papillomavirus types 16 and 18 in healthy women with cytologically
- negative pap smear in Iran. Indian Journal of Pathology and Microbiology. 2010;53(4):681.
- 438 10. Sasieni P, Castanon A. Call and recall cervical screening programme: Screening interval
- and age limits. Current Diagnostic Pathology. 2006;12(2):114-26.
- 440 11. Wakeham K, Kavanagh K, Cuschieri K, Millan D, Pollock KG, Bell S, et al. HPV status
- and favourable outcome in vulvar squamous cancer. International journal of cancer.
- 442 2017;140(5):1134-46.
- 12. Dillner J, Kjaer SK, Wheeler CM, Sigurdsson K, Iversen OE, Hernandez-Avila M, et al.
- 444 Four year efficacy of prophylactic human papillomavirus quadrivalent vaccine against low grade
- 445 cervical, vulvar, and vaginal intraepithelial neoplasia and anogenital warts: randomised
- controlled trial. BMJ (Clinical research ed). 2010;341:c3493.
- 447 13. Leung W, Chan P, Lau K, Ho K. The prevalence of human papilloma virus in the anal
- 448 region of male Chinese attendees in three public sexually transmitted disease clinics in Hong
- Kong. Hong Kong J Dermatol Venereol. 2011;19(1):6-13.
- 450 14. Lo Cigno I, Calati F, Albertini S, Gariglio M. Subversion of host innate immunity by
- human papillomavirus oncoproteins. Pathogens. 2020;9(4):292.
- 452 15. Anhang R, Goodman A, Goldie SJ. HPV communication: review of existing research and
- recommendations for patient education. CA: A Cancer Journal for Clinicians. 2004;54(5):248-
- 454 59.

- 455 16. Ciavattini A, Carpini GD, Giannella L, Del Fabro A, Banerji V, Hall G, et al. An online
- 456 survey on emotions, impact on everyday life, and educational needs of women with HPV
- positivity or abnormal Pap smear result. Medicine. 2021;100(45).
- 458 17. Waller J, McCaffery K, Nazroo J, Wardle J. Making sense of information about HPV in
- 459 cervical screening: a qualitative study. British journal of cancer. 2005;92(2):265-70.
- 460 18. Bennett KF, Waller J, Ryan M, Bailey JV, Marlow LA. Concerns about disclosing a
- high-risk cervical human papillomavirus (HPV) infection to a sexual partner: a systematic review
- and thematic synthesis. BMJ sexual & reproductive health. 2021;47(1):17-26.
- 463 19. McBride E, Marlow LA, Forster AS, Ridout D, Kitchener H, Patnick J, et al. Anxiety and
- distress following receipt of results from routine HPV primary testing in cervical screening: The
- psychological impact of primary screening (PIPS) study. International journal of cancer.
- 466 2020;146(8):2113-21.
- 467 20. Dodd RH, Mac OA, McCaffery KJ. Women's experiences of the renewed National
- 468 Cervical Screening Program in Australia 12 months following implementation: a qualitative
- study. BMJ open. 2020;10(7):e039041.
- 470 21. O'Connor M, O'Donovan B, Waller J, Céilleachair AÓ, Gallagher P, Martin C, et al.
- 471 Communicating about HPV in the context of head and neck cancer: A systematic review of
- 472 quantitative studies. Patient education and counseling. 2020;103(3):462-72.
- 22. Denny-Smith T, Bairan A, Page MC. A survey of female nursing students' knowledge,
- 474 health beliefs, perceptions of risk, and risk behaviors regarding human papillomavirus and
- cervical cancer. Journal of the American Academy of Nurse practitioners. 2006;18(2):62-9.
- 476 23. Schadewaldt V. How to use this appraisal tool 2015 [Available from:
- 477 <u>www.reache.files.wordpress.com/2010/03/cross-sectional-appraisal-tool.</u>
- 478 24. Njau B, Covin C, Lisasi E, Damian D, Mushi D, Boulle A, et al. A systematic review of
- 479 qualitative evidence on factors enabling and deterring uptake of HIV self-testing in Africa. BMC
- 480 Public Health. 2019;19(1):1289.
- 481 25. Bahri N, Latifnejad Roudsari R. A Critical Appraisal of Research Evidence on Iranian
- Women's Attitude towards Menopause. The Iranian Journal of Obstetrics, Gynecology and
- 483 Infertility. 2015;18(178):1-11.
- 484 26. Alsous MM, Ali AA, Al-Azzam SI, Jalil MHA, Al-Obaidi HJ, Al-Abbadi EI, et al.
- 485 Knowledge and awareness about human papillomavirus infection and its vaccination among
- women in Arab communities. Scientific reports. 2021;11(1):1-8.
- 487 27. Ulasan S, NAHRAWI N, MUSTAFA WA, KANAFIAH SNAM. Knowledge of Human
- 488 Papillomavirus (HPV) and cervical cancer among Malaysia Residents: A review. Sains
- 489 Malaysiana. 2020;49(7):1687-95.
- 490 28. Underwood SM, Ramsay-Johnson E, Browne L, Caines N, Dean A, Duval S, et al. What
- women in the United States Virgin Islands still want and need to know about HPV, cervical
- cancer, and condom use. Journal of National Black Nurses' Association: JNBNA. 2010;21(1):25.
- 493 29. Marlow LA, Wardle J, Waller J, Grant N. Human papillomavirus (HPV) information
- 494 needs: a theoretical framework. Journal of Family Planning and Reproductive Health Care.
- 495 2009;35(1):29-33.
- 496 30. Marlow L, Forster AS, McBride E, Rockliffe L, Kitchener H, Waller J. Information
- 497 needs among women taking part in primary HPV screening in England: a content analysis. BMJ
- 498 open. 2020;10(12):e044630.

- 499 31. Symmons SM, Waller J, McBride E. Testing positive for Human Papillomavirus (HPV)
- at primary HPV cervical screening: A qualitative exploration of women's information needs and
- preferences for communication of results. Preventive Medicine Reports. 2021;24:101529.
- Verhoeven V, Baay MF, Baay PE, Lardon F, Van Royen P, Vermorken JB. Everything you always wanted to know about HPV (but could not ask your doctor). Patient education and
- 504 counseling. 2010;81(1):101-5.
- 505 33. Milbury K, Rosenthal DI, El-Naggar A, Badr H. An exploratory study of the
- 506 informational and psychosocial needs of patients with human papillomavirus-associated
- oropharyngeal cancer. Oral oncology. 2013;49(11):1067-71.
- 508 34. Qaderi K, Mirmolaei ST, Geranmayeh M, Farnam F, Hasani SS. 'Does HPV affect my
- 509 fertility?'Reproductive concerns of HPV-positive women: a qualitative study. Reproductive
- 510 Health. 2021;18(1):1-11.
- 511 35. McCaffery K, Irwig L. Australian women's needs and preferences for information about
- 512 human papillomavirus in cervical screening. Journal of Medical Screening. 2005;12(3):134-41.
- 513 36. Anhang R, Wright Jr TC, Smock L, Goldie SJ. Women's desired information about
- 514 human papillomavirus. Cancer. 2004;100(2):315-20.
- 515 37. Lee S-J, Yun H-J, Lee K-H, Kim C-J, Park J-S. What questions do people ask on a
- 516 Human Papillomavirus website? a comparative analysis of public and private questions.
- 517 International journal of medical sciences. 2012;9(2):142.
- 518 38. Garland SM, Quinn MA. How to manage and communicate with patients about HPV?
- International Journal of Gynecology & Obstetrics. 2006;94:S106-S12.
- 520 39. Qaderi K, Geranmayeh M, Farnam F, Hasani SS, Mirmolaei ST. Iranian HPV-positive
- Women's Needs and Preferences in Relation to Receiving Health Services: A Qualitative Study.
- 522 2021.
- 523 40. Chen H, Zhang X, Wang W, Zhang R, Du M, Shan L, et al. Effect of an educational
- 524 intervention on human papillomavirus (HPV) knowledge and attitudes towards HPV vaccines
- among healthcare workers (HCWs) in Western China. Human Vaccines & Immunotherapeutics.
- 526 2021;17(2):443-50.
- 527 41. López N, Garcés-Sánchez M, Panizo MB, de la Cueva IS, Artés MT, Ramos B, et al.
- 528 HPV knowledge and vaccine acceptance among European adolescents and their parents: a
- 529 systematic literature review. Public health reviews. 2020;41(1):1-24.
- Head KJ, Imburgia TM, Zimet GD, Shew ML. Women's understanding of their pap and
- HPV test results: implications for patient–provider communication. Journal of Communication in
- 532 Healthcare. 2017;10(1):37-46.
- 533 43. Nagpal TS, Blake J. Stigma as a Barrier to Care for Human Papillomavirus: A
- 534 Commentary Summarizing Literature and Presenting Recommendations for Health Care
- 535 Providers. Journal of Obstetrics and Gynaecology Canada. 2020;42(10):1276-8.
- 536 44. Gold D. The psychosocial care needs of patients with HPV-related head and neck cancer.
- Otolaryngologic Clinics of North America. 2012;45(4):879-97.
- 538 45. Friedman AL, Shepeard H, editors. Making sense of HPV and Pap test results: What do
- women need and want to know? Findings from CDC qualitative research with women, ages 30-
- 540 65 years. The 2008 National STD Prevention Conference; 2008.




Figure 1: Flow diagram of study selection

Table1: Characteristics of studies in the review

No	Reference	Country	Aim	Participants (n)	Study design	Analysis	Disclosure outcomes report
1	Sophie Mulcahy Symmons, et al.2021(31)	England	investigate women's	■ 30 women ■ age 24 to 63 ■ HPV positive	■ In-depth semi-structured interview ■ Face-to-face interviews ■ 15 women with highly educated and 15 with low-educated women. ■ The women participated in the interview on average 35.5 days after receiving their HPV test results.	Data were coded using qualitative analysis software NVivo 12	Summary of information needs in general and among people with higher and lower education including: Get test results using letters Not knowing the test result before receiving the test result Poor knowledge about HPV Try and search to interpret the test result Content and structure of the received letter Request more information about HPV Request personalization of the sent letter Prioritize the test result by first mentioning the natural cytological result and then the HPV-related result Explain about the next follow-up and additional tests Search for information after receiving the test result Talk to your doctor about the test result Search for information after receiving the test result Search for information about the test result Receive appropriate information along with providing test results Introducing trusted websites

	ı	1	I	T .		I	
							 In the group of people with higher education, people preferred to receive Supportive information. But in the group of people with lower education in terms of understanding and receiving information Questions about test interpretation ✓ Be explicitly informed about the risk of uterine cancer. ✓ Ask for information about the cause of HPV and how it is transmitted ✓ The effect of HPV on gender and sexual partner ✓ The effect of HPV on fertility ✓ Lack of clear association between normal cytology and HPV positive test ✓ Lack of a clear link between HPV and other sexually transmitted diseases ✓ How to share the test result with your partner and sexual
							partner
2	Laura Marlow, et al.2020(30)	England	The aim of this study was to investigate more information needs among women undergoing primary HPV screening.	• 381 women • Age d 24-65 who have registered one or more responses.	 Qualitative study The postal questionnaire was completed immediately, 6 and 12 months after receiving the results of the human papillomavirus (HPV) test. Ask the participant an openended question entitled "Do you have an unanswered question about cervical screening and HPV testing?" 	content	A total of 921 women who underwent initial HPV screening returned their original questionnaire. A total of 507 free texts were registered (base = 239/931, 6 months = 762/110 and 12 months = 537/68). Results as: Reaction and understanding of test results (45%) 170 ✓ Emotional response ✓ Trust the results ✓ Meaning of results ✓ Impact on sexuality

			1					
						Was asked.		✓ Lack of general
								understanding
								✓ Epidemiology of HPV
								✓ Cause of HPV
								✓ Prevention / treatment of
								HPV
								✓ HPV vaccination
3	Qaderi, et	IRAN	The sim of this study was to	40 H	IPV	semi-structured interviews	aantant	The subject and subcategory of
3		IKAN	The aim of this study was to		1P V	semi-structured interviews	content	
	al.2020(39)		evaluate the needs and	positive			analysis	this study include:
			experiences of HPV positive	women				1. Communication and
			women receiving health					counseling skills
			services.					A. Discussion about HPV
								B. Emotional support and
								acceptance of the disease
								C. Provide recommendations
								D. Clinical considerations
								2. Commitment to professional
								principles
								A. Gain the trust of patients
								B. Avoid financial misconduct
								C. Assertiveness and privacy
								D. Clinical considerations
								3. Providers' knowledge about
								HPV
								A. Follow the screening
								instructions
								B. Avoid misconceptions
								C. Adopt multilateral approaches
								are a series and a series approaches
								Other needs and questions of
								patients include:
								✓ Search for information on
								HPV types, transmission,
]
								spread and removal of the
								virus from the body
								✓ Need to request information
								about other sexually
								transmitted diseases, the
								potential benefits of the HPV
								vaccine, and the risk of HPV-

				1			
							related cancers Questions about HPV cofactors for cervical cancer in women with abnormal cytology Participants look for ways to improve their safety and sexual health
4	Kathrin Milbury et al.2013(33)	U.S.A	The aim of this study was to evaluate the information and psychosocial needs of patients with oral and pharyngeal squamous cell carcinoma in HPV-positive individuals and to identify the psychosocial challenges associated with HPV-positive cancer.	HPV men and women are positive with the OPSCC and have relationships	■ Exploratory study ■ Use open-ended questions such as "What information and knowledge would you like to know about HPV and related cancer"? ■ Questionnaires used include: Gritz 8-item questionnaire to assess past and present nicotine use and drug interactions ■ 10 item questionnaire (AUDIT) consumption assessment	analyzed using SPSS v19	The results of this study include: 66% of patients were aware of their HPV status but only 35% of them considered HPV to be a possible cause of cancer. Most patients report their HPV status to their partner. 41% discussed virus transmission and only 23% were aware of potential risks and preventive measures. 39% of patients sought to talk to an oncologist for information on HPV issues, and 58% sought it from other sources. More than a third of them were interested in learning more about HPV. Women are interested in receiving any information. 18% want more information on how HPV causes cancer. 15% wanted information on HPV vaccination (especially whether vaccinating their children to prevent cancer), 10% wanted information on how to prevent transmission to their partner, and 10% wanted to know if there was

							any cure for Is there HPV or
							not?
							The findings suggest that the
							diagnosis of HPV + OPSCC
							may have important
							psychosocial consequences including stigma, self-blame,
							and social interaction problems
5	Dorothy	U.S.A	The aim of this study was to		Qualitative Study		✓ Women with HPV need
	Gold, et	0.S.A	understand a framework for	_	Quantative Study	_	psychosocial care at different
	al.2012(44)		describing and recognizing the				stages of the disease.
	a1.2012(44)		psychosocial factors and				✓ Shock, uncertainty, and fear
			challenges that HNC patients				of deformed limbs following
			face, including those with				genital warts, dysfunction, or
			HPV-related tumors during				disability. Patients with HNC
			their disease stages from				face physical and
			diagnosis, treatment, and				psychological challenges
			recovery to survival., it				during illness.
			shows.				✓ There is a high potential for
							psychosocial disorders,
							including anxiety and
							depression, during ongoing
							care, including survival.
							✓ Patients with HNC associated
							with human papillomavirus
							(HPV) are at risk for
							psychological stress due to
							their demographic
							characteristics as well as the
							viral cause of their tumors. Failure to meet psychosocial
							Failure to meet psychosocial needs can complicate the
							course of treatment and
							recovery.
6	SungJong		Identify the questions that	Of the 3062	Comparative Study	Chi-square	10 categories for topics and
	Lee1, et	Korea	women ask on the human	subjects who	Data on all questions raised	test.	sample questions with
	al.2012(37)		papillomavirus-related	visited the	between March 2004 and July	Statistical	different topics are shown in.
			website.	HPV website,	2011 were collected and	analysis was	
			(www.hpvkorea.org)	2330 asked	analyzed.	performed	visited the HPV website,
				general	All questions on the HPV	using SAS	2,330 asked general questions

T	T	1			Т	
			questions and	website were in Korean and in	program,	and 732 asked private
			732 asked	free text.	version 8.0.	questions.
			private	At the time of drafting, the text	P <0.05	
			questions.	was being translated from	values were	public and private questions
				Korean to English.	defined as	showed a statistical difference
				Data on all questions raised	significant.	between the two groups (p
				between March 2004 and July		<0.001).
				2011 were collected and		From 2004 to 2011, the most
				analyzed.		common questions are as
				All questions on the HPV		follows: Frequency and
				website were in Korean and in		percentage of each question
				free text.		are as follows.
				At the time of drafting, the text		1.Treatment of HPV and
				was being translated from		cervical dysplasia 1156(37.8)
				Korean to English.		Is HPV treatable? What
				_		treatment is recommended for
						HPV infection?
						What is the best treatment for
						my condition (for example,
						cervical dysplasia?)
						2. How HPV is transmitted 684
						(22.3)
						Ask for information about the
						routes of transmission and
						duration of HPV infection
						Do other people become infected
						with HPV by sharing towels or
						bowls of soup?
						Have I got HPV infection from
						my partner?
						Does skin warts spread to other
						parts of the body than to the
						genitals?
						Is genital warts transmitted to
						the baby during breastfeeding?
						3. HPV regression 481(15.7)
						How long does it take for an
						HPV infection to go away on its
						own?
						Does a negative HPV DNA test
	<u> </u>					

							really indicate that the HPV
							virus has been removed from the
							body?
							4 .Genital warts 174(5.7)
							Why do genital warts recur even
							after successful treatment?
							If a small lump is found in the
							genital area, is it a genital wart?
							5Risk factors for cervical
							cancer 147(4.8)
							What is the risk of cervical
							cancer after HPV infection?
							What is the risk of cervical
							cancer due to cervical dysplasia?
							6. HPV vaccination 116(3.8)
							How effective is HPV
							vaccination?
							Which company vaccine is most
							effective in preventing HPV?
							When can the vaccine be used?
							7. Prevention of HPV 90(2.9)
							How to prevent HPV infection?
							Are Diets Good for HPV
							Prevention?
							Can condoms always prevent
							HPV infection?
							8. Vaginal discharge 86 (2.8)
							Is vaginal discharge associated
							with symptoms of HPV
							infection?
							9. HPV and pregnancy 81(2.6)
							Is HPV transmitted from mother
							to baby during pregnancy?
							10.Laryngeal cancer and HPV
							47(1.5)
							If a lump or lump is felt in the
							mouth, is this laryngeal cancer
							associated with HPV?
7	Sandra	U.S.A	The aim of this study was to	4 single	cross-sectional, mixed methods,	Content	Women age 18 to 44 years.
'	Millon		obtain the required	women	exploratory survey design was		or Sexual behavior, chronic
		l	1 Toganou	1	- Processed and and and and and and and and and an	1	a de la companya de l

	1.0	1 1 1 1 1	1::	LOCAL TIPLE
Underwood	information about HPV,	used in this study	qualitative	infections, HPV, are risk factors
,etal.2010	cervical cancer and condom		stdy	for cervical cancer.
(28)	use.		Quantitative	76.8% (352) of women reported
			data analysis	having had sexual activity in the
			of SPSS	last six months. Among them,
			using	62.2 (219) reported having one
			descriptive	sexual partner, 15.6% (55)
			and	reported having two sexual
			inferential	partners and 22.2% (78) reported
			statistics	having three or more partners in
				the past six months.
				Floors and basements include:
				Cause of HPV
				Where does HPV come from?
				How does one actually get it?
				How many different types of
				HPV are there? Do all types of
				HPV cause cancer? Do some
				types of HPV cause warts?
				How is HPV diagnosed in a
				sexual partner? What are HPV
				lesions like?
				Why cervical cancer is called a
				sexually transmitted disease?
				Can I get HPV infection without
				having sex?
				Can I get HPV from contact with
				strangers?
				Is HPV transmitted from person
				to person by kissing? Can it be
				transmitted to the throat?
				Who is most at risk for HPV
				infection?
				Does having more than one
				sexual partner increase a
				person's risk of contracting
				HPV?
				Does HPV increase with sexual
				activity?
				Where does the infection enter
		<u> </u>		

		the body?
		Cause of cervical cancer
		What is the main cause of
		cervical cancer?
		Is cervical cancer inherited? If
		we do not have a family history,
		what is the cause of this disease?
		Who is most at risk for cervical
		cancer?
		Does having more than one
		sexual partner increase the risk
		of cervical cancer?
		What is cervical cancer like?
		Does anyone diagnosed with
		HPV infection get cervical
		cancer?
		Signs and symptoms of HPV
		infection
		How do we know if we have
		been exposed to HPV or become infected with HPV?
		What are the signs and symptoms
		of HPV infection?
		Signs and symptoms of cervical
		cancer
		What are the symptoms of
		cervical cancer? Are there any
		early warning signs?
		Does cervical cancer cause pain?
		HPV and cervical cancer
		screening
		How easy is it for doctors to
		diagnose cervical cancer?
		Is a Pap smear really enough to
		diagnose?
		Can HPV be detected before
		cancer develops?
		In addition to annual
		examinations, what kind of
		examination should we do?

8	Veronique Verhoeven, et al.2010(32)	Belgium	Determine the specific information needs of people searching the Internet for information on human papillomavirus (HPV).	527 emails were received in which 713 separate questions were identified and	Data on all questions raised between March 2004 and July 2011 were collected and analyzed. All questions on the HPV website were in Korean and in	qualitative content analysis	HPV vaccine Is there a vaccine to prevent HPV? How effective is the vaccine? Is the vaccine safe? Why is the vaccine so expensive? Where can I go to get the vaccine in the community? Where can I get more information about the HPV vaccine? HPV and cervical cancer risk management What can be done to reduce the risk of HPV infection and cervical cancer? What can we do to prevent girls and younger women from being exposed? What can older women do to reduce their risk of HPV and cervical cancer? All questions are categorized in one of the following topics In descending order: transfer (n = 162) Vaccine (n = 98) Natural history (n = 70)
8	Verhoeven, et	Belgium	information needs of people searching the Internet for information on human	were received in which 713 separate questions were identified and	between March 2004 and July 2011 were collected and analyzed. All questions on the HPV website were in Korean and in	content	risk of HPV infection and cervical cancer? What can we do to prevent girls and younger women from being exposed? What can older women do to reduce their risk of HPV and cervical cancer? All questions are categorized in one of the following topics In descending order: transfer (n = 162) Vaccine (n = 98) Natural history (n = 70)
				answered.	free text. At the time of writing the manuscript, the text was translated from Korean to English. Website address: (www.ua.ac.be/hpvinfo)		 Defective circle (n=60) (meaning: the belief that two Partners with HPV infection continue to infect each other, so it is not possible to clear the virus. Diagnosis of HPV in men (n=52) Diagnosis of HPV in women (n=41) Treatment in women (n=40) Commune period (n=34)

		Pregnancy / fertility (n = 27)
		■ Genital warts (n = 23)
		 Loyalty and unfaithfulness (n
		=21)
		■ Treatment in men (n=19)
		• Symptoms of HPV infection
		(n=10)
		1 .When did I get the virus?
		when? Did my partner take it
		from a former relationship or
		betray me?
		2. Is the virus transmitted
		through towels, dirty toilet seats,
		pets?
		3. Can children be infected with
		HPV?
		4. What can be done to prevent
		HPV transmission?
		5. When diagnosed with HPV in
		a couple. What are the
		consequences for his wife and
		sexual partner? Do you have a
		sexual partner? Should he test?
		6. I have HPV and my husband
		will probably get it too. Won't we
		constantly infect each other?
		7. When I got HPV. What to say
		to my partner, ex-partner, and my
		lover. To?
		8 .I have HPV. Does this fact
		affect my sex life?
		9.Does HPV infection jeopardize
		my chances of getting pregnant?
		10. I just got my first HPV
		vaccine. Should I abstain from
		sex until after the third injection?
		Can I have sex while using a
		condom?
		11.I was diagnosed with HPV. Is
		vaccination good for me?

	l women	Face-to-face interview	thematic	The questions were in 6
	8-53 yesrs	The samples were entered	framework	themes:
09(29) old	ld	into the study using the	analysis	Identity (for example, what
		snowball technique		are the symptoms?)
				Does HPV have symptoms?
				Is it really common in the UK?
				Cause (for example, how do
				you get HPV?)
				How to fight HPV?
				Can HPV be transmitted in a
				way other than sexually?
				Is HPV more common in young
				people?
				Are women who are multiple
				partners and sex workers more at
				risk?
				Does drinking alcohol increase
				the risk of infection?
				Schedule (for example, how
				long does it take?)
				How long does it take for HPV
				to become cervical cancer?
				How long can the virus remain
				dormant?
				Consequences (for example,
				does it always cause the
				cervix?)
				How many cases of HPV
				become cervical cancer?
				Is HPV the only cause of
				cervical cancer?
				What other cancers are caused
				by this virus?
				Can I get pregnant with this
				virus?
				Does HPV have other effects on
				the body?
				Are Genital Warts and Cancer
				Related?
				How many women get cervical

	cancer?
	How is cervical cancer treated
	with treatment?
	Can a man get HPV?
	Does it affect men?
	Should the sexual partner be
	notified if the HPV test is
	positive?
	Treatment control (e.g. Can you prevent infection?)
	Is there a way to protect yourself against HPV?
	Does using a condom protect
	against the virus? What is the treatment for HPV?
	What does the treatment of
	cellular changes involve?
	How long does it take for this
	virus to clear the body?
	Why do some people clean it
	with their immune system, but
	others do not?
	How often should you go for a checkup?
	At what age should screening for
	the virus begin?
	Methods: HPV testing/
	vaccination
	What does the [HPV] test
	include?
	Can I get an HPV test from a
	doctor?
	How long does it take to
	determine the result of an HPV
	test?
	Is HPV included in the STI test?
	Is Pap smear different from
	HPV?
	How effective is the vaccine?

							vaccine? What are the long-term effects of the vaccine? How does the vaccine work, is it a bit of a virus? Can the virus be vaccinated? Is the vaccine for prevention or treatment? How many vaccinations should be given? How many years have they been working on the vaccine? Who has the vaccine been tested on? Who has tested the vaccine? Will the vaccine be suitable for everyone? Is the vaccination only for a certain age group? When is the vaccine available? Will the vaccination be free for everyone? Does the vaccine also protect against warts? \Should I have a Pap smear and test after the injection?
10	Allison L et al.2008(45)	African- America n	This study aims to understand the results of HPV and Pap smear test entitled: What do women with HPV need and want to know? Was performed	 A total of ten focus groups (N = 90) Age 30-45 years; 46-65 years Language English/Spanish 	 Qualitative Study Participants' knowledge about cervical cancer and HPV and the reactions of each group were recorded with more than 2 observers / notes. 	The results were analyzed using a note-based strategy with three independent judges.	According to the study, participants were aware of HPV and cervical cancer, but did not have the necessary knowledge. They were confused about the nature of the disease, the severity, the transmission, the types, the treatment, and the instructions for the HPV test, and needed to have complete information because they stated that this reduced their anxiety and caused them to follow up. Do

					the necessary. Many questions about your sexual partner and how to deal with it If you have the virus, several concepts of HPV for women are contradictory and must be carefully explained to ensure that patients understand their test results. Do not experience unnecessary
					fear, anxiety or stigma; And return for follow-up. Recommendations are provided
					to meet these patient needs.
11 Suzanne M. et al.2006(38)	Australia	Purpose, how to manage and communicate with patients about HPV?	-		The needs of the community in relation to HPV What is HPV and how do we get it? How long does the infection stay in the body and does it lead to cancer? How is HPV diagnosed? Despite a normal Pap smear, is there no cure for the virus? Who should do the HPV DNA test and when? HPV can cause cancer: Are other cancers caused by these microorganisms? Does taking OCP pills affect this disease? Should it be stopped? How long does taking the pill cause uterine cancer? I have had this partner for many years. Now that I'm positive for HPV, does that mean she've had sex with someone else?
					✓I have had the many years. positive for Fred mean she've

Г				I		I	П		a low wish and a high will
									a low-risk and a high-risk type?
									✓ Do I have to tell my partner?
									✓ Do I have to quit smoking?
									✓ If my spouse and I were in
									the beginning of a virgin
									relationship, how can I have
									an abnormal HPV or Pap smear?
F	12	KirstenMc	Australia	The aim of this study was to	19 women	Qualitative	face-to-face	Framework	1. Women's perception of HPV
	12	Caffery,	Australia	determine the information	with HPV	interview	race-to-race	Analysis	✓ Knowledge of HPV before
		etal.2005		needs of women about the role	infection were	interview		7 111d1 y 515	current diagnosis
		(35)		of human papillomavirus	selected after				✓ The patient's current
		, ,		(HPV) in cervical cancer and	routine				understanding of HPV
				how to prevent cervical	cervical				2. Required information about
				cancer.	screening.				HPV
									✓ High-risk types of HPV against genital warts
									✓ Sexual transmission: Where
									does the virus come from?
									✓ Impact on sexual partner,
									future partners and disease
									transmission between
									partners ✓ Outbreak of HPV,
									information on recurrence
									and recovery
									✓ Information on management
									and treatment options
									✓ Consequences of cancer risk
									and fertility
									✓ Lack of appropriate and available information
									✓ Key information that women
									reported as reliable
									information
									✓ High prevalence of HPV
									✓ How the disease goes away
									on its own
L									✓ Creating cancer following

						infection with the virus ✓ Absence of genital warts and symptoms of the disease 3. How to present HPV results ✓ Using the phone ✓ Using letters ✓ During the consultation 4. Physician communication method ✓ Notification style ✓ Amount of information 5. HPV information retrieval experiences
13	Rebecca Anhang, et al.2004(15)	USA	Determining the relevance of HPV: Reviewing existing research and recommendations for educating patients	-	Qualitative Study	According to this study, information and psychological needs were among the important needs of individuals, which include: I How and when did I get HPV? Does HPV affect pregnancy or the baby? Can a person get HPV orally or by hand? How can I get tested for HPV? How can HPV be prevented from being given or received? Can partners re-infect each other? Does HPV cause cervical cancer? What should I tell my partner about HPV? What are the best HPV treatment options? Looking for psychological

				 Problems such as: Anxiety Regret Angry Fear of cancer Concerns about loss of reproductive functions Fear of negative reactions from friends, family or sexual partners Concerns about partner infidelity or enmity with a partner are thought to be a partner as a thought to be a partner are thought to be a partner as a a partner a
				person are thought to be a source of infection Changes in body image Reduce intimate activities
580				

583

585

590

592 593

597

Table2:CASP (Critical Appraisal Skills Programme) qualitative checklist scores for total included studies.

Tuoiez.erioi (enticui rip	praisa	CASP criterion						rade	Total	Quality of		
	nt of aim	qualitative	research	ting strategy	in data	relationship esearcher and ts	considerations	data analysis	statement of	e of research	Score	study
	Clear statement	Suitable methodology	Appropriate design	Proper recruiting	Adequacy collection	Adequate relationship between researcher and participants	Ethical cc	Rigor in	Clear state findings	Overall value		
Sophie Mulcahy Symmons, et al.2021											19	Moderate
Laura Marlow, et al.2020											20	High
Qaderi, et al.2020											20	High
Dorothy Gold, et al.2012											14	Low
Veronique Verhoeven, et al.2010											20	High
Laura A V Marlow,2009											20	High
KirstenMcCaffery, etal.2005											20	High
Rebecca Anhang, et al.2004											17	Moderate

Table3: CASP (Critical Appraisal Skills Programme) Crossectional checklist scores for total included studies.

CASP criterion											
	Profile of	Evaluation	Study	Results	Total	Quality					
	participants	tool	design		Score	of					
						study					
Kathrin Milbury et al.2013					18	High					
SungJong Lee1, et al.2012					18	High					
Sandra Millon Underwood, et al.2010					18	High					