

1 **A comparative analysis of depression between pregnant and non-pregnant adolescents in a**
2 **southwestern town in Nigeria**

3 Authors: Julianah T. MOSANYA¹, Temilade A. ADEGBITE², Kazeem O. ADEBAYO³, Bolaji
4 E. EGBEWALE⁴, Kayode T. IJADUNOLA⁵

5 Affiliation: 1. Department of Psychiatry, Osun State University (UNIOSUN) Teaching Hospital,
6 Osogbo, Osun State, Nigeria

7 2. Department of Psychiatry, Lagos University Teaching Hospital, Lagos, Lagos
8 State, Nigeria

9 3. Department of Psychiatry, Osun State University, Osogbo, Osun State, Nigeria

10 4. Department of Community Medicine, Ladoke Akintola University of Technology,
11 Ogbomoso, Oyo State, Nigeria

12 5. Department of Community Health, Obafemi Awolowo University, Ile-Ife, Osun
13 State, Nigeria

14 Corresponding author: J.T. Mosanya, temdad2@yahoo.com

15 **Abstract**

16 **Background:** Adolescence constitutes a risk factor for mental health problems, and this may be
17 further complicated by pregnancy. The rate of adolescent pregnancy is still extremely high in the
18 sub-Saharan Africa including Nigeria. Pregnancy and mental health problems during
19 adolescence constitute double vulnerability for negative outcomes for the adolescents and their
20 offspring.

21 **Methodology:** The study was cross-sectional in design and it compared prevalence of depression
22 and associated factors among pregnant and non-pregnant adolescents. It was conducted in
23 Osogbo metropolis, Osun State, Southwest, Nigeria. The study population comprised pregnant
24 adolescents (aged 15-19 years) attending antenatal care (ANC) in selected formal and informal
25 health facilities. Non-pregnant adolescents who were equally attending services at the facilities
26 were recruited as the control group. Information was obtained from the adolescents with the use
27 of a structured questionnaire and data was analysed with IBM-SPSS version 21 software.

28 **Results:** Three hundred and thirty-four respondents (167 per group) were involved in the study;
29 the pregnant adolescents had a mean age (\pm SD) of 17.92 (\pm 1.13) years while the non-pregnant
30 adolescents had a mean age of 17.70 (\pm 1.23) years. The prevalence of depression among the
31 pregnant adolescents was 8.4% while that of the non-pregnant adolescents was 3.0%. The result
32 showed a statistically significant association between pregnancy status and depression among the
33 adolescents ($p= 0.033$). Living arrangement was the only socio-demographic variable that had
34 significant relationship with depression among the pregnant adolescents while living arrangement
35 and employment status had significant relationships with depression among the non-pregnant
36 adolescents. History of mental illness, childhood sexual abuse and anxiety symptoms showed
37 significant relationship with depression among pregnant adolescents, however, only anxiety
38 symptoms showed significant relationship with depression among non-pregnant adolescents.

39 **Conclusion:** The study concluded that the prevalence of depression is significantly higher
40 among pregnant adolescents with similarities and differences in the factors associated with
41 depression in the two groups.

42

43 **Keywords: Pregnant Adolescents, Depression, psychosocial, childhood physical abuse,**
44 **childhood sexual abuse**

45 **Introduction**

46 The challenges of the developmental phase of adolescence are enormous and constitute risk for
47 various mental health problems including depression. Depression is a common mental health
48 disorder among adolescents (1), with equal incidence in males and females before puberty.
49 However, there is a steady increase in the incidence in females after puberty till adulthood, when
50 the risk doubles in them. (2). The vulnerability of females to depression beginning in adolescence
51 (3) and the impact of an unidentified and untreated illness remain a challenge globally (4).

52 According to the World Health Organization, about 16 million girls aged 15-19 years and 1 million
53 girls under 15 years give birth annually and most live in the low/middle-income countries (5).The
54 burden of pregnancy or parenting during adolescence constitutes a risk for mental health problems
55 in addition to the developmental risks of adolescence (6,7). Pregnancy and mental health problems
56 in adolescents therefore constitute “double vulnerability” for negative outcomes in them and the
57 future generation.

58 The poor mental health of the mothers has a potential to exert enormous impact on them, the
59 unborn child and their infants after delivery. A pregnant woman with poor mental health is more
60 likely to attend antenatal clinics poorly, comply poorly with treatment and subject herself to
61 practices that may be detrimental to her and the foetus (7–9), consequently, impairing the success
62 of public health interventions targeted at reduction of both maternal and neonatal morbidity and
63 mortality. The two most important stages in human growth and development- the foetal period
64 and adolescence are at stake in the context of adolescent pregnancy, especially when complicated

65 by mental disorders. Poor attention to this vulnerable group of adolescents has a potential to set
66 up a vicious cycle of poverty and mental health problems in them and the next generation (10).
67 Depression has been recognized as one of the most common perinatal mental health disorders
68 among women including adolescent mothers. Despite its high prevalence, significance and
69 availability of effective treatment options, it is usually undetected and untreated (6,11). Many
70 studies on depression among adolescents have considered the school population extensively in
71 both developed and developing countries (3,12–16). There are however fewer studies on
72 depression among the pregnant adolescents and most of the available studies were conducted in
73 the developed nations (17,18) . The prevalence reported in some of these studies varied widely
74 with rates as low as 1.6% (19) and as high as 68% (20), depending on various factors including
75 the population, diagnostic criteria used, and age range of the respondents.

76 Commonly reported prevalence of current episode of depression among adolescents ranges
77 between 2-10% (21–24). Although some studies in high-income countries showed that
78 adolescent mothers are at a higher risk of depressive disorders compared with adult mothers
79 (25,26) and non-pregnant adolescent girls (27), others refuted the evidence (28,29). However,
80 there is a dearth of literature on both antepartum and postpartum mental disorders among
81 adolescent girls in the developing countries where the burden of adolescent pregnancy is highest.
82 This study aims to determine and compare the prevalence of depression and its relationship with
83 sociodemographic and psychosocial variables between pregnant and non-pregnant adolescents in
84 a south-western town in Nigeria.

85 **Materials And Methods**

86 **Study setting and Participants**

87 We employed a cross-sectional analytical design where comparison was made between the
88 prevalence and correlates of depression among pregnant and non-pregnant adolescents aged 15 to
89 19 years. The participants were recruited from formal and informal health care facilities in Osogbo,
90 the capital town of Osun State in south-west, Nigeria.

91 The pregnant adolescents were recruited from the public primary and secondary health care
92 facilities as well as informal health facilities where they were attending antenatal care (ANC).
93 Non-pregnant adolescents who attended the same facilities for one health reason or the other at the
94 time of the study were recruited as the control group. Adolescents who were acutely ill or had
95 chronic medical conditions were excluded from the study.

96 Simple random sampling method (balloting) was used to select two Primary Health Care Centres
97 (PHC) from each of Osogbo and Olorunda Local Government areas in Osogbo town. The only
98 secondary health facility in the town, the State Specialist Hospital, was also included (making a
99 total of five formal health facilities). The secondary facility was included because health care
100 services are free as obtainable in the PHCs. Two traditional / faith homes offering services that
101 included antenatal care were also selected by snowballing method from each of the local
102 governments as there was no sampling frame for this group. In total, nine facilities were included
103 in the study.

104 Consecutive pregnant adolescents were recruited serially and the controls were also recruited from
105 the same facilities; for formal health services, they were recruited from the general outpatient
106 section while for the informal services, they were recruited among clients that came to receive care
107 for other reasons besides pregnancy in the same facility. They were also recruited serially and
108 matched for age of the pregnant adolescents, using frequency matching techniques.

109

110 **Sample size determination**

111 Minimum sample size for each group required for this study was calculated using the formula for
112 comparing two independent proportions (30).

113

$$114 \quad n = \frac{(Z_{\alpha/2} + Z_{\beta})^2 * \{P_1(1 - P_1) + P_2(1 - P_2)\}}{(P_1 - P_2)^2}$$

115

116 Where:

117 n = sample size in each group

118 $Z_{\alpha/2}$ = Standard normal deviate corresponding to confidence level; at confidence level of 95%, Z_{α}
119 = 1.96 for a two tailed test

120 Z_{β} = Standard normal deviate corresponding to (1-Power); at power of 80%, $Z_{\beta} = 0.84$

121 P_1 is the estimate of prevalence of depression in the pregnant adolescents

122 P_2 is the estimate of prevalence of depression in the non-pregnant adolescents

123 P_1 and P_2 were derived from an earlier study in Brazil where 26.3% of pregnant adolescents had
124 depression compared with 13.6% of the non-pregnant adolescents (27).

125 $P_1 = 0.263$

126 $P_2 = 0.136$

127
$$n = \frac{(1.96 + 0.84)^2 * \{0.263(0.737) + 0.136(0.864)\}}{(0.263 - 0.136)^2}$$

128
$$n = \frac{(7.84) * (0.19381 + 0.117504)}{0.016129}$$

129
$$= 151.33$$

130
$$= 152$$

131 Ten percent was added to the sample size to make up for non-response $(152 + 15) = 167$

132 One hundred and sixty-seven respondents were recruited for each group in this study.

133

134 **Data Collection**

135 Data was obtained from the respondents by means of a structured questionnaire. The questionnaire
136 was administered to them after explaining the purpose of the research to them and obtaining
137 informed consent/ assent. This was done after they had received care at the facility for the day.
138 An identification sticker was placed on the hospital card of the patients that have been interviewed
139 to avoid repeat sampling. The researcher ensured that there was privacy for the respondents while
140 completing the questionnaire. The first section had questions on socio-demographic data, sexual
141 history, whether pregnancy was planned and disposition to it. The other sections contained
142 measures for depression, anxiety, social support, childhood sexual and physical abuse and recent
143 abuse.

144

145 **Measures**

146 **Depression:** The Beck Depression Inventory-II (BDI-II), which was used to assess depression in
147 this study, is a popular screening instrument which has been validated and used previously among
148 adolescents. It was adapted to the diagnostic criteria of major depressive episode of the Diagnostic
149 and Statistical Manual of Mental Disorders-fourth edition (DSM-IV). It measures depressive
150 symptomatology as well as assess the severity of depression in people who are thirteen years and
151 above. It has 21 items, each item is scored 0 to 3 with possible total score of 0-63. A score of 0-16
152 is considered normal while 17-20, 21-30, and 31 and above are considered borderline, moderate
153 and severe depression respectively Those who scored 21 and above were regarded as depressed
154 because this cut-off has been reported previously to correlate significantly with clinical depression
155 (31).

156 **Anxiety:** Patient Health Questionnaire for Depression and Anxiety-4 (PHQ-4): PHQ-4 combined
157 both the two item Patient Health Questionnaire (PHQ-2) and the Generalized Anxiety Disorder
158 (GAD-2). The scales were derived from the original eight and seven items PHQ-8 and GAD-7
159 respectively. PHQ-8 is used for screening for depression while GAD-7 is used for screening for
160 anxiety. The first two of four items on PHQ-4 screened for anxiety and the last two screen for
161 depression. Each item is rated on a scale of 0-3. Scores for each of the subscale is derived by
162 adding scores for items on each subscales. Score ≥ 3 on each of the scale suggests that disorder.
163 The scale for anxiety is used in this study as a measure of anxiety

164 **Social Support:** OSLO Social Support Questionnaire) has three items for assessing perceived
165 social support, Minimum and maximum score obtainable on the scale are 3 and 14 respectively.

166 Higher score denotes higher level of perceived social support. Scores of 3-8 is considered “poor
167 social support”, 9-11 “moderate support and 12-14 “strong support”.

168 **Childhood Physical and Sexual Abuse:** Sexual and Physical Abuse Questionnaire (SPAQ) used
169 in this study is a nine-item questionnaire that was designed as a screening tool for sexual and
170 physical abuse in childhood. For this questionnaire, the operational definition of sexual and
171 physical abuse involved having actual contact for sexual abuse, while presence of physical harm
172 like bruises is required for physical abuse (32).

173 **Abuse Assessment Screen (AAS):** Abuse assessment screen is a five-item screening tool for
174 detection of domestic violence, an individual screens positive for abuse once any of the items is
175 answered in the affirmative. The instrument was used as a measure of current/recent abuse in this
176 study.

177 All instruments were translated to Yoruba and back translated to English to ensure content validity.

178 **Statistical Analysis**

179 Data were analyzed with the Statistical Package for Social Sciences (SPSS) version 21.
180 Appropriate descriptive statistics such as percentages, means and standard deviation were used to
181 summarize the data depending on the type of variable. Chi-square test was used to determine the
182 relationship between the depression and the independent variables and significance level was set
183 at $p < 0.05$.

184 **Ethical Considerations**

185 Ethical approval was obtained from the Institute of Public Health, Obafemi Awolowo University,
186 Ile-Ife, Osun State. Permission was sought from the head of the selected health facilities and the

187 doctors whose patients were involved in the research. Informed consent was obtained from the
188 pregnant adolescents or parents/guardians. All pregnant adolescents gave informed consent
189 irrespective of their age because they were considered “matured minors” (33). The
190 parents/guardians of those that were less than 18 years old were required to give consent while the
191 adolescents gave assent after which they were enrolled for participation. The questionnaire was
192 administered to the adolescents privately. Privacy and confidentiality of all participants were
193 ensured during and after data collection.

194

195 **Results**

196 **Socio-demographic characteristics of pregnant and non-pregnant adolescents**

197 Three hundred and thirty-four respondents (167 per group) were involved in the study. The mean
198 (\pm SD) age of all the respondents was 17.81 (\pm 1.18) years. There was no significant difference
199 between the mean age of both groups of respondents ($t = 1.670$, $p = 0.096$). The pregnant
200 adolescents had a mean age (\pm SD) of 17.92 (\pm 1.13) years while the non-pregnant adolescents had
201 a mean age of 17.70 (\pm 1.23) years.

202 Thirty-nine percent of the pregnant adolescents compared with 84% of the non-pregnant
203 adolescents were students and the difference was statistically significant ($\chi^2 = 77.24$, $p = 0.001$).

204 Thirty seven percent of the non-pregnant adolescents compared with 11% of the pregnant ones
205 had attained tertiary education and the difference was statistically significant ($\chi^2 = 29.525$,
206 $p = 0.001$). Concerning employment status, 25% of the pregnant adolescents compared with 5% of
207 the non-pregnant adolescents were employed. Although a higher proportion of pregnant
208 adolescents were married (25%) compared with the non-pregnant ones (13%), the proportions of

209 both groups that were never married were high at 72% and 84% respectively. Concerning living
210 arrangements of the respondents, 77 (46%) pregnant adolescents compared with 127 (76%) non-
211 pregnant ones lived with their parents. However, significantly higher proportions of the pregnant
212 ones (29%) lived with a partner (husband or boyfriend) compared with the non-pregnant ones
213 (6%), $p < 0.001$. About 65% of the parents of the pregnant adolescents compared with 90.4% of
214 the parents of the non-pregnant adolescents were married and the difference was statistically
215 significant ($\chi^2 = 31.839$, $p = 0.001$). Likewise, 18% and 4.8% of mothers of the pregnant
216 adolescents compared with 43.7% and 18.0% of the mothers of the non-pregnant ones had tertiary
217 and postgraduate education respectively ($\chi^2 = 58.445$, $p = 0.001$). The pattern was also similar for
218 the educational attainment of fathers of the respondents. (Table 1).

219 **Table 1: Sociodemographic Characteristics of Pregnant and Non-pregnant Adolescents**

220

221 **Psychosocial characteristics of pregnant and non-pregnant adolescents**

222 The proportion of those that signified that they have had history of mental illness was 5.4% among
223 the pregnant adolescents compared with 1.8% of the non-pregnant peers, the difference was not
224 statistically significant ($p = 0.139$).

225 Twenty-three (13.8%) of the pregnant adolescents compared with 30 (18.0%) of the non-pregnant
226 adolescents had poor social support ($p = 0.084$). Twenty-five (15%) and 33 (19.8%) of the pregnant
227 adolescents compared with 16 (9.6%) and 30 (18.0%) of non-pregnant adolescents had history of
228 childhood physical and sexual abuse respectively, but the difference in proportions were not
229 statistically significant ($p = 0.133$ and 0.675). Similarly, the proportions of those exposed to recent
230 abuse was comparable between the two groups ($p = 0.082$). Fewer, 13 (7.8%) of the pregnant

231 adolescents compared with 24 (14.4%) of the non-pregnant adolescents screened positive for
232 anxiety, but the difference did not reach statistical significance ($p= 0.055$) (Table 2).

233 **Table 2: Psychosocial Variables of Pregnant and Non-pregnant Adolescents**

234 **Planning status of pregnancy and its disposition among pregnant adolescents**

235 Figure 1 and 2 shows the planning status of the pregnancy and their disposition to it among
236 pregnant adolescents.

237 **Figure 1: Planning Status of the Pregnancies of the Pregnant Adolescents**

238 **Figure 2: Disposition of Pregnant Adolescents in Osogbo towards their** 239 **Pregnancies**

240 **Depression**

241 The mean BDI score (which assessed depression among the respondents) of the pregnant
242 adolescents was significantly higher, 8.32 (± 8.17) than that of the non-pregnant adolescents (6.71
243 (± 7.85) ($p < 0.001$). Overall, 19 (5.7%) of the respondents were categorized as being depressed, 14
244 (8.4%) among the pregnant adolescents and 5 (3.0%) among the non-pregnant adolescents. The
245 proportion of pregnant adolescents categorized as depressed was significantly higher than that of
246 the non-pregnant adolescents ($p = 0.033$).

247

248 **Relationship between depression and sociodemographic and Psychosocial variables among** 249 **the respondents**

250 Table 3 shows the relationship between depression and selected sociodemographic variables
251 among pregnant and non-pregnant adolescents. Living arrangement was the only variable that had
252 significant relationship with depression among the pregnant adolescents while living arrangement
253 and employment status had significant relationships with depression among the non-pregnant
254 adolescents.

255 **Table 3: Relationship between depression and socio-demographic variables in pregnant**
256 **and non-pregnant adolescents**

257
258 Table 4 shows the relationship between depression and psychosocial variables among pregnant
259 and non-pregnant adolescents. Among pregnant adolescents, 33% of those with positive history of
260 mental illness compared with only 7% of those with negative history of mental illness had
261 depression and the difference was statistically significant (Fisher's Exact Test, $p=0.029$).
262 Similarly, a higher proportion of those who were exposed to childhood sexual abuse (18%)
263 compared to those who were not exposed (6%) had depression ($p=0.023$). Conversely, exposure
264 to childhood physical abuse had no significant relationship with depression among the pregnant
265 adolescent. Also, social support and current abuse had no relationship with depression status.
266 Similarly, whether pregnancy was planned or not as well as disposition to the pregnancy showed
267 no relationship with depression among them. Anxiety however showed significant relationship
268 with depression as significantly higher proportions of those that screened positive for anxiety
269 (38.5%) compared with those that were negative (5.8%) had depression ($p=0.001$).

270 **Table 4: Relationship between depression and psychosocial variables in pregnant and non-**
271 **pregnant adolescents**

272 Among non-pregnant adolescents, the history of mental illness, social support, childhood physical
273 as well as sexual abuse had no relationship with depression. Current experience of abuse also had
274 no relationship with depression. However, screening positive for anxiety showed significant
275 relationship with depression with 12.5% of those with anxiety compared with 1.4% of those
276 without anxiety having depression, $p=0.022$.

277 **DISCUSSION**

278 The prevalence of depression among the pregnant adolescents (8.4%) was significantly higher than
279 that of the non-pregnant adolescents (3.0%), $p=0.033$, this pattern followed the previous findings
280 of pregnant adolescents having a higher risk of depression than their non-pregnant counterparts
281 (27). The higher prevalence in the pregnant adolescents as found in this study may be due to the
282 stress and/or physiological changes of pregnancy and social discrimination associated with teenage
283 pregnancy (34). It may also be that the presence of depression actually predisposed to pregnancy
284 in the adolescents as earlier studies suggested, that depression predisposes the adolescents to high-
285 risk sexual behaviour which may increase the incidence of pregnancy in them and may account
286 for the increased prevalence of depression among the pregnant adolescents (35,36). This finding
287 highlights the importance of paying close attention to the mental health of adolescents in general
288 and the pregnant ones in particular.

289 This observed relationship between studentship status and depression could make one to consider
290 the possibility that being a student is protective against depression or that those who were not
291 students had other risk factors like poverty that increased the risk of depression in them. On the
292 one hand, it is possible that the presence of depressive disorder in these other groups predisposed
293 them to dropping out of school, thereby looking for other things to do beyond schooling. On the

294 other hand, depression in them could be a psychological reaction to dropping out of school which
295 was informed by reasons out of their control. There may be need to watch out for depression in
296 adolescents who are out of school. This finding is in agreement with a study in India that found a
297 significantly higher prevalence of depression among female adolescents that dropped out of school
298 (19).

299 Concerning the relationship of depression with living arrangements in this study, the proportion
300 of respondents who stayed alone or with partners (boyfriend or husbands) who had depression was
301 significantly higher than those who stayed with their parents or grandparents ($p < 0.001$). This
302 finding suggests that living with parents or grandparents may be a protective factor against
303 depression in this age group rather than living alone or living with a partner. Furthermore, living
304 with partner, especially when unmarried may put significant emotional strain on those who were
305 practicing it, therefore increasing the risk of depression in them since the practice is not culturally
306 acceptable in the environment where the study was carried out. Beyond this, the social support
307 available for those living alone or with partners may not be as good as those living with their
308 parents or grandparents among this population, thereby putting significant strain on their emotional
309 health. The finding of an association of depression with few socio-demographic factors in this
310 study is contrary to the finding of an earlier study that found no such association in pregnant
311 adolescents (37).

312 Among the psychosocial factors examined, three factors; the past history of mental illness, anxiety
313 and childhood sexual abuse had a significant relationship with depression. The past history of
314 mental illness having a significant relationship with depression among the pregnant adolescents is
315 not surprising because depression and different other mental illnesses are chronic illnesses that are
316 prone to relapse, so anyone who has had an episode could have further episodes (38). The fact that

317 there is no significant relationship between past history of mental illness and depression among
318 the non-pregnant adolescents could then suggest that there is a relationship between mental illness
319 and poor reproductive behaviour as suggested in previous studies (39), which could have
320 predisposed the pregnant adolescents to the pregnancy in the first place. The type of mental illness
321 that those with history of mental illness had previously was however not documented in this study,
322 so it is not clear whether the past mental illness they had was depression or not.

323 The finding of a relationship between depression and childhood sexual abuse but not with
324 childhood physical abuse among the pregnant adolescents in this study is surprising, since they are
325 both forms of violence. It therefore raises the question as to whether sexual abuse has more
326 emotional impact on the victims than physical abuse. In addition, one may want to know the
327 contribution of sexual abuse to the occurrence of pregnancy as previous researches suggested (40).
328 The finding of an association between depression and childhood sexual abuse in this study is in
329 agreement with extant literature (41,42). Presence of anxiety symptoms had significant
330 relationship with depression, this further confirms the strong possibility of depression co-existing
331 with other psychological illnesses, especially anxiety symptoms/disorders as previous studies have
332 documented. It is interesting to note that neither planned status of pregnancy nor disposition to the
333 pregnancy had significant relationship with depression among the pregnant adolescents. This is at
334 variance with findings among adults where unplanned pregnancy showed relationship with
335 depression (43,44). This suggests that pregnancy in adolescence may be associated with emotional
336 disturbance irrespective of the planned status or disposition of the adolescent to the pregnancy.

337 **Study strengths and Limitations**

338 The strength of the study includes the focus on antenatal depression which has been less studied
339 compared with postnatal depression. Furthermore, the inclusion of a comparison group and the use

340 of structured instruments to assess the dependent and independent variables. However, there are
341 few limitations, which include our inability to select the informal health facilities involved in this
342 study randomly because effort made to get the sampling frame for the informal services was not
343 successful. Also, screening instrument was used to assess depression as against structured
344 diagnostic interview.

345 **Conclusion**

346 This study concludes that the prevalence of depression is significantly higher among the pregnant
347 adolescents compared with the non-pregnant adolescents. Living arrangement is the only socio-
348 demographic factor that has a significant relationship with depression among pregnant adolescents
349 while employment status in addition to living arrangement showed significant relationship among
350 non-pregnant adolescents. On the other hand, history of mental illness, childhood sexual abuse and
351 anxiety were the psychosocial factors with significant relationship with depression among
352 pregnant adolescents and only anxiety among non-pregnant adolescents.

353 **References**

- 354 1. Thapar A, Collishaw S, Pine DS, Thapar AK. Depression in adolescence. *The Lancet*.
355 2012;379(9820):1056–67.
- 356 2. Parker G, Brotchie H. Gender differences in depression. *International Review of*
357 *Psychiatry*. 2010 Oct 1;22(5):429–36.
- 358 3. Sajjadi H, Kamal SHM, Rafiey H, Vameghi M, Forouzan AS, Rezaei M. A Systematic
359 Review of the Prevalence and Risk Factors of Depression among Iranian Adolescents. *Glob J*
360 *Health Sci*. 2013 May;5(3):16–27.

- 361 4. Field T, Diego M, Sanders C. Adolescent depression and risk factors. *Adolescence*.
362 2001;36(143):491–8.
- 363 5. WHO. Adolescent Pregnancy: Fact sheet. 2014; Available from:
364 <http://www.who.int/mediacentre/factsheets/fs364/en/>
- 365 6. Mitsuhiro SS, Chalem E, Barros MM, Guinsburg R, Laranjeira R. Teenage pregnancy:
366 use of drugs in the third trimester and prevalence of psychiatric disorders. *Revista Brasileira de*
367 *Psiquiatria*. 2006;28:122–5.
- 368 7. Mitsuhiro SS, Chalem E, Moraes Barros MC, Guinsburg R, Laranjeira R. Brief report:
369 Prevalence of psychiatric disorders in pregnant teenagers. *Journal of Adolescence*.
370 2009;32(3):747–52.
- 371 8. Marcus SM. Depression during pregnancy: rates, risks and consequences. *Can J Clin*
372 *Pharmacol*. 2009;16(1):15–22.
- 373 9. Patel V, Rahman A, Jacob KS, Hughes M. Effect of maternal mental health on infant
374 growth in low income countries: new evidence from South Asia. *BMJ*. 2004;328(7443):820–3.
- 375 10. Plant DT, Barker ED, Waters CS, Pawlby S, Pariante CM. Intergenerational transmission
376 of maltreatment and psychopathology: the role of antenatal depression. *Psychological Medicine*.
377 2013;43(3):519–28.
- 378 11. Chalem E, Mitsuhiro SS, Manzolli P, Barros MCM, Guinsburg R, Sass N, et al.
379 Underdetection of Psychiatric Disorders During Prenatal Care: A Survey of Adolescents in Sao
380 Paulo, Brazil. *Journal of Adolescent Health*. 2012;50(1):93–6.

- 381 12. Adewuya AO, Ola BA, Aloba OO. Prevalence of major depressive disorders and a
382 validation of the beck depression inventory among Nigerian adolescents. *European Child and*
383 *Adolescent Psychiatry*. 2007;16(5):287–92.
- 384 13. Eisenberg D, Gollust SE, Golberstein E, Hefner JL. Prevalence and Correlates of
385 Depression, Anxiety, and Suicidality Among University Students. *American Journal of*
386 *Orthopsychiatry*. 2007 Oct 1;77(4):534–42.
- 387 14. Eskin M, Ertekin K, Harlak H, Dereboy Ç. Prevalence of and Factors Related to
388 Depression in High School Students. *Turkish journal of psychiatry*. 2008;19(4).
- 389 15. Sooky Z, Sharifi KH, Tagharrobi Z. The depression prevalence and its related factors in
390 teenagers in Kashan-Iran 2006. *European Psychiatry*. 2007 Mar 1;22:S246.
- 391 16. Unsal A, Ayranci U. Prevalence of Students With Symptoms of Depression Among High
392 School Students in a District of Western Turkey: An Epidemiological Study. *J School Health*.
393 2008 May;78(5):287–93.
- 394 17. Brown JD, Harris SK, Woods ER, Buman MP, Cox JE. Longitudinal study of depressive
395 symptoms and social support in adolescent mothers. *Maternal and Child Health Journal*.
396 2012;16(4):894–901.
- 397 18. Coelho FM, Pinheiro RT, Silva RA, Quevedo Lde A, Souza LD, Castelli RD. Major
398 depressive disorder during teenage pregnancy: socio-demographic, obstetric and psychosocial
399 correlates. *Revista brasileira de psiquiatria (Sao Paulo, Brazil: 1999) [Internet]*. 2013;35.
400 Available from: <http://dx.doi.org/10.1016/j.rbp.2012.03.006>

- 401 19. Nair M, Paul MK, John R. Prevalence of depression among adolescents. *The Indian*
402 *Journal of Pediatrics*. 2004;71(6):523–4.
- 403 20. Logsdon MC, Cross R, Williams B, Simpson T. Prediction of Postpartum Social Support
404 and Symptoms of Depression in Pregnant Adolescents: A Pilot Study. *The Journal of School*
405 *Nursing*. 2004 Feb 1;20(1):36–42.
- 406 21. Hoffmann F, Petermann F, Glaeske G, Bachmann CJ. Prevalence and Comorbidities of
407 Adolescent Depression in Germany. *Zeitschrift für Kinder- und Jugendpsychiatrie und*
408 *Psychotherapie*. 2012 Oct 29;40(6):399–404.
- 409 22. Mojtabai R, Olfson M, Han B. National Trends in the Prevalence and Treatment of
410 Depression in Adolescents and Young Adults. *Pediatrics* [Internet]. 2016 Dec 1 [cited 2019 Nov
411 15];138(6). Available from: <https://pediatrics.aappublications.org/content/138/6/e20161878>
- 412 23. Munhoz TN, Santos IS, Matijasevich A. Depression among Brazilian adolescents: A
413 cross-sectional population-based study. *Journal of Affective Disorders*. 2015;175:281–6.
- 414 24. Rey JM, Sawyer MG, Clark JJ, Baghurst PA. Depression among Australian adolescents.
415 *Medical Journal of Australia*. 2001;175(1):19–23.
- 416 25. Kim THM, Connolly JA, Tamim H. The effect of social support around pregnancy on
417 postpartum depression among Canadian teen mothers and adult mothers in the maternity
418 experiences survey. *BMC Pregnancy and Childbirth*. 2014;14(1):162–162.
- 419 26. Siegel RS, Brandon AR. Adolescents, Pregnancy, and Mental Health. *Journal of Pediatric*
420 *and Adolescent Gynecology*. 2014;27(3):138–50.

- 421 27. Freitas GVS, Cais CFS, Stefanello S, Botega NJ. Psychosocial conditions and suicidal
422 behavior in pregnant teenagers. *European Child & Adolescent Psychiatry*. 2008;17(6):336–336.
- 423 28. Biello KB, Sipsma HL, Kershaw T. Effect of Teenage Parenthood on Mental Health
424 Trajectories: Does Sex Matter? *American Journal of Epidemiology*. 2010;172(3):279–87.
- 425 29. Troutman BR, Cutron CE. Nonpsychotic postpartum depression among adolescent
426 mothers. *J Abnorml Psychol [Internet]*. 1990;99. Available from: [https://doi.org/10.1037/0021-](https://doi.org/10.1037/0021-843X.99.1.69)
427 843X.99.1.69
- 428 30. Charan J, Biswas T. How to calculate sample size for different study designs in medical
429 research? *Indian journal of psychological medicine*. 2013;35(2):121.
- 430 31. Steer RA, Scholl TO, Beck AT. Revised Beck Depression Inventory Scores of Inner-City
431 Adolescents: Pre- and Postpartum. *Psychol Rep*. 1990 Feb;66(1):315–20.
- 432 32. Kooiman CG, Ouwehand AW, Kuile MM. The Sexual and Physical Abuse Questionnaire
433 (SPAQ) A screening instrument for adults to assess past and current experiences of abuse. *Child*
434 *abuse & neglect*. 2002;26:939–53.
- 435 33. Folayan MO, Haire B, Harrison A, Odetoyingbo M, Fatusi O, Brown B. Ethical issues in
436 adolescents' sexual and reproductive health research in Nigeria. *Developing world bioethics*.
437 2015;15(3):191–8.
- 438 34. Wiemann CM, Rickert VI, Berenson AB, Volk RJ. Are pregnant adolescents stigmatized
439 by pregnancy? *J Adolesc Health*. 2005 Apr;36(4):352.e1-8.

- 440 35. Barnet B, Liu J, DeVoe M. Double jeopardy: Depressive symptoms and rapid subsequent
441 pregnancy in adolescent mothers. *Archives of Pediatrics & Adolescent Medicine*.
442 2008;162(3):246–52.
- 443 36. Hallfors DD, Waller MW, Ford CA, Halpern CT, Brodish PH, Iritani B. Adolescent
444 Depression and Suicide Risk Association with Sex and Drug Behavior. *American Journal of*
445 *Preventive Medicine* 2004. 2004;27(3).
- 446 37. Villanueva LA, Pérez-Fajardo MM, Fernando LI. [Sociodemographic factors associated
447 with depression in pregnant adolescents]. *Ginecol Obstet Mex*. 2000 Apr;68:143–8.
- 448 38. Monroe SM, Harkness KL. Is depression a chronic mental illness? *Psychological*
449 *Medicine*. 2012 May;42(5):899–902.
- 450 39. Lundberg P, Rukundo G, Ashaba S, Thorson A, Allebeck P, Östergren P-O, et al. Poor
451 mental health and sexual risk behaviours in Uganda: A cross-sectional population-based study.
452 *BMC Public Health*. 2011;11(1):125–125.
- 453 40. Noll JG, Shenk CE, Putnam KT. Childhood Sexual Abuse and Adolescent Pregnancy: A
454 Meta-analytic Update. *Journal of Pediatric Psychology*. 2009;34(4):366–78.
- 455 41. Fergusson DM, Horwood LJ, Lynskey MT. Childhood Sexual Abuse and Psychiatric
456 Disorder in Young Adulthood: II. Psychiatric Outcomes of Childhood Sexual Abuse. *Journal of*
457 *the American Academy of Child & Adolescent Psychiatry*. 1996 Oct 1;35(10):1365–74.
- 458 42. Weiss EL, Longhurst JG, Mazure CM. Childhood Sexual Abuse as a Risk Factor for
459 Depression in Women: Psychosocial and Neurobiological Correlates. *American Journal of*
460 *Psychiatry*. 1999 Jun;156(6):816–28.

461 43. Faisal-Cury A, Menezes PR, Quayle J, Matijasevich A. Unplanned pregnancy and risk of
462 maternal depression: secondary data analysis from a prospective pregnancy cohort. *Psychology,*
463 *Health & Medicine.* 2017 Jan 2;22(1):65–74.

464 44. Ege E, Timur S, Zincir H, Geçkil E, Sunar-Reeder B. Social support and symptoms of
465 postpartum depression among new mothers in Eastern Turkey. *Journal of Obstetrics and*
466 *Gynaecology Research.* 2008;34(4):585–93.

467

468

1 **A comparative analysis of depression between pregnant and non-pregnant adolescents in a**
 2 **southwestern town in Nigeria**

3 **TABLES AND FIGURES**

4 **Table 1: Sociodemographic Characteristics of Pregnant and Non-pregnant Adolescents**

Variable	Pregnant (n=167) Frequency (%)	Non-pregnant (n=167) Frequency (%)	Total (n=334) Frequency(%)	Statistics
Age (years)				
15	7 (4.2)	10 (6.0)	17 (5.1)	$\chi^2 = 3.553$ df = 4 p = 0.470
16	17 (10.2)	23 (13.8)	40 (12.0)	
17	21 (12.6)	28 (16.8)	49 (14.7)	
18	60 (35.9)	52 (31.1)	112 (33.5)	
19	62 (37.1)	54 (32.3)	116 (34.7)	
Employment status				
Employed	41(24.6)	8 (4.8)	49 (14.7)	$\chi^2 = 77.24$ df = 3 p = 0.001
Unemployed	29 (17.4)	14 (8.4)	43 (12.9)	
Student	65 (38.8)	141 (84.4)	206 (61.7)	
Apprentice	32 (19.2)	4 (2.4)	36 (10.7)	
Religion				
Christianity	57 (34.1)	80 (47.9)	137 (41.0)	$\chi^2 = 7.061$

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Islam	108 (64.7)	84 (50.3)	192 (57.5)	df =2
Traditional	2 (1.2)	3 (1.8)	5 (1.5)	p = 0.029
Educational attainment				
Primary	36 (21.6)	22 (13.2)	58 (17.4)	$\chi^2 = 29.525$
Secondary	107 (64.1)	81 (48.5)	188 (56.3)	df =3
Tertiary	70 (41.4)	51 (36.8)	80 (24.0)	p = 0.001
No formal	5 (3.0)	3 (1.8)	8 (2.4)	
Marital status				
Never married	120 (71.9)	141 (84.4)	261 (78.1)	$\chi^2 = 9.420$
Married	41 (24.6)	22 (13.2)	63 (18.9)	df =3
Separated/divorced	2 (1.2)	0 (0)	2 (0.6)	p = 0.024
Cohabiting	4 (2.4)	4 (2.4)	8 (2.4)	
Living arrangement				
Partner	49 (29.3)	10 (6.0)	59 (17.7)	$\chi^2 = 55.329$
Parents	77 (46.1)	127 (76.0)	204 (61.1)	df =3
Grandparents	34 (20.4)	11 (6.6)	45 (13.5)	p = 0.001
Alone	7 (4.2)	19 (11.4)	26 (7.8)	

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

5

6

7

8

9

10 **Table 2: Psychosocial Variables of Pregnant and Non-pregnant Adolescents**

Variable	Pregnant (n=167) Freq (%)	Non-pregnant (n=167) Freq (%)	Total (n=334) Freq (%)	Statistics
History of mental illness				Fisher's Exact P=0.139
Yes	9 (5.4)	3 (1.8)	12 (3.6)	
No	158 (94.6)	164 (98.2)	322 (96.4)	
Social support category				$\chi^2 = 4.954$ df = 2 p = 0.084
Poor	23 (13.8)	30 (18.0)	53 (15.9)	
Intermediate	108 (64.7)	88 (52.7)	196 (58.7)	
Strong	36 (21.6)	49 (29.3)	85 (25.4)	
History of CPA*				$\chi^2 = 2.252$ df = 1 p = 0.133
Present	25 (15)	16 (9.6)	41 (12.3)	
Absent	142 (85.0)	151 (90.4)	293 (87.7)	
History of CSA**				$\chi^2 = 0.176$

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).


Present	33 (19.8)	30 (18.0)	63 (18.9)	df = 1
Absent	134 (80.2)	137 (82.0)	271 (81.1)	p = 0.675
Recent Abuse				$\chi^2 = 3.024$
Absent	130 (77.8)	116 (69.5)	246 (73.7)	df = 1
Present	37 (22.2)	51 (30.5)	88 (26.3)	p = 0.082
Anxiety				$\chi^2 = 3.678$
Negative	154 (92.2)	143 (85.6)	297 (88.9)	df = 1
Positive	13 (7.8)	24 (14.4)	37 (11.1)	p = 0.055

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

11 *CPA- childhood physical abuse, **CSA- childhood sexual abuse

12

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).




13

14 **Figure 1: Planning Status of the Pregnancies of the Pregnant Adolescents**

15

16

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).



17

18 **Figure 2: Disposition of Pregnant Adolescents in Osogbo towards their**
19 **Pregnancies**

20

21 **Table 3: Relationship between depression and socio-demographic variables in pregnant**
22 **and non-pregnant adolescents**

Pregnant adolescents	Non-pregnant adolescents
----------------------	--------------------------

Variable	Not Depressed (n=153) Freq (%)	Depressed (n=14) Freq (%)	Total (n=167) Freq (%)	Statistics	Not Depressed (n=162) Freq (%)	Depressed (n=5) Freq (%)	Total (n=167) Freq (%)
Marital Status							
Married	38 (92.7)	3 (7.3)	41 (100.0)	Fisher's Exact P=1.000	21 (95.5)	1 (4.5)	22 (100.0)
Not married	115 (91.3)	11 (8.7)	126 (100.0)		141 (97.2)	4 (2.8)	145 (100.0)
Educational attainment							
<Secondary	37(90.2)	4 (9.8)	41 (100.0)	Fisher's Exact P=0.748	25(100.0)	0 (0.0)	25 (100.0)
≥Secondary	116 (92.1)	10 (7.9)	126(100.0)		137 (96.5)	5 (3.5)	142(100.0)
Employment Status							
Employed	38 (92.7)	3 (7.3)	41 (100.0)	χ ² =0.814* df=3 p=0.846	6 (75.0)	2 (25.0)	8 (100.0)
Unemployed	27 (93.1)	2 (6.9)	29 (100.0)		13 (92.9)	1 (7.1)	14 (100.0)
Student	60 (92.3)	5 (7.7)	65 (100.0)		140(99.3)	1 (0.7)	141 (100.0)
Apprentice	28 (87.5)	4 (12.5)	32 (100.0)		3 (75.0)	1 (25.0)	4 (100.0)
Religion							
Christianity	53 (93.0)	4 (7.0)	57 (100.0)	χ ² =2.507* df =2	77 (96.3)	3 (3.8)	80 (100.0)
Islam	99 (91.7)	9 (8.3)	108 (100.0)		82 (97.6)	2 (2.4)	84 (100.0)

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22233375>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Traditional	1 (50.0)	1 (50.0)	2 (100.0)	p=0.286	3(100.0)	0 (0.0)	3 (100.0)
Living arrangement							
Partner	46 (93.9)	3 (6.1)	49 (100.0)	χ^2	9 (90.9)	1(10.0)	10(100.0)
Parents	71 (92.2)	6 (7.8)	77 (100.0)	=12.903*	126 (99.2)	1 (0.8)	127 (100.0)
Grandparents	33 (97.1)	1 (2.9)	34 (100.0)	df =3	11 (100.0)	0(0.0)	11 (100.0)
Alone	3 (42.9)	4 (57.1)	7 (100.0)	p=0.005	16 (84.2)	3 (15.8)	19 (100.0)
Parent's Marital Status							
				Fisher's			
Married	98 (90.7)	10 (9.3)	108 (100.0)	Exact	146 (96.7)	5 (3.3)	151 (100.0)
Not Married	55 (93.2)	4 (6.8)	59 (100.0)	P=0.772	16 (100.0)	0 (0.0)	16 (100.0)

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

23

24

25

26

27

28 **Table 4 :Relationship between depression and psychosocial variables in pregnant and non-**
 29 **pregnant adolescents**

	Pregnant adolescents	Non-pregnant adolescents
--	----------------------	--------------------------

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Variable	Not Depressed (n=153) Freq (%)	Depressed (n=14) Freq (%)	Total (n=167) Freq (%)	Statistics	Not Depressed (n=162) Freq (%)	Depressed (n=5) Freq (%)	Total (n=167) Freq (%)
Hx of mental illness							
Yes	159 (97.0)	5 (3.0)	164 (100.0)	Fisher's Exact P=0.029	3 (100.0)	0 (0.0)	3 (100.0)
No	147 (93.0)	11 (7.0)	158 (100.0)				
Social support							
Poor	20 (87.0)	3 (13.0)	23 (100.0)	$\chi^2=4.954$ df=2 p=0.611	28 (93.3)	2 (6.7)	30 (100.0)
Intermediate	99 (91.7)	9 (8.3)	108 (100.0)		87 (98.8)	1 (1.1)	88 (100.0)
Strong	34 (94.4)	2 (5.6)	36 (100.0)		47 (95.9)	2 (4.1)	49 (100.0)
Hx of CPA*							
Absent	131 (92.3)	11 (7.7)	142 (100.0)	Fisher's Exact p=0.443	147 (97.4)	4 (2.6)	151 (100.0)
Present	22 (88.0)	3 (12.0)	25 (100.0)		15 (93.8)	1 (6.3)	16 (100.0)
Hx of CSA++							
Absent	126 (94.0)	8 (6.0)	134 (100.0)	$\chi^2=5.141$ df=1	134 (97.8)	3 (2.2)	137 (100.0)
Present	27 (81.8)	6 (18.2)	33 (100.0)		28 (93.3)	2 (6.7)	30 (100.0)

				p=0.023			
Recent Abuse							
Absent	118 (90.8)	12 (9.2)	130 (100.0)	Fisher's	113 (97.4)	3 (2.6)	116 (100.0)
Present	35 (94.6)	2 (5.4)	37 (100.0)	Exact	49 (96.1)	2 (3.9)	51 (100.0)
				p =0.737			
Anxiety							
Negative	145 (94.2)	9 (5.8)	144 (100.0)	$\chi^2 =16.606$	141 (98.6)	2 (1.4)	143 (100.0)
Positive	8 (61.5)	5(38.5)	13 (100.0)	df = 1	21 (87.5)	3 (12.5)	24 (100.0)
				p=0.001			
Pregnancy*+ status							
				$\chi^2 =0.137$			
Planned	62 (92.5)	5 (7.5)	67 (100.0)	df = 1			
Not planned	90 (90.9)	9 (9.1)	99 (100.0)	p=0.711	-----	-----	-----
Pregnancy disposition^s							
Happy	69 (93.2)	5 (6.8)	74 (100/0)	χ^2			
Sad	47 (87.0)	7 (13.0)	54 (100.0)	=2.035**	-----	-----	-----
Indifferent	35 (94.6)	2 (5.4)	37 (100.0)	df = 2			
				p=0.362			

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.01.22273317>; this version posted April 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

30 ***CPA- childhood physical abuse; ++CSA-childhood sexual abuse; **likelihood ratio χ^2**

31 ***+n=166 ; ^sn=165**

32