Health conditions associated with sexual assault in a large hospital population

Allison M. Lake, BSc[1,2], Slavina B. Goleva, PhD[1,2,3], Lauren R. Samuels, PhD[4], Laura M.

Carpenter, PhD[5], Lea K. Davis, PhD[1,2,3,6,7,8]

1 – Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Nashville, TN

2 – Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical

Center, Nashville, TN

3 – Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville,

Tennessee

4 – Department of Biostatistics, Vanderbilt University, Nashville, TN

5 – Department of Sociology, Vanderbilt University Medical Center, Nashville, TN

6 – Department of Psychiatry and Behavioral Sciences, Vanderbilt University Medical Center,

Nashville, TN

7 – Department of Medicine, Vanderbilt University Medical Center, Nashville, TN

8 – Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN

Abstract

Objective: To develop a clinical informatics approach to identify patients with a history of

sexual assault and to characterize the clinical risk factors and comorbidities of this population in

a sex-stratified manner.

Methods: We developed and applied a keyword-based approach to clinical notes to identify

patients with a history of sexual assault in the Vanderbilt University Medical Center (VUMC)

electronic health record from 1989 to 2021. Using a phenome-wide association study (PheWAS),

we then examined diagnoses that co-occurred with evidence of sexual assault. We also examined

whether sex assigned at birth modified any of these associations.

Results: Our keyword-based algorithm achieved a positive predictive value of 90.4%, as

confirmed by manual patient chart review. Out of 1,703 diagnoses tested across all subgroup

analyses, we identified a total of 465 associated with sexual assault, many of which have been

previously observed in the literature. Interaction analysis revealed 55 sex-differential phenotypic

associations.

Conclusions: In a large hospital setting, disclosures of sexual assault were associated with

increased rates of hundreds of health conditions.

<u>Introduction</u>

Sexual assault, which includes rape, forced penetration of another person, sexual

coercion, and unwanted sexual contact, is a major public health and human rights concern,

estimated to affect 44% of women and 25% of men in the United States during their lifetimes[1].

Rape alone has an astonishing lifetime prevalence of 20% in women[1]. Sexual assault has

immediate and wide-ranging health consequences, including direct physical injuries and adverse

effects on reproductive and mental health[2]. Sexual assault is also associated with debilitating

long-term health outcomes. Survivors of sexual assault exhibit increased rates of psychiatric

conditions including post-traumatic stress disorder (PTSD), anxiety, depression, sleep disorders,

eating disorders, and suicide attempts[3]. Furthermore, sexual assault is associated with lifetime

diagnoses of multiple functional and somatic disorders, including gastrointestinal symptoms,

chronic pain, and functional seizures[4].

The ability to identify patients with a history of sexual assault is crucial for understanding

health consequences related to sexual assault. Large studies are needed to understand the unique

healthcare needs of this population. However, the challenges inherent in sexual assault research

make this a daunting task. The highly sensitive and often stigmatized nature of sexual assault can

leave many patients and research participants unwilling to disclose these experiences.

Stigmatization and other cultural factors may also lead survivors to reject the classification of

their experiences as sexual assault[5] (cited in[6]). Furthermore, research on male survivors is

relatively limited, likely in part due to differential reporting of sexual assault between men and

women[7]. Given the sex differences widely observed for common mental health conditions, it is

plausible that the effects of sexual assault on mental health differ between men and women.

Despite the importance of this line of research, sex differences in sexual assault-related health

effects are understudied, and few robust sex differences have been identified[3,4,7].

Research utilizing electronic health records (EHRs) has the potential to overcome some

limitations of traditional epidemiological approaches. However, few EHR-based studies of

sexual assault and associated health sequelae exist[8,9]. In the present study, we utilized both

billing codes and key phrases to identify patients with a history of sexual assault in a large

hospital setting. Our novel key phrase approach, which mines clinical note text for matches to

certain relevant phrases, identified over 14,000 patients who have experienced sexual assault out

of over 3 million patients, a twofold increase over the 6,998 patients identified using billing

codes alone. We then characterized the medical phenome of a subset of this patient population

and identified hundreds of associated clinical phenotypes, including many related to psychiatric

conditions. Finally, we demonstrated the sex-differentiated nature of dozens of clinical

associations, addressing a key gap in sexual assault research.

Methods

Sample and data description

This study included 3,376,424 de-identified patients who received care at VUMC, an

academic medical center in Nashville, TN, between the years 1989 and 2021. Demographic

information, ICD-9 and ICD-10 codes, and clinical notes were obtained from the medical record

for use in subsequent analyses. No age restrictions were applied. For all association analyses, we

retained only individuals meeting "medical home" criteria as described[10] to limit the study

population to patients receiving regular care at VUMC and reduce nonrandom missingness

between cases and controls. This was accomplished by restricting our study population to

individuals with at least 5 ICD-9 or ICD-10 codes, of any type, on different days over a period of

at least 3 years. We further limited the study population to individuals with recorded male or

female sex assigned at birth, and with at least one recorded BMI. This study was reviewed and

approved by the VUMC institutional review board (IRB #212285) and exempted from informed

consent requirements as non-human subjects research using deidentified medical data.

Sexual assault case/control algorithms

Although ICD codes do exist for rape, sexual assault, and sexual abuse, these are

typically reserved for care immediately after an assault and are infrequently used to indicate a

history of sexual abuse or assault. Across the entire cohort, only 6,998 patient charts included

ICD codes for sexual assault (0.21%), likely an extreme underestimate given sexual assault

prevalence estimates[1]. To improve detection of patients with a history of sexual assault, we

developed an algorithm to search the unstructured free text from clinical notes associated with

patient charts for matches to specific phrases. These clinical notes consisted of admission notes,

ancillary reports, discharge summaries, emergency department notes, inpatient notes, notes not

otherwise classified, nursing reports, outpatient notes, pathology reports, and problem lists.

In developing the algorithm, we first identified charts containing relevant ICD codes

and/or matches to an initial set of keywords with obvious relevance (e.g., "rape" and "sexual

assault"). We then performed an exploratory chart review and identified an expanded set of

relevant keywords and exclusion phrases. This subsequent "phase 1" algorithm identified

individuals meeting criteria for disclosure of sexual assault using the chart-review-derived set of

key phrases as well as relevant ICD codes (**Supplemental Table 1**). We then performed a

comprehensive manual review of 25 charts meeting these criteria to further refine our keyword

search terms accordingly. This included identifying commonly used phrasing to indicate the

presence of a history of sexual assault (i.e., inclusion) or the absence of a history of sexual

assault (i.e., exclusion).

Our final "phase 2" keyword-based algorithm consisted of 18 inclusion phrases (e.g.,

"history of sexual assault", "hx of rape") and 12 exclusion phrases (i.e., "denies history of sexual

abuse", "no hx of sexual assault") to identify patients who disclosed sexual assault

(**Supplemental Table 1**). To allow comparisons between keyword-based and ICD-based

approaches, ICD codes were not used in this final keyword-based algorithm. A patient was

considered a case if their chart contained a match to at least one inclusion phrase and contained

no matches to any exclusion phrases. Otherwise, the patient was considered a control.

To benchmark our keyword-based algorithm, we performed additional analyses using

only ICD codes to identify patients with positive and negative histories of sexual assault. In the

ICD-code-based approach, charts were defined as cases if they contained at least 1 of 25 ICD-9

and ICD-10 codes pertaining to sexual assault (**Supplemental Table 2**), and as controls

otherwise. We did not exclude individuals meeting both ICD-based and keyword-based criteria.

Chart review

To evaluate our final "phase 2" keyword-based algorithm, AML conducted a manual

review of 52 randomly selected charts identified by the sexual assault algorithm and calculated

the positive predictive value of the algorithm. The chart reviews were conducted within a subset

of patients who are also enrolled in the VUMC biobank, BioVU. These patients tend to have

longer health records (median 9.99 years among biobank participants compared to median of

1.15 years for the entire VUMC-EHR) and thus present an optimal setting for validating a

phenotyping algorithm that relies on the presence of key phrases in clinical notes. Charts were

identified as true positives if they clearly contained a description of the patient's reported history

of sexual assault.

Phecode-based analysis

The presence or absence of a history of sexual assault was used as the primary

independent variable in a PheWAS. The medical phenome was categorized by phecodes—single

codes representing clinical phenotypes—that were generated by aggregating related billing codes

into hierarchical code families[11]. Using the PheWAS package[12] (version 0.99) in R[13], we mapped

ICD-9 and ICD-10 codes to 1,817 phecodes using Phecode Map version 1.2

(phewascatalog.org)[14,15]. We conducted a sex-combined PheWAS and sex-stratified (i.e.,

separately in males and females) PheWAS. For each phenotype tested in the sex-combined

analysis, we required a minimum of 200 records, with at least 100 records for both males and

females. Accordingly, for the sex-stratified analyses, we required a minimum of 100 records per

phecode. For each phecode analyzed (n=1482, 1658, and 1527 for sex-combined, female-only,

and male-only, respectively), we fitted a multiple logistic regression model with sexual assault

history (binary: positive or negative) as the primary independent variable and EHR-reported

race, ethnicity, median age of all visits in an individual's medical record, median body mass

index (BMI) of all visits in an individual's medical record, and the log-transformed mean

number of ICD codes per day in the record as covariates. Sex-combined analyses additionally

included sex as a covariate.

For the 354 phenotypes that were significantly associated with sexual assault in at least

one sex in the sex-stratified analyses and that had at least 100 records for each sex, we conducted

interaction analyses to formally examine the moderating effect of sex on phenotypic associations

with sexual assault. These analyses used models identical to those used in the sex-combined

analyses, with the addition of a sex-by-sexual-assault interaction term.

For each association analysis, we applied a Bonferroni correction (p= 0.05/total number

of phecodes tested) to correct for the number of phecodes tested, using a threshold of corrected

p<0.05 for an association to be considered phenome-wide-significant. Although Bonferroni

correction is likely to be overly conservative given the correlation inherent in related diagnostic

codes, the large sample size counteracts this loss of power.

Results

Sexual assault prevalence in the electronic health record

Across our entire cohort (3,376,424 patients), the keyword-based approach identified

14,496 individuals (0.43%) with documentation of sexual assault, whereas only 6,998 cases

(0.21%) (with an overlap of 2,928 individuals) received ICD codes for sexual assault. After

applying medical-home criteria to our cohort (see *Methods*), we removed 11 individuals with

unknown sex and 154,804 individuals with missing BMI, resulting in a total sample of 833,185

individuals. Among these individuals, 9,333 were designated as cases by the keyword-based

algorithm (0.94%, **Table 1**), and 4,422 by the ICD-code-based algorithm (0.45%, **Supplemental**

Table 3), with an overlap of 2,014 individuals. Patients with keyword- or ICD-based

documentation of sexual assault were predominantly female, and younger on average than the

remainder of the medical-home population (**Table 1, Supplemental Table 3**).

Sexual assault case-control algorithm performance

Of the 52 patient charts identified as cases by the sexual assault case-control algorithm

and randomly selected to assess the positive predictive value (PPV) of the algorithm, 47 were

found to be true positives (PPV=90.4%, 95% confidence interval [CI] 78.2%-96.4%). In the 5/52

charts identified as false positives, inclusion phrases were present either as negation of sexual

assault history (e.g., "denies a h/o sexual abuse") or in reference to someone other than the

patient (e.g., "[sibling] was sexually abused").

History of sexual assault is associated with hundreds of clinical phenotypes

Our sex-combined PheWAS identified 386 out of 1,482 medical and psychiatric

conditions as significantly overrepresented (Bonferroni-adjusted α=3.34e-05) among patients

disclosing sexual assault. Of the 50 associations with lowest p-values, 40 were psychiatric

disorders, including schizophrenia (OR=86.36, 95% CI=76.31, 97.75), suicide or self-inflicted

injury (OR=86.33, 95% CI=74.82, 99.6), depression (OR=27.5, 95% CI=25.57, 29.57), and

PTSD (OR=122.81, 95% CI=113.47, 132.91, **Figure 1, Supplemental Table 4**). These top

associations also included phecodes relating to personality disorders: "personality disorders"

(OR=107.35, 95% CI=94.51, 121.92) and "antisocial/borderline personality disorder"

(OR=129.17, 95% CI=111.73, 149.32). In total, 64 of the 71 tested phecodes pertaining to

psychiatric disorders were significantly associated with sexual assault.

The 386 statistically significant associations also included many non-psychiatric health

conditions. Some of these associations plausibly represent immediate physical consequences of

sexual assault, including urinary tract infection (OR=1.79, 95% CI=1.68, 1.91), sexually

transmitted infection (OR=4.6, 95% CI=3.85, 5.5), and contusion (OR=2.19, 95% CI=1.9, 2.53),

Figure 2, Supplemental Table 4). Many of the non-psychiatric clinical associations with the

largest effect sizes pertained to poisonings, including poisoning by psychotropic agents

(OR=18.13, 95% CI=15.55, 21.13) and poisoning by analgesics, antipyretics, or antirheumatics

(OR=5.35, 95% CI=4.69, 6.09).

Survivors of sexual assault frequently suffer from disorders involving functional and

somatic symptoms including chronic pelvic pain and chronic gastrointestinal symptoms[4]. In our

analysis, the broad "somatoform disorder" phecode is strongly associated with sexual assault

(OR=53.84, 95% CI=46.35, 62.55). We additionally note multiple clinical associations related to

functional and somatic symptom disorders, including abdominal pain (OR=2.18, 95% CI=2.07,

2.29), chronic (nonspecific) pain (OR=2.24, 95% CI=2.09, 2.41), unspecified myalgia and

myositis (OR=2.31, 95% CI=2.13, 2.5), and cervicalgia (OR=1.75, 95% CI=1.62, 1.88). We also

identify a range of urinary symptoms associated with sexual assault, consistent with prior studies

linking sexual assault and genitourinary symptoms[16,17], including dysuria (OR=2.11, 95%

CI=1.95, 2.29), urinary frequency (OR=2.18, 95% CI=1.95, 2.43), urinary retention (OR=2.67,

95% CI=2.33, 3.06), urinary incontinence (OR=1.97, 95% CI=1.78, 2.19), and cystitis

(OR=2.35, 95% CI=2.06, 2.68).

Several phenome-wide-significant phecodes identified in our analysis pertained to

seizures and epilepsy, including the broad "epilepsy, recurrent seizures, convulsions" phecode

and all phecodes classified under it. Given the previously identified relationship between sexual

assault and the development of functional seizures, we investigated whether these associations

were partially explained by patients experiencing functional seizures as classified by our

previously published phenotyping algorithm[10]. Indeed, we observed a high level of overlap

between seizure-related phecodes and functional seizure status (**Supplemental Table 5**).

Furthermore, when we included functional seizure status as a covariate in the logistic regression

models, all associations between sexual assault and seizure-related phecodes were at least

partially attenuated (**Supplemental Table 6**).

Sex-differential phenotypic association analyses

Sex differences in prevalence, age of onset, and presentation of many neuropsychiatric

disorders are well characterized in the literature[18,19]. We analyzed our cohort for sex-differential

phenotypic associations to determine whether a history of sexual assault is modified by sex

assigned at birth. First, we first performed sex-stratified PheWAS of sexual assault disclosure

across all phecodes with sufficient case counts (**Supplemental Table 4,** "male_only" and

"female_only" sheets). We identified 410 phecodes significantly associated with sexual assault

in at least one of the analyses, 332 of which were also significant in the sex-combined analysis.

Many of the significant sex-stratified associations were specific to one sex. We identified

three male-specific associations with sexual assault: viral hepatitis A (OR=7.93, 95% CI=3.2,

19.64), testicular hypofunction (OR=2.25, 95% CI=1.73, 2.93), and testicular dysfunction

(OR=2.24, 95% CI=1.72, 2.91). The female-specific associations were diverse and included

infectious diseases, genitourinary conditions, and psychiatric disorders. Focusing on the

psychiatric associations, we note that sexual assault is strongly associated with the broad "mental

disorders during/after pregnancy" phecode (OR=28.77, 95% CI=24.6, 33.66). We also observe a

strong association between sexual assault and the female-predominant diagnosis of dissociative

disorder (OR=113.77, 95% CI=86.15, 150.24), which recapitulates previous findings that the

majority of patients with this disorder have a history of abuse or neglect[20-21] (cited in [22]).

Of the 410 significant sex-stratified associations, the 354 phecodes with sufficient case

counts in both sexes were tested in an interaction analysis examining the moderating effect of

sex on sexual assault comorbidities. Of these, 54 phecodes demonstrated significant differences

between males and females (Bonferroni-adjusted α=1.41e-04, **Figure 3, Supplemental Table 4**

["sex_interaction" sheet]), 32 of which pertained to psychiatric conditions, including depression,

anxiety disorders, schizophrenia, and PTSD. Interestingly, in all 32 sex-differential psychiatric

associations with sexual assault, the effect sizes were significantly greater in males than in

females.

Among the non-psychiatric associations, those stronger in females than in males included

"sprains and strains," gastrointestinal symptoms ("abdominal pain;" "nausea and vomiting"),

urinary retention, urinary tract infection, and seizure-related phecodes ("epilepsy, recurrent

seizures, convulsions;" "convulsions"). Non-psychiatric associations that were stronger in males

included anal and rectal conditions, HIV-related phecodes, herpes simplex infection, and

unspecified congenital anomalies. Several sex-differential associations with non-psychiatric

phecodes exhibit opposite effects in men and women. For example, hypotension, unspecified

bacterial infection, muscle weakness, and cough were observed more frequently in females

affected by sexual assault than in females not affected, but the reverse was true for males.

Conversely, colorectal cancer was observed more frequently in males with a sexual assault

history than in those without, but the reverse was true for females.

Benchmarking of keyword algorithm using ICD-code-based case-selection

To determine whether the associations identified by a keyword-based case-control

definition are consistent with those identified by a simpler ICD-code-based approach, we

repeated the PheWAS using ICD codes to classify patients as positive or negative for a history of

sexual assault (see *Methods*). Of the 386 phenome-wide-significant sex-combined associations

identified using the keyword-based algorithm, 289 were also nominally associated (p<.05) using

the ICD-code-based algorithm. Furthermore, among phecodes identified as phenome-wide-

significant by the keyword-based approach, log ORs from the two approaches were strongly

correlated (Pearson R≥0.84, **Supplemental Figure 1, Supplemental Table 7**).

Discussion

Epidemiological studies have established that individuals with a history of sexual assault

have increased risk of developing both immediate and long-term physical and mental health

conditions[2–4,7,8,22–24]. We aimed to replicate and extend these findings using a novel clinical

informatics-based phenotyping approach in a large hospital setting. Our approach ascertained

substantially more patients than did diagnostic codes alone, yielding improved power to study

the medical phenomes of sexual assault survivors. This study not only confirmed the utility of

our approach but also produced novel insights into the consequences of sexual assault for males

and females.

Our phenotyping approach identified hundreds of clinical phenotypes significantly associated

with sexual assault, including a preponderance of associations with psychiatric conditions such

as schizophrenia, depression, PTSD, and suicidal behaviors. We observed several associations

between sexual assault and phenotypes related to functional and somatic symptom disorders

including gastrointestinal symptoms, chronic pain, and seizures. These results are consistent with

findings from the few epidemiological studies examining the impact of sexual violence on the

development of functional and somatic symptom disorders[4,8]. Nevertheless, they remain largely

understudied and warrant further investigation.

We noted several phenotypes associated with sexual assault pertaining to toxic

ingestions, such as poisoning by psychotropic agents and analgesics. Psychotropic agents

including antidepressants, as well as commonly used analgesics such as paracetamol, are used

frequently in suicide attempts[25,26]. Consistent with this, our analysis demonstrated a strong

association between sexual assault and suicide and self-harm. Given the high overlap between

toxic ingestions and documentation of suicidal behavior in our data (**Supplemental Table 8**), the

associations between sexual assault and toxic ingestions are likely in part driven by suicidal

behavior. Furthermore, when we conditioned on the "suicide or self-inflicted injury" phecode,

these associations were diminished (**Supplemental Table 9**).

Finally, we addressed a gap in the sexual assault literature by leveraging our approach to

study sex differences in risk factors and comorbidity patterns among survivors of sexual assault.

We found that dissociative disorder, which occurs predominantly in women, is strongly

associated with sexual assault in our cohort, recapitulating the observation that most women with

the disorder have a history of abuse or neglect[20–22]. Our interaction analysis revealed that

abdominal pain, nausea and vomiting, epilepsy, recurrent seizures, and convulsions are more

commonly observed in female than male survivors of sexual assault. While we cannot

conclusively assert that these conditions represent functional or somatic symptom disorders,

prior work linking sexual assault to this class of disorders[4] makes this interpretation plausible.

Furthermore, we showed that the seizure-related associations are partially attenuated when

conditioning on functional seizures case-control status, the phenotyping algorithm for which we

recently published[10].

Interestingly, most of the sex-differential psychiatric associations with sexual assault were

stronger in men than in women. This finding should be interpreted in light of possible differential

reporting and recall bias between males and females who have experienced sexual assault. For

example, if males tend to report experiences of sexual assault less frequently than females, it is

possible that the men who do disclose such a history in a medical setting are those who are

experiencing especially severe health consequences as a result of their trauma. In this scenario,

observed clinical associations with sexual assault would appear stronger in males than in

females, which is what we observe for many sex-differential phenotypes in our analysis (in

particular, psychiatric conditions).

<u>Limitations</u>

The cross-sectional design of our study prohibits us from drawing conclusions about causal

or temporal relationships between sexual assault and associated phenotypes. Some of the

phenotypic associations we identified, including phecodes relating to congenital anomalies and

neurodevelopmental disorders, are likely to have preceded sexual assault and could be

interpreted as risk factors for experiencing sexual assault later in life. However, many of the

identified associations are consistent with health sequelae of sexual assault identified in

longitudinal case-control and cohort studies.

Secondly, our analysis is subject to the ascertainment bias inherent in EHR-based medical

phenome studies. It is difficult to extrapolate our results, which are derived from a cohort of

individuals seeking medical care, to the broader population. For example, survivors of sexual

assault who frequently interact with the healthcare system may be more likely to receive a

greater number of medical diagnoses than those who do not seek or have limited access to care.

Nevertheless, it is critical that health care systems are prepared to anticipate the care needs of

survivors of sexual assault to provide high quality primary preventative care. Moreover, we have

attempted to account for this bias by including a measure of healthcare utilization as a covariate

in our analyses.

Thirdly, the small number of cases identified by our keyword-based algorithm relative to

sexual assault prevalence estimates[1] suggests that, though it is an improvement over the use of

ICD codes alone, our algorithm is still misclassifying many cases as controls and

underestimating the prevalence of sexual assault in the VUMC patient population. This

underestimation likely stems from ubiquitous underreporting of sexual assault in healthcare

settings. For example, only approximately one third of women with injuries due to rape seek any

type of medical care[27]. In the context of this misclassification, our analyses of clinical

associations with sexual assault are likely to be overly conservative. While an overly

conservative analysis introduces type II error, it does not reduce confidence in the positive

clinical associations we have identified.

<u>Public health implications</u>

Sexual assault is an urgent public health concern with both immediate and enduring

physical and mental health consequences. Our study provides novel evidence supporting a strong

relationship between sexual assault and psychiatric conditions, including functional and somatic

symptom disorders. In our cohort, the associations between sexual assault history and psychiatric

conditions are stronger in males than in females, which may reflect reporting bias and/or sex-

differential biopsychological responses to trauma. Conversely, associations relating to

gastrointestinal symptoms and seizures, which may represent manifestations of functional and

somatic symptom disorders, were more commonly observed in female than male survivors of

sexual assault.

Our analysis, in context of prior epidemiological studies, suggests that patients who have

experienced sexual assault are at risk for developing wide-ranging medical and psychiatric

comorbidities. Yet, sexual assault continues to be vastly underreported in healthcare settings.

Further research is needed to understand the conditions under which patients feel safe in

disclosing a history of sexual assault trauma. Furthermore, safe, compassionate, and effective

protocols are needed for providers to routinely screen survivors for a variety of physical and

mental health conditions, both immediately and longitudinally.

<u>Figure and table legends</u>

Table 1: Demographics of patients classified as sexual assault cases or controls by our keyword-
based phenotyping algorithm. To assess case-control differences in demographic variables, two-
sided t-tests were performed for continuous variables (record-median age, record-median BMI,
log(mean records per day)), and chi-square tests were performed for binary variables (sex,
white/non-Hispanic). Log(mean records per day) is a measure of the density of diagnostic codes
across an individual's health record.

Figure 1: Psychiatric phenotypic associations with sexual assault. Log10-scale forest plots
displaying odds ratios with 95% confidence intervals for adjusted associations between sexual
assault and all tested psychiatric phecodes in sex-combined analyses. Associations achieving
statistical significance are colored green. Related phecodes are grouped together (for example,
the "Antisocial/borderline personality DO" phecode [301.2] is related to its parent phecode
"Personality DOs" [301]). Red dashed line indicates an OR of 1. (Note: DO=disorder).

Figure 2: Non-psychiatric associations with sexual assault. PheWAS plot from sex-combined
analysis displaying –log10(p-value) for all non-psychiatric phecodes analyzed. Dashed red line
indicates significance threshold (Bonferroni-corrected $p<0.05$).

Figure 3: History of sexual assault increases odds of clinical phenotypes in a sex-differential
manner. For each of the 55 phenotypes with significant sexual-assault-by-sex interaction effects,
log10-scale odds ratio (OR) and 95% confidence interval is plotted separately by sex for non-
psychiatric phecodes (left) and psychiatric phecodes (right). The ORs correspond to the sex-
stratified associations between sexual assault and each phenotype. Associations are grouped by
related phenotypes. Red dashed line indicates an OR of 1. (Note: DO=disorder).

Supplemental Table 1. Criteria used in developing algorithm to identify patients with a history
of sexual assault. Patients were identified using natural language processing of clinical notes
using various inclusion phrases and exclusion phrases, shown here. The "Phase 1" case
definition additionally included patients with matches to relevant ICD-9 and ICD-10 codes and
yielded a positive predictive value (PPV) of 64% derived from 25 manually reviewed patient
charts. The ICD codes used for the Phase 1 algorithm were all codes listed in Supplemental

Table 2 except three codes (E960.1, Z04.41, Z04.42) which were identified later in the analysis

process. False positives were used to refine the search terms, resulting in the final or "Phase 2"

algorithm. The Phase 2 algorithm relies solely on natural language processing and does not take

ICD codes into account. The PPV of the Phase 2 algorithm, based on manual review of 52 charts,

was 90.4%.

Supplemental Table 2 ICD-9 and ICD-10 codes used to identify sexual assault patients in the

ICD-code-based algorithm. * indicates codes from the ICD-9 code set; all other codes listed are

ICD-10.

Supplemental Table 3 Demographics of patients classified as sexual assault cases or controls

based only on relevant ICD codes. To assess case-control differences in demographic variables,

two-sided t-tests were performed for continuous variables (record-median age, record-median

BMI, log(mean records per day)), and chi-square tests were performed for binary variables (sex,

white/non-Hispanic). Log(mean records per day) is a measure of the density of diagnostic codes

across an individual's health record.

Supplemental Table 4 PheWAS summary statistics for sex-combined, male-only, female-only,

and sexual-assault-by-sex interaction analyses. Here, the keyword-based sexual assault case

definition was applied. In the sex_interaction tab, ORs, confidence intervals, and p-values

derived from the sex-stratified analyses are re-displayed (OR.m, p.value.f, etc.). (OR=odds ratio,

CI=confidence interval, ".f" = female-only analysis, ".m" = male-only analysis, ".int" =

interaction analysis.)

Supplemental Table 5 Contingency tables demonstrating co-incidence of seizure-related

phecodes and functional seizure cases (functional seizure phenotyping algorithm described in [10]).

For a given phenotype, "NA" corresponds to individuals excluded from case-control analyses for

that phenotype (see [11]).

Supplemental Table 6 Sex-combined associations between sexual assault and seizure-related

phecodes after conditioning on functional seizures case-control status (functional seizure

phenotyping algorithm described in [10]). Here, the keyword-based sexual assault case definition

was applied. (OR=odds ratio, CI=confidence interval.)

Supplemental Figure 1. Correlation scatterplots of log ORs from PheWAS on keyword-based

(y-axis) and ICD-code-based (x-axis) case definitions. Log ORs are plotted for keyword-based

associations meeting Bonferroni significance threshold. Results are reported separately for sex-

combined, male-only, and female-only analyses.

Supplemental Table 7 PheWAS summary statistics for sex-combined, male-only, and female-

only analyses. Here, the ICD-code-based sexual assault case definition was applied. (OR=odds

ratio, CI=confidence interval, ".f" = female-only analysis, ".m" = male-only analysis.)

Supplemental Table 8 Contingency tables demonstrating co-incidence of toxic ingestions and

suicidal behavior. For a given phenotype, "NA" corresponds to individuals excluded from case-

control analyses for that phenotype (see [11]).

Supplemental Table 9 Sex-combined associations between sexual assault and toxic ingestions

after conditioning on the "suicide or self-inflicted injury" phecode. Here, the keyword-based

sexual assault case definition was applied. (OR=odds ratio, CI=confidence interval.)

<u>**References**</u>

1. Smith SG, Zhang X, Basile KC, et al. *The National Intimate Partner and Sexual Violence*
Survey (NISVS): 2015 Data Brief – Updated Release. National Center for Injury Prevention
and Control, Centers for Disease Control and Prevention; 2018. Accessed August 30, 2020.
https://www.cdc.gov/violenceprevention/datasources/nisvs/2015NISVSdatabrief.html

2. Jina R, Thomas LS. Health consequences of sexual violence against women. *Best Pract Res*
Clin Obstet Gynaecol. 2013;27(1):15-26. doi:10.1016/j.bpobgyn.2012.08.012

3. Chen LP, Murad MH, Paras ML, et al. Sexual Abuse and Lifetime Diagnosis of Psychiatric
Disorders: Systematic Review and Meta-analysis. *Mayo Clin Proc*. 2010;85(7):618-629.
doi:10.4065/mcp.2009.0583

4. Paras ML, Murad MH, Chen LP, et al. Sexual Abuse and Lifetime Diagnosis of Somatic
Disorders: A Systematic Review and Meta-analysis. *JAMA*. 2009;302(5):550-561.
doi:10.1001/jama.2009.1091

5. Dartnall E, Jewkes R. Sexual violence against women: the scope of the problem. *Best Pract*
Res Clin Obstet Gynaecol. 2013;27(1):3-13. doi:10.1016/j.bpobgyn.2012.08.002

6. Oram S, Khalifeh H, Howard LM. Violence against women and mental health. *Lancet*
Psychiatry. 2017;4(2):159-170. doi:10.1016/S2215-0366(16)30261-9

7. Rhodes AE, Boyle MH, Tonmyr L, et al. Sex Differences in Childhood Sexual Abuse and
Suicide-Related Behaviors. *Suicide Life Threat Behav*. 2011;41(3):235-254.
doi:10.1111/j.1943-278X.2011.00025.x

8. Young-Wolff KC, Sarovar V, Klebaner D, Chi F, McCaw B. Changes in Psychiatric and
Medical Conditions and Healthcare Utilization Following a Diagnosis of Sexual Assault: A
Retrospective Cohort Study. *Med Care*. 2018;56(8):649-657.
doi:10.1097/MLR.0000000000000930

9. Brignone E, Gundlapalli AV, Blais RK, et al. Increased Health Care Utilization and Costs
Among Veterans With a Positive Screen for Military Sexual Trauma. *Med Care*.
2017;55:S70. doi:10.1097/MLR.0000000000000767

10. Goleva SB, Lake AM, Torstenson ES, Haas KF, Davis LK. Epidemiology of Functional
Seizures Among Adults Treated at a University Hospital. *JAMA Netw Open*.
2020;3(12):e2027920. doi:10.1001/jamanetworkopen.2020.27920

11. Denny JC, Ritchie MD, Basford MA, et al. PheWAS: demonstrating the feasibility of a
phenome-wide scan to discover gene–disease associations. *Bioinformatics*.
2010;26(9):1205-1210. doi:10.1093/bioinformatics/btq126

12. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for
phenome-wide association studies in the R environment. *Bioinformatics*. 2014;30(16):2375-
2376. doi:10.1093/bioinformatics/btu197

13. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing; 2020. https://www.R-project.org/

14. Denny JC, Bastarache L, Ritchie MD, et al. Systematic comparison of phenome-wide
association study of electronic medical record data and genome-wide association study
data. *Nat Biotechnol*. 2013;31(12):1102-1111. doi:10.1038/nbt.2749

15. Wu P, Gifford A, Meng X, et al. Mapping ICD-10 and ICD-10-CM Codes to Phecodes:
Workflow Development and Initial Evaluation. *JMIR Med Inform*. 2019;7(4):e14325.
doi:10.2196/14325

16. Mayson BE, Teichman JMH. The relationship between sexual abuse and interstitial
cystitis/painful bladder syndrome. *Curr Urol Rep*. 2009;10(6):441-447.
doi:10.1007/s11934-009-0070-3

17. Lalchandani P, Lisha N, Gibson C, Huang AJ. Early Life Sexual Trauma and Later Life
500   Genitourinary Dysfunction and Functional Disability in Women. *J Gen Intern Med*.
501   2020;35(11):3210-3217. doi:10.1007/s11606-020-06118-0

502   18. Arnett AB, Pennington BF, Willcutt EG, DeFries JC, Olson RK. Sex differences in ADHD
503   symptom severity. *J Child Psychol Psychiatry*. 2015;56(6):632-639.
504   doi:10.1111/jcpp.12337

505   19. Brody DJ, Pratt LA, Hughes JP. *Prevalence of Depression among Adults Aged 20 and over:*
506   *United States, 2013–2016*. National Center for Health Statistics; 2018. Accessed August 30,
507   2020. https://www.cdc.gov/nchs/products/databriefs/db303.htm

508   20. Kluft RP. An Update on Multiple Personality Disorder. *Psychiatr Serv*. 1987;38(4):363-373.
509   doi:10.1176/ps.38.4.363

510   21. Ross CA, Miller SD, Bjornson L, Reagor P, Fraser GA, Anderson G. Abuse histories in 102
511   cases of multiple personality disorder. *Can J Psychiatry Rev Can Psychiatr*. 1991;36(2):97-
512   101. doi:10.1177/070674379103600204

513   22. Teicher MH, Samson JA. Childhood Maltreatment and Psychopathology: A Case for
514   Ecophenotypic Variants as Clinically and Neurobiologically Distinct Subtypes. *Am J*
515   *Psychiatry*. 2013;170(10):1114-1133. doi:10.1176/appi.ajp.2013.12070957

516   23. Finestone HM, Stenn P, Davies F, Stalker C, Fry R, Koumanis J. Chronic pain and health
517   care utilization in women with a history of childhood sexual abuse. *Child Abuse Negl*.
518   2000;24(4):547-556. doi:10.1016/S0145-2134(00)00112-5

519   24. Kilpatrick DG, Ruggiero KJ, Acierno R, Saunders BE, Resnick HS, Best CL. Violence and
520   risk of PTSD, major depression, substance abuse/dependence, and comorbidity: results
521   from the National Survey of Adolescents. *J Consult Clin Psychol*. 2003;71(4):692-700.
522   doi:10.1037/0022-006x.71.4.692

523   25. Hawton K, Bergen H, Simkin S, et al. Toxicity of antidepressants: rates of suicide relative to
524   prescribing and non-fatal overdose. *Br J Psychiatry*. 2010;196(5):354-358.
525   doi:10.1192/bjp.bp.109.070219

526   26. Sarchiapone M, Mandelli L, Iosue M, Andrisano C, Roy A. Controlling Access to Suicide
527   Means. *Int J Environ Res Public Health*. 2011;8(12):4550-4562.
528   doi:10.3390/ijerph8124550

529   27. Full Report of the Prevalence, Incidence, and Consequences of Violence Against Women:
530   Findings From the National Violence Against Women Survey. National Institute of Justice.
531   Accessed November 10, 2021. https://nij.ojp.gov/library/publications/full-report-
532   prevalence-incidence-and-consequences-violence-against-women

<u>Acknowledgements</u>

Research reported in this publication was supported by NIGMS of the National Institutes of

Health under award number T32GM007347 and Grant No. R01 HG011405, partially funded by

the Office of Research on Women's Health, Office of the Director, NIH and the National Human

Genome Research Institute. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the Office of Research on Women's Health, the

National Human Genome Research Institute, or the NIH.

Odds ratio with 95% confidence interval

Odds ratio with 95% confidence interval

	Case (N=9333)	Control (N=823852)	Test statistic	P-value
Sex				
Female	7610 (81.5%)	467617 (56.8%)	χ^2=2311.26	<0.001
Male	1723 (18.5%)	356235 (43.2%)		
Record-median age				
Median [Min, Max]	28.0 [0, 86.0]	37.5 [0, 89.0]	t=-39.47	<0.001
Record-median BMI				
Median [Min, Max]	26.4 [11.3, 59.2]	25.8 [10.0, 60.0]	t=18.04	<0.001
White and non-Hispanic				
No	2436 (26.1%)	207663 (25.2%)	χ^2=3.87	0.0492
Yes	6897 (73.9%)	616189 (74.8%)		
Log(mean records per day)				
Median [Min, Max]	-3.44 [-7.16, 0.699]	-4.09 [-7.68, 1.37]	t=53.07	<0.001