Ambient Temperature and Dengue Hospitalisation in Brazil over a 10-year period, 2010-2019: a times series analysis

Rafael Lopes Paixão da Silva[1,2,3,4], Xavier Basagaña[2,3,4], Leonardo S. L. Bastos[5], Fernando A. Bozza[6], Otavio T. Ranzani[2,3,4,7]

1 - Instituto de Física Teórica, IFT, UNESP, São Paulo, Brazil
2 - Barcelona Institute for Global Health, ISGlobal, Barcelona, Spain
3 - Universitat Pompeu Fabra, UPF, Barcelona, Spain
4- CIBER Epidemiología y Salud Pública, CIBERESP, Madrid, Spain
5 - Department of Industrial Engineering (DEI), Pontifical Catholic University of Rio de Janeiro (PUC-Rio), Rio de Janeiro, RJ, Brazil
6 - National Institute of Infectious Disease Evandro Chagas (INI), Oswaldo Cruz Foundation (FIOCRUZ), Rio de Janeiro, RJ, Brazil
7 - Pulmonary Division, Heart Institute (InCor), HCFMUSP, Faculdade de Medicina da Universidade de São Paulo, Brazil

Corresponding author:
Rafael Lopes Paixão da Silva (rafael.lp.silva@unesp.br)
IFT-Unesp, R. Dr. Bento Teobaldo Ferraz, 271, 01140-070 São Paulo, Brazil

Otavio T. Ranzani (otavio.ranzani@isglobal.org)
ISGlobal, Dr. Aiguader, 88, 08003 Barcelona, Spain

Word count: 2,669.

Summary:

Background: Climate factors are known to influence seasonal patterns of dengue transmission. However, little is known about the effect of extreme heat on the severity of dengue infection, such as hospital admission. We aimed to quantify the effect of ambient temperature on dengue hospitalisation risk in Brazil.

Methods: We retrieved daily dengue hospitalisation counts by each of 5,570 municipalities across the 27 states of Brazil from 1st of January 2010 to 31st of December 2019, from the Brazilian Public Hospital Admission System ("SIH"). We obtained average daily ambient temperature for each municipality from the ERA5-land product reanalysis. We combined distributed lag non-linear models with time stratified design model framework to pool an estimate for dose-response and lag-response structures for the association of Dengue hospitalisation relative risk (RR) and temperature. We estimated the overall dengue hospitalisation RR for the whole country as well as for each of the five macro-regions by meta-analysing state level estimates.

Findings: 579,703 hospital admissions due to dengue occurred over the 10 years period of 2010 to 2019. We observed a positive association between high temperatures and high risk of hospitalisation for Brazil and each of the five macro-regions. The overall RR for dengue hospitalisation was at the 50th percentile of temperature distribution 1·40 (95% IC 1·27-1·54) and at 95th percentile of temperature the RR was 1·50 (1·38-1·66) for Brazil, relative to the minimum temperature, which was the one with the lowest risk. We also found lagged effects of heat on hospitalisation, particularly observed on the same day (lag 0) both at the 50th percentile and 95th.

Interpretation: High temperatures are associated with an increase in the risk of hospitalisation by dengue infection. These findings may guide preparedness and mitigation policies during dengue season outbreaks, particularly on the health-care demand.

Funding: Conselho Nacional de Pesquisa, Coordenação Nacional de Aperfeiçoamento de Pessoal, Institut de Salud Carlos III.

Keywords: Dengue; severe Dengue; temperature; heatwave; hospital

Research in context

Evidence before this study Temperature can have an effect of aggravating a dengue infection progressing to severe disease requiring hospitalisation. On February 12, 2022, we searched PudMed for the terms "dengue", "severe or hospital" and "temperature or heat". Although there is extensive literature on the association of temperature and number of dengue cases, we did not find any study quantifying the association of ambient temperature and hospitalisation by dengue. Quantitative evidence on this association, and how the extreme high temperature augments the risk of hospitalisation by dengue is needed to guide plans of mitigation and preparedness.

Added value of this study Our findings show that heat can aggravate an on-going infection of Dengue virus leading to hospitalisation. Average temperatures above 23.96º Celsius degrees, the 50th percentile of temperature distribution over Brazil, represents a risk of over 1.3x to the Minimum Hospitalisation Temperature. An average temperature of 28.68º Celsius degrees, the 95th percentile of temperature distribution, represents a risk of hospitalisation of over 1.4x to the Minimum Hospitalisation Temperature. The South region, which does not have a high incidence of Dengue cases and a colder climate compared to other regions, has a temperature-related risk of hospitalisation of greater magnitude compared to other regions.

Implications of all the available evidence Heat exposure plays an important role to the public health policy for mitigating dengue virus burden on the population. Hospitalisation is costly to any health system and can be more impacting on fragile or unprepared health systems as frequently found in countries affected by dengue. Due to its hyperendemicity and its seasonality, dengue can be predicted in mid to long term allowing for better designed early-warnings systems to mitigation measures.

INTRODUCTION

Dengue fever has been a major global seasonal endemic disease, mainly present in the tropics, with an estimated burden over ~390 million infections per year.[1–4] Approximately one-quarter of these infections manifest as clinical or subclinical disease.[1] Brazil is one of the most affected countries by Dengue, that has caused over ~20·9 million cases of infections in the last 20 years, since the compulsory universal notification to the Brazilian health systems was established.[5,6] Additionally, it is estimated that the Brazilian Universal Health System (SUS) spent more than USD 159 million in the treatment and assistance to Dengue cases and USD 10 million on severe Dengue between 2000 and 2015.[5] The impressive burden of Dengue in Brazil is also observed in several others low-income and middle-income countries.[7]

Dengue incidence is influenced by climate variables due to mosquito life cycle and human behaviour changes, particularly with rainfall and temperature.[1,6,7] However, there is lacking evidence for the association of ambient temperature and the frequency of severe dengue cases. The pathophysiology of Dengue severity involves dehydration and coagulation disorders, both conditions that can be aggravated by ambient temperature.[8,9]

Based on the extensive literature on the association of temperature and all-cause hospitalizations,[2,3,5,6] we hypothesised that high ambient temperatures are associated with a

higher risk of hospitalisation due to Dengue. We aimed to evaluate the association between ambient temperature and Dengue hospitalisation in Brazil, analysing the period from 2010 to 2019.

METHODS

Setting

Brazil has about 211 million inhabitants distributed over 8·5 million km² area, divided into 26 states and the country capital, the federal district.[10] These 27 states are grouped in 5 administrative macro-regions: North, Northeast, Center-West, Southeast, and South. The country is located in a tropical region and has 3 main Köppen climate types and 12 subtypes.[11]

We conducted a time-stratified regression analysis to evaluate the association between ambient temperature and Dengue hospitalisation in Brazil. The unit of analysis was municipality (n=5,570) of residence.

Data sources and definitions

Dengue hospitalizations

We used the Brazilian Hospital Admission System (SIH), a nationwide database that comprises individual level data of all hospitalizations covered by the Universal Healthcare System (SUS) in Brazil. We defined a dengue hospitalisation by the ICD-10 codes 'A90', 'A91', 'A97', 'A970', 'A971', 'A972', 'A979'.[12] We build daily time series by date of hospitalisation for each municipality of residence. We excluded dengue hospitalizations from three municipalities that are placed in small islands, because there was no ambient temperature for them. This filtering resulted in a loss of 153 (0.03%) cases in total, for the whole period of the 10 years. The final

time series aggregation by municipalities encompasses a total of 579,703 hospital admissions due to Dengue over the 10 years period of 2010 to 2019.

Ambient temperature

We used the gridded (0.1º x 0.1º) 2 metres daily temperature taken from reanalysis products (ERA5-Land) freely available from the Coppernicus Climate Service through the Climate date Store.[13] The original data was an hourly gridded dataset over the extension of Brazil and encompassing the whole period of 10 years. We estimated the daily mean temperature for each municipality by calculating the daily temperature weighted to the municipality area. All these weighted mean areas were done with the 'exactextractr' R package.[14]

Data Analysis

We studied the association between daily dengue hospitalizations counts time-series and ambient daily temperature using the Distributed Lag Nonlinear Model (DLNM) framework.[15,16] We used the two-stage approach, first fitting generalised linear models at Brazilian State level. In the second stage we pooled the estimates using a meta-analysis at the country level and its five macro-regions.[17–19] At the first stage, we fitted a Conditional Poisson Model for the aggregated time series of Dengue hospitalisation counts.[20] The model strata was a month-city term[21]. We report the overall Relative Risk (RR) compared to the Minimum Hospitalisation Temperature (MHT) point for each state. MHT is the temperature with minimum hospitalisations according to the model. We also reported the lagged effects at the 50th and 95th percentiles. The second-stage was one meta-analysis for the whole country and another for each of the macro administrative regions. The meta-analysis for the whole country was conducted aggregating all the coefficients fitted by each state and their covariance matrix in a multivariate meta-analysis random effects

model. The meta-analysis for each macro-region used the coefficients and the covariance matrix of the states within a given region. From the meta-analysis, we estimated the overall cumulative RR curve, relative to MHT and the lag-effects at 50th and 95th percentile.

First-stage:

In the first-stage, we ran the following model for each state:

$$g\left[E\left(Y_{it}\right)\right]=\xi_{si}+f\left(x_{it},l\right)+\sum_{k=1}^{K} s_k\left(t\right)$$

Where Y_{it} is the daily count of hospitalisations on the *i-th* municipality, ξ_{si} the dow-month-municipality strata term conditioned out. $f(x_{it}, l)$ which is the bi-dimensional exposure-lag-response distributed lag non-linear model for mean temperature by each day of delay, until 21 days of lags. The cross-basis is parametrized with natural splines, with 2 knots equally spaced on the exposure-response structure and 3 knots equally spaced on the log transformed scale for the lag-response structure. The last term is the long-term trend model choice for temperature trend along the whole period, a natural spline with 7 degrees of freedom by each year on the whole period.

Second-stage:

Whole Country and Regions:

In the meta-analysis we did the aggregation of the data generated by the first-stage analysis on the whole country and by each of the five macro regions. We ran a multivariate meta-analysis with random effects, so from the i-*th* state-level estimate, θ_i, we have the following equation:

$$\theta_i \vee u_i \sim N_k\left(\theta_i+u_i, S_i\right), u_i \sim N_k\left(0, \psi\right)$$

is the estimate for the i-th location, u_i the random effects for the coefficients of this study with S_j intra-location covariance matrix of coefficient estimates, and ψ the between-location covariance matrix. For the whole country meta-analysis, the coefficients θ_i were taken from the 27 states,

while in the macro regions meta-analysis the coefficients θ_i were only taken from states pertaining to the macro region.

We ran two sensitivity analyses with different splines parametrizations for each of the exposure-response and lag-response structures. On the first sensitivity analysis, we parameterized the cross-basis with 3 knots (one extra knot compared to the main analysis) equally spaced on the exposure-response structure and again 3 knots equally spaced on the log transformed scale for the lag-response structure. In the second sensitivity analysis we parameterized the cross-basis with 2 knots equally spaced on the dose-response structure and 4 knots placed on the days 1, 2, 7, 14 for the lag-response structure.

All the analyses were run on R statistical Software, version 4.1.2.

Role of the funding source

The funders had no role in any decision about the manuscript. All authors had full access to all the data in the study and RLPS and OTR verified the data, and all authors approved the final version of the manuscript for publication.

RESULTS

Climate data description

A descriptive table (**Table S1**) of the mean daily temperature for each state, macro regions and the whole country over the ten years period can be found in the supplementary material. The overall temperature range across locations and years covered from -0·09º Celsius to 34·8º

Celsius. **Figure 1** gives the distribution of the mean daily temperature for each state.

Figure 1. Mean temperature density distribution over each state. The colours represent each macro administrative region.

Table 1. Characteristics of Dengue Hospitalisation cases

Characteristics Variables	Brazil, N = 579,703	North, N = 75,304	Northeast, N = 224,085	Center-West, N = 100,056	Southeast, N = 155,405	South, N = 24,853
Age, median [p25, p75], years	30 (15, 49)	27 (15, 44)	25 (12, 45)	36 (20, 53)	34 (17, 54)	37 (21, 55)
Age Categories, years						
0 to 1	8,752 (1.5%)	1,309 (1.7%)	4,457 (2.0%)	1,058 (1.1%)	1,726 (1.1%)	202 (0.8%)
1 to 9	73,507 (13%)	9,401 (12%)	37,354 (17%)	8,438 (8.4%)	16,620 (11%)	1,694 (6.8%)
10 to 17	91,786 (16%)	12,562 (17%)	41,991 (19%)	12,155 (12%)	22,171 (14%)	2,907 (12%)
18 to 39	192,819 (33%)	29,153 (39%)	72,769 (32%)	34,330 (34%)	48,009 (31%)	8,558 (34%)
40 to 59	126,059 (22%)	14,863 (20%)	39,640 (18%)	26,825 (27%)	38,268 (25%)	6,463 (26%)
60 to 79	71,395 (12%)	6,737 (8.9%)	22,314 (10.0%)	14,656 (15%)	23,494 (15%)	4,194 (17%)
80+	15,385 (2.7%)	1,279 (1.7%)	5,560 (2.5%)	2,594 (2.6%)	5,117 (3.3%)	835 (3.4%)
Self-Reported Race						
Black	12,467 (3.1%)	1,042 (2.3%)	3,946 (2.7%)	1,187 (1.9%)	5,760 (4.9%)	532 (2.6%)
Brown	244,869 (62%)	39,925 (87%)	120,476 (81%)	34,973 (55%)	45,113 (38%)	4,382 (21%)
Indigenous	1,092 (0.3%)	306 (0.7%)	116 (<0.1%)	595 (0.9%)	62 (<0.1%)	13 (<0.1%)
White	129,031 (33%)	3,893 (8.5%)	19,700 (13%)	24,396 (39%)	65,746 (55%)	15,296 (74%)
Yellow	9,165 (2.3%)	700 (1.5%)	4,003 (2.7%)	2,116 (3.3%)	1,991 (1.7%)	355 (1.7%)
(Missing)	183,079	29,438	75,844	36,789	36,733	4,275
Sex						
Female	310,308 (54%)	38,225 (51%)	121,175 (54%)	54,916 (55%)	82,550 (53%)	13,442 (54%)
Male	269,395 (46%)	37,079 (49%)	102,910 (46%)	45,140 (45%)	72,855 (47%)	11,411 (46%)
Hospital length of stay, median [p25-p75], days	3.00 (2.00, 4.00)	2.00 (2.00, 3.00)	3.00 (2.00, 4.00)	2.00 (2.00, 3.00)	3.00 (2.00, 4.00)	2.00 (2.00, 3.00)
Outcome						
Death	3,436 (0.6%)	281 (0.4%)	958 (0.4%)	548 (0.5%)	1,510 (1.0%)	139 (0.6%)

Cases data description:

Table 1 gives a characterization of the data summarised by each macro administrative region. In

the overall data the mean age for a hospitalisation by Dengue was 30 years old. There were

differences in mean age of hospitalisation by region, ranging from 25 years old in the Northeast

region to 37 years old in the South region. The sex ratio was representative of the sex ratio

estimates form the national statistics. The crude in-hospital mortality was in general low, with a

mortality rate of 0·6% of the total hospitalizations, ranging from 0·4% in the North and

Northeast regions to 1·0% in the Southeast region. **Figure S1** gives a visual description of the

time series for Dengue Hospitalisation for each region.

First-stage Results:

The cumulative overall lags RR curves to the MHT on each state level, for the whole period of

analysis, 2010 to 2019 are given in the supplementary material (**Figure S2**). The majority of the

states show an increasing risk of hospitalisation with high temperatures.

Second-stage Results:

Figure 2A and **3**, presents the cumulative overall RR accounting for the 21 days of lags, derived

from the meta-analysis of the association with temperature and hospitalisation, for the whole

country and for each macro-region. The overall curve (**Figure 2A**) shows that the risk of dengue

hospitalisation is minimal for the lowest temperature, and steadily increases up to more than a

40% increase at the highest temperatures. **Figure 2B** and 2**C**, present the lag effects for the 50th

(23·96º C) and 95th (28·68º C) percentiles of mean temperature distribution for Brazil, showing

that the association is mainly observed on the same day (lag 0), with a greater than 2-fold

increase in risk. The lag effects for each macro-region are shown on supplementary material,

Figure S3, S4, S5, S6 and S7. A summary of the RR and MHT for Brazil and each macro-region

is shown in **Table 2**.

Figure 2. A) Cumulative overall lags RR for the relationship between Dengue hospitalisation

and mean temperature, relative to the MHT. Vertical traced lines mark the 50th (Blue) and 95th

(Red) percentile of the temperature distribution. The grey shade represents the 95% confidence

interval derived from the meta-analysis. B) Lag effect to the RR to the MHT of Hospitalisation

due to Dengue on the 50th (23·96º C) percentile of temperature. C) Lag effect to the RR to the

MHT of hospitalisation due to Dengue on the 95th (28·68º C) percentile of temperature.

Figure 3. Cumulative overall lags RR for the relationship between Dengue hospitalisation and mean temperature, relative to the MHT. Each panel is the result of a meta-analysis model run over the estimates for all states that pertains to a given macro administrative region. Order by region is given by latitude extent.

Table 2. Dengue hospitalisation relative risk by Brazil and each macro-region			
Unit	MHT (ºC)	RR (CI 95%) 50th (23·96º C)	RR (CI 95%) 95th (28·68ºC)
Brazil	10·8	1·404 (1·274-1·548)	1·491 (1·337-1·664)
North	22·8	1·153 (1·026-1·295)	1·256 (1·039-1·518)
Northeast	19·7	1·196 (1·091-1·312)	1·255 (1·059-1·488)
Center-West	17	1·303 (1·093-1·553)	1·460 (1·198-1·779)
Southeast	13·7	1·353 (1·171-1·563)	1·498 (1·298-1·729)
South	7·4	2·723 (1·447-5·125)	2·508 (1·362-4·618)
* Relative Risk to the Minimum Hospitalisation Temperature of hospitalisation due Dengue			
** Minimum Hospitalisation Temperature, on Celsius degree			

Sensitivity analyses

Overall, the sensitivity analysis results were comparable with the main analysis. From the meta-analyses, on the 50th percentile for whole Brazil aggregation the RR was 1·629 (1·276-2·079) and 1·405 (1·274-1·549), for the first and second sensitivity analysis parametrization, respectively. On the 95th percentile they were 1·780 (1·354-2·339) and 1·495 (1·344-1·663), respectively. These results are shown on **Table S2** and **Figures S8** and **S9**, in the supplementary material.

Discussion:

We ran a two-stage time stratified design study to assess the association between hospitalisation by Dengue disease and ambient temperature. Our main findings are that there is a higher risk of being hospitalised by Dengue as the temperature gets higher. In a quantitative way, in general for Brazil, after 20 degrees Celsius, the risk is 1·2x higher than for the minimum temperature

observed, which coincides with the MHT. This is the cumulative effects of lags up to 21 days before the date of hospitalisation. We found a stronger immediate effect on the RR to the hospitalisation. We found the same patterns over each macro region of Brazil for the general association between RR hospitalisation and ambient temperature.

In comparison with the literature, we see that, in general, higher temperature is related to an augmented RR of mortality and hospitalisation by any causes[2,3,22,23]. There is scarce literature on the association between temperatures and hospitalisation by an infectious disease. Our study is the first, to our knowledge, to measure an association between temperature and hospitalisation by dengue directly.[23,24] We did not observe a typical U-shaped or J-shaped curves observed in the majority of studies looking for all-cause and cause-specific hospitalisations.[24,25] Some specific features of Dengue could explain this difference. First, its progression to severe disease is highly linked with factors that would be aggravated as the temperature increases. Second, the observed days with relatively low temperature are not frequent in Brazil and the incidence of the Dengue disease, of any severity, is much lower during cold months, therefore limiting the power to observe a potential effect of cold temperatures on Dengue hospitalisations. This reflects an additional challenge of studying climate variables and vector-borne infectious diseases severity, among others.[26]

Overall in the country and in each of the macro regions, the extreme heat had an effect of increasing the severity of cases, which evolved to a necessity of a hospitalisation. Quantitatively, the South region had the highest RR to its own MHT overall regions, with 1·5x higher risk on

the 95th percentile of its distribution of temperatures. This might be explained by the fact that the

South region has the lowest MHT and the wider range of temperature in comparison to all other

regions. The difference observed for the South could also reflect differences in the mosquito life

cycle, human behaviour and Dengue mitigation strategies[1,27,28]. A direct comparison between

regions is not straightforward because of regional disparities. Although Brazil has a universal

access health system (SUS), which has great disparities over geographical distribution[29]. Thus,

we might expect that regions with more hospital beds per population, such as South and

Southeast, would hospitalise milder cases compared with regions with fewer hospital beds. This

is also important during massive Dengue outbreaks, where the threshold for hospitalisation might

change according to beds availability.

Several mechanisms can play a role in the observed overall association and lag effects. The

immediate effect generating the first peak on lag-0 is expected and observed in the majority of

studies evaluating temperature and heat waves and hospitalizations and deaths.[23,25,30–33] It reflects

an immediate worsening of the clinical condition due to high temperature. This sharp increase on

lag-0 reinforces that our results is a direct effect of ambient temperature on hospitalisation risk,

rather than the indirect effect of temperature on mosquito activity and increased risk of dengue

incidence, association usually present in the scale of months[6]. Another mechanism is that Dengue

has a natural evolution of 1 to 3 days of incubation period, and more 5 to 7 days to complete

clearance. If an individual is exposed to higher temperatures during the viral infection evolution

this can lead to more severe infection, as seen a potential late lag-effect on some regions. Finally,

the decrease on the RR following the immediate effect could be due to a depletion of susceptible

("harvesting effect")[31,32]. This mechanism is reported for the lag between temperature and deaths,

but also has been hypothesised in another study evaluating temperature and other seasonal infectious diseases such as hand, foot and mouth disease[34]. If harvesting was the only explanation for the observed association, then the increase in hospitalizations observed on the hot day would just represent an advancement of a few days, because of heat, of hospitalizations that would have occurred anyway a few days later in the absence of heat. In that case, the overall RR cumulated over all lags will be close to 1. In our study, we observe important increases in the RR cumulated over three weeks, suggesting that these represent extra hospitalisations that would have not occurred in the absence of high temperatures. Interestingly, the South and Center-West regions present significant associations at later lags, see **Figures S5** and **S7**. When varying the points for the knots in a sensitivity analysis, lag-effects were less prominent, increasing the likelihood of collinearity explaining this finding.[35]

Our study has some strengths. We evaluated a nationwide database providing a 10 year time-series of Dengue hospitalizations in a LMIC exposed to a wide temperature range. Ambient temperature from reanalysis products from ERA5-Land are the state-of-the-art in climatology,[32,33] as the framework applied for data analysis, accounting for dose-response and lag-response structures and correlated daily data on temperature[21,36]. This has great generalizability to many other LMIC countries with great burden of Dengue and provides support to public health measures.

Our study has limitations to be mentioned. First, we did not evaluate other factors that could modify the high temperature effects, such as green space, urbanisation and relative humidity. In the same extension, we did not evaluate individual factors, such as age and sex, that could show different effects of temperature and risk of Dengue hospitalization in vulnerable populations.

Second, we obtained estimates for 27 states and pooled them for Brazil and each corresponding macro-regions. States from the same macro-region will have similar climate as well a similar Dengue incidence. This guarantees that they were likely comparing states with similar conditions for mosquito transmission as well hospitalisations conditions.[27] Although this analysis respected administrative divisions, it might not be generalizable to some municipalities. Finally, we considered hospitalisation due to Dengue as a surrogate of severe Dengue without evaluation of clinical severity using defined clinical criteria.

Conclusion

We found an increased risk of Dengue hospitalisation due high temperatures in Brazil for a ten years period, 2010-2019. This increased risk varies over the whole extension of Brazil and between macro-regions. With the findings presented here we hope the climate variability of Brazil can be taken into account when public health measures are designed to curb and mitigate the impacts of Dengue on the population.

Acknowledgements:

The authors thank Carles Milà for the advice on ERA5-Land data management. The authors also thank the research funding agencies: the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior -- Brazil (Finance Code 001 to RLPS), Conselho Nacional de Desenvolvimento Científico e Tecnológico -- Brazil (grant number: 141698/2018-7 to RLPS). OTR is funded by a Sara Borrell fellowship (CD19/00110) from the Instituto de Salud Carlos III. We acknowledge

support from the Spanish Ministry of Science and Innovation through the Centro de Excelencia Severo Ochoa 2019-2023 programme and from the Generalitat de Catalunya through the Centres de Recerca de Catalunya (CERCA) programme. The funding agencies had no role in the conceptualization of the study.

Role of the funding source

Contributors

RLPS, OTR and XB participated in the design and concept of the study. RLPS analysed the data. RLPS wrote the first version of the manuscript. XB, LSLB, FAB and OTR revised the manuscript with important intellectual contributions. XB and OTR supervised the study. All authors had full access to all data in the study, participated in data interpretation, revised the manuscript, and approved the final version of the manuscript for publication. RLPS and OTR verified the underlying data.

Declaration of interests

We declare no competing interests.

Data Availability Statement:

All data used in this study are publicly available. Hospitalisation data is available at

http://sihd.datasus.gov.br/principal/index.php, and temperature at

https://cds.climate.copernicus.eu/#!/home. The code to reproduce this analysis is available at:

https://github.com/rafalopespx/dengue_t2m_severity_paper

References:

Bhatt S, Gething PW, Brady OJ, *et al.* The global distribution and burden of dengue. *Nature* 2013; **496**: 504–7.
Martínez-Solanas È, Basagaña X. Temporal changes in the effects of ambient temperatures on hospital admissions in Spain. *PLOS ONE* 2019; **14**: e0218262.
Seah A, Aik J, Ng L-C, Tam CC. The effects of maximum ambient temperature and heatwaves on dengue infections in the tropical city-state of Singapore – A time series analysis. *Sci Total Environ* 2021; **775**: 145117.
Guzman MG, Harris E. Dengue. *The Lancet* 2015; **385**: 453–65.
Godói IP, Da Silva LVD, Sarker AR, *et al.* Economic and epidemiological impact of dengue illness over 16 years from a public health system perspective in Brazil to inform future health policies including the adoption of a dengue vaccine. *Expert Rev Vaccines* 2018; **17**: 1123–33.
Lowe R, Lee SA, O'Reilly KM, *et al.* Combined effects of hydrometeorological hazards and urbanisation on dengue risk in Brazil: a spatiotemporal modelling study. *Lancet Planet Health* 2021; **5**: e209–19.
Suaya JA, Shepard DS, Siqueira JB, *et al.* Cost of dengue cases in eight countries in the Americas and Asia: a prospective study. *Am J Trop Med Hyg* 2009; **80**: 846–55.
Burattini M, Lopez L, Coutinho F, *et al.* Age and regional differences in clinical presentation and risk of hospitalization for dengue in Brazil, 2000-2014. *Clinics* 2016; **71**: 455–63.
Werneck GL, Macias AE, Mascarenas C, *et al.* Comorbidities increase in-hospital mortality in dengue patients in Brazil. *Mem Inst Oswaldo Cruz* 2018; **113**. DOI:10.1590/0074-02760180082.
Brasil | Cities and States | IBGE. https://www.ibge.gov.br/en/cities-and-states.html (accessed Feb 2, 2022).
Alvares CA, Stape JL, Sentelhas PC, de Moraes Gonçalves JL, Sparovek G. Köppen's climate classification map for Brazil. *Meteorol Z* 2013; **22**: 711–28.
Coelho GE, Leal PL, Cerroni M de P, Simplicio ACR, Siqueira JB. Sensitivity of the Dengue Surveillance System in Brazil for Detecting Hospitalized Cases. *PLoS Negl Trop Dis* 2016; **10**: e0004705.
Muñoz-Sabater J, Dutra E, Agustí-Panareda A, *et al.* ERA5-Land: a state-of-the-art global reanalysis dataset for land applications. *Earth Syst Sci Data* 2021; **13**: 4349–83.
Fast Extraction from Raster Datasets using Polygons. https://isciences.gitlab.io/exactextractr/index.html (accessed Feb 2, 2022).
Gasparrini A, Armstrong B, Kenward MG. Distributed lag non-linear models. *Stat Med* 2010; **29**: 2224–34.
Gasparrini A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J Stat Softw* 2011; **43**: 1–20.
Gasparrini A, Armstrong B. Reducing and meta-analysing estimates from distributed lag non-linear models. *BMC Med Res Methodol* 2013; **13**: 1.
Jackson D, Riley R, White IR. Multivariate meta-analysis: Potential and promise. *Stat Med* 2011; **30**: 2481–98.

Gasparrini A, Armstrong B, Kenward MG. Multivariate meta-analysis for non-linear and other multi-parameter associations. *Stat Med* 2012; **31**: 3821–39.

Armstrong BG, Gasparrini A, Tobias A. Conditional Poisson models: a flexible alternative to conditional logistic case cross-over analysis. *BMC Med Res Methodol* 2014; **14**: 122.

Gasparrini A. The Case Time Series Design. *Epidemiology* 2021; **32**: 829–37.

Jacobson L da SV, Oliveira BFA de, Schneider R, Gasparrini A, Hacon S de S. Mortality Risk from Respiratory Diseases Due to Non-Optimal Temperature among Brazilian Elderlies. *Int J Environ Res Public Health* 2021; **18**: 5550.

Pudpong N, Hajat S. High temperature effects on out-patient visits and hospital admissions in Chiang Mai, Thailand. *Sci Total Environ* 2011; **409**: 5260–7.

Wang B, Chai G, Sha Y, Zha Q, Su Y, Gao Y. Impact of ambient temperature on cardiovascular disease hospital admissions in farmers in China's Western suburbs. *Sci Total Environ* 2021; **761**: 143254.

Vaidyanathan A, Saha S, Vicedo-Cabrera AM, *et al.* Assessment of extreme heat and hospitalizations to inform early warning systems. *Proc Natl Acad Sci* 2019; **116**: 5420–7.

Imai C, Armstrong B, Chalabi Z, Mangtani P, Hashizume M. Time series regression model for infectious disease and weather. *Environ Res* 2015; **142**: 319–27.

Lee SA, Economou T, de Castro Catão R, Barcellos C, Lowe R. The impact of climate suitability, urbanisation, and connectivity on the expansion of dengue in 21st century Brazil. *PLoS Negl Trop Dis* 2021; **15**: e0009773.

Brady OJ, Gething PW, Bhatt S, *et al.* Refining the Global Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. *PLoS Negl Trop Dis* 2012; **6**: e1760.

Noronha KVM de S, Guedes GR, Turra CM, *et al.* Pandemia por COVID-19 no Brasil: análise da demanda e da oferta de leitos hospitalares e equipamentos de ventilação assistida segundo diferentes cenários. *Cad Saúde Pública* 2020; **36**: e00115320.

Royé D, Íñiguez C, Tobías A. Comparison of temperature–mortality associations using observed weather station and reanalysis data in 52 Spanish cities. *Environ Res* 2020; **183**: 109237.

Schwartz J, Samet JM, Patz JA. Hospital admissions for heart disease: the effects of temperature and humidity. *Epidemiol Camb Mass* 2004; **15**: 755–61.

Saha MV, Davis RE, Hondula DM. Mortality Displacement as a Function of Heat Event Strength in 7 US Cities. *Am J Epidemiol* 2014; **179**: 467–74.

Morral-Puigmal C, Martínez-Solanas È, Villanueva CM, Basagaña X. Weather and gastrointestinal disease in Spain: A retrospective time series regression study. *Environ Int* 2018; **121**: 649–57.

Yi X, Chang Z, Zhao X, Ma Y, Liu F, Xiao X. The temporal characteristics of the lag-response relationship and related key time points between ambient temperature and hand, foot and mouth disease: A multicity study from mainland China. *Sci Total Environ* 2020; **749**: 141679.

Basagaña X, Barrera-Gómez J. Reflection on modern methods: visualizing the effects of collinearity in distributed lag models. *Int J Epidemiol* 2021; : dyab179.

de SE, Bundo M, Ragettli MS, *et al.* Nationwide Analysis of the Heat- and Cold-Related Mortality Trends in Switzerland between 1969 and 2017: The Role of Population Aging. *Environ Health Perspect*; **130**: 037001.