Ambient Temperature and Dengue Hospitalisation in Brazil:

A 10-year period two-stage case-design time series analysis

Rafael Lopes[1,2], Xavier Basagaña[2], Leonardo S. L. Bastos[3], Fernando A. Bozza[4], Otavio T. Ranzani[2,5]

1 - Instituto de Física Teórica - IFT, UNESP, São Paulo, Brazil
2 - Barcelona Institute for Global Health, Universitat Pompeu Fabra, CIBER Epidemiología y Salud Pública, Barcelona, Spain
3 - Department of Industrial Engineering (DEI), Pontifical Catholic University of Rio de Janeiro (PUC-Rio), Rio de Janeiro, RJ, Brazil
4 - National Institute of Infectious Disease Evandro Chagas (INI), Oswaldo Cruz Foundation (FIOCRUZ), Rio de Janeiro, RJ, Brazil
5 - Pulmonary Division, Heart Institute (InCor), HCFMUSP, Faculdade de Medicina da Universidade de São Paulo, Brazil

Corresponding author:

Otavio T. Ranzani (otavio.ranzani@isglobal.org)

ISGlobal, Dr. Aiguader, 88, 08003 Barcelona, Spain

Conflicts of interest: None.

Word count: 3,492

Keywords: Dengue; severe Dengue; temperature; heatwave; hospital; temperature

26   **Abstract (**270/300 words**)**
27
Climate factors are known to influence seasonal patterns of dengue transmission. However, little

is known about the effect of extremes of temperature on the severity of dengue infection, such as

hospital admission. We aimed to quantify the effect of ambient temperature on dengue

hospitalisation risk in Brazil. We retrieved daily dengue hospitalisation counts by each of 5,565

municipalities across the 27 states of Brazil from 1st of January 2010 to 31st of December 2019,

from the Brazilian Public Hospital Admission System ("SIH"). We obtained average daily

ambient temperature for each municipality from the ERA5-land product reanalysis. We

combined distributed lag non-linear models with time stratified design model framework to pool

a relative risk (RR) estimate for dose-response and lag-response structures for the association of

temperature and Dengue hospitalisation. We estimate the overall dengue hospitalisation RR for

the whole country as well for each of the five macro-regions by meta-analysing state level

estimates. 579,703 hospital admissions due to dengue occurred over the 10 years' period of 2010

to 2019. We observed a positive association between high temperatures and high risk of

hospitalisation for Brazil. The overall RR for dengue hospitalisation was 2.185 (95% CI 1.457-

3.276) at the 50th percentile of temperature and 2.385 (95% CI 1.556-3.655) at 95th percentile of

temperature for Brazil. We also found lag effects of heat on hospitalisation, particularly an

immediate augmented risk to hospitalisation, both at the 50th percentile and 95th. High

temperatures are associated with an increase in the risk of hospitalisation by dengue. These

findings may guide preparedness and mitigation policies during dengue season outbreaks,

particularly on the health-care demand.

48
49
50
51

1. **Introduction**

Dengue fever has been a major global seasonal endemic disease, present largely in the tropics,

with an estimated burden over ~390 million infections per year (Bhatt et al., 2013; Guzman and

Harris, 2015). Approximately one quarter of these infections manifest as clinical or subclinical

disease (Bhatt et al., 2013). Brazil is one of the most affected countries by dengue, that has

caused over ~20.9 millions cases of infections in the last 20 years, since the compulsory

universal notification to the Brazilian health systems (Godói et al., 2018; Lowe et al., 2021).

Additionally, it is estimated that the Brazilian Universal Health system (SUS) spent more than

USD 159 millions in the treatment and assistance to dengue cases and USD 10 million on severe

dengue between 2000 and 2015 (Godói et al., 2018). The impressive burden of dengue in Brazil

is also observed in several others low-income and middle-income countries (Suaya et al., 2009).

Dengue incidence is influenced by climate variables, particularly with temperature and rainfall,

due to mosquito life cycle and human behaviour changes, resulting in increased transmission risk

(Bhatt et al., 2013; Campbell et al., 2015; Colón-González et al., 2021; Lee et al., 2021; Lowe et

al., 2021; Suaya et al., 2009; Wibawa et al., 2024). A systematic review and meta-analysis

published in 2023 pooled 106 studies evaluating the association between high temperature and

heatwaves with dengue incidence (Damtew et al., 2023). There was sufficient evidence to show

increased relative risk for high temperatures (pooled RR 1.13 (95% CI, 1.11-1.16, for 1°C

increase in temperature), however limited evidence for heatwave events (pooled RR 1.08, 95%

CI, 0.95-1.23). The majority of articles used a monthly time resolution of the exposure, usually

exploring lag effects up to one year. The magnitude of the association reported seemed to depend

on the climate zone, showing a greater incidence risk in the tropical monsoon and humid

subtropical climate zones (Damtew et al., 2023). In this scenario, projections show that dengue

incidence, as well as other mosquito-borne diseases such as Zika, can have increased epidemic

potential with the global rising temperature due to climate change (Colón-González et al., 2021;

Van Wyk et al., 2023).

Dengue fever can present with increasing severity, requiring hospital admission and the support

of vital organs (Paz-Bailey et al., 2024). The main mechanisms involved in dengue severity

comprises dehydration and coagulation disorders (Burattini et al., 2016; Paz-Bailey et al., 2024;

Werneck et al., 2018), both conditions which can be aggravated by ambient temperature (Kenney

et al., 2014; Schneider et al., 2017). Although the literature is clear regarding the association

between temperature and dengue transmission, there is a lack of evidence regarding the

association between ambient temperature and dengue severity. Based on the reported association

of short-term extremes of temperature and all-cause hospitalizations (Martínez-Solanas and

Basagaña, 2019; Pudpong and Hajat, 2011; Vaidyanathan et al., 2019), we hypothesised that

short-term ambient temperature has a positive association with the risk of hospitalisation due to

dengue. We aimed to evaluate the association between ambient temperature and dengue

hospitalisation in Brazil, analysing the period from 2010 to 2019.

2. **Methods**

2.1 **Study design**

We conducted a two-stage case-design time series analysis to evaluate the association between

ambient temperature and dengue hospitalisation in Brazil. The unit of analysis was municipality

of residence.

2.2 **Study area**

Brazil is located in South America and has ~211 million inhabitants distributed over an area of
8.5 million km². The country is divided into 26 states and the country capital, a federal district
(Brazilian Institute of Geography and Statistics - IBGE, n.d.). These 27 states are grouped in 5
administrative macro-regions: North (7 states), Northeast (9 states), Center-West (4 states),
Southeast (4 states) and South (3 states). The country is located in a tropical region and has 3
main Köppen climate types and 12 subtypes (Alvares et al., 2013).

2.3 **Outcome and covariates**

Our outcome is a dengue hospitalization event. We used the Brazilian Hospital Admission
System (SIH), a nationwide database that comprises individual level data of all hospitalizations
covered by the Universal Healthcare System (SUS) in Brazil. We defined a dengue
hospitalisation by the following ICD-10 codes: 'A90', 'A91', 'A97', 'A970', 'A971', 'A972', 'A979'
(Coelho et al., 2016). We build daily time series aggregating the number of events by date of
hospitalisation for each municipality of residence.

We used the number of dengue cases as a covariate. We used the National Dengue Surveillance
System (SINAN-Dengue), that receives any notification for a suspected or confirmed dengue
case. We selected the confirmed cases of dengue and build daily times times series aggregating
the number of cases by date of symptoms onset at each municipality. To add this covariate in the
model in a sensitivity analysis, we derived the 7-days moving average of the dengue cases series,
an estimated time of progression from symptoms to hospitalization.

Both databases are publicly accessible and following ethically agreed principles on open data,

the use of this data did not require ethical approval in Brazil, according to the Brazilian Ethics

Resolution n° 510/2016.

2.4 **Temperature exposure assessment**

Our exposure of interest is the daily average temperature at each Brazilian municipality. To

derive it, we used the hourly 2-metres temperature (gridded 0.1º x 0.1º) from reanalysis products

(ERA5-Land), freely available by the Copernicus Climate Service through the Climate date Store

(Muñoz-Sabater et al., 2021). We estimated the daily mean temperature for each municipality by

calculating the mean daily temperature of each grid cell and weighting it to the municipality area.

All these weighted mean areas were done with the 'exactextractr' R package (Baston, Daniel,

2023).

Regarding the performance of the temperature estimated from ERA5-Land reanalysis in Brazil,

one study evaluated the agreement between ERA5-Land and monitoring stations for monthly

temperature averages, using 12 automatic stations for the period 2011-2020 from one Brazilian

state in the Northeast region (Pernambuco state - PE), with an average R-square of 0.92 (Araújo

et al., 2022). For this study, we performed a validation comparing the daily average of

temperature from monitoring stations at the municipal level with the ERA5-Land estimates

described before. We used data from 389 stations (269 automatic and 120 manual) from the

National Institute of Meteorology (National Institute of Meteorology (INMET), 2024), with at

least 90% of days with complete data during the period. The analysis of 340 municipalities

covering the 27 states and approximately 65 million individuals showed a Pearson correlation

coefficient of 0.94, R-square of 0.90 and RMSE of 1.54°C. Further information is available in

the supplementary material.

2.5 **Data Analysis**

We used a two-stage approach in the times-series analysis. At the first stage, we fitted

Conditional Poisson Models for each Brazilian state (Armstrong et al., 2014; Gasparrini, 2021).

At the second stage, we pooled the 27 estimates using a multivariate meta-analysis (Gasparrini et

al., 2012; Gasparrini and Armstrong, 2013; Jackson et al., 2011).

2.5.1 **First-stage analysis**

At the first-stage, we run the following model for each state:

$g[E(Y_{ijt})] = \xi_{ijs} + f(x_{tij}, l, \theta_j) + \sum_{j=i}^{J} s_k(t; \gamma_j)$

Where Y_{ijt} is the daily count of cases by the *j-th* state on the *i-th* municipality, ξ_{ijs} the month-

municipality strata term conditioned out, $f(x_{tij}, l, \theta_j)$ is the bi-dimensional exposure-lag-response

distributed lag non-linear cross-basis for the mean temperature on the *j-th* state by each day of

delay, until 21 days of lags (Gasparrini et al., 2015; Martínez-Solanas and Basagaña, 2019). The

cross-basis is parametrized with natural splines, with 2 knots equally spaced on the dose-

response structure and 3 knots equally spaced on the log transformed scale for the lag-response

structure. The last term is the long-term trend model choice for temperature trend along the

whole period, a natural spline with 7 degrees of freedom for each year over the whole period.

2.5.2. **Second-stage analysis**

We ran a multivariate meta-analysis with random effects, so from the *j-th* state-level study, θ_j, we

have the following equation:

172    $$\theta_i | u_i \sim N_k(\theta_i + u_i, S_i), u_i \sim N_k(0, \psi)$$

Where θ is the study to estimate, with u_i the random effects for the coefficients of this study

together with S_j covariance matrix of intra state studies and ψ covariance matrix of between

states studies. We run one meta-analysis for the whole country and another for each of the macro

administrative regions separately. For the whole country meta-analysis, the θ studies were taken

from all the 27 states, and for the macro regions meta-analyses, θ was only taken from states

pertaining to the same macro region.

2.5.3 **Sensitivity analyses**

We ran three sensitivity analyses to explore the DLNM parametrization and to account for the

number of dengue cases. On the first sensitivity analysis, we parameterized the cross-basis with 3

knots equally spaced on the dose-response structure and, as in the main analysis, 3 knots equally

spaced on the log transformed scale for the lag-response structure. The second sensitivity

analysis we parameterized the cross-basis with 2 knots equally spaced on the dose-response

structure as the main analysis, however with 4 knots placed on the days 1, 2, 7, 14 for the lag-

response structure. On the third sensitivity analysis we adjusted the first stage models adding the

7-days moving average of the times series of confirmed dengue cases, considering the average

time between incubation period and time to hospitalization.

2.5.4 **Reporting**

We report the overall cumulative relative risk (RR) compared to the Minimum Hospitalisation

Temperature (MHT) point. MHT is defined as the temperature which has an overall RR equal to

1. We also report RR for the lag effects up to 21 days at the 50th and 95th percentiles of the

temperature distribution.

All the analyses were run on R Software, version 4.1.2.

3. **Results**

3.1 **Climate data description**

A descriptive table (**Table S1**) of the mean daily temperature for each state, macro regions and

the whole country over the ten years period is shown in the supplementary material. For the

whole period of ten years, it varied from -0.09 ºC to 34.8 ºC. **Figure 1** gives the daily mean

temperature distribution across all states.

Figure 1. **Mean temperature density distribution over each state**. On y-axis is each of the

state acronyms and on x-axis is the range of mean temperature in degree Celsius. The height of

each curve is the cumulative density of mean temperature at that specific degree. The colours are

given by each macro administrative region. The inset map gives the location of each of the state

code at Brazilian geography.

203   **Table 1. Characteristics description of Dengue Hospitalisation data**

Variables	N	Brazil (N=579,703)	North (N=75,304)	Northeast (N=224,085)	Center-West (N=100,056)	Southeast (N=155,405)	South (N=24,853)
Mean Age (IQR), years	579,703	30 (15, 49)	27 (15, 44)	25 (12, 45)	36 (20, 53)	34 (17, 54)	37 (21, 55)
Age Categories, n (%) years	579,703						
0 to 1		8,752 (1.5%)	1,309 (1.7%)	4,457 (2.0%)	1,058 (1.1%)	1,726 (1.1%)	202 (0.8%)
1 to 9		73,507 (13%)	9,401 (12%)	37,354 (17%)	8,438 (8.4%)	16,620 (11%)	1,694 (6.8%)
10 to 17		91,786 (16%)	12,562 (17%)	41,991 (19%)	12,155 (12%)	22,171 (14%)	2,907 (12%)
18 to 39		192,819 (33%)	29,153 (39%)	72,769 (32%)	34,330 (34%)	48,009 (31%)	8,558 (34%)
40 to 59		126,059 (22%)	14,863 (20%)	39,640 (18%)	26,825 (27%)	38,268 (25%)	6,463 (26%)
60 to 79		71,395 (12%)	6,737 (8.9%)	22,314 (10.0%)	14,656 (15%)	23,494 (15%)	4,194 (17%)
80+		15,385 (2.7%)	1,279 (1.7%)	5,560 (2.5%)	2,594 (2.6%)	5,117 (3.3%)	835 (3.4%)
Self-Reported Race, n (%)	396,624						
Black		12,467 (3.1%)	1,042 (2.3%)	3,946 (2.7%)	1,187 (1.9%)	5,760 (4.9%)	532 (2.6%)
Pardo		244,869 (62%)	39,925 (87%)	120,476 (81%)	34,973 (55%)	45,113 (38%)	4,382 (21%)
Indigenous		1,092 (0.3%)	306 (0.7%)	116 (<0.1%)	595 (0.9%)	62 (<0.1%)	13 (<0.1%)
White		129,031 (33%)	3,893 (8.5%)	19,700 (13%)	24,396 (39%)	65,746 (55%)	15,296 (74%)
Asian		9,165 (2.3%)	700 (1.5%)	4,003 (2.7%)	2,116 (3.3%)	1,991 (1.7%)	355 (1.7%)
(Missing)		183,079	29,438	75,844	36,789	36,733	4,275
Sex, n (%)	579,703						
Female		310,308 (54%)	38,225 (51%)	121,175 (54%)	54,916 (55%)	82,550 (53%)	13,442 (54%)
Male		269,395 (46%)	37,079 (49%)	102,910 (46%)	45,140 (45%)	72,855 (47%)	11,411 (46%)
Duration of Hospitalisation, median (IQR), days	579,703	3 (2, 4)	2 (2, 3)	3 (2, 4)	2 (2, 3)	3 (2, 4)	2 (2, 3)
In-hospital mortality, n (%)	579,703	3,436 (0.6%)	281 (0.4%)	958 (0.4%)	548 (0.5%)	1,510 (1.0%)	139 (0.6%)

3.2 **Dengue hospitalization data description**

We excluded dengue hospitalizations from five small municipalities (e.g, small islands), because

there was no ambient temperature for them. This filtering resulted in a loss of 153 (0.03%)

hospitalized cases, for the whole period of the 10 years. The final time series aggregation by

municipalities encompasses a total of 579,703 hospital admissions due to dengue over the 10

years' period of 2010 to 2019 in 5,565 municipalities.

Table 1 gives a characterization of the data summarised by each macro administrative region.

Overall, the mean age for a hospitalisation by dengue was 30 years old. Between regions it can

be seen a difference of 12 years old to the mean age of hospitalisation, being the mean age of

hospitalisation of 37 years old for the South region and 25 years old for the Northeast region.

The sex ratio is typically representative of the sex ratio estimates from the national statistics. The

crude in-hospital mortality in general is low, with mortality rate of 0.6% of the total

hospitalizations, and there are some discrepancies between regions. The Southeast has a rate of

1.0% of the hospitalizations with the highest death outcome rate. The North and Northeast with

lowest rate of mortality with an average of 0.4% rate of deaths to the total hospitalizations.

Figure S1 gives a visual description of the time series for dengue hospitalisation for each region.

3.4 **First-stage Results**

The cumulative RR over all lags compared to the MHT on each state level, for the whole period

of analysis, 2010 to 2019 are given in the supplementary material (**Figure S2**).

3.5 **Second-stage Results**

Figure 2 and **3**, presents the cumulative RR over all lags compared to the MHT derived from the

meta-analysis, for the whole country and for each macro-region. **Figure 2B** and **2C**, presents the

lag effects for the 50th (23.96º C) and 95th (28.68º C) percentiles of mean temperature

distribution for Brazil. The lag effects for each macro-region is shown on supplementary

material, **Figure S3, S4, S5, S6** and **S7**. A summary of the RR and MHT for Brazil and each

macro-region is shown in **Table 2**.

Figure 2. **Pooled cumulative relative risk for the whole country meta-analysis.** A)

Cumulative RR over all lags for a Dengue hospitalisation relative to the MHT. Vertical traced

lines mark the 50th (Blue) and 95th (Red) percentile of the temperature distribution. The grey

shade ribbon is 95% confidence interval derived from the meta-analysis. B) Lag effect to the RR

to the MHT of Hospitalisation due to Dengue at the 50th (23.96º C) percentile of temperature. C)

Lag effect to the RR to the MHT of hospitalisation due to Dengue on the 95th (28.68º C)

percentile of temperature. MHT: Minimum Hospitalisation Temperature; RR: relative risk.

Figure 3. Pooled cumulate relative risk to each of the macro region meta-analysis.

Cumulative overall lags RR to the MHT curves to temperature by region. Each panel is given by

a meta-analysis model run over all the states coefficients and covariance matrices that pertains to

a given macro administrative region. Order by region is given by latitude extent. MHT:

Minimum Hospitalisation Temperature; RR: relative risk.

253   Table 2. **Dengue hospitalisation relative risk (RR) over Brazil and each macro-region**

Unit	MHT (ºC)	RR (IC 95%) 50th (23·96º C)	RR (IC 95%) 95th (28·68º C)
Brazil	10.8	2.185 (1.457-3.276)	2.385 (1.556-3.655)
North	30.8	1.139 (0.707-1.836)	1.048 (0.811-1.355)
Northeast	19.7	1.660 (1.116-2.468)	1.547 (0.860-2.784)
Center-West	17	2.393 (1.394-4.110)	2.420 (1.365-4.289)
Southeast	13.7	3.335 (1.605-6.929)	4.930 (2.567-9.468)
South	16	1.119 (0.580-2.159)	2.253 (0.592-8.569)

RR: Relative Risk to the Minimum Hospitalisation Temperature of hospitalisation due Dengue

MHT: Minimum Hospitalisation Temperature, on Celsius degree

3.6 **Sensitivity analyses**

From the meta-analyses, on the 50th percentile for whole Brazil the RR was 2.080 (95% CI

1.210-3.578) and 2.315 (95% CI 1.558-3.439), for the first and second sensitivity analyses

parametrizations, respectively. After adjusting for the number of dengue cases (third sensitivity

analysis), the RR on the 50th percentile for whole Brazil was 1.991 (95% CI 1.382-2.867).

Overall, the sensitivity analyses result for each macro-region were comparable with the main

analysis at the 50th percentile (**Figure 4**) and at the 95th percentile (Figure S8), except for the

South region. There was an increased risk for dengue hospitalization in the South in the three

sensitivity analyses, particularly for the 95th percentile, although with high imprecision (RR

4.830, 95% CI, 1.34617.331). Additional results are shown on **Table S2** and **Figures S9**, **S10,**

S11, and **S12**, in the supplementary material.

Figure 4. Dengue hospitalization relative risk by Brazil and each macro-region: main and
sensitivity analyses at 50th percentile of temperature

270
*MHT: Minimum Hospitalisation Temperature, on Celsius degree; RR: relative risk

4. **Discussion**

We ran a two-stage time stratified design study to the association between ambient temperature

and hospitalisation due to Dengue. Our main finding is an increased risk of being hospitalised by

Dengue as the temperature increases. In a quantitative way, in general for Brazil, above 20

degrees Celsius, the RR to the MHT is 1.2x higher. This is the cumulative effects of lags up to 21

days before the date of hospitalisation. We found a stronger immediate effect on the RR to the

hospitalisation. We found the same patterns for each macro region of Brazil, but the North and

South regions.

In comparison with the literature, we see that, in general, higher temperature is related to an

augmented RR of all-cause and cause-specific hospitalisation and mortality (Jacobson et al.,

2021; Martínez-Solanas and Basagaña, 2019; Pudpong and Hajat, 2011; Silveira et al., 2019;

Wang et al., 2021; Zhao et al., 2019). There is scarce literature on the association between

temperatures and hospitalisation by an infectious disease. Our study is the first, to our

knowledge, to measure an association between temperature and hospitalisation caused by

dengue. Interestingly, we did not observe a typical U-shape curve increase on the risk for

hospitalization, as commonly observed for studies looking for all-cause and cause-specific

hospitalisations, except for the South region, where the association was unclear. The pattern we

observed has been reported for non-infectious causes in the US and China (Vaidyanathan et al.,

2019; Wang et al., 2021). Some specific features of dengue could explain this difference. First,

its progression to severe disease is highly linked with factors that would be aggravated as the

temperature increases, such dehydration and capillary leak syndrome. Second, the incidence of

the dengue disease, of any severity, is much lower during cold months, therefore limiting the

power to observe a potential effect of cold temperatures on Dengue hospitalizations. This reflects

an additional challenge when studying climate variables and vector-borne infectious diseases

severity, among others (Imai et al., 2015).

Overall, in the country and each of the macro regions, the extreme heat (above 20 Celsius

degrees) has an effect of increasing the severity of cases, which evolves to a necessity of a

hospitalisation. The association for the North and South regions are unclear and a direct

comparison between regions is not straightforward because of regional disparities as well as

climate disparities. Although Brazil has a universal access health system, SUS has great

disparities over geographical distribution (Noronha et al., 2020). Thus, we might expect that

regions with more hospital beds per population, such as South and Southeast, would hospitalise

milder cases compared with regions with fewer hospital beds. This is also important during

massive dengue outbreaks, where the threshold for hospitalisation might change according to

beds availability and age distribution of cases.

Several mechanisms can play a role in the observed overall association and lag effects. The

immediate effect generating the first peak on lag-0 is expected and observed in the majority of

studies evaluating temperature and heat waves and hospitalizations and deaths (Morral-Puigmal

et al., 2018; Pudpong and Hajat, 2011; Royé et al., 2020; Saha et al., 2014; Schwartz et al., 2004;

Vaidyanathan et al., 2019). It reflects an immediate worsening of the clinical condition due to

high temperature over the course of an infection. This sharp increase on lag-0 reinforces that our

results is a direct effect of ambient temperature on hospitalisation risk, rather than the indirect

effect of temperature on mosquito activity which increases risk of dengue infection, association

usually present in the scale of months (Lowe et al., 2021). Even though, the sensitivity analysis

when adding the 7-day rolling mean of mild cases as a covariate in the model shows that the

effect of temperature on our model is reduced from 2.185 to 1.991 at the 50[th] and from 2.385 to

2.221 at the 95[th] percentile of temperature (**Table S2**). This sensitivity analysis suggests, as

before, that the augmented risk for hospitalization by dengue with increased temperature it is a

direct effect of heat on individuals already infected by dengue. This finding shows that high

temperature seems have a short- and a long-term effect on dengue burden, being the first

particularly on severity and the second on infection.

We hypothesize that the natural evolution of 1 to 3 days of incubation period from an infection,

plus 5 to 7 days to the complete clearance of the viral infection, can be affected by heat exposure

based on the lagged effects observed in our results. If an individual is exposed to higher

temperatures during the viral infection evolution this can lead to more severe infection, as seen

on lagged-effect around 5 to 7 days before hospitalization. Additionally, the RR decrease right

after the lag0-1 effect could be explained by depletion of susceptibles ("harvesting effect") (Saha

et al., 2014; Schwartz et al., 2004). This mechanism is reported for the lag between temperature

and deaths, but also has been hypothesised in another study evaluating temperature and other

seasonal infectious diseases such as hand, foot and mouth disease (Yi et al., 2020). Importantly,

the lag-effect varies over regions, for instance, the South region has a greater delayed lag-effect

than other regions. However, when varying the knots placement in a sensitivity analysis, lag-

effects were less prominent, increasing the likelihood of collinearity explaining this finding

(Basagaña and Barrera-Gómez, 2022).

Our study has some strengths. We evaluated a nationwide database providing a 10-year time-

series of dengue hospitalizations in a LMIC exposed to a wide temperature range. We used

ambient temperature from reanalysis products from ERA5-Land (Royé et al., 2020), which might

increase the generalizability of our results to many other LMIC countries as countries in the

regions which dengue has been expanding during last years. Finally, we used the DLNM

approach, accounting for dose-response and lag-response structures and correlated daily data on

temperature (De Schrijver et al., 2022; Gasparrini, 2021).

Our study has limitations to be mentioned. We did not evaluate other factors that could modify

the high temperature effects, such as green space, urbanisation and relative humidity . In the

same extension, we did not evaluate individual factors, such as age and sex, that could show

different effects of temperature and risk of dengue hospitalization in vulnerable populations.

Second, we obtained estimates for 27 states and pooled them for Brazil and each corresponding

macro-regions. States from the same macro-region will have similar climate as well as a similar

dengue incidence. This choice of aggregation allows to comparing states with probably similar

transmission conditions to the mosquito as well hospitalizations conditions (Lee et al., 2021).

However, this may introduce bias to the analysis and do not allow for generalizations to some

municipalities within each state or to border line areas. Finally, we might have exposure

misclassification by using the ERA-5-land reanalysis, particularly for the North region based on

our validation. Nevertheless, the use of daily average temperature from ERA5-Land reanalysis

has been applied in several epidemiological analysis (Alahmad et al., 2023; Kephart et al., 2022),

with two studies showing minimal differences in estimates when comparing the exposure from

ERA5-Land reanalysis with the observed weather station data for mortality (Mistry et al., 2022;

Royé et al., 2020). One of these studies included 18 municipalities from Brazil in the period

1997-2011, with comparable estimates of excess mortality due to cold (2.83, 95% CI, 2.29-3.38

for station versus 2.90, 95% CI, 2.07-3.69 for ERA5-Land) and to heat (0.73, 95% CI, 0.47-0.99

for station versus 0.70, 95% CI, -0.14-1.50 for ERA5-Land) (Mistry et al., 2022). Thus, we did

not expect meaningful impact on the results due to the potential exposure misclassification.

5. **Conclusion**

In conclusion, we observed an association between high temperature and increased risk of

dengue hospitalization in Brazil. That association is mainly driven by an immediate effect of

heat, and varied over the five Brazilian macro-regions, being unclear at the North and South

regions. This study adds to the gap in knowledge showing a short-term effect of temperature on

dengue severity, in addition to the well-known association between long-term temperature and

dengue infection reported in the literature.

Acknowledgements:

The authors also thank the research funding agencies: the Coordenação de Aperfeiçoamento de

Pessoal de Nível Superior -- Brazil (Finance Code 001 to RLPS), Conselho Nacional de

Desenvolvimento Científico e Tecnológico -- Brazil (grant number: 141698/2018-7 to RLPS).

OTR is funded by a Sara Borrell fellowship (CD19/00110) from the Instituto de Salud Carlos III.

We acknowledge support from the grant CEX2018-000806-S funded by MCIN/AEI/

10.13039/501100011033, and support from the Generalitat de Catalunya through the CERCA

Program. The funding agencies had no role in the conceptualization of the study.

Data Availability Statement:

All data used in this study are publicly available. Hospitalisation data is available at

http://sihd.datasus.gov.br/principal/index.php, and temperature at

https://cds.climate.copernicus.eu/#!/home. The code to reproduce this analysis is available at:

https://github.com/rafalopespx/dengue_t2m_severity_paper

Author Contribution Statement:

Rafael Lopes: Conceptualization, Data curation, Formal analysis, Funding acquisition,
Methodology, Software, Visualization, Writing – original draft

Xavier Basagaña: Methodology, Supervision, Validation, Writing - review & editing

Leonardo S. L. Bastos: Data curation, Writing - review & editing

Fernando A. Bozza: Writing - review & editing

Otavio T. Ranzani: Conceptualization, Data curation, Formal analysis, Funding acquisition,
Methodology, Supervision, Validation, Writing - review & editing

References

Alahmad, B., Khraishah, H., Royé, D., Vicedo-Cabrera, A.M., Guo, Y., Papatheodorou, S.I., Achilleos, S., Acquaotta, F., Armstrong, B., Bell, M.L., Pan, S.-C., De Sousa Zanotti Stagliorio Coelho, M., Colistro, V., Dang, T.N., Van Dung, D., De' Donato, F.K., Entezari, A., Guo, Y.-L.L., Hashizume, M., Honda, Y., Indermitte, E., Íñiguez, C., Jaakkola, J.J.K., Kim, H., Lavigne, E., Lee, W., Li, S., Madureira, J., Mayvaneh, F., Orru, H., Overcenco, A., Ragettli, M.S., Ryti, N.R.I., Saldiva, P.H.N., Scovronick, N., Seposo, X., Sera, F., Silva, S.P., Stafoggia, M., Tobias, A., Garshick, E., Bernstein, A.S., Zanobetti, A., Schwartz, J., Gasparrini, A., Koutrakis, P., 2023. Associations Between Extreme Temperatures and Cardiovascular Cause-Specific Mortality: Results From 27 Countries. Circulation 147, 35–46. https://doi.org/10.1161/CIRCULATIONAHA.122.061832

Alvares, C.A., Stape, J.L., Sentelhas, P.C., De Moraes Gonçalves, J.L., Sparovek, G., 2013. Köppen's climate classification map for Brazil. metz 22, 711–728. https://doi.org/10.1127/0941-2948/2013/0507

Araújo, C.S.P.D., Silva, I.A.C.E., Ippolito, M., Almeida, C.D.G.C.D., 2022. Evaluation of air temperature estimated by ERA5-Land reanalysis using surface data in Pernambuco, Brazil. Environ Monit Assess 194, 381. https://doi.org/10.1007/s10661-022-10047-2

Armstrong, B.G., Gasparrini, A., Tobias, A., 2014. Conditional Poisson models: a flexible alternative to conditional logistic case cross-over analysis. BMC Med Res Methodol 14, 122. https://doi.org/10.1186/1471-2288-14-122

Basagaña, X., Barrera-Gómez, J., 2022. Reflection on modern methods: visualizing the effects of collinearity in distributed lag models. International Journal of Epidemiology 51, 334–344. https://doi.org/10.1093/ije/dyab179

Baston, Daniel, 2023. exactextractr: Fast Extraction from Raster Datasets using Polygons. R package.

Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., Drake, J.M., Brownstein, J.S., Hoen, A.G., Sankoh, O., Myers, M.F., George, D.B., Jaenisch, T., Wint, G.R.W., Simmons, C.P., Scott, T.W., Farrar, J.J., Hay, S.I., 2013. The global distribution and burden of dengue. Nature 496, 504–507. https://doi.org/10.1038/nature12060

Brazilian Institute of Geography and Statistics - IBGE, n.d. 2010 Population Census. [WWW Document]. URL https://www.ibge.gov.br/en/statistics/social/income-expenditure-and-consumption/18391-2010-population-census.html?edicao=19720&t=sobre (accessed 4.27.21).

Burattini, M.N., Lopez, L.F., Coutinho, F.A.B., Siqueira, J.B., Homsani, S., Sarti, E., Massad, E., 2016. Age and regional differences in clinical presentation and risk of hospitalization for dengue in Brazil, 2000-2014. Clinics 71, 455–463. https://doi.org/10.6061/clinics/2016(08)08

Campbell, K.M., Haldeman, K., Lehnig, C., Munayco, C.V., Halsey, E.S., Laguna-Torres, V.A., Yagui, M., Morrison, A.C., Lin, C.-D., Scott, T.W., 2015. Weather Regulates Location, Timing, and Intensity of Dengue Virus Transmission between Humans and Mosquitoes. PLoS Negl Trop Dis 9, e0003957. https://doi.org/10.1371/journal.pntd.0003957

Coelho, G.E., Leal, P.L., Cerroni, M.D.P., Simplicio, A.C.R., Siqueira, J.B., 2016. Sensitivity of the Dengue Surveillance System in Brazil for Detecting Hospitalized Cases. PLoS Negl Trop Dis 10, e0004705. https://doi.org/10.1371/journal.pntd.0004705

Colón-González, F.J., Sewe, M.O., Tompkins, A.M., Sjödin, H., Casallas, A., Rocklöv, J.,
Caminade, C., Lowe, R., 2021. Projecting the risk of mosquito-borne diseases in a
warmer and more populated world: a multi-model, multi-scenario intercomparison
modelling study. The Lancet Planetary Health 5, e404–e414.
https://doi.org/10.1016/S2542-5196(21)00132-7
Damtew, Y.T., Tong, M., Varghese, B.M., Anikeeva, O., Hansen, A., Dear, K., Zhang, Y.,
Morgan, G., Driscoll, T., Capon, T., Bi, P., 2023. Effects of high temperatures and
heatwaves on dengue fever: a systematic review and meta-analysis. eBioMedicine 91,
104582. https://doi.org/10.1016/j.ebiom.2023.104582
De Schrijver, E., Bundo, M., Ragettli, M.S., Sera, F., Gasparrini, A., Franco, O.H., Vicedo-
Cabrera, A.M., 2022. Nationwide Analysis of the Heat- and Cold-Related Mortality
Trends in Switzerland between 1969 and 2017: The Role of Population Aging. Environ
Health Perspect 130, 037001. https://doi.org/10.1289/EHP9835
Gasparrini, A., 2021. The Case Time Series Design. Epidemiology 32, 829–837.
https://doi.org/10.1097/EDE.0000000000001410
Gasparrini, A., Armstrong, B., 2013. Reducing and meta-analysing estimates from distributed lag
non-linear models. BMC Med Res Methodol 13, 1. https://doi.org/10.1186/1471-2288-
13-1
Gasparrini, A., Armstrong, B., Kenward, M.G., 2012. Multivariate meta-analysis for non-linear
and other multi-parameter associations. Stat Med 31, 3821–3839.
https://doi.org/10.1002/sim.5471
Gasparrini, A., Guo, Y., Hashizume, M., Lavigne, E., Zanobetti, A., Schwartz, J., Tobias, A.,
Tong, S., Rocklöv, J., Forsberg, B., Leone, M., De Sario, M., Bell, M.L., Guo, Y.-L.L.,
Wu, C., Kan, H., Yi, S.-M., De Sousa Zanotti Stagliorio Coelho, M., Saldiva, P.H.N.,
Honda, Y., Kim, H., Armstrong, B., 2015. Mortality risk attributable to high and low
ambient temperature: a multicountry observational study. The Lancet 386, 369–375.
https://doi.org/10.1016/S0140-6736(14)62114-0
Godói, I.P., Da Silva, L.V.D., Sarker, A.R., Megiddo, I., Morton, A., Godman, B., Alvarez-
Madrazo, S., Bennie, M., Guerra-Junior, A.A., 2018. Economic and epidemiological
impact of dengue illness over 16 years from a public health system perspective in Brazil
to inform future health policies including the adoption of a dengue vaccine. Expert
Review of Vaccines 17, 1123–1133. https://doi.org/10.1080/14760584.2018.1546581
Guzman, M.G., Harris, E., 2015. Dengue. The Lancet 385, 453–465.
https://doi.org/10.1016/S0140-6736(14)60572-9
Imai, C., Armstrong, B., Chalabi, Z., Mangtani, P., Hashizume, M., 2015. Time series regression
model for infectious disease and weather. Environmental Research 142, 319–327.
https://doi.org/10.1016/j.envres.2015.06.040
Jackson, D., Riley, R., White, I.R., 2011. Multivariate meta-analysis: potential and promise. Stat
Med 30, 2481–2498. https://doi.org/10.1002/sim.4172
Jacobson, L.D.S.V., Oliveira, B.F.A.D., Schneider, R., Gasparrini, A., Hacon, S.D.S., 2021.
Mortality Risk from Respiratory Diseases Due to Non-Optimal Temperature among
Brazilian Elderlies. IJERPH 18, 5550. https://doi.org/10.3390/ijerph18115550
Kenney, W.L., Craighead, D.H., Alexander, L.M., 2014. Heat Waves, Aging, and Human
Cardiovascular Health. Medicine & Science in Sports & Exercise 46, 1891–1899.
https://doi.org/10.1249/MSS.0000000000000325
Kephart, J.L., Sánchez, B.N., Moore, J., Schinasi, L.H., Bakhtsiyarava, M., Ju, Y., Gouveia, N.,

Caiaffa, W.T., Dronova, I., Arunachalam, S., Diez Roux, A.V., Rodríguez, D.A., 2022.
City-level impact of extreme temperatures and mortality in Latin America. Nat Med 28,
1700–1705. https://doi.org/10.1038/s41591-022-01872-6
Lee, S.A., Economou, T., De Castro Catão, R., Barcellos, C., Lowe, R., 2021. The impact of
climate suitability, urbanisation, and connectivity on the expansion of dengue in 21st
century Brazil. PLoS Negl Trop Dis 15, e0009773.
https://doi.org/10.1371/journal.pntd.0009773
Lowe, R., Lee, S.A., O'Reilly, K.M., Brady, O.J., Bastos, L., Carrasco-Escobar, G., De Castro
Catão, R., Colón-González, F.J., Barcellos, C., Carvalho, M.S., Blangiardo, M., Rue, H.,
Gasparrini, A., 2021. Combined effects of hydrometeorological hazards and urbanisation
on dengue risk in Brazil: a spatiotemporal modelling study. The Lancet Planetary Health
5, e209–e219. https://doi.org/10.1016/S2542-5196(20)30292-8
Martínez-Solanas, È., Basagaña, X., 2019. Temporal changes in the effects of ambient
temperatures on hospital admissions in Spain. PLoS ONE 14, e0218262.
https://doi.org/10.1371/journal.pone.0218262
Mistry, M.N., Schneider, R., Masselot, P., Royé, D., Armstrong, B., Kyselý, J., Orru, H., Sera,
F., Tong, S., Lavigne, É., Urban, A., Madureira, J., García-León, D., Ibarreta, D., Ciscar,
J.-C., Feyen, L., De Schrijver, E., De Sousa Zanotti Stagliorio Coelho, M., Pascal, M.,
Tobias, A., Multi-Country Multi-City (MCC) Collaborative Research Network, Alahmad,
B., Abrutzky, R., Saldiva, P.H.N., Correa, P.M., Orteg, N.V., Kan, H., Osorio, S.,
Indermitte, E., Jaakkola, J.J.K., Ryti, N., Schneider, A., Huber, V., Katsouyanni, K.,
Analitis, A., Entezari, A., Mayvaneh, F., Michelozzi, P., de'Donato, F., Hashizume, M.,
Kim, Y., Diaz, M.H., De La Cruz Valencia, C., Overcenco, A., Houthuijs, D., Ameling,
C., Rao, S., Seposo, X., Nunes, B., Holobaca, I.-H., Kim, H., Lee, W., Íñiguez, C.,
Forsberg, B., Åström, C., Ragettli, M.S., Guo, Y.-L.L., Chen, B.-Y., Colistro, V.,
Zanobetti, A., Schwartz, J., Dang, T.N., Van Dung, D., Guo, Y., Vicedo-Cabrera, A.M.,
Gasparrini, A., 2022. Comparison of weather station and climate reanalysis data for
modelling temperature-related mortality. Sci Rep 12, 5178.
https://doi.org/10.1038/s41598-022-09049-4
Morral-Puigmal, C., Martínez-Solanas, È., Villanueva, C.M., Basagaña, X., 2018. Weather and
gastrointestinal disease in Spain: A retrospective time series regression study.
Environment International 121, 649–657. https://doi.org/10.1016/j.envint.2018.10.003
Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G.,
Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D.G., Piles,
M., Rodríguez-Fernández, N.J., Zsoter, E., Buontempo, C., Thépaut, J.-N., 2021. ERA5-
Land: a state-of-the-art global reanalysis dataset for land applications. Earth Syst. Sci.
Data 13, 4349–4383. https://doi.org/10.5194/essd-13-4349-2021
National Institute of Meteorology (INMET), 2024. Metheorological database, INMET. [WWW
Document]. URL https://bdmep.inmet.gov.br/# (accessed 2.19.24).
Noronha, K.V.M.D.S., Guedes, G.R., Turra, C.M., Andrade, M.V., Botega, L., Nogueira, D.,
Calazans, J.A., Carvalho, L., Servo, L., Ferreira, M.F., 2020. Pandemia por COVID-19
no Brasil: análise da demanda e da oferta de leitos hospitalares e equipamentos de
ventilação assistida segundo diferentes cenários. Cad. Saúde Pública 36, e00115320.
https://doi.org/10.1590/0102-311x00115320
Paz-Bailey, G., Adams, L.E., Deen, J., Anderson, K.B., Katzelnick, L.C., 2024. Dengue. The
Lancet 403, 667–682. https://doi.org/10.1016/S0140-6736(23)02576-X

Pudpong, N., Hajat, S., 2011. High temperature effects on out-patient visits and hospital
admissions in Chiang Mai, Thailand. Science of The Total Environment 409, 5260–5267.
https://doi.org/10.1016/j.scitotenv.2011.09.005
Royé, D., Íñiguez, C., Tobías, A., 2020. Comparison of temperature–mortality associations using
observed weather station and reanalysis data in 52 Spanish cities. Environmental
Research 183, 109237. https://doi.org/10.1016/j.envres.2020.109237
Saha, M.V., Davis, R.E., Hondula, D.M., 2014. Mortality Displacement as a Function of Heat
Event Strength in 7 US Cities. American Journal of Epidemiology 179, 467–474.
https://doi.org/10.1093/aje/kwt264
Schneider, A., Rückerl, R., Breitner, S., Wolf, K., Peters, A., 2017. Thermal Control, Weather,
and Aging. Curr Envir Health Rpt 4, 21–29. https://doi.org/10.1007/s40572-017-0129-0
Schwartz, J., Samet, J.M., Patz, J.A., 2004. Hospital Admissions for Heart Disease: The Effects
of Temperature and Humidity. Epidemiology 15, 755–761.
https://doi.org/10.1097/01.ede.0000134875.15919.0f
Silveira, I.H., Oliveira, B.F.A., Cortes, T.R., Junger, W.L., 2019. The effect of ambient
temperature on cardiovascular mortality in 27 Brazilian cities. Science of The Total
Environment 691, 996–1004. https://doi.org/10.1016/j.scitotenv.2019.06.493
Suaya, J.A., Shepard, D.S., Siqueira, J.B., Martelli, C.T., Lum, L.C.S., Tan, L.H., Kongsin, S.,
Jiamton, S., Garrido, F., Montoya, R., Armien, B., Huy, R., Castillo, L., Caram, M., Sah,
B.K., Sughayyar, R., Tyo, K.R., Halstead, S.B., 2009. Cost of dengue cases in eight
countries in the Americas and Asia: a prospective study. Am J Trop Med Hyg 80, 846–
855.
Vaidyanathan, A., Saha, S., Vicedo-Cabrera, A.M., Gasparrini, A., Abdurehman, N., Jordan, R.,
Hawkins, M., Hess, J., Elixhauser, A., 2019. Assessment of extreme heat and
hospitalizations to inform early warning systems. Proc. Natl. Acad. Sci. U.S.A. 116,
5420–5427. https://doi.org/10.1073/pnas.1806393116
Van Wyk, H., Eisenberg, J.N.S., Brouwer, A.F., 2023. Long-term projections of the impacts of
warming temperatures on Zika and dengue risk in four Brazilian cities using a
temperature-dependent basic reproduction number. PLoS Negl Trop Dis 17, e0010839.
https://doi.org/10.1371/journal.pntd.0010839
Wang, B., Chai, G., Sha, Y., Zha, Q., Su, Y., Gao, Y., 2021. Impact of ambient temperature on
cardiovascular disease hospital admissions in farmers in China's Western suburbs.
Science of The Total Environment 761, 143254.
https://doi.org/10.1016/j.scitotenv.2020.143254
Werneck, G.L., Macias, A.E., Mascarenas, C., Coudeville, L., Morley, D., Recamier, V.,
Guergova-Kuras, M., Puentes-Rosas, E., Baurin, N., Toh, M.-L., 2018. Comorbidities
increase in-hospital mortality in dengue patients in Brazil. Mem. Inst. Oswaldo Cruz 113.
https://doi.org/10.1590/0074-02760180082
Wibawa, B.S.S., Wang, Y.-C., Andhikaputra, G., Lin, Y.-K., Hsieh, L.-H.C., Tsai, K.-H., 2024.
The impact of climate variability on dengue fever risk in central java, Indonesia. Climate
Services 33, 100433. https://doi.org/10.1016/j.cliser.2023.100433
Yi, X., Chang, Z., Zhao, X., Ma, Y., Liu, F., Xiao, X., 2020. The temporal characteristics of the
lag-response relationship and related key time points between ambient temperature and
hand, foot and mouth disease: A multicity study from mainland China. Science of The
Total Environment 749, 141679. https://doi.org/10.1016/j.scitotenv.2020.141679
Zhao, Q., Li, S., Coelho, M.S.Z.S., Saldiva, P.H.N., Hu, K., Huxley, R.R., Abramson, M.J., Guo,

Y., 2019. The association between heatwaves and risk of hospitalization in Brazil: A
nationwide time series study between 2000 and 2015. PLoS Med 16, e1002753.
https://doi.org/10.1371/journal.pmed.1002753

Supplementary Material

Ambient Temperature and Dengue Hospitalisation in Brazil: A 10-year period two-stage case-design time series analysis

Lopes R, et al.

Table of Contents

ERA5-Land validation

We performed a validation comparing the daily average of temperature from monitoring stations at the municipal level with the ERA5-Land estimates. We used data from 389 stations (269 automatic and 120 manual) from the National Institute of Meteorology (INMET), with at least 90% of days with complete data during the period. This validation has a total of 1,221,051 days evaluated, from 340 municipalities covering the 27 states and the 5 macro-regions of Brazil.

	R-Pearson	R-square	RMSE
Overall	0.94	0.90	1.54
Distribution across 340 municipalities			
Mean ± SD	0.89 ± 0.1	0.81 ± 0.1	1.02 ± 0.3
Median [p25-p75]	0.92 [0.86-0.95]	0.85 [0.74-0.90]	1.01 [0.85-1.17]

Spatial distribution of R-square for each of the 340 municipalities. The shade areas are the municipality areas. N stands for North (n=39), NE for Northeast (n=95), CW for Center-West (n=42), SE for Southeast (n=103) and S for South (n=61).

Spatial distribution of RMSE for each of the 340 municipalities. The shade areas are the municipality areas. N stands for North (n=39), NE for Northeast (n=95), CW for Center-West (n=42), SE for Southeast (n=103) and S for South (n=61).

Table S1. Descriptive distribution of 2m height Mean Temperature

Brazil, Region, State	Minimum	5th	25th	50th	75th	95th	Maximum
Brazil	-0.09	15.39	20.92	23.96	26.08	28.68	34.8
North	13.99	23.91	25.15	26.13	27.39	29.32	34.32
AC	13.99	22.81	24.35	25.13	26.01	27.49	31.01
AM	17.98	24.18	25.09	25.81	26.68	28.39	33.06
AP	21.98	23.72	24.73	25.65	27.14	28.59	30.5
PA	21.7	24.24	25.38	26.31	27.53	28.99	32.59
RO	14.11	23.34	24.67	25.52	26.6	28.59	31.61
RR	20.04	22.88	24.29	25.33	26.67	28.82	32.24
TO	19.6	24.17	25.58	26.71	28.03	30.27	34.32
Northeast	14.04	21.49	24.16	25.8	27.24	29.54	34.39
AL	18.6	21.76	23.52	25.02	26.32	27.9	32.78
BA	14.04	19.99	22.31	24.08	25.73	27.8	33.99
CE	19.09	23.82	25.58	26.77	27.81	29.16	32.41
MA	21.75	24.53	25.79	26.84	28.09	29.89	33.29
PB	19.17	22.36	24.19	25.49	26.74	29.15	32.08
PE	17.59	20.77	22.95	24.35	25.76	27.84	32.71
PI	20.71	24.44	26.07	27.55	29.3	31.41	34.39
RN	20.18	24.02	25.56	26.6	27.66	29.29	31.86
SE	19.77	22.17	23.74	25.35	26.53	27.77	32.25
Center-West	6.09	20.68	23.34	24.88	26.41	28.96	34.3
DF	15.96	19.42	21.01	22.23	23.54	25.87	29.58
GO	11.3	20.99	22.97	24.4	26	28.76	34.18
MS	6.09	16.95	22.42	24.71	26.46	28.81	34.26
MT	10.6	22.55	24.46	25.61	26.89	29.31	34.3
Southeast	4.69	16.13	19.7	22.13	24.31	27.03	34.8
ES	10.78	17.22	20.39	22.49	24.66	26.82	31.46
MG	6.71	16.15	19.47	21.76	23.82	26.62	34.54
SP	4.69	15.88	19.96	22.66	24.83	27.45	34.8
RJ	9.78	16.75	19.79	22.12	24.45	27.2	32.25
South	-0.09	11.03	16.52	19.92	22.76	26.02	33.03
PR	0.67	13.25	18.19	21.27	23.78	26.78	32.42
RS	0.77	9.99	15.71	19.33	22.35	25.7	33.03
SC	-0.09	10.8	15.86	18.98	21.73	24.89	31.13

Figure S1. Time series of Dengue hospitalisation by macro administrative region of Brazil. Colour by year, the data covers a period of 10 years, from the whole epidemiological year of 2010 to the whole epidemiological year of 2019.

Figure S2. Cumulative relative risk over all the lags compared to the MHT on each state level, for the whole period of analysis, 2010 to 2019 (First stage, Main Analysis)

The curve is plotted in red lines and the 95% confidence interval generated from the fitted model is given by the grey shaded ribbon around it. The title of each subplot is the abbreviations for the name of each state.

Figure S3. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT at the North Region. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Main analysis)

The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S4. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT at the Northeast Region. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Main analysis)

The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S5. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT at the Center-West Region. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Main analysis)

Meta-analysis Results, Center-West Region

The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S6. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT at the Southeast Region. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Main analysis)

Meta-analysis Results, Southeast Region

The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S7. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT at the South Region. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Main analysis)

The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Table S2. Dengue hospitalisation relative risk by Brazil and each macro-region: main and sensitivity analyses

	MHT (°C)	RR (95% CI), 50th	RR (95% CI), 95th
Main analysis			
Brazil	10.8	2.185 (1.457-3.276)	2.385 (1.556-3.655)
North	30.8	1.139 (0.707-1.836)	1.048 (0.811-1.355)
Northeast	19.7	1.660 (1.116-2.468)	1.547 (0.860-2.784)
Center-West	17	2.393 (1.394-4.110)	2.420 (1.365-4.289)
Southeast	13.7	3.335 (1.605-6.929)	4.930 (2.567-9.468)
South	16	1.119 (0.580-2.159)	2.253 (0.592-8.569)
Sensitivity analysis 1			
Brazil	10.8	2.080 (1.210-3.578)	2.285 (1.152-4.531)
North	30.8	1.297 (0.865-1.945)	1.061 (0.798-1.412)
Northeast	19.7	1.981 (1.179-3.329)	1.983 (0.978-4.020)
Center-West	17	2.176 (1.110-4.265)	2.122 (1.058-4.255)
Southeast	13.7	3.175 (1.865-5.403)	4.358 (2.699-7.038)
South	7.4	4.160 (0.557-31.070)	9.125 (1.308-63.655)
Sensitivity analysis 2			
Brazil	10.8	2.315 (1.558-3.439)	2.501 (1.639-3.819)
North	30.8	1.162 (0.791-1.707)	1.058 (0.824-1.359)
Northeast	19.7	2.051 (1.282-3.284)	2.151 (1.180-3.921)
Center-West	17	2.140 (1.330-3.443)	1.972 (1.172-3.317)
Southeast	13.7	3.156 (1.630-6.112)	3.813 (2.601-5.588)
South	11	2.433 (0.875- 6.761)	5.140 (2.135-12.371)
Sensitivity analysis 3			
Brazil	10.8	1.991 (1.382-2.867)	2.221 (1.512-3.263)
North	22.8	1.091 (0.811-1.469)	1.098 (0.736-1.640)
Northeast	19.7	1.707 (1.163-2.507)	1.735 (1.074-2.803)
Center-West	17	1.688 (1.278-2.228)	1.734 (1.145-2.626)
Southeast	13.7	2.212 (1.344-3.638)	2.553 (1.634-3.989)
South	7.4	2.347 (0.653- 8.440)	4.830 (1.346-17.331)

RR: Relative Risk to the Minimum Hospitalisation Temperature of hospitalisation due Dengue
MHT: Minimum Hospitalisation Temperature, on Celsius degree

Figure S8. Dengue hospitalisation relative risk by Brazil and each macro-region: main and sensitivity analyses forest plot at 95th percentile of temperature

Brazil	MHT*	RR (95% CI)
Main analysis	10.8	2.385 (1.556-3.655)
Sensitivity 1	10.8	2.285 (1.152-4.531)
Sensitivity 2	10.8	2.501 (1.639-3.819)
Sensitivity 3	10.8	2.221 (1.512-3.263)
North		
Main analysis	30.8	1.048 (0.811-1.355)
Sensitivity 1	30.8	1.061 (0.798-1.412)
Sensitivity 2	30.8	1.058 (0.824-1.359)
Sensitivity 3	22.8	1.098 (0.736-1.640)
Northeast		
Main analysis	19.7	1.547 (0.860-2.784)
Sensitivity 1	19.7	1.983 (0.978-4.020)
Sensitivity 2	19.7	2.151 (1.180-3.921)
Sensitivity 3	19.7	1.735 (1.074-2.803)
Center-West		
Main analysis	17	2.420 (1.365-4.289)
Sensitivity 1	17	2.122 (1.058-4.255)
Sensitivity 2	17	1.972 (1.172-3.317)
Sensitivity 3	17	1.734 (1.145-2.626)
Southeast		
Main analysis	13.7	4.930 (2.567-9.468)
Sensitivity 1	13.7	4.358 (2.699-7.038)
Sensitivity 2	13.7	3.813 (2.601-5.588)
Sensitivity 3	13.7	2.553 (1.634-3.989)
South		
Main analysis	16	2.253 (0.592-8.569)
Sensitivity 1	7.4	9.125 (1.308-63.655
Sensitivity 2	11	5.140 (2.135-12.371
Sensitivity 3	7.4	4.830 (1.346-17.331

Relative Risk (95% CI)

*MHT: Minimum Hospitalisation Temperature, on Celsius degree

Figure S9. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT in Brazil. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Sensitivity Analysis 1)

Sensitivity Analysis 1: parametrization of dose-response: 3 knots equally spaced; lag-response: 3 knots equally spaced at the log-scale. Vertical traced lines mark the 50th (Blue) and 95th (Red) percentile of the temperature distribution. The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S10. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT in Brazil. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Sensitivity Analysis 2)

Sensitivity Analysis 2: parametrization of dose-response: 2 knots equally spaced; lag-response: 4 knots at days 1, 2, 7 and 14 from date of hospitalization. Vertical traced lines mark the 50th (Blue) and 95th (Red) percentile of the temperature distribution. The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.

Figure S11. A) Cumulative relative risk over all lags for a Dengue hospitalisation compared to the MHT in Brazil. B) Lag effect of the RR on the 50th percentile of temperature. C) Lag effect of the RR on the 95th percentile of temperature (Sensitivity Analysis 3)

Sensitivity Analysis 3: parametrization of dose-response: 2 knots equally spaced; lag-response: lag-response: 3 knots equally spaced at the log-scale; covariate: 7-day moving average of confirmed dengue cases. Vertical traced lines mark the 50th (Blue) and 95th (Red) percentile of the temperature distribution. The grey shade (A) is 95% confidence interval, as the error bars (B and C), derived from the meta-analysis.