Ambient Temperature and Dengue Hospitalisation in Brazil: A 10-year period two-stage case-design time series analysis

Rafael Lopes[1,2], Xavier Basagaña[2], Leonardo S. L. Bastos[3], Fernando A. Bozza[4], Otavio T. Ranzani[2,5]

1 - Instituto de Física Teórica - IFT, UNESP, São Paulo, Brazil
2 - Barcelona Institute for Global Health, Universitat Pompeu Fabra, CIBER Epidemiología y Salud Pública, Barcelona, Spain
3 - Department of Industrial Engineering (DEI), Pontifical Catholic University of Rio de Janeiro (PUC-Rio), Rio de Janeiro, RJ, Brazil
4 - National Institute of Infectious Disease Evandro Chagas (INI), Oswaldo Cruz Foundation (FIOCRUZ), Rio de Janeiro, RJ, Brazil
5 - Pulmonary Division, Heart Institute (InCor), HCFMUSP, Faculdade de Medicina da Universidade de São Paulo, Brazil

Corresponding author:

Otavio T. Ranzani (otavio.ranzani@isglobal.org)

ISGlobal, Dr. Aiguader, 88, 08003 Barcelona, Spain

Conflicts of interest: None.

Word count: 2,850

Keywords: Dengue; severe Dengue; temperature; heatwave; hospital; temperature

Abstract (270/300 words**)**

Climate factors are known to influence seasonal patterns of dengue transmission. However, little is known about the effect of extremes of temperature on the severity of dengue infection, such as hospital admission. We aimed to quantify the effect of ambient temperature on dengue hospitalisation risk in Brazil. We retrieved daily dengue hospitalisation counts by each of 5,565 municipalities across the 27 states of Brazil from 1st of January 2010 to 31st of December 2019, from the Brazilian Public Hospital Admission System ("SIH"). We obtained average daily ambient temperature for each municipality from the ERA5-land product reanalysis. We combined distributed lag non-linear models with time stratified design model framework to pool a relative risk (RR) estimate for dose-response and lag-response structures for the association of temperature and Dengue hospitalisation. We estimate the overall dengue hospitalisation RR for the whole country as well for each of the five macro-regions by meta-analysing state level estimates. 579,703 hospital admissions due to dengue occurred over the 10 years' period of 2010 to 2019. We observed a positive association between high temperatures and high risk of hospitalisation for Brazil. The overall RR for dengue hospitalisation was 2.185 (95% CI 1.457-3.276) at the 50th percentile of temperature and 2.385 (95% CI 1.556-3.655) at 95th percentile of temperature for Brazil. We also found lag effects of heat on hospitalisation, particularly an immediate augmented risk to hospitalisation, both at the 50th percentile and 95th. High temperatures are associated with an increase in the risk of hospitalisation by dengue. These findings may guide preparedness and mitigation policies during dengue season outbreaks, particularly on the health-care demand.

Introduction

Dengue fever has been a major global seasonal endemic disease, present largely in the tropics, with an estimated burden over ~390 million infections per year (Bhatt et al., 2013; Guzman and Harris, 2015). Approximately one quarter of these infections manifest as clinical or subclinical disease (Bhatt et al., 2013). Brazil is one of the most affected countries by Dengue, that has caused over ~20·9 millions cases of infections in the last 20 years, since the compulsory universal notification to the Brazilian health systems (Godói et al., 2018; Lowe et al., 2021). Additionally, it is estimated that the Brazilian Universal Health system (SUS) spent more than USD 159 millions in the treatment and assistance to Dengue cases and USD 10 million on severe Dengue between 2000 and 2015 (Godói et al., 2018). The impressive burden of Dengue in Brazil is also observed in several others low-income and middle-income countries (Suaya et al., 2009).

Dengue incidence is influenced by climate variables, due to mosquito life cycle and human behaviour changes, particularly with rainfall and temperature (Bhatt et al., 2013; Colón-González et al., 2021; Lee et al., 2021; Lowe et al., 2021; Suaya et al., 2009). However, there is lacking evidence for the association of ambient temperature and dengue severity. The pathophysiology of Dengue severity involves dehydration and coagulation disorders, both conditions that can be aggravated by ambient temperature (Burattini et al., 2016; Werneck et al., 2018).

Based on the reported association of temperature and all-cause hospitalizations (Martínez-Solanas and Basagaña, 2019; Pudpong and Hajat, 2011; Vaidyanathan et al., 2019), we hypothesised that ambient temperature has a positive association with the risk of hospitalisation due to Dengue. We aimed to evaluate the association between ambient temperature and Dengue hospitalisation in Brazil, analysing the period from 2010 to 2019.

Methods

Brazil has about 211 million inhabitants distributed over 8·5 million km² area, is divided into 26 states and the country capital, a federal district (Brazilian Institute of Geography and Statistics - IBGE, n.d.). These 27 states are grouped in 5 administrative macro-regions: North, Northeast, Center-West, Southeast and South. The country is located in a tropical region and has 3 main Köppen climate types and 12 subtypes (Alvares et al., 2013).

We conducted a time-stratified regression analysis to evaluate the association between ambient temperature and Dengue hospitalisation in Brazil. The unit of analysis was municipality of residence. We considered municipalities boundaries as in 2010 census.

We used publicly accessible data and following ethically agreed principles on open data, the use of this data did not require ethical approval in Brazil, according to the Brazilian ethics Resolution n° 510/2016.

Data sources and definitions:

Dengue hospitalizations: We used the Brazilian Hospital Admission System (SIH), a nationwide database that comprises individual level data of all hospitalizations covered by the Universal Healthcare System (SUS) in Brazil. We defined a dengue hospitalisation by the ICD-10 codes 'A90', 'A91', 'A97', 'A970', 'A971', 'A972', 'A979' (Coelho et al., 2016). We build daily time series by date of hospitalisation for each municipality of residence. We excluded dengue hospitalizations from five small municipalities (e.g, small islands), because there was no ambient temperature for them. This filtering resulted in a loss of 153 (0.03%) cases in total, for the whole period of the 10 years. The final time series aggregation by municipalities encompasses a total of

579,703 hospital admissions due to Dengue over the 10 years' period of 2010 to 2019 in 5,565 municipalities.

Number of dengue cases: We used the National Dengue Surveillance System (SINAN-Dengue) to retrieve a daily times series by date of symptoms onset of confirmed dengue cases at each municipality.

Ambient temperature: We used the gridded (0.1º x 0.1º) 2 metres daily temperature taken from reanalysis products (ERA5-Land) freely available by the Coppernicus Climate Service through the Climate date Store (Muñoz-Sabater et al., 2021). The original data was an hourly gridded dataset over the extension of Brazil and encompassing the whole period of 10 years. We estimated the daily mean temperature for each municipality by calculating the mean daily temperature of each grid cell and weighting it to the municipality area. All these weighted mean areas were done with the 'exactextractr' R package (Baston, Daniel, 2023).

Data Analysis:

We study the association between dengue hospitalizations counts time-series and ambient daily temperature using the Distributed Lag Nonlinear Model (DLNM) framework (Gasparrini, 2011; Gasparrini et al., 2010). We use a two-stage approach, first fitting generalised linear models at each Brazilian State and second pooling the estimates using a multivariate meta-analysis (Gasparrini et al., 2012; Gasparrini and Armstrong, 2013; Jackson et al., 2011). At the first stage, we fitted a Conditional Poisson Model for the aggregated time series of Dengue hospitalisation counts (Armstrong et al., 2014). The model strata was a month-city term (Gasparrini, 2021). We report the overall Relative Risk (RR) compared to the Minimum Hospitalisation Temperature

(MHT) point for each state. MHT is defined as the temperature which has an overall RR equal to 1. We also report the lag effects at the 50th and 95th percentiles. The second-stage was one meta-analysis for the whole country and another for each of the macro administrative regions separately. The meta-analysis for the whole country was conducted aggregating all the coefficients fitted on each state and their covariance matrix in a multivariate meta-analysis random effects model. The meta-analysis for each macro-region was using the coefficients and the covariance matrix of the states within a given region. From the meta-analysis we estimated the overall cumulative curve of the RR, relative to MHT and the lag-effects at 50th and 95th percentile.

First-stage:

At the first-stage, we run the following model for each state:

$$g[E(Y_{ijt})] = \xi_{ijs} + f(x_{tij}, l, \theta_j) + \sum_{j=i}^{J} s_k(t; \gamma_j)$$

Where Y_{ijt} is the daily count of cases by the *j-th* state on the *i-th* municipality, ξ_{ijs} the month-city strata term conditioned out. $f(x_{tij}, l, \theta_j)$ which is the bi-dimensional exposure-lag-response distributed lag non-linear cross-basis for the mean temperature on the *j-th* state by each day of delay, until 21 days of lags. The cross-basis is parametrized with natural splines, with 2 knots equally spaced on the dose-response structure and 3 knots equally spaced on the log transformed scale for the lag-response structure. The last term is the long-term trend model choice for temperature trend along the whole period, a natural spline with 7 degrees of freedom for each year over the whole period.

Second-stage:

Whole Country and Regions:

We ran a multivariate meta-analysis with random effects, so from the *j-th* state-level study, θ$_j$, we have the following equation:

$$\theta_i | u_i \sim N_k(\theta_i + u_i, S_i), u_i \sim N_k(0, \psi)$$

Where θ is the study to estimate, with u_i the random effects for the coefficients of this study together with S_j covariance matrix of intra state studies and ψ covariance matrix of between states studies. For the whole country meta-analysis, the θ studies were taken from all the 27 states, and for the macro regions meta-analysis θ was only taken from states pertaining to the same macro region.

We ran three sensitivity analyses to cover the DLNM parametrization and by adjusting for the number of confirmed dengue cases. On the first sensitivity analysis, we parameterized the cross-basis with 3 knots equally spaced on the dose-response structure and, as in the main analysis, 3 knots equally spaced on the log transformed scale for the lag-response structure. The second sensitivity analysis we parameterized the cross-basis with 2 knots equally spaced on the dose-response structure as the main analysis, however with 4 knots placed on the days 1, 2, 7, 14 for the lag-response structure. On the third sensitivity analysis we adjusted the first stage models adding the 7-days moving average of the times series of confirmed dengue cases, considering the average time between incubation period and time to hospitalization.

All the analyses were run on R Software, version 4.1.2.

Results:

Climate data description:

A descriptive table (**Table S1**) of the mean daily temperature for each state, macro regions and

the whole country over the ten years period is shown in the supplementary material. For the whole period of ten years, it varied from -0.09 ℃ to 34.8 ℃. **Figure 1** gives the daily mean temperature distribution across all states.

Figure 1. **Mean temperature density distribution over each state**. On y-axis is each of the state acronyms and on x-axis is the range of mean temperature in degree Celsius. The height of each curve is the cumulative density of mean temperature at that specific degree. The colours are given by each macro administrative region. The inset map gives the location of each of the state code at Brazilian geography.

Table 1. Characteristics description of Dengue Hospitalisation data

Variables	N	Brazil (N=579,703)	North (N=75,304)	Northeast (N=224,085)	Center-West (N=100,056)	Southeast (N=155,405)	South (N=24,853)
Mean Age (IQR), years	579,703	30 (15, 49)	27 (15, 44)	25 (12, 45)	36 (20, 53)	34 (17, 54)	37 (21, 55)
Age Categories, n (%) years	579,703						
0 to 1		8,752 (1.5%)	1,309 (1.7%)	4,457 (2.0%)	1,058 (1.1%)	1,726 (1.1%)	202 (0.8%)
1 to 9		73,507 (13%)	9,401 (12%)	37,354 (17%)	8,438 (8.4%)	16,620 (11%)	1,694 (6.8%)
10 to 17		91,786 (16%)	12,562 (17%)	41,991 (19%)	12,155 (12%)	22,171 (14%)	2,907 (12%)
18 to 39		192,819 (33%)	29,153 (39%)	72,769 (32%)	34,330 (34%)	48,009 (31%)	8,558 (34%)
40 to 59		126,059 (22%)	14,863 (20%)	39,640 (18%)	26,825 (27%)	38,268 (25%)	6,463 (26%)
60 to 79		71,395 (12%)	6,737 (8.9%)	22,314 (10.0%)	14,656 (15%)	23,494 (15%)	4,194 (17%)
80+		15,385 (2.7%)	1,279 (1.7%)	5,560 (2.5%)	2,594 (2.6%)	5,117 (3.3%)	835 (3.4%)
Self-Reported Race, n (%)	396,624						
Black		12,467 (3.1%)	1,042 (2.3%)	3,946 (2.7%)	1,187 (1.9%)	5,760 (4.9%)	532 (2.6%)
Pardo		244,869 (62%)	39,925 (87%)	120,476 (81%)	34,973 (55%)	45,113 (38%)	4,382 (21%)
Indigenous		1,092 (0.3%)	306 (0.7%)	116 (<0.1%)	595 (0.9%)	62 (<0.1%)	13 (<0.1%)
White		129,031 (33%)	3,893 (8.5%)	19,700 (13%)	24,396 (39%)	65,746 (55%)	15,296 (74%)
Asian		9,165 (2.3%)	700 (1.5%)	4,003 (2.7%)	2,116 (3.3%)	1,991 (1.7%)	355 (1.7%)
(Missing)		183,079	29,438	75,844	36,789	36,733	4,275
Sex, n (%)	579,703						
Female		310,308 (54%)	38,225 (51%)	121,175 (54%)	54,916 (55%)	82,550 (53%)	13,442 (54%)
Male		269,395 (46%)	37,079 (49%)	102,910 (46%)	45,140 (45%)	72,855 (47%)	11,411 (46%)
Duration of Hospitalisation, median (IQR), days	579,703	3 (2, 4)	2 (2, 3)	3 (2, 4)	2 (2, 3)	3 (2, 4)	2 (2, 3)
In-hospital mortality, n (%)	579,703	3,436 (0.6%)	281 (0.4%)	958 (0.4%)	548 (0.5%)	1,510 (1.0%)	139 (0.6%)

Dengue hospitalization data description:

Table 1 gives a characterization of the data summarised by each macro administrative region. Overall, the mean age for a hospitalisation by dengue was 30 years old. Between regions it can be seen a difference of 12 years old to the mean age of hospitalisation, being the mean age of hospitalisation of 37 years old for the South region and 25 years old for the Northeast region. The sex ratio is typically representative of the sex ratio estimates from the national statistics. The crude in-hospital mortality in general is low, with mortality rate of 0.6% of the total hospitalizations, and there are some discrepancies between regions. The Southeast has a rate of 1.0% of the hospitalizations with the highest death outcome rate. The North and Northeast with lowest rate of mortality with an average of 0.4% rate of deaths to the total hospitalizations. **Figure S1** gives a visual description of the time series for dengue hospitalisation for each region.

First-stage Results:

The cumulative RR over all lags compared to the MHT on each state level, for the whole period of analysis, 2010 to 2019 are given in the supplementary material (**Figure S2**).

Second-stage Results:

Figure 3 and **4**, presents the cumulative RR over all lags compared to the MHT derived from the meta-analysis, for the whole country and for each macro-region. **Figure 3B** and **3C**, presents the lag effects for the 50th (23.96º C) and 95th (28.68º C) percentiles of mean temperature distribution for Brazil. The lag effects for each macro-region is shown on supplementary material, **Figure S3, S4, S5, S6** and **S7**. A summary of the RR and MHT for Brazil and each macro-region is shown in **Table 2**.

Figure 3. **Pooled cumulative relative risk for the whole country meta-analysis.** A)

Cumulative RR over all lags for a Dengue hospitalisation relative to the MHT. Vertical traced

lines mark the 50th (Blue) and 95th (Red) percentile of the temperature distribution. The grey

shade ribbon is 95% confidence interval derived from the meta-analysis. B) Lag effect to the RR

to the MHT of Hospitalisation due to Dengue at the 50th (23.96º C) percentile of temperature. C)

Lag effect to the RR to the MHT of hospitalisation due to Dengue on the 95th (28.68º C)

percentile of temperature. MHT: Minimum Hospitalisation Temperature; RR: relative risk.

Figure 4. Pooled cumulate relative risk to each of the macro region meta-analysis.

Cumulative overall lags RR to the MHT curves to temperature by region. Each panel is given by

a meta-analysis model run over all the states coefficients and covariance matrices that pertains to

a given macro administrative region. Order by region is given by latitude extent. MHT:

Minimum Hospitalisation Temperature; RR: relative risk.

Table 2. **Dengue hospitalisation relative risk (RR) over Brazil and each macro-region**

Unit	MHT (ºC)	RR (IC 95%) 50th (23·96º C)	RR (IC 95%) 95th (28·68º C)
Brazil	10.8	2.185 (1.457-3.276)	2.385 (1.556-3.655)
North	30.8	1.139 (0.707-1.836)	1.048 (0.811-1.355)
Northeast	19.7	1.660 (1.116-2.468)	1.547 (0.860-2.784)
Center-West	17	2.393 (1.394-4.110)	2.420 (1.365-4.289)
Southeast	13.7	3.335 (1.605-6.929)	4.930 (2.567-9.468)
South	16	1.119 (0.580-2.159)	2.253 (0.592-8.569)

RR: Relative Risk to the Minimum Hospitalisation Temperature of hospitalisation due Dengue

MHT: Minimum Hospitalisation Temperature, on Celsius degree

Sensitivity analyses

Overall, the sensitivity analyses results were comparable with the main analysis (**Figure 5**).

From the meta-analyses, on the 50th percentile for whole Brazil the RR was 2.080 (95% CI

1.210-3.578) and 2.315 (95% CI 1.558-3.439), for the first and second sensitivity analyses

parametrizations, respectively. After adjusting for the number of dengue cases, the RR on the

50th percentile for whole Brazil was 1.991 (95% CI 1.382-2.867). Additional results are shown

on **Table S2** and **Figures S8**, **S9, S10,** and **S11**, in the supplementary material.

Figure 3. Dengue hospitalization relative risk by Brazil and each macro-region: main and sensitivity analyses at 50th percentile of temperature

Brazil	MHT*	RR (95% CI)
Main analysis	10.8	2.185 (1.457-3.276)
Sensitivity 1	10.8	2.080 (1.210-3.578)
Sensitivity 2	10.8	2.315 (1.558-3.439)
Sensitivity 3	10.8	1.991 (1.382-2.867)
North		
Main analysis	30.8	1.139 (0.707-1.836)
Sensitivity 1	30.8	1.297 (0.865-1.945)
Sensitivity 2	30.8	1.162 (0.791-1.707)
Sensitivity 3	22.8	1.091 (0.811-1.469)
Northeast		
Main analysis	19.7	1.660 (1.116-2.468)
Sensitivity 1	19.7	1.981 (1.179-3.329)
Sensitivity 2	19.7	2.051 (1.282-3.284)
Sensitivity 3	19.7	1.707 (1.163-2.507)
Center-West		
Main analysis	17	2.393 (1.394-4.110)
Sensitivity 1	17	2.176 (1.110-4.265)
Sensitivity 2	17	2.140 (1.330-3.443)
Sensitivity 3	17	1.688 (1.278-2.228)
Southeast		
Main analysis	13.7	3.335 (1.605-6.929)
Sensitivity 1	13.7	3.175 (1.865-5.403)
Sensitivity 2	13.7	3.156 (1.630-6.112)
Sensitivity 3	13.7	2.212 (1.344-3.638)
South		
Main analysis	16	1.119 (0.580-2.159)
Sensitivity 1	7.4	4.160 (0.557-31.070
Sensitivity 2	11	2.433 (0.875- 6.761)
Sensitivity 3	7.4	2.347 (0.653- 8.440)

Relative Risk (95% CI)

*MHT: Minimum Hospitalisation Temperature, on Celsius degree; RR: relative risk

Discussion

We ran a two-stage time stratified design study to the association between ambient temperature and hospitalisation due to Dengue. Our main finding is an increased risk of being hospitalised by Dengue as the temperature increases. In a quantitative way, in general for Brazil, above 20 degrees Celsius, the RR to the MHT is 1.2x higher. This is the cumulative effects of lags up to 21 days before the date of hospitalisation. We found a stronger immediate effect on the RR to the hospitalisation. We found the same patterns for each macro region of Brazil, but the North and South regions.

In comparison with the literature, we see that, in general, higher temperature is related to an augmented RR of all-cause and cause-specific hospitalisation and mortality (Jacobson et al., 2021; Martínez-Solanas and Basagaña, 2019; Pudpong and Hajat, 2011; Silveira et al., 2019; Wang et al., 2021; Zhao et al., 2019). There is scarce literature on the association between temperatures and hospitalisation by an infectious disease. Our study is the first, to our knowledge, to measure an association between temperature and hospitalisation caused by dengue. Interestingly, we did not observe a typical U-shape curve increase on the risk for hospitalization, as commonly observed for studies looking for all-cause and cause-specific hospitalisations, except for the South region, where the association was unclear. The patter we observed has been reported for non-infectious causes in the US and China (Vaidyanathan et al., 2019; Wang et al., 2021). Some specific features of dengue could explain this difference. First, its progression to severe disease is highly linked with factors that would be aggravated as the temperature increases, such dehydration and capillary leak syndrome. Second, the incidence of

the dengue disease, of any severity, is much lower during cold months, therefore limiting the power to observe a potential effect of cold temperatures on Dengue hospitalizations. This reflects an additional challenge when studying climate variables and vector-borne infectious diseases severity, among others (Imai et al., 2015).

Overall, in the country and each of the macro regions, the extreme heat (above 20 Celsius degrees) has an effect of increasing the severity of cases, which evolves to a necessity of a hospitalisation. The association for the North and South are unclear and a direct comparison between regions is not straightforward because of regional disparities as well as climate disparities. Although Brazil has a universal access health system, SUS has great disparities over geographical distribution (Noronha et al., 2020). Thus, we might expect that regions with more hospital beds per population, such as South and Southeast, would hospitalise milder cases compared with regions with fewer hospital beds. This is also important during massive Dengue outbreaks, where the threshold for hospitalisation might change according to beds availability and age distribution.

Several mechanisms can play a role in the observed overall association and lag effects. The immediate effect generating the first peak on lag-0 is expected and observed in the majority of studies evaluating temperature and heat waves and hospitalizations and deaths (Morral-Puigmal et al., 2018; Pudpong and Hajat, 2011; Royé et al., 2020; Saha et al., 2014; Schwartz et al., 2004; Vaidyanathan et al., 2019). It reflects an immediate worsening of the clinical condition due to high temperature during an infection. This sharp increase on lag-0 reinforces that our results is a direct effect of ambient temperature on hospitalisation risk, rather than the indirect effect of temperature on mosquito activity and increased risk of dengue incidence, association usually present in the scale of months (Lowe et al., 2021). Even though, the sensitivity analysis when

adding the 7-day rolling mean of mild cases as a covariate in the model shows that the effect of temperature on our model is reduced from 2.185 to 1.991 at the 50[th] and from 2.385 to 2.221 at the 95[th] percentile of temperature (**Table S2**).

Another mechanism that Dengue has is a natural evolution of 1 to 3 days of incubation period, and more 5 to 7 days to complete clearance of the viral infection. If an individual is exposed to higher temperatures during the viral infection evolution this can lead to more severe infection, as seen a potential late lag-effect on some regions. Finally, the decrease on the RR following the immediate effect could be due to a depletion of susceptible ("harvesting effect") (Saha et al., 2014; Schwartz et al., 2004). This mechanism is reported for the lag between temperature and deaths, but also has been hypothesised in another study evaluating temperature and other seasonal infectious diseases such as hand, foot and mouth disease (Yi et al., 2020). Importantly, the lag-effect varies over regions, for instance, the South region has a delayed lag-effect (**Figure S7**). When varying points for the knots in a sensitivity analysis, lag-effects were less prominent, increasing the likelihood of collinearity explaining this finding (Basagaña and Barrera-Gómez, 2022).

Our study has some strengths. We evaluated a nationwide database providing a 10-year time-series of Dengue hospitalizations in a LMIC exposed to a wide temperature range. Ambient temperature from reanalysis products from ERA5-Land are the state-of-the-art in climatology (Royé et al., 2020), as the framework applied for data analysis, accounting for dose-response and lag-response structures and correlated daily data on temperature (De Schrijver et al., 2022; Gasparrini, 2021). This has great generalizability to many other LMIC countries as countries in

the regions which dengue has been expanding during last years. The present study adds to the literature dengue burden and provides sources to better support to the public health measures, showing increased risk of hospitalization in warmer days.

Our study has limitations to be mentioned. First, we did not evaluate other factors that could modify the high temperature effects, such as green space, urbanisation and relative humidity. In the same extension, we did not evaluate individual factors, such as age and sex, that could show different effects of temperature and risk of dengue hospitalization in vulnerable populations. Second, we obtained estimates for 27 states and pooled them for Brazil and each corresponding macro-regions. States from the same macro-region will have similar climate as well as a similar dengue incidence. This choice of aggregation allows to comparing states with probably similar transmission conditions to the mosquito as well hospitalizations conditions (Lee et al., 2021). However, this may introduce bias to the analysis and do not allow for generalizations to some municipalities within each state or to border line areas.

Conclusion

We found an increased RR to the Dengue hospitalisation due high temperatures in Brazil during a ten years period, 2010-2019. This RR varies over the whole extension of Brazil and between macro-regions. With the findings presented here we hope the climate variability of Brazil and the present climate emergency can be taken into account when public health measures are designed to curb and mitigate the impacts of dengue on the population.

Acknowledgements:

The authors also thank the research funding agencies: the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior -- Brazil (Finance Code 001 to RLPS), Conselho Nacional de Desenvolvimento Científico e Tecnológico -- Brazil (grant number: 141698/2018-7 to RLPS). OTR is funded by a Sara Borrell fellowship (CD19/00110) from the Instituto de Salud Carlos III. We acknowledge support from the grant CEX2018-000806-S funded by MCIN/AEI/ 10.13039/501100011033, and support from the Generalitat de Catalunya through the CERCA Program. The funding agencies had no role in the conceptualization of the study.

Data Availability Statement:

All data used in this study are publicly available. Hospitalisation data is available at

http://sihd.datasus.gov.br/principal/index.php, and temperature at

https://cds.climate.copernicus.eu/#!/home. The code to reproduce this analysis is available at:

https://github.com/rafalopespx/dengue_t2m_severity_paper

Author Contribution Statement:

Rafael Lopes: Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Software, Visualization, Writing – original draft

Xavier Basagaña: Methodology, Supervision, Validation, Writing - review & editing

Leonardo S. L. Bastos: Data curation, Writing - review & editing

Fernando A. Bozza: Writing - review & editing

Otavio T. Ranzani: Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Supervision, Validation, Writing - review & editing

References

Alvares, C.A., Stape, J.L., Sentelhas, P.C., De Moraes Gonçalves, J.L., Sparovek, G., 2013. Köppen's climate classification map for Brazil. metz 22, 711–728. https://doi.org/10.1127/0941-2948/2013/0507

Armstrong, B.G., Gasparrini, A., Tobias, A., 2014. Conditional Poisson models: a flexible alternative to conditional logistic case cross-over analysis. BMC Med Res Methodol 14, 122. https://doi.org/10.1186/1471-2288-14-122

Basagaña, X., Barrera-Gómez, J., 2022. Reflection on modern methods: visualizing the effects of collinearity in distributed lag models. International Journal of Epidemiology 51, 334–344. https://doi.org/10.1093/ije/dyab179

Baston, Daniel, 2023. exactextractr: Fast Extraction from Raster Datasets using Polygons. R package.

Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., Drake, J.M., Brownstein, J.S., Hoen, A.G., Sankoh, O., Myers, M.F., George, D.B., Jaenisch, T., Wint, G.R.W., Simmons, C.P., Scott, T.W., Farrar, J.J., Hay, S.I., 2013. The global distribution and burden of dengue. Nature 496, 504–507. https://doi.org/10.1038/nature12060

Brady, O.J., Gething, P.W., Bhatt, S., Messina, J.P., Brownstein, J.S., Hoen, A.G., Moyes, C.L., Farlow, A.W., Scott, T.W., Hay, S.I., 2012. Refining the Global Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. PLoS Negl Trop Dis 6, e1760. https://doi.org/10.1371/journal.pntd.0001760

Brazilian Institute of Geography and Statistics - IBGE, n.d. 2010 Population Census. [WWW Document]. URL https://www.ibge.gov.br/en/statistics/social/income-expenditure-and-consumption/18391-2010-population-census.html?edicao=19720&t=sobre (accessed 4.27.21).

Burattini, M.N., Lopez, L.F., Coutinho, F.A.B., Siqueira, J.B., Homsani, S., Sarti, E., Massad, E., 2016. Age and regional differences in clinical presentation and risk of hospitalization for dengue in Brazil, 2000-2014. Clinics 71, 455–463. https://doi.org/10.6061/clinics/2016(08)08

Coelho, G.E., Leal, P.L., Cerroni, M.D.P., Simplicio, A.C.R., Siqueira, J.B., 2016. Sensitivity of the Dengue Surveillance System in Brazil for Detecting Hospitalized Cases. PLoS Negl Trop Dis 10, e0004705. https://doi.org/10.1371/journal.pntd.0004705

Colón-González, F.J., Sewe, M.O., Tompkins, A.M., Sjödin, H., Casallas, A., Rocklöv, J., Caminade, C., Lowe, R., 2021. Projecting the risk of mosquito-borne diseases in a warmer and more populated world: a multi-model, multi-scenario intercomparison modelling study. The Lancet Planetary Health 5, e404–e414. https://doi.org/10.1016/S2542-5196(21)00132-7

De Schrijver, E., Bundo, M., Ragettli, M.S., Sera, F., Gasparrini, A., Franco, O.H., Vicedo-Cabrera, A.M., 2022. Nationwide Analysis of the Heat- and Cold-Related Mortality Trends in Switzerland between 1969 and 2017: The Role of Population Aging. Environ Health Perspect 130, 037001. https://doi.org/10.1289/EHP9835

Gasparrini, A., 2021. The Case Time Series Design. Epidemiology 32, 829–837. https://doi.org/10.1097/EDE.0000000000001410

Gasparrini, A., 2011. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. J Stat Softw 43, 1–20.

Gasparrini, A., Armstrong, B., 2013. Reducing and meta-analysing estimates from distributed lag

non-linear models. BMC Med Res Methodol 13, 1. https://doi.org/10.1186/1471-2288-13-1

Gasparrini, A., Armstrong, B., Kenward, M.G., 2012. Multivariate meta-analysis for non-linear and other multi-parameter associations. Stat Med 31, 3821–3839. https://doi.org/10.1002/sim.5471

Gasparrini, A., Armstrong, B., Kenward, M.G., 2010. Distributed lag non-linear models. Stat Med 29, 2224–2234. https://doi.org/10.1002/sim.3940

Godói, I.P., Da Silva, L.V.D., Sarker, A.R., Megiddo, I., Morton, A., Godman, B., Alvarez-Madrazo, S., Bennie, M., Guerra-Junior, A.A., 2018. Economic and epidemiological impact of dengue illness over 16 years from a public health system perspective in Brazil to inform future health policies including the adoption of a dengue vaccine. Expert Review of Vaccines 17, 1123–1133. https://doi.org/10.1080/14760584.2018.1546581

Guzman, M.G., Harris, E., 2015. Dengue. The Lancet 385, 453–465. https://doi.org/10.1016/S0140-6736(14)60572-9

Imai, C., Armstrong, B., Chalabi, Z., Mangtani, P., Hashizume, M., 2015. Time series regression model for infectious disease and weather. Environmental Research 142, 319–327. https://doi.org/10.1016/j.envres.2015.06.040

Jackson, D., Riley, R., White, I.R., 2011. Multivariate meta-analysis: potential and promise. Stat Med 30, 2481–2498. https://doi.org/10.1002/sim.4172

Jacobson, L.D.S.V., Oliveira, B.F.A.D., Schneider, R., Gasparrini, A., Hacon, S.D.S., 2021. Mortality Risk from Respiratory Diseases Due to Non-Optimal Temperature among Brazilian Elderlies. IJERPH 18, 5550. https://doi.org/10.3390/ijerph18115550

Lee, S.A., Economou, T., De Castro Catão, R., Barcellos, C., Lowe, R., 2021. The impact of climate suitability, urbanisation, and connectivity on the expansion of dengue in 21st century Brazil. PLoS Negl Trop Dis 15, e0009773. https://doi.org/10.1371/journal.pntd.0009773

Lowe, R., Lee, S.A., O'Reilly, K.M., Brady, O.J., Bastos, L., Carrasco-Escobar, G., De Castro Catão, R., Colón-González, F.J., Barcellos, C., Carvalho, M.S., Blangiardo, M., Rue, H., Gasparrini, A., 2021. Combined effects of hydrometeorological hazards and urbanisation on dengue risk in Brazil: a spatiotemporal modelling study. The Lancet Planetary Health 5, e209–e219. https://doi.org/10.1016/S2542-5196(20)30292-8

Martínez-Solanas, È., Basagaña, X., 2019. Temporal changes in the effects of ambient temperatures on hospital admissions in Spain. PLoS ONE 14, e0218262. https://doi.org/10.1371/journal.pone.0218262

Morral-Puigmal, C., Martínez-Solanas, È., Villanueva, C.M., Basagaña, X., 2018. Weather and gastrointestinal disease in Spain: A retrospective time series regression study. Environment International 121, 649–657. https://doi.org/10.1016/j.envint.2018.10.003

Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D.G., Piles, M., Rodríguez-Fernández, N.J., Zsoter, E., Buontempo, C., Thépaut, J.-N., 2021. ERA5-Land: a state-of-the-art global reanalysis dataset for land applications. Earth Syst. Sci. Data 13, 4349–4383. https://doi.org/10.5194/essd-13-4349-2021

Noronha, K.V.M.D.S., Guedes, G.R., Turra, C.M., Andrade, M.V., Botega, L., Nogueira, D., Calazans, J.A., Carvalho, L., Servo, L., Ferreira, M.F., 2020. Pandemia por COVID-19 no Brasil: análise da demanda e da oferta de leitos hospitalares e equipamentos de ventilação assistida segundo diferentes cenários. Cad. Saúde Pública 36, e00115320.

https://doi.org/10.1590/0102-311x00115320

Pudpong, N., Hajat, S., 2011. High temperature effects on out-patient visits and hospital admissions in Chiang Mai, Thailand. Science of The Total Environment 409, 5260–5267. https://doi.org/10.1016/j.scitotenv.2011.09.005

Royé, D., Íñiguez, C., Tobías, A., 2020. Comparison of temperature–mortality associations using observed weather station and reanalysis data in 52 Spanish cities. Environmental Research 183, 109237. https://doi.org/10.1016/j.envres.2020.109237

Saha, M.V., Davis, R.E., Hondula, D.M., 2014. Mortality Displacement as a Function of Heat Event Strength in 7 US Cities. American Journal of Epidemiology 179, 467–474. https://doi.org/10.1093/aje/kwt264

Schwartz, J., Samet, J.M., Patz, J.A., 2004. Hospital Admissions for Heart Disease: The Effects of Temperature and Humidity. Epidemiology 15, 755–761. https://doi.org/10.1097/01.ede.0000134875.15919.0f

Silveira, I.H., Oliveira, B.F.A., Cortes, T.R., Junger, W.L., 2019. The effect of ambient temperature on cardiovascular mortality in 27 Brazilian cities. Science of The Total Environment 691, 996–1004. https://doi.org/10.1016/j.scitotenv.2019.06.493

Suaya, J.A., Shepard, D.S., Siqueira, J.B., Martelli, C.T., Lum, L.C.S., Tan, L.H., Kongsin, S., Jiamton, S., Garrido, F., Montoya, R., Armien, B., Huy, R., Castillo, L., Caram, M., Sah, B.K., Sughayyar, R., Tyo, K.R., Halstead, S.B., 2009. Cost of dengue cases in eight countries in the Americas and Asia: a prospective study. Am J Trop Med Hyg 80, 846–855.

Vaidyanathan, A., Saha, S., Vicedo-Cabrera, A.M., Gasparrini, A., Abdurehman, N., Jordan, R., Hawkins, M., Hess, J., Elixhauser, A., 2019. Assessment of extreme heat and hospitalizations to inform early warning systems. Proc. Natl. Acad. Sci. U.S.A. 116, 5420–5427. https://doi.org/10.1073/pnas.1806393116

Wang, B., Chai, G., Sha, Y., Zha, Q., Su, Y., Gao, Y., 2021. Impact of ambient temperature on cardiovascular disease hospital admissions in farmers in China's Western suburbs. Science of The Total Environment 761, 143254. https://doi.org/10.1016/j.scitotenv.2020.143254

Werneck, G.L., Macias, A.E., Mascarenas, C., Coudeville, L., Morley, D., Recamier, V., Guergova-Kuras, M., Puentes-Rosas, E., Baurin, N., Toh, M.-L., 2018. Comorbidities increase in-hospital mortality in dengue patients in Brazil. Mem. Inst. Oswaldo Cruz 113. https://doi.org/10.1590/0074-02760180082

Yi, X., Chang, Z., Zhao, X., Ma, Y., Liu, F., Xiao, X., 2020. The temporal characteristics of the lag-response relationship and related key time points between ambient temperature and hand, foot and mouth disease: A multicity study from mainland China. Science of The Total Environment 749, 141679. https://doi.org/10.1016/j.scitotenv.2020.141679

Zhao, Q., Li, S., Coelho, M.S.Z.S., Saldiva, P.H.N., Hu, K., Huxley, R.R., Abramson, M.J., Guo, Y., 2019. The association between heatwaves and risk of hospitalization in Brazil: A nationwide time series study between 2000 and 2015. PLoS Med 16, e1002753. https://doi.org/10.1371/journal.pmed.1002753