medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Title page

Psychosocial predictors of COVID-19 vaccine uptake among pregnant women: a cross-sectional study in Greece

Petros Galanis, Assistant Professor, Clinical Epidemiology Laboratory, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Irene Vraka, MD, PhD, Department of Radiology, P & A Kyriakou Children's Hospital, Athens, Greece

Aglaia Katsiroumpa, RN, MSc, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Olympia Konstantakopoulou, Economist, MSc, PhD, Center for Health Services Management and Evaluation, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Olga Siskou, Assistant Professor, University of Piraeus, Piraeus, Greece

Eleftheria Zogaki, Midwife, MSc, Faculty of Midwifery, Technological Education Institution of Western Macedonia, Kozani, Greece

Daphne Kaitelidou, Professor, Center for Health Services Management and Evaluation, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Corresponding author: Petros Galanis, Assistant Professor, Clinical Epidemiology Laboratory, Faculty of Nursing, National and Kapodistrian University of Athens, 123 Papadiamantopoulou street, GR-11527, Athens, Greece, e-mail: pegalan@nurs.uoa.gr, https://orcid.org/0000-0002-1308-5782

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Abstract

Background: Unvaccinated pregnant women with symptomatic COVID-19 have been found to have a higher risk of iatrogenic preterm births, intensive care unit admission, and invasive ventilation.

Objective: To estimate the vaccination rate of pregnant women against the COVID-19 and to evaluate psychosocial factors associated with vaccine uptake among them.

Methods: We conducted an anonymous cross-sectional study with a convenience sample in Greece from December 2021 to March 2022. We measured socio-demographic data of pregnant women, COVID-19-related vaccination status, worry about the side effects of COVID-19 vaccines, trust in COVID-19 vaccines, and COVID-19-related stress.

Results: The study population included 812 pregnant women with a mean age of 31.6 years. Among the pregnant women, 58.6% had received a COVID-19 vaccine. The most important reasons that pregnant women were not vaccinated were doubts about the safety and effectiveness of COVID-19 vaccines (31.4%), fear that COVID-19 vaccines could be harmful to fetus (29.4%), and fear of adverse side effects of COVID-19 vaccines (29.4%). Increased danger and contamination fears, increased fears about economic consequences, and higher levels of trust in COVID-19 vaccines were related with COVID-19 vaccine uptake. On the other hand, increased compulsive checking and reassurance seeking and increased worry about the adverse side effects of COVID-19 vaccines reduced the likelihood of pregnant women being vaccinated against the COVID-19.

Conclusions: An understanding of the psychosocial factors associated with COVID-19 vaccine uptake in pregnant women is paramount to persuade women to get vaccinated against the COVID-19. There is a need for targeted educational campaigns to increase knowledge about COVID-19 vaccines and reduce COVID-19 vaccine hesitancy in pregnancy.

Key words: COVID-19; pregnant women; vaccine uptake; predictors; Greece

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Introduction

Unvaccinated pregnant women with symptomatic COVID-19 are at increased risk for iatrogenic preterm births, intensive care unit admission, and invasive ventilation (Khalil et al., 2020; Lokken et al., 2021; Vousden et al., 2021). Also, risk of death for unvaccinated women with symptomatic COVID-19 during pregnancy is higher than non pregnant women with symptomatic COVID-19 (Zambrano et al., 2020). Moreover, unvaccinated pregnant women have been found to have a higher risk of hospital admission for COVID-19 than vaccinated pregnant women (Iacobucci, 2021).

Pregnant women were excluded from early randomized controlled trials, leading to the lack of safety and efficacy data (Pogue et al., 2020; Polack et al., 2020). However, the most recent data are rather encouraging for vaccinated pregnant women. In particular, COVID-19 vaccine research finds that (1) abortion rate, adverse pregnancy, and adverse neonatal outcomes are similar in vaccinated and non-vaccinated pregnant women, (2) anti-SARS-CoV-2 immunoglobulins provide immunity to the newborns, and (3) COVID-19 vaccines do not cause vaccine-related adverse events (Falsaperla et al., 2021; Fu et al., 2022; Garg et al., 2021; Mendoza et al., 2020; Rawal et al., 2022). Thus, several organizations worldwide, now recommend COVID-19 vaccination for pregnant women and women who are trying to become pregnant, or who might become pregnant in the future to prevent severe maternal morbidity and adverse birth outcomes (American College of Obstetricians and Gynecologist, 2021; Centers for Disease Control and Prevention, 2021).

Until now, only a handful of studies investigated the determinants of COVID-19 vaccine uptake among pregnant women (Blakeway et al., 2022; Hosokawa et al., 2022; Razzaghi et al., 2021; Siegel et al., 2021; UK Health Security Agency, 2021). Moreover, only one study searched for psychosocial predictors of COVID-19 vaccine uptake among pregnant women, e.g. trust in COVID-19 vaccines, fear of COVID-19, worry about the adverse side effects of the COVID-19 vaccines, etc. (Siegel et al., 2021). So far, research has focused on the impact of the socio-demographic characteristics on the decision of pregnant women to be vaccinated against the COVID-19. Thus, the aim of our study was to estimate the rate of vaccinated pregnant women against the COVID-19 and to study psychosocial predictors of COVID-19 vaccine uptake.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Methods

Study design and participants

We conducted a cross-sectional study with a convenience sample in Greece. Recruitment of pregnant women began about one year after the Greek government has offered a free COVID-19 vaccine to all adults. In particular, we collected data from December 2021 to March 2022. We used several approaches to collect our data. Firstly, we used Google forms to create an anonymous version of the study questionnaire and then we disseminated it through social media, e.g. Facebook. Moreover, we sent the questionnaire via e-mail to all our contacts targeting in pregnant women. At the same time, we asked pregnant women who completed the study questionnaire to invite to our study any other pregnant women they might have known. Thus, a snowball sampling method was applied. Women were considered eligible for the study if they were pregnant aged 18 or older and indicated that they could complete the study questionnaire in Greek. Data were anonymous and confidential and participation in the survey was voluntary. We informed the pregnant women that they could cease to participate at any stage of the study should they wish to do so.

Considering a low effect size (odds ratio=1.4), the precision level as 5% (alpha level), and the power as 95%, a minimum sample size of 564 pregnant women was required. Nevertheless, we strove to recruit a larger number of pregnant women to further decrease random error.

The Ethics Committee of Department of Nursing, National and Kapodistrian University of Athens approved the study protocol (reference number; 370, 02-09-2021).

Questionnaires

Firstly, we collected the following socio-demographic data of pregnant women: age (continuous), marital status (singles, married, in a couple relationship without marriage, or divorced), educational level (elementary school, high school, or university degree), financial status (very poor, poor, average, good, or very good), physical health (very poor, poor, average, good, or very good), children aged 18 or

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

younger (no or yes), chronic disease (no or yes), previous COVID-19 diagnosis (no or yes), and family members/friends with previous COVID-19 diagnosis (no or yes).

Also, we used four questions to measure pregnant women' vaccination status and COVID-19-related vaccination status. In particular, we asked the pregnant women (1) if they have received the vaccination against the seasonal influenza (no or yes), (2) if they have received the vaccination against the COVID-19 (no or yes), and (3) if they have received the vaccination against the COVID-19 before or during pregnancy (no or yes). Moreover, we asked unvaccinated pregnant women to state the main reason for this denial (doubts about the safety and effectiveness of COVID-19 vaccines, fear of adverse side effects of COVID-19 vaccines, low self-perceived threat regarding COVID-19, previous COVID-19 diagnosis, fear due to a chronic disease, family doctor' recommendation due to physical health of pregnant women, religious issues, or fear that the COVID-19 vaccination could be harmful to the fetus).

We measured pregnant women's worry about the side effects of COVID-19 vaccines with a single item: "I am worried about the side effects that COVID-19 vaccines can have". Also, we measured pregnant women's trust in COVID-19 vaccines with a single item: "I trust the COVID-19 vaccines". Answers in both cases were indicated on a scale ranging from 0 (totally disagree) to 10 (totally agree).

We used the 36-item COVID Stress Scales (CSS) to measure COVID-19-related stress (Taylor et al., 2020). In particular, we used the Greek version of the CSS (Galanis, Vraka, Katsiroumpa, et al., 2022). The questionnaire consists of five factors: (1) Danger and contamination fears, (2) fears about economic consequences, (3) xenophobia, (4) compulsive checking and reassurance seeking, and (5) traumatic stress symptoms about COVID-19. Responses for the first three factors are indicated on a 5-point Likert scale ranging from 0 (not at all) to 4 (extremely), and for the last two factors are indicated on a 5-point Likert scale ranging from 0 (never) to 4 (almost always). For each factor, a stress score is calculated by averaging the answers to all items of the factor. Thus, score for each factor ranges from 0 to 4 with higher scores reflect greater stress. In our study, Cronbach's alpha was 0.93, 0.94, 0.95, 0.88, and 0.93 for the five factors respectively.

Statistical analysis

We use numbers and percentages to present categorical variables and mean and standard deviation to present continuous variables. Socio-demographic data of

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

pregnant women, COVID-19-related stress, worry about the adverse side effects of COVID-19 vaccines, and trust in COVID-19 vaccines were the independent variables in our study. The outcome variable was COVID-19 vaccination uptake among pregnant women, measured through "yes/no" answers. Thus, we used logistic regression analysis to identify the predictors of COVID-19 vaccine uptake. We performed univariate and multivariate logistic regression analysis, calculating unadjusted and adjusted odds ratios respectively. Also, we calculated 95% confidence intervals (CI), and two-sided p-values. A p-value<0.05 considering as statistically significant. Statistical analysis was performed with the Statistical Package for Social Sciences software (IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.).

Results

Socio-demographic characteristics

The study population included 812 pregnant women. Mean age of the pregnant women was 31.6 years, while 35.3% were expecting their first child. The majority of the participants were married or in a couple relationship without marriage (89.7%). Among pregnant women, 48.3% defined their financial status as good/very good. Almost 15% of the pregnant women suffered from a chronic disease and 90.5% defined their health status as good/very good. Twenty-four point one percent of pregnant women and 79.3% of their family members/friends were diagnosed with COVID-19. Detailed socio-demographic characteristics of the pregnant women are shown in Table 1.

COVID-19-related vaccination status

Among the pregnant women in our study, 58.6% had received a COVID-19 vaccine. Among the vaccinated women, 45.6% had received a COVID-19 vaccine during pregnancy. The most important reasons that pregnant women were not vaccinated were doubts about the safety and effectiveness of COVID-19 vaccines (31.4%), fear that COVID-19 vaccines could be harmful to my fetus (29.4%), and fear of adverse side effects of COVID-19 vaccines (29.4%). In our sample, 24.1% of the pregnant women had received a flu vaccine during 2021. COVID-19-related vaccination status of the pregnant women is presented in Table 2.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

COVID-19-related stress

The highest levels of COVID-19-related stress were due to danger and contamination fears and then to xenophobia, compulsive checking and reassurance seeking, fears about economic consequences, and traumatic stress symptoms about COVID-19 was low. The level of danger and contamination fears was moderate for both vaccinated and unvaccinated pregnant women, while the level of (1) fears about economic consequences, (2) xenophobia, (3) compulsive checking and reassurance seeking, and (4) traumatic stress symptoms about COVID-19 was low. The mean level of danger and contamination fears, fears about economic consequences, xenophobia, compulsive checking and reassurance seeking, and traumatic stress symptoms about COVID-19 among the vaccinated pregnant women is higher than that of the unvaccinated pregnant women. COVID-19-related stress of pregnant women according to COVID-19 vaccination status is shown in Table 3.

Predictors of COVID-19 vaccines uptake

Multivariate logistic regression analysis identified that five psychosocial factors and seven socio-demographic characteristics affect pregnant women decision to receive a COVID-19 vaccine. The multivariate logistic regression analysis for COVID-19 vaccine uptake among the pregnant women revealed that the independent variables explained 55.6% of the variance in this outcome. Predictors of COVID-19 vaccines uptake are shown in Table 4.

Regarding psychosocial factors, increased danger and contamination fears, increased fears about economic consequences, and higher levels of trust in COVID-19 vaccines were related with pregnant women's COVID-19 vaccine uptake. On the other hand, increased compulsive checking and reassurance seeking and increased worry about the adverse side effects of COVID-19 vaccines reduced the likelihood of pregnant women being vaccinated against the COVID-19.

Regarding socio-demographic characteristics, singles women, women that already had at least one child aged <18 years old, women that defined their health status as very poor/poor/moderate, women without a previous COVID-19 diagnosis, women with family members/friends with previous COVID-19 diagnosis, and women who were vaccinated against the seasonal influenza had a greater probability to be vaccinated against the COVID-19. Moreover, increased age was significantly associated with higher COVID-19 vaccine uptake among pregnant women.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Discussion

To the best of our knowledge this is the first study that investigates psychosocial predictors of COVID-19 vaccine uptake among pregnant women with a valid questionnaire. Siegel et al. 2021 studied several psychosocial predictors but without using a valid questionnaire. In a sample of pregnant women in Greece, we found that 58.6% of them were vaccinated against the COVID-19 during pregnancy. A recent meta-analysis found that the overall proportion of vaccinated pregnant women against the COVID-19 is 27.5% (Galanis, Vraka, Siskou, Konstantakopoulou, Katsiroumpa, & Kaitelidou, 2022b). Also, it is interesting to highlight the fact that the vaccination rate is much higher in Israel (43.3%) than in the USA (27.3%) and other countries (12.8%). The higher vaccination rate in our study may be due to the fact that the studies in the meta-analysis were conducted from December 2020 to September 2021, while our study was conducted from December 2021 to March 2022. Evidence on the safety and efficacy of the COVID-19 vaccines is constantly increasing which may also increase the confidence of pregnant women in vaccines.

We found that the main reasons that pregnant women were not vaccinated were doubts about the safety and effectiveness of COVID-19 vaccines, fear that COVID-19 vaccines could be harmful to my fetus, and fear of adverse side effects. This finding is confirmed by the literature, since several studies have revealed women's fear that COVID-19 vaccines can cause problems in fertility, pregnancy, and breastfeeding (Nguyen et al., 2021; Schrading et al., 2021; Xu et al., 2021). Moreover, pregnant women are more worried about lack of safety data in pregnancy, fetal effects, vaccine adverse side effects, and rushed development of vaccines (Siegel et al., 2021; Skjefte et al., 2021). Moreover, worry in the general population about safety, efficacy, and side-effects of COVID-19 vaccines is related with hesitancy in COVID-19 vaccines uptake (Freeman et al., 2020).

According to our multivariate analysis, several psychosocial factors affect pregnant women decision to receive a COVID-19 vaccine. In particular, increased danger, contamination fears, and fears about economic consequences were related with COVID-19 vaccine uptake among pregnant women. Siegel et al. (2021) confirm our finding, since they found that vaccinated pregnant women are more afraid of the COVID-19 during pregnancy than unvaccinated pregnant women. Moreover, vaccinated pregnant women believe that if they are infected they are at risk for getting very sick (Siegel et al., 2021). Increased fear of pregnant women in our study may

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

also be due to the fact that they have family members/friends with previous COVID-19 diagnosis. Siegel et al. (2021) confirm this result since vaccinated pregnant women are more likely to know hospitalized COVID-19 patients. In general, fear of COVID-19 is associated with good prevention practices in both pregnant women and the general population (Ahorsu et al., 2022; Masjoudi et al., 2022; W/Mariam et al., 2021; Wong et al., 2020; Yıldırım et al., 2021). Probably, individuals who fear to acquire and spread the COVID-19 could better adopt a protective health behavior against the COVID-19 and comply with preventive measures recommendations. Therefore, fear appears to promote preventive measures during the COVID-19 pandemic, e.g. frequent hand-washing, physical distancing, face masks, and vaccination.

On the other hand, we found that high levels of COVID-19-related stress and more specifically the stress caused by the compulsive checking and reassurance seeking reduce the COVID-19 vaccine uptake among the pregnant women. Pregnant women experience moderate to high levels of stress during the COVID-19 pandemic (Ayaz et al., 2020; Mappa et al., 2020; Stepowicz et al., 2020). Particular attention should be focused on pregnant women since COVID-19-related stress is associated with depressive symptoms (King et al., 2021). Higher depressive symptoms in pregnancy may have negative effects on infants' health, e.g. disruptions of pro- and anti-inflammatory markers, infant stress responses, shortened breastfeeding period, and poorer bonding between mothers and their infants (Edvinsson et al., 2017; Figueiredo et al., 2014; Gustafsson et al., 2018; Osborne et al., 2018; Rossen et al., 2016). Moreover, there is a negative association between perceived stress and self-care behaviors among pregnant women (Masjoudi et al., 2022). Negative self-care behaviors among pregnant women with high levels of COVID-19-related stress could explain the low vaccination rate among those women since self-care behaviors are purposeful actions that individuals take on to improve their health (Masjoudi et al., 2020; Stearns et al., 2000).

Also, we found that higher levels of trust in COVID-19 vaccines were associated with higher COVID-19 vaccine uptake among pregnant women. Literature confirms this finding, since Siegel et al. (2021) found that trust in COVID-19 vaccines effectiveness for women and newborns is associated with increased vaccine uptake. Also, they recognized that unvaccinated pregnant women are less likely to trust vaccine developers. Moreover, general distrust is related with hesitancy in COVID-19 vaccines uptake in the general population (Freeman et al., 2020). It is noteworthy that

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

the mistrust in the COVID-19 vaccines is higher among individuals from ethnic minorities, both in the general population and among pregnant women (Green et al., 2021; Paul et al., 2021; Taubman – Ben Ari et al., 2022). Another important issue is that several studies found a positive relation between trust in COVID-19 vaccines and parents' willingness to vaccinate their children against the COVID-19 (Galanis, Vraka, Siskou, Konstantakopoulou, Katsiroumpa, & Kaitelidou, 2022a; Skjefte et al., 2021; Yılmaz & Sahin, 2021; Zhang et al., 2020).

Our study identified several socio-demographic predictors of COVID-19 vaccination uptake among pregnant women. Among others, we found that pregnant women who received a flu vaccine had also a greater probability to be vaccinated against the COVID-19. Systematic reviews have already shown that previous seasonal influenza vaccination history is a strong predictive factor for individuals to accept a COVID-19 vaccine, both for themselves and their children (Galanis, Vraka, et al., 2021; Galanis, Vraka, Siskou, Konstantakopoulou, Katsiroumpa, & Kaitelidou, 2022a; Li et al., 2021; Luo et al., 2021; Shamshirsaz et al., 2021). Moreover, recent studies expand this evidence by confirming that COVID-19 vaccination uptake is more common among individuals that are already vaccinated against the seasonal influenza (Doran et al., 2022; Galanis, Moisoglou, et al., 2021; Galanis, Vraka, Siskou, Konstantakopoulou, Katsiroumpa, Moisoglou, et al., 2022; Gilboa et al., 2021; Shariff et al., 2022). Influenza vaccination during the COVID-19 pandemic is crucial since may have a protective role against the COVID-19 pandemic by reducing SARS-CoV-2 infection rate, hospitalization, severity of COVID-19, admission to intensive care unit and mortality (Wang et al., 2021; Zeynali Bujani et al., 2021).

Older age of pregnant women is another predictor of COVID-19 vaccination uptake. Our findings confirm the evidence that older age is associated with increased likelihood of acceptance and uptake of a COVID-19 vaccine among pregnant women (Razzaghi et al., 2021; Skjefte et al., 2021; Stuckelberger et al., 2021; UK Health Security Agency, 2021). Probably, older pregnant women experience more fear of COVID-19 since the evidence shows that pregnancy at advanced maternal age is a strong predictor for adverse outcomes, e.g. neonatal intensive care unit admission, low birthweight babies, worse Apgar scores, spontaneous miscarriage, cesarean deliveries, and pre-eclampsia (Glick et al., 2021; Pinheiro et al., 2019). Moreover, it is well known that older age increases adverse outcomes in COVID-19 patients, such as hospitalization, intensive care unit admission and mortality (Mehraeen et al., 2020;

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Sepandi et al., 2020; Yanez et al., 2020). This fear that pregnant women experience against the COVID-19 is confirmed by two more finding in our study. In particular, we found that pregnant women that defined their health status as poor and women without a previous COVID-19 diagnosis were more likely to get vaccinated against the COVID-19. In a similar way, Blakeway et al. (2022) found that pre-gestational diabetes mellitus is associated with COVID-19 vaccination uptake among pregnant women. Higher levels of fear about contracting COVID-19 increase the intention of individuals to accept a COVID-19 vaccine (Caserotti et al., 2021; Ward et al., 2020). For instance, intention of people to accept a COVID-19 vaccine is higher during the lockdown periods when the self-perceived COVID-19 vulnerability is higher (Brooks et al., 2020; Moccia et al., 2020). Moreover, past COVID-19 patients are less likely to be vaccinated, since they perceive a low risk and feel protected against the COVID-19 (Pacella-LaBarbara et al., 2021).

Limitations

We should note a number of limitations in our study. Firstly, we conducted a cross-sectional study and therefore we are unable to establish a causal mechanism between psychosocial factors and COVID-19 vaccination uptake among pregnant women. Second, we relied on a convenience sample of pregnant women that cannot be considered representative of the population of pregnant women in Greece. For instance, the educational level of the participants in our study was high, while the participation rate of migrants was probably low since the questionnaire was only in Greek language. Furthermore, we used a valid questionnaire to measure psychosocial pattern of pregnant women but our data were based on self-reported measures which may introduce information bias due to tendency of participants to seek for social desirability. In addition, there are also other psychosocial factors that could affect pregnant women decision to receive a COVID-19 vaccine, e.g. anxiety, depression, quality of life, etc. Moreover, we did not measure some possible confounders, such as ethnicity, gestational week, at-risk pregnancy, vaccine type (mRNA or viral vector vaccine) that was offered to pregnant women, number of people in household, employment status (housewife or employed), and work location (working in person or working remotely).

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Conclusions

Our study is the first to assess psychosocial predictors of COVID-19 vaccines uptake among pregnant women with a valid instrument. This study is very timely due to the ongoing high COVID-19 case rates globally and the known increased risks of COVID-19 in pregnant women. An understanding of the psychosocial factors associated with increased COVID-19 vaccine uptake in pregnant women could be helpful for policy makers and healthcare professionals in their efforts to persuade women to get vaccinated against the COVID-19. Furthermore, we found that several socio-demographic characteristics of pregnant women affect their decision to receive a COVID-19 vaccine. Our findings emphasize the need for a more sensitive approach in the attempt to encourage vaccination uptake in pregnancy, especially in younger women, women with a good self-perceived health status, women with a previous COVID-19 diagnosis, and women who are not vaccinated against the seasonal influenza. Thus, public educational campaigns to encourage COVID-19 vaccination in the population of pregnant women should be targeted taking into account different concerns, needs, and attitudes. Mass vaccination of pregnant women is paramount to reduce COVID-19 vaccine hesitancy in pregnancy, halt spread of SARS-CoV-2 and reduce adverse outcomes in mothers and newborns.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 1. Socio-demographic characteristics of the pregnant women.

Variable	**N**	**%**
Marital status		
Singles	84	10.3
Married or in a couple relationship without marriage	728	89.7
Divorced	0	0
Age (years)[a]	31.6	4.6
Children <18 years old		
No	525	64.7
Yes	287	35.3
Educational level		
Elementary school	7	0.9
High school	210	25.9
University degree	595	73.3
Self-perceived financial status		
Very poor	7	0.9
Poor	28	3.4
Moderate	385	47.4
Good	336	41.4
Very good	56	6.9
Self-perceived health status		
Very poor	0	0
Poor	0	0
Moderate	77	9.5
Good	462	56.9
Very good	273	33.6
Chronic disease		
No	693	85.3
Yes	119	14.7
Previous COVID-19 diagnosis		
No	616	75.9
Yes	196	24.1

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Family members/friends with previous COVID-19 diagnosis		
No	168	20.7
Yes	644	79.3

[a] mean, standard deviation

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 2. COVID-19-related vaccination status in the pregnant women**.**

Variable	**N**	**%**
COVID-19 vaccination uptake		
No	336	41.4
Yes	476	58.6
COVID-19 vaccination uptake before pregnancy		
No	217	45.6
Yes	259	54.4
Seasonal influenza vaccination in 2021		
No	616	75.9
Yes	196	24.1
Reasons for decline of pregnant women to receive a COVID-19 vaccine		
I have doubts about the safety and effectiveness of COVID-19 vaccines	112	31.4
I am afraid of adverse side effects of COVID-19 vaccines	105	29.4
I believe that even I get infected with COVID-19, nothing bad will happen to me	14	3.9
I have already been diagnosed with COVID-19 and the vaccine will not be beneficial for me	14	3.9
I am afraid because I suffer from a chronic disease	7	2.0
I am afraid that COVID-19 vaccines could be harmful to my fetus	105	29.4

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 3. COVID-19-related stress of pregnant women according to COVID-19 vaccination status.

Scale	Mean	Standard deviation
Danger and contamination fears		
Unvaccinated	2.01	1.06
Vaccinated	2.34	1.02
Fears about economic consequences		
Unvaccinated	0.51	0.82
Vaccinated	0.65	1.05
Xenophobia		
Unvaccinated	0.97	1.24
Vaccinated	1.04	1.27
Compulsive checking and reassurance seeking		
Unvaccinated	0.87	0.84
Vaccinated	0.92	0.97
Traumatic stress symptoms about COVID-19		
Unvaccinated	0.40	0.74
Vaccinated	0.45	0.74

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 4. Univariate and multivariate logistic regression analysis with COVID-19 vaccine uptake among the pregnant women as the dependent variable (reference: COVID-19 vaccine denial).

Variable	Unadjusted OR (95% CI)	P-value	Adjusted OR (95% CI)[a]	P-value
Marital status (singles vs. married)	1.45 (0.91 – 2.36)	0.12	2.43 (1.09 – 5.43)	**0.03**
Age (years)	1.07 (1.04 – 1.10)	<0.001	1.09 (1.04 – 1.15)	**0.001**
Children aged <18 years old (yes vs. no)	1.36 (1.01 – 1.83)	0.04	1.84 (1.19 – 2.83)	**0.006**
Educational level (University degree vs. high school)	2.75 (1.99 – 3.79)	<0.001	1.08 (0.65 – 1.77)	0.77
Self-perceived financial status (very good/good vs. moderate/poor/very poor)	1.18 (0.89 – 1.56)	0.24	0.91 (0.59 – 1.39)	0.67
Self-perceived health status (very poor/poor/moderate vs. good/ very good)	1.26 (0.78 – 2.05)	0.35	3.17 (1.49 – 6.72)	**0.003**
Chronic disease (yes vs. no)	1.35 (0.90 – 2.03)	0.15	0.95 (0.52 – 1.73)	0.85
Previous COVID-19 diagnosis (no vs. yes)	1.31 (0.95 – 1.81)	0.10	1.69 (1,04 – 2.78)	**0.04**
Family members/friends with previous COVID-19 diagnosis (yes vs. no)	2.39 (1.69 – 3.38)	<0.001	1.82 (1.09 – 3.05)	**0.02**
Seasonal influenza vaccination in 2021 (yes vs. no)	8.72 (5.40 – 14.09)	<0.001	17.89 (8.51 – 37.64)	**<0.001**
Danger and contamination fears	1.35 (1.18 – 1.54)	<0.001	1.38 (1.07 – 1.78)	**0.013**
Fears about economic consequences	1.16 (0.99 – 1.35)	0.05	1.48 (1.09 – 2.01)	**0.01**
Xenophobia	1.04 (0.93 – 1.17)	0.47	1.07 (0.85 – 1.35)	0.54
Compulsive checking and reassurance seeking	1.07 (0.91 – 1.24)	0.41	0.46 (0.35 – 0.62)	**<0.001**
Traumatic stress symptoms about COVID-19	1.09 (0.89 – 1.32)	0.38	1.19 (0.86 – 1.66)	0.28
Worry about the side effects of COVID-19 vaccines	0.84 (0.81 – 0.88)	<0.001	0.73 (0.67 – 0.78)	**<0.001**
Trust in COVID-19 vaccines	1.36 (1.29 – 1.44)	<0.001	1.53 (1.41 – 1.66)	**<0.001**

An odds ratio <1 indicates a negative association, while an odds ratio >1 indicates a positive association. Bold values indicate statistically significant relations in multivariate model.

CI: confidence interval; OR: odds ratio

[a] R^2 for the final multivariate model was 55.6%

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

References

Ahorsu, D. K., Imani, V., Lin, C.-Y., Timpka, T., Broström, A., Updegraff, J. A., Årestedt, K., Griffiths, M. D., & Pakpour, A. H. (2022). Associations Between Fear of COVID-19, Mental Health, and Preventive Behaviours Across Pregnant Women and Husbands: An Actor-Partner Interdependence Modelling. *International Journal of Mental Health and Addiction*, *20*(1), 68–82. https://doi.org/10.1007/s11469-020-00340-x

American College of Obstetricians and Gynecologist. (2021). *ACOG and SMFM Recommend COVID-19 Vaccination for Pregnant Individuals*. https://www.acog.org/news/newsreleases/ 2021/07/acog-smfm-recommend-covid-19-vaccination-for-pregnant-individuals

Ayaz, R., Hocaoğlu, M., Günay, T., Yardımcı, O. devrim, Turgut, A., & Karateke, A. (2020). Anxiety and depression symptoms in the same pregnant women before and during the COVID-19 pandemic. *Journal of Perinatal Medicine*, *48*(9), 965–970. https://doi.org/10.1515/jpm-2020-0380

Blakeway, H., Prasad, S., Kalafat, E., Heath, P. T., Ladhani, S. N., Le Doare, K., Magee, L. A., O'Brien, P., Rezvani, A., von Dadelszen, P., & Khalil, A. (2022). COVID-19 vaccination during pregnancy: Coverage and safety. *American Journal of Obstetrics and Gynecology*, *226*(2), 236.e1-236.e14. https://doi.org/10.1016/j.ajog.2021.08.007

Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *The Lancet*, *395*(10227), 912–920. https://doi.org/10.1016/S0140-6736(20)30460-8

Caserotti, M., Girardi, P., Rubaltelli, E., Tasso, A., Lotto, L., & Gavaruzzi, T. (2021). Associations of COVID-19 risk perception with vaccine hesitancy over time for Italian residents. *Social Science & Medicine*, *272*, 113688. https://doi.org/10.1016/j.socscimed.2021.113688

Centers for Disease Control and Prevention. (2021). *COVID-19 vaccines while pregnant or breastfeeding*. https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/pregnancy.html

Doran, J., Seyidov, N., Mehdiyev, S., Gon, G., Kissling, E., Herdman, T., Suleymanova, J., Rehse, A. P. C., Pebody, R., Katz, M. A., & Hagverdiyev, G.

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

(2022). Factors associated with early uptake of COVID-19 vaccination among healthcare workers in Azerbaijan, 2021. *Influenza and Other Respiratory Viruses*. https://doi.org/10.1111/irv.12978

Edvinsson, Å., Bränn, E., Hellgren, C., Freyhult, E., White, R., Kamali-Moghaddam, M., Olivier, J., Bergquist, J., Boström, A. E., Schiöth, H. B., Skalkidou, A., Cunningham, J. L., & Sundström-Poromaa, I. (2017). Lower inflammatory markers in women with antenatal depression brings the M1/M2 balance into focus from a new direction. *Psychoneuroendocrinology*, *80*, 15–25. https://doi.org/10.1016/j.psyneuen.2017.02.027

Falsaperla, R., Leone, G., Familiari, M., & Ruggieri, M. (2021). COVID-19 vaccination in pregnant and lactating women: A systematic review. *Expert Review of Vaccines*, *20*(12), 1619–1628. https://doi.org/10.1080/14760584.2021.1986390

Figueiredo, B., Canário, C., & Field, T. (2014). Breastfeeding is negatively affected by prenatal depression and reduces postpartum depression. *Psychological Medicine*, *44*(5), 927–936. https://doi.org/10.1017/S0033291713001530

Freeman, D., Loe, B. S., Chadwick, A., Vaccari, C., Waite, F., Rosebrock, L., Jenner, L., Petit, A., Lewandowsky, S., Vanderslott, S., Innocenti, S., Larkin, M., Giubilini, A., Yu, L.-M., McShane, H., Pollard, A. J., & Lambe, S. (2020). COVID-19 vaccine hesitancy in the UK: The Oxford coronavirus explanations, attitudes, and narratives survey (Oceans) II. *Psychological Medicine*, 1–15. https://doi.org/10.1017/S0033291720005188

Fu, W., Sivajohan, B., McClymont, E., Albert, A., Elwood, C., Ogilvie, G., & Money, D. (2022). Systematic review of the safety, immunogenicity, and effectiveness of COVID-19 vaccines in pregnant and lactating individuals and their infants. *International Journal of Gynaecology and Obstetrics: The Official Organ of the International Federation of Gynaecology and Obstetrics*, *156*(3), 406–417. https://doi.org/10.1002/ijgo.14008

Galanis, P., Moisoglou, I., Vraka, I., Siskou, O., Konstantakopoulou, O., Katsiroumpa, A., & Kaitelidou, D. (2021). Predictors of COVID-19 Vaccine Uptake in Healthcare Workers: A Cross-sectional Study in Greece. *Journal of Occupational & Environmental Medicine*, *Publish Ahead of Print*. https://doi.org/10.1097/JOM.0000000000002463

Galanis, P., Vraka, I., Fragkou, D., Bilali, A., & Kaitelidou, D. (2021). Intention of healthcare workers to accept COVID-19 vaccination and related factors: A systematic

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

review and meta-analysis. *Asian Pacific Journal of Tropical Medicine*, *14*(12), 543. https://doi.org/10.4103/1995-7645.332808

Galanis, P., Vraka, I., Katsiroumpa, A., Konstantakopoulou, O., Siskou, O., Zogaki, E., & Kaitelidou, D. (2022). *Cross-cultural adaptation and validation of the "COVID Stress Scales" in Greek* [Preprint]. Public and Global Health. https://doi.org/10.1101/2022.02.28.22271615

Galanis, P., Vraka, I., Siskou, O., Konstantakopoulou, O., Katsiroumpa, A., & Kaitelidou, D. (2022a). Willingness, refusal and influential factors of parents to vaccinate their children against the COVID-19: A systematic review and meta-analysis. *Preventive Medicine*, *157*, 106994. https://doi.org/10.1016/j.ypmed.2022.106994

Galanis, P., Vraka, I., Siskou, O., Konstantakopoulou, O., Katsiroumpa, A., & Kaitelidou, D. (2022b). *Uptake of COVID-19 vaccines among pregnant women: A systematic review and meta-analysis* [Preprint]. Public and Global Health. https://doi.org/10.1101/2022.04.01.22273296

Galanis, P., Vraka, I., Siskou, O., Konstantakopoulou, O., Katsiroumpa, A., Moisoglou, I., & Kaitelidou, D. (2022). Predictors of Real-World Parents' Acceptance to Vaccinate Their Children Against the COVID-19. *Acta Medica Lituanica*, *29*(1), 4. https://doi.org/10.15388/Amed.2021.29.1.4

Garg, I., Shekhar, R., Sheikh, A. B., & Pal, S. (2021). COVID-19 Vaccine in Pregnant and Lactating Women: A Review of Existing Evidence and Practice Guidelines. *Infectious Disease Reports*, *13*(3), 685–699. https://doi.org/10.3390/idr13030064

Gilboa, M., Tal, I., Levin, E. G., Segal, S., Belkin, A., Zilberman-Daniels, T., Biber, A., Rubin, C., Rahav, G., & Regev-Yochay, G. (2021). Coronavirus disease 2019 (COVID-19) vaccination uptake among healthcare workers. *Infection Control and Hospital Epidemiology*, 1–6. https://doi.org/10.1017/ice.2021.421

Glick, I., Kadish, E., & Rottenstreich, M. (2021). Management of Pregnancy in Women of Advanced Maternal Age: Improving Outcomes for Mother and Baby. *International Journal of Women's Health*, *Volume 13*, 751–759. https://doi.org/10.2147/IJWH.S283216

Green, M. S., Abdullah, R., Vered, S., & Nitzan, D. (2021). A study of ethnic, gender and educational differences in attitudes toward COVID-19 vaccines in Israel – implications for vaccination implementation policies. *Israel Journal of Health Policy Research*, *10*(1), 26. https://doi.org/10.1186/s13584-021-00458-w

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Gustafsson, H. C., Sullivan, E. L., Nousen, E. K., Sullivan, C. A., Huang, E., Rincon, M., Nigg, J. T., & Loftis, J. M. (2018). Maternal prenatal depression predicts infant negative affect via maternal inflammatory cytokine levels. *Brain, Behavior, and Immunity*, *73*, 470–481. https://doi.org/10.1016/j.bbi.2018.06.011

Hosokawa, Y., Okawa, S., Hori, A., Morisaki, N., Takahashi, Y., Fujiwara, T., Nakayama, S. F., Hamada, H., Satoh, T., & Tabuchi, T. (2022). The Prevalence of COVID-19 Vaccination and Vaccine Hesitancy in Pregnant Women: An Internet-based Cross-sectional Study in Japan. *Journal of Epidemiology*, JE20210458. https://doi.org/10.2188/jea.JE20210458

Iacobucci, G. (2021). Covid-19 and pregnancy: Vaccine hesitancy and how to overcome it. *BMJ (Clinical Research Ed.)*, *375*, n2862. https://doi.org/10.1136/bmj.n2862

Khalil, A., von Dadelszen, P., Draycott, T., Ugwumadu, A., O'Brien, P., & Magee, L. (2020). Change in the Incidence of Stillbirth and Preterm Delivery During the COVID-19 Pandemic. *JAMA*. https://doi.org/10.1001/jama.2020.12746

King, L. S., Feddoes, D. E., Kirshenbaum, J. S., Humphreys, K. L., & Gotlib, I. H. (2021). Pregnancy during the pandemic: The impact of COVID-19-related stress on risk for prenatal depression. *Psychological Medicine*, 1–11. https://doi.org/10.1017/S003329172100132X

Li, M., Luo, Y., Watson, R., Zheng, Y., Ren, J., Tang, J., & Chen, Y. (2021). Healthcare workers' (HCWs) attitudes and related factors towards COVID-19 vaccination: A rapid systematic review. *Postgraduate Medical Journal*, postgradmedj-2021-140195. https://doi.org/10.1136/postgradmedj-2021-140195

Lokken, E. M., Huebner, E. M., Taylor, G. G., Hendrickson, S., Vanderhoeven, J., Kachikis, A., Coler, B., Walker, C. L., Sheng, J. S., Al-Haddad, B. J. S., McCartney, S. A., Kretzer, N. M., Resnick, R., Barnhart, N., Schulte, V., Bergam, B., Ma, K. K., Albright, C., Larios, V., … Washington State COVID-19 in Pregnancy Collaborative. (2021). Disease severity, pregnancy outcomes, and maternal deaths among pregnant patients with severe acute respiratory syndrome coronavirus 2 infection in Washington State. *American Journal of Obstetrics and Gynecology*, *225*(1), 77.e1-77.e14. https://doi.org/10.1016/j.ajog.2020.12.1221

Luo, C., Yang, Y., Liu, Y., Zheng, D., Shao, L., Jin, J., & He, Q. (2021). Intention to COVID-19 vaccination and associated factors among health care workers: A

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

systematic review and meta-analysis of cross-sectional studies. *American Journal of Infection Control*, *49*(10), 1295–1304. https://doi.org/10.1016/j.ajic.2021.06.020

Mappa, I., Distefano, F. A., & Rizzo, G. (2020). Effects of coronavirus 19 pandemic on maternal anxiety during pregnancy: A prospectic observational study. *Journal of Perinatal Medicine*, *48*(6), 545–550. https://doi.org/10.1515/jpm-2020-0182

Masjoudi, M., Aslani, A., Khazaeian, S., & Fathnezhad-Kazemi, A. (2020). Explaining the experience of prenatal care and investigating the association between psychological factors with self-care in pregnant women during COVID-19 pandemic: A mixed method study protocol. *Reproductive Health*, *17*(1), 98. https://doi.org/10.1186/s12978-020-00949-0

Masjoudi, M., Aslani, A., Seifi, M., Khazaeian, S., & Fathnezhad-Kazemi, A. (2022). Association between perceived stress, fear and anxiety of COVID 19 with self-care in pregnant women: A cross-sectional study. *Psychology, Health & Medicine*, *27*(2), 289–300. https://doi.org/10.1080/13548506.2021.1894344

Mehraeen, E., Karimi, A., Barzegary, A., Vahedi, F., Afsahi, A. M., Dadras, O., Moradmand-Badie, B., Seyed Alinaghi, S. A., & Jahanfar, S. (2020). Predictors of mortality in patients with COVID-19–a systematic review. *European Journal of Integrative Medicine*, *40*, 101226. https://doi.org/10.1016/j.eujim.2020.101226

Mendoza, M., Garcia□Ruiz, I., Maiz, N., Rodo, C., Garcia□Manau, P., Serrano, B., Lopez□Martinez, R., Balcells, J., Fernandez□Hidalgo, N., Carreras, E., & Suy, A. (2020). Pre□eclampsia□like syndrome induced by severe COVID□19: A prospective observational study. *BJOG: An International Journal of Obstetrics & Gynaecology*, *127*(11), 1374–1380. https://doi.org/10.1111/1471-0528.16339

Moccia, L., Janiri, D., Pepe, M., Dattoli, L., Molinaro, M., De Martin, V., Chieffo, D., Janiri, L., Fiorillo, A., Sani, G., & Di Nicola, M. (2020). Affective temperament, attachment style, and the psychological impact of the COVID-19 outbreak: An early report on the Italian general population. *Brain, Behavior, and Immunity*, *87*, 75–79. https://doi.org/10.1016/j.bbi.2020.04.048

Nguyen, L. H., Joshi, A. D., Drew, D. A., Merino, J., Ma, W., Lo, C.-H., Kwon, S., Wang, K., Graham, M. S., Polidori, L., Menni, C., Sudre, C. H., Anyane-Yeboa, A., Astley, C. M., Warner, E. T., Hu, C. Y., Selvachandran, S., Davies, R., Nash, D., … Chan, A. T. (2021). *Racial and ethnic differences in COVID-19 vaccine hesitancy and uptake* [Preprint]. Epidemiology. https://doi.org/10.1101/2021.02.25.21252402

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Osborne, S., Biaggi, A., Chua, T. E., Du Preez, A., Hazelgrove, K., Nikkheslat, N., Previti, G., Zunszain, P. A., Conroy, S., & Pariante, C. M. (2018). Antenatal depression programs cortisol stress reactivity in offspring through increased maternal inflammation and cortisol in pregnancy: The Psychiatry Research and Motherhood – Depression (PRAM-D) Study. *Psychoneuroendocrinology*, *98*, 211–221. https://doi.org/10.1016/j.psyneuen.2018.06.017

Pacella-LaBarbara, M. L., Park, Y. L., Patterson, P. D., Doshi, A., Guyette, M. K., Wong, A. H., Chang, B. P., & Suffoletto, B. P. (2021). COVID-19 Vaccine Uptake and Intent Among Emergency Healthcare Workers: A Cross-Sectional Survey. *Journal of Occupational & Environmental Medicine*, *63*(10), 852–856. https://doi.org/10.1097/JOM.0000000000002298

Paul, E., Steptoe, A., & Fancourt, D. (2021). Attitudes towards vaccines and intention to vaccinate against COVID-19: Implications for public health communications. *The Lancet Regional Health - Europe*, *1*, 100012. https://doi.org/10.1016/j.lanepe.2020.100012

Pinheiro, R. L., Areia, A. L., Mota Pinto, A., & Donato, H. (2019). Advanced Maternal Age: Adverse Outcomes of Pregnancy, A Meta-Analysis. *Acta Médica Portuguesa*, *32*(3), 219. https://doi.org/10.20344/amp.11057

Pogue, K., Jensen, J. L., Stancil, C. K., Ferguson, D. G., Hughes, S. J., Mello, E. J., Burgess, R., Berges, B. K., Quaye, A., & Poole, B. D. (2020). Influences on Attitudes Regarding Potential COVID-19 Vaccination in the United States. *Vaccines*, *8*(4), E582. https://doi.org/10.3390/vaccines8040582

Polack, F. P., Thomas, S. J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J. L., Pérez Marc, G., Moreira, E. D., Zerbini, C., Bailey, R., Swanson, K. A., Roychoudhury, S., Koury, K., Li, P., Kalina, W. V., Cooper, D., Frenck, R. W., Hammitt, L. L., … C4591001 Clinical Trial Group. (2020). Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *The New England Journal of Medicine*, *383*(27), 2603–2615. https://doi.org/10.1056/NEJMoa2034577

Rawal, S., Tackett, R. L., Stone, R. H., & Young, H. N. (2022). COVID-19 Vaccination among Pregnant People in the U.S.: A Systematic Review. *American Journal of Obstetrics & Gynecology MFM*, 100616. https://doi.org/10.1016/j.ajogmf.2022.100616

Razzaghi, H., Meghani, M., Pingali, C., Crane, B., Naleway, A., Weintraub, E., Kenigsberg, T. A., Lamias, M. J., Irving, S. A., Kauffman, T. L., Vesco, K. K., Daley,

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

M. F., DeSilva, M., Donahue, J., Getahun, D., Glenn, S., Hambidge, S. J., Jackson, L., Lipkind, H. S., … Patel, S. A. (2021). COVID-19 Vaccination Coverage Among Pregnant Women During Pregnancy—Eight Integrated Health Care Organizations, United States, December 14, 2020–May 8, 2021. *MMWR. Morbidity and Mortality Weekly Report*, *70*(24), 895–899. https://doi.org/10.15585/mmwr.mm7024e2

Rossen, L., Hutchinson, D., Wilson, J., Burns, L., A Olsson, C., Allsop, S., J Elliott, E., Jacobs, S., Macdonald, J. A., & Mattick, R. P. (2016). Predictors of postnatal mother-infant bonding: The role of antenatal bonding, maternal substance use and mental health. *Archives of Women's Mental Health*, *19*(4), 609–622. https://doi.org/10.1007/s00737-016-0602-z

Schrading, W. A., Trent, S. A., Paxton, J. H., Rodriguez, R. M., Swanson, M. B., Mohr, N. M., Talan, D. A., Project COVERED Emergency Department Network, Bahamon, M., Carlson, J. N., Chisolm□Straker, M., Driver, B., Faine, B., Galbraith, J., Giordano, P. A., Haran, J. P., Higgins, A., Hinson, J., House, S., … Weber, K. D. (2021). Vaccination rates and acceptance of SARS□CoV□2 vaccination among U.S. emergency department health care personnel. *Academic Emergency Medicine*, *28*(4), 455–458. https://doi.org/10.1111/acem.14236

Sepandi, M., Taghdir, M., Alimohamadi, Y., Afrashteh, S., & Hosamirudsari, H. (2020). Factors Associated with Mortality in COVID-19 Patients: A Systematic Review and Meta-Analysis. *Iranian Journal of Public Health*. https://doi.org/10.18502/ijph.v49i7.3574

Shamshirsaz, A. A., Hessami, K., Morain, S., Afshar, Y., Nassr, A. A., Arian, S. E., Asl, N. M., & Aagaard, K. (2021). Intention to Receive COVID-19 Vaccine during Pregnancy: A Systematic Review and Meta-analysis. *American Journal of Perinatology*, a-1674-6120. https://doi.org/10.1055/a-1674-6120

Shariff, S. Z., Richard, L., Hwang, S. W., Kwong, J. C., Forchuk, C., Dosani, N., & Booth, R. (2022). COVID-19 vaccine coverage and factors associated with vaccine uptake among 23□247 adults with a recent history of homelessness in Ontario, Canada: A population-based cohort study. *The Lancet. Public Health*, *7*(4), e366–e377. https://doi.org/10.1016/S2468-2667(22)00037-8

Siegel, M. R., Lumbreras-Marquez, M. I., James, K., McBay, B. R., Gray, K. J., Schantz-Dunn, J., Diouf, K., & Goldfarb, I. T. (2021). *Perceptions and Attitudes Towards COVID-19 Vaccination Amongst Pregnant and Postpartum Individuals* [Preprint]. Obstetrics and Gynecology. https://doi.org/10.1101/2021.12.17.21267997

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Skjefte, M., Ngirbabul, M., Akeju, O., Escudero, D., Hernandez-Diaz, S., Wyszynski, D. F., & Wu, J. W. (2021). COVID-19 vaccine acceptance among pregnant women and mothers of young children: Results of a survey in 16 countries. *European Journal of Epidemiology*, *36*(2), 197–211. https://doi.org/10.1007/s10654-021-00728-6

Stearns, S., Bernard, S., Fasick, S., Schwartz, R., Konrad, T., Ory, M., & DeFriese, G. (2000). The economic implications of self-care: The effect of lifestyle, functional adaptations, and medical self-care among a national sample of Medicare beneficiaries. *American Journal of Public Health*, *90*(10), 1608–1612. https://doi.org/10.2105/AJPH.90.10.1608

Stepowicz, A., Wencka, B., Bieńkiewicz, J., Horzelski, W., & Grzesiak, M. (2020). Stress and Anxiety Levels in Pregnant and Post-Partum Women during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, *17*(24), 9450. https://doi.org/10.3390/ijerph17249450

Stuckelberger, S., Favre, G., Ceulemans, M., Nordeng, H., Gerbier, E., Lambelet, V., Stojanov, M., Winterfeld, U., Baud, D., Panchaud, A., & Pomar, L. (2021). SARS-CoV-2 Vaccine Willingness among Pregnant and Breastfeeding Women during the First Pandemic Wave: A Cross-Sectional Study in Switzerland. *Viruses*, *13*(7), 1199. https://doi.org/10.3390/v13071199

Taubman – Ben□Ari, O., Weiss, E., Abu□Sharkia, S., & Khalaf, E. (2022). A comparison of COVID-19 vaccination status among pregnant Israeli Jewish and Arab women and psychological distress among the Arab women. *Nursing & Health Sciences*, nhs.12929. https://doi.org/10.1111/nhs.12929

Taylor, S., Landry, C. A., Paluszek, M. M., Fergus, T. A., McKay, D., & Asmundson, G. J. G. (2020). Development and initial validation of the COVID Stress Scales. *Journal of Anxiety Disorders*, *72*, 102232. https://doi.org/10.1016/j.janxdis.2020.102232

UK Health Security Agency. (2021). *COVID-19 vaccine surveillance report week 47*. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1036047/Vaccine_surveillance_report_- _week_47.pdf

Vousden, N., Bunch, K., Morris, E., Simpson, N., Gale, C., O'Brien, P., Quigley, M., Brocklehurst, P., Kurinczuk, J. J., & Knight, M. (2021). The incidence, characteristics and outcomes of pregnant women hospitalized with symptomatic and asymptomatic SARS-CoV-2 infection in the UK from March to September 2020: A national cohort

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

study using the UK Obstetric Surveillance System (UKOSS). *PloS One*, *16*(5), e0251123. https://doi.org/10.1371/journal.pone.0251123

Wang, R., Liu, M., & Liu, J. (2021). The Association between Influenza Vaccination and COVID-19 and Its Outcomes: A Systematic Review and Meta-Analysis of Observational Studies. *Vaccines*, *9*(5), 529. https://doi.org/10.3390/vaccines9050529

Ward, J. K., Alleaume, C., Peretti-Watel, P., Peretti-Watel, P., Seror, V., Cortaredona, S., Launay, O., Raude, J., Verger, P., Beck, F., Legleye, S., L'Haridon, O., & Ward, J. (2020). The French public's attitudes to a future COVID-19 vaccine: The politicization of a public health issue. *Social Science & Medicine*, *265*, 113414. https://doi.org/10.1016/j.socscimed.2020.113414

W/Mariam, T. G., Kassie, B. A., Asratie, M. H., & Abate, A. T. (2021). The Effects of Fear and Knowledge of COVID-19 on Preventive Practice Among Pregnant Women Who Attend Antenatal Care in Northwest Ethiopia, 2020: Institution-Based Cross-Sectional Study. *International Journal of Women's Health*, *Volume 13*, 95–100. https://doi.org/10.2147/IJWH.S286088

Wong, L. P., Hung, C.-C., Alias, H., & Lee, T. S.-H. (2020). Anxiety symptoms and preventive measures during the COVID-19 outbreak in Taiwan. *BMC Psychiatry*, *20*(1), 376. https://doi.org/10.1186/s12888-020-02786-8

Xu, B., Gao, X., Zhang, X., Hu, Y., Yang, H., & Zhou, Y.-H. (2021). Real-World Acceptance of COVID-19 Vaccines among Healthcare Workers in Perinatal Medicine in China. *Vaccines*, *9*(7), 704. https://doi.org/10.3390/vaccines9070704

Yanez, N. D., Weiss, N. S., Romand, J.-A., & Treggiari, M. M. (2020). COVID-19 mortality risk for older men and women. *BMC Public Health*, *20*(1), 1742. https://doi.org/10.1186/s12889-020-09826-8

Yıldırım, M., Geçer, E., & Akgül, Ö. (2021). The impacts of vulnerability, perceived risk, and fear on preventive behaviours against COVID-19. *Psychology, Health & Medicine*, *26*(1), 35–43. https://doi.org/10.1080/13548506.2020.1776891

Yılmaz, M., & Sahin, M. K. (2021). Parents' willingness and attitudes concerning the COVID□19 vaccine: A cross□sectional study. *International Journal of Clinical Practice*, *75*(9). https://doi.org/10.1111/ijcp.14364

Zambrano, L. D., Ellington, S., Strid, P., Galang, R. R., Oduyebo, T., Tong, V. T., Woodworth, K. R., Nahabedian, J. F., Azziz-Baumgartner, E., Gilboa, S. M., Meaney-Delman, D., & CDC COVID-19 Response Pregnancy and Infant Linked Outcomes Team. (2020). Update: Characteristics of Symptomatic Women of Reproductive Age

medRxiv preprint doi: https://doi.org/10.1101/2022.04.06.22273526; this version posted April 10, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

with Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status - United States, January 22-October 3, 2020. *MMWR. Morbidity and Mortality Weekly Report*, *69*(44), 1641–1647. https://doi.org/10.15585/mmwr.mm6944e3

Zeynali Bujani, M., Behnampour, M., Rahimi, N., Safari, T., Khazaei Feizabad, A., Hossein Sarbazi, A., Baniasadi, M., Rezaei, N., & Ansari Moghaddam, A. (2021). The Effect of Influenza Vaccination on COVID-19 Morbidity, Severity and Mortality: Systematic Review and Meta-Analysis. *The Malaysian Journal of Medical Sciences: MJMS*, *28*(6), 20–31. https://doi.org/10.21315/mjms2021.28.6.3

Zhang, K. C., Fang, Y., Cao, H., Chen, H., Hu, T., Chen, Y. Q., Zhou, X., & Wang, Z. (2020). Parental Acceptability of COVID-19 Vaccination for Children Under the Age of 18 Years: Cross-Sectional Online Survey. *JMIR Pediatrics and Parenting*, *3*(2), e24827. https://doi.org/10.2196/24827