Triangulating Causality Between Childhood Obesity and Neurobehavior: Twin and Longitudinal Evidence

Leonard Konstantin Kulisch[a,b], Kadri Arumäe[a], Daniel A. Briley[c], Uku Vainik[a,d,e]

[a] Institute of Psychology, University of Tartu, 50409 Tartu, Estonia

[b] Leipzig University, 04109 Leipzig, Germany

[c] Department of Psychology, University of Illinois at Urbana-Champaign, Champaign, IL 61820, USA

[d] Institute of Genomics, University of Tartu, 50409 Tartu, Estonia

[e] Montreal Neurological Institute, McGill University, Montreal, QC H3A 2B4, Canada

Corresponding author: Uku Vainik ukuvainik@gmail.com

ORCID Identifiers: LKK 0000-0002-9754-0411, KA 0000-0001-9950-904X, DB 0000-0001-6344-597X, UV 0000-0002-9375-9520

Author contributions: LKK, UV designed research; LKK, UV, DB performed research; LKK analyzed data; and LKK, KA, UV wrote the paper.

The authors declare no conflict of interest.

Abstract

Childhood obesity has been linked with brain functioning and morphology. The causal directions of these relationships remain mostly untested. We filled this gap by using the Adolescent Brain Cognitive Development study cohort comprising 11,875 children aged 9-10. Cross-sectionally, body mass percentile correlated with measures of personality, psychopathology, eating behavior, cognition, cortical thickness, cortical surface area, and cortical volume. Effects were then aggregated by neurobehavioral domain for causal analyses. Direction of Causation twin modeling was used to test the direction of each relationship. Findings were validated by longitudinal analyses. Twin modeling suggested causal effects of childhood obesity on eating behavior (β=.26), cognition (β=.05), cortical thickness (β=.15), and cortical surface area (β=.07). Personality/psychopathology (β=.09) and eating behavior (β=.16) appeared to causally influence childhood obesity. Longitudinal evidence broadly supported these findings. Results regarding cortical volume were inconsistent. Childhood obesity had considerable causal effects on brain functioning and morphology, supporting the brain inflammation theory.

Main Text

 Introduction

 Childhood obesity is a growing health issue with every fifth child in the U.S. having obesity
 and these numbers expected to rise[1]. High body weight as indexed by body mass index (BMI)
 tends to persist into adulthood and is associated with individual health risks and increased
 financial burden to the health and labor system[2,3]. Since all food intake happens through
 behaviors, such as choosing and eating it[4], it is hoped that examining the association between
 obesity and neurobehavioral factors like behavior, cognition, and brain structure could aid in
 designing interventions that prevent or reduce childhood obesity.

 Many studies have found links between obesity and such neurobehavioral factors and
 obesity. Regarding self- and parent-reported traits, correlations have been found between
 pediatric weight status and mental health issues like depression[5] as well as various eating
 behaviors[6]. Child and adolescent obesity have also been found to correlate with behavioral
 and self-reported measures of impulsivity[7], motivation, and cognitive functioning[8]. Regarding
 brain morphology, associations between childhood obesity and cortical thickness have been
 found in diverse brain regions. Many studies have reported association to areas within the
 frontal lobe[9–14]. Yet, other brain regions may also play a role[15]. In participants with higher
 obesity, cortical surface area was observed to be increased in the left rostral middle frontal
 gyrus and the right superior frontal gyrus[10] while others found cortical volume to be altered in
 frontal and temporal brain regions[16–18]. Some of these associations have already been
 replicated within the Adolescent Brain Cognitive Development (ABCD) study dataset[19–25] due
 to its open accessibility.

 However, it is unclear if obesity may have causal associations with these
 neurobehavioral domains and aspects of brain morphology and, if so, what the direction of
 these associations may be. A recent study in adults showed that neurobehavioral correlates of
 obesity are largely heritable[26] and follow-up genetic analyses with two different methods
 suggest that obesity may cause variance in personality traits[27]. In childhood, obesity is
 assumed to be causally influenced by eating, lifestyle, and socio-cultural behaviors and to
 have causal effects on cognitive abilities[28] and brain morphology[10,18]. Psychopathology is
 conceptualized to have a reciprocal relationship with child weight status[28]. However, few
 studies have tested these causal assumptions. Longitudinal evidence suggests causal links
 from inhibitory control[29], reward responsiveness[30], prosociality[31], depression[32,33],
 aggression[34], and psychopathology in general[35] to childhood obesity. BMI has additionally
 been found to influence internalizing problems[36] and depression[33] suggesting a reciprocal

relationship between childhood obesity and depressive symptoms. Eating behaviors like emotional eating[37] and picky eating[38] were found to longitudinally influence childhood obesity. Longitudinal studies found a link from child weight status to academic attainment[39] but no evidence for any causal relations to intelligence[40]. Cortical surface area predicted subsequent BMI changes during adolescence[41]. An experimental study observing brain morphology changes following bariatric surgery found a an effect of childhood obesity on cortical thickness[12]. Genetically informed analyses revealed directional effects from polygenic scores for BMI to ADHD, heavy smoking[42], and overeating[43], as well as cortical volume[44]. Schizophrenia and anorexia nervosa genetic propensity were identified to predict childhood obesity[42]. Another genetically informed study found a reciprocal causal relationship between BMI and disordered eating[45].

In this study, we sought to triangulate causality in well-known correlates of childhood obesity. First, cross-sectional obesity-neurobehavior associations were validated using the large ABCD study dataset. We then applied twin and longitudinal designs to analyze these correlations from a causal perspective. Two causal inference methods with different assumptions on separate subsets of the data can aid in establishing robust causal effects[46].

Results

Personality/psychopathology and eating behavior factors

For an overview of the association between childhood obesity and personality/psychopathology, the features (i.e., items of personality and psychopathology tests) within the factor were summarized into their respective sum scales according to test manuals. In the subset of singletons, obesity was positively correlated to drive, reward responsiveness [50], social problems, somatic complaint, aggressive behavior, and being depressed [51]. A negative correlation was found with sensation seeking[52]. The strongest associations were between obesity and measures of psychopathology (Fig. 1).

Correlation analyses were conducted between the percentage of the sex- and age-specific 95th BMI percentile and the eating-specific items "overeating" and "doesn't eat well" of the Child Behavior Checklist[51] which formed the eating behavior factor. Both eating-related and parent-report items had considerable correlations with obesity. A strong positive correlation was found with "overeating" and a weaker negative correlation with "doesn't eat well" (Fig. 1). The feature "doesn't eat well" was also represented in the somatic complaints score. In following analyses, eating-specific items were excluded from the personality/psychopathology factor.

--- Fig. 1 here ---

Cognition

Cognitive test performance consistently showed negative correlations with obesity in the

singleton subset. Exceptions were inhibitory control, perception speed, and pattern

comparison[53] where no significant associations occurred. The obesity-cognition association

was strongest for the Wechsler matrix reasoning task (Fig. 1)[54].

Brain morphology factors

Analyses of singletons revealed positive associations of posterior cortical thickness in the

occipital and parietal lobe. Moreover, a positive correlation was also found with the precentral

region of interest within the frontal lobe. Negative associations between cortical thickness and

obesity were found for most regions of interest within the frontal lobe as well as superior

temporal and temporal pole regions within the temporal lobe. Higher correlations for the right

compared to the left hemisphere were discovered in some frontal and parietal brain regions

(Fig. 2).

Fewer meaningful associations were found for cortical surface area and volume.

Surface area-obesity associations were present in the isthmus cingulate, lateral orbitofrontal,

medial orbitofrontal, rostral middle frontal, insula, and temporal pole regions of interest (Fig.

2) while volume-obesity associations were found with the isthmus cingulate, rostral middle

frontal, frontal pole, insula, posterior cingulate, superior temporal, and temporal pole regions

of interest (Fig. 2). See Fig. 3 for mapped visualization of obesity-morphology correlations.

--- Fig. 2 here ---

--- Fig. 3 here ---

Poly-phenotype scores

Poly-phenotype scores (PPS)[26] were calculated to summarize the total association of each

neurobehavioral factor with obesity. Obesity-PPS correlations in the twin sample were r=.09,

p<.001, n=2469 for personality/psychopathology, r=.36, p<.001, n=2475 for eating behavior,

r=.04, p=.02, n=2296 for cognition, r=.14, p<.001, n=2391 for cortical thickness, r=.06,

p<.01, n=2391 for cortical surface area, and r=.07, p<.001, n=2391 for cortical volume (for

full correlation matrix see Supplementary Table S4). The coefficient for the

personality/psychopathology factor containing mainly mental health items is very close to the

meta-analytic estimate found for the childhood obesity-depression correlation of r=.08[5]. A

meta-analysis investigating various eating behaviors detected effects between r=|.08| and

r=|.22| with a lower estimate for emotional overeating (r=.15) than in the present study[6]. In

contrast, a previous meta-analysis of associations between childhood obesity and cognitive

domains found a greater overall effect size (r=|.24|) than this analysis[8]. A previous analysis of

obesity-PPS correlations among 895 adults yielded a similar coefficient for personality

(r=.08) but stronger associations between obesity and cognition (r=.16) as well as cortical

thickness (r=.26)[26].

Direction of Causation

Direction of Causation (DoC) twin modeling was conducted to investigate the causal paths

building the observed obesity-PPS correlations. In DoC, the relationships of heritability

variance components of two phenotypes are structurally modeled[56]. Four models were

compared: obesity to PPS, PPS to obesity, reciprocal, and Cholesky (i.e., no causal paths).

The heritability of obesity was 89.5% with the remaining 10.5% being due to unique

experiences of a co-twin. PPS heritability varied between 6.6% (eating behavior) and 57.5%

(cortical thickness, see Supplementary Tables S5-6).

Obesity was found to have unidirectional causal effects on cognition (β=.05, CI=[.00,

.09], n_{Pairs}=1056), cortical thickness (β=.15, CI=[.11, .20], n_{Pairs}=1138), and cortical surface

area (β=.07, CI=[.03, .11], n_{Pairs}=1138). Contrarily, the personality/psychopathology (β=.09,

CI=[.05, .12], n_{Pairs}=1213) and cortical volume (β=.07, CI=[.04, .11], n_{Pairs}=1138) PPS were

found to have unidirectional causal effects on obesity. The reciprocal causation model was

found the be best fitting for eating behavior ($\beta_{Obesity-to-PPS}$ =.26, $CI_{Obesity-to-PPS}$ =[.20, .32], $\beta_{PPS-to-}$

$_{Obesity}$ =.16, $CI_{PPS-to-Obesity}$ =[.11, .20], n_{Pairs}=1218).

--- Fig. 4 here ---

Cross-lagged panel models

To validate DoC results, longitudinal analyses using cross-lagged panel models (CLPM) were

performed. CLPMs estimate autoregressive paths which reflect stability of each measured

trait over time (i.e., *a* paths) by structural modeling. Moreover, paths from previous measures

of one trait to later measures of another trait are modeled. CLPMs refer to these as causal

paths or *c* paths. However, they may not truly reflect causality[61]: CLPMs test the ability of

one trait's past levels to predict later levels of another trait, but such relationships can be

driven by unobserved third variables as well as causal influences.

Using the preferred random-intercept CLPM, meaningful *c* paths were found from

childhood obesity to eating behavior (β=.11, CI=[.03, .19], n=3497) and from

personality/psychopathology to childhood obesity (β=.04, CI=[.02, .07], n=3482). Traditional

CLPMs found *c* paths from obesity to all neurobehavioral factors:

personality/psychopathology (β=.07, CI=[.04, .10], n=3416), eating behavior (β=.22, CI=[.18,

.26], n=3442), cognition (β=-.04, CI=[-.08, -.01], n=2952), cortical thickness (β=.08, CI=[.05,

.11], n=3215), cortical surface area (β=.02, CI=[.01, .03], n=3215), and cortical volume

(β=.02, CI=[.00, .03], n=3215). *C* paths also emerged from personality/psychopathology

(β=.03, CI=[.01, .05], n=3416) and eating behavior (β=.05, CI=[.02, .08], n=3442) to obesity.

Complete CLPM path estimates are shown in Supplementary Table S8.

The directional relationships found in DoC twin modeling were fully replicated except

causal effects of cortical volume and eating behavior (only traditional CLPM) on childhood

obesity. Additional longitudinal paths emerged from childhood obesity to

personality/psychopathology (only traditional CLPM) and cortical volume (Fig. 4). The

negative longitudinal childhood obesity-cognition effect implies that higher percentage of the

sex- and age-specific 95[th] BMI percentile at age nine to ten is associated with higher cognitive

test scores at age eleven to twelve.

Discussion

The purpose of the present study was to use multiple causal inference methods to understand

associations between childhood obesity and neurobehavioral factors. Replicating previous

cross-sectional findings, childhood obesity at age nine to ten was found to be associated to

measures of personality/psychopathology, eating behavior, cognition, cortical thickness,

cortical surface area, and cortical volume. Regarding the direction of these relationships, the

data supported causal influences of obesity on eating behavior, cognition, cortical thickness,

and cortical surface area across analysis methods. Personality/psychopathology showed

consistent causal and longitudinal effects on obesity. A reciprocal relationship may drive the associations between obesity and eating behavior. However, a random-intercept CLPM found only longitudinal effects from obesity to eating behavior. Both reciprocal obesity-eating behavior models (i.e., DoC and traditional CLPM) implied the latter to be the stronger path, too. For the association between obesity and personality/psychopathology, only the methodologically weaker traditional CLPM suggested a reciprocal relationship. Conflicting findings leave the relationship between obesity and cortical volume unclear.

Together, the results of this multi-method study in a large sample highlight the importance of obesity as a driving force behind individual differences in brain functioning and morphology during childhood.

In recent studies, deficits in cognitive abilities of obese adults are attributed to an obesity associated inflammation in the brain[62]. Even though less pronounced, this effect can be partly observed in children, too[63]. The beginning inflammation in brains of obese children may explain the small effect from obesity to cognition in children aged 9-10. Obesity-triggered inflammation may explain accelerated brain aging in obese adults. This results in neurodegeneration which can be observed in cortical thinning. Albeit the effect being most pronounced in adulthood, similar degeneration processes may drive the effect of obesity on cortical thickness and surface area in the present study[64]. Reduced cortical thickness or surface area in frontal brain areas associated with executive functioning could mediate effects of obesity on several tests of cognition.

However, a longitudinal correlation from increased obesity to better cognitive functioning was observed using a traditional CLPM. The previous negative association disappeared at age 11-12. Changes in the association between obesity and intelligence across childhood were previously described implying that negative effects of obesity can potentially be compensated in certain developmental periods[65]. However, the positive effect could not be replicated across methods. Further research is necessary to map this effect.

On the other hand, potential self- and parent-reported contributors to childhood obesity (i.e., eating behavior, psychopathology, impulsivity, and motivation) were identified. Of them, strongest effects could be summarized as increased approach behavior (sensation-seeking, drive, reward responsiveness), which may lead to unhealthy food choices that maximize short term satisfaction[66]. Besides food quality, also food quantity could be affected. The reduced ability to stop eating when hunger is satisfied may lead to overeating[67]. The other domain seems to be stress-related psychopathology. Children with increased mental health problems may attempt to regulate adverse emotions by stress eating, which may contribute to

weight gain[68]. Current evidence favors focusing on interventions aiding reducing stress and/or

try to manage increased sensation seeking.

Eating behavior was another causal contributor to obesity, which seems trivial. On the

other hand, the reverse effect of obesity on eating behavior was even stronger. While rarely

considered, such association is consistent with the model, where obesity increases fat free

mass which in turn increases food intake, which then maintains obesity[69]. Fat-free mass and

food intake have been linked in adolescents[70], and genetic obesity predicts future childhood

eating behaviors[43]. Therefore, studies intervening on eating behaviors should also consider

reverse causality.

The results are limited by the use of %BMI_{p95} as a proxy for obesity and the

assumption of obesity being continuous while the majority of participants were not obese. We

explored relative fat mass as an alternative obesity indicator but discarded due to lack of

additional information[71]. The PPS method limits the results by ignoring causal relationships

of individual features within a factor, but this was offset by gain in statistical power. DoC

results are influenced by measurement reliability in a way that DoC tends to model the more

reliable variable to have causal effects on the less reliable variable. Reliability could be lower

for behavioral[72] and cognitive factors[73] which could contribute to the finding that obesity had

causal effects on eating behavior and cognition. Personality/psychopathology and cognitive

test results can be compromised by intrapersonal components like response bias or motivation

which are irrelevant for the directly observed measures of childhood obesity and brain

morphology. At the same time, previous analyses showed that explicitly modeling reliability

did not change DoC conclusions[27]. Future studies using different causal inference methods

should build complementary evidence[27].

In summary, the present study provides evidence that childhood obesity may influence

rather than be influenced by the various aspects of brain functioning and morphology that

have been found to differ between children with and without obesity. Therefore, obesity may

play an important role in brain development during childhood. The present findings are hoped

to inspire further investigation of the underlaying mechanisms behind these associations and

development of more effective obesity interventions.

Methods

Participants

The baseline, one-year and two-year follow-ups of the ABCD study[47] were used for data

analysis. The study cohort includes 11875 children aged 9-10 years at baseline. After applying

typical exclusion criteria to the available cases, the final sample comprised 7016 singletons

and 2511 siblings (for exclusion criteria see Supplementary Table S1; for demographics see

Supplementary Table S2). Singletons were participants with unique family-IDs. Siblings were

participants who had the same family-ID as one other participant within study waves and had

information about their respective relationship available. Singleton data were used for cross-

sectional analyses of associations between childhood obesity and neurobehavioral indicators

as well as longitudinal analyses of those relationships. Sibling data were the basis for behavior

genetics analyses.

Measures

The percentage of the sex- and age-specific 95th BMI percentile was calculated for each

participant following guidelines for childhood obesity research[48]. BMI is often used in

research on obesity in adulthood, but the dependency of its distribution on sex and age during

childhood renders it impractical for childhood obesity studies. For grouping of different

neurobehavioral aspects, each variable of interest recorded in the ABCD study (called *feature*)

was assigned to one of six neurobehavioral factors. These were personality/psychopathology,

eating behavior, cognition, cortical thickness, cortical surface area, and cortical volume. All

analyses are controlled for gender at birth, age, race, Hispanic ethnicity, handedness, puberty

status and study site. Additionally, in analyses with brain morphology measures the individual

mean or total of the respective measure was controlled for. This procedure was chosen to

maintain consistency regarding controls for brain measures. Multiple control variables were

correlated to childhood obesity (see Supplementary Figure S1). Brain morphology measures

needed to successfully pass ABCD quality control. False discovery rate correction[49] was

applied when screening for features within a neurobehavioral factor. See Supplementary

Table S3 and https://nda.nih.gov/general-query.html for further information on the measures.

Poly-phenotype scores

The PPS method is adapted from polygenic scores and helps to reduce measurement noise as

well as multiple testing of redundant measures [e.g., 55]. PPSs are calculated by summing the

products of each feature's measure (i.e., personality/psychopathology/eating behavior item

scores, cognitive test scores, and thickness in mm/surface area in mm² /volume in mm³ of

cortical regions of interest) with its correlation with the sex- and age-specific 95[th] BMI

percentile within a factor. The personality/psychopathology behavior factor used personality

and psychopathology test items (without eating/weight-related items) and their specific

correlation opposed to the sum scales used for preliminary correlation analyses. To avoid

overfitting, correlations were obtained from the singleton subset and applied to measures

within the subset of siblings to generate six PPSs for each participant in the sibling subset.

These PPSs informed preliminary correlation and twin modeling analyses. Ten-fold cross-

validation within the singleton sample was later used to generate PPSs in an independent

sample for longitudinal analysis (see Supplementary Table S3).

Twin modeling

DoC twin modeling builds on the ACE heritability model which decomposes observed traits

into additive genetic (A), common environment (C), and unique environment (E) variance

components by comparing mono- and dizygotic twin pairs (non-twin siblings were treated

equal to dizygotic twins)[57].If one trait has a causal influence another trait, then the A, C, and E

variance components of this first trait should be proportionally represented in the variance of

the second. In this way, DoC models can be specified for obesity causally influencing PPS, or

PPS causally influencing obesity. A reciprocal model can be defined by allowing both causal

paths. Removing any causal paths leads to the Cholesky model where observed correlations

are driven by a third variable[56]. For each obesity-PPS association, the best-fitting model was

chosen by comparing model fit with regards to model complexity using Chi-squared and

Akaike information criterion (see Supplementary Table S7). For both, lower values indicate

better model fit.

Longitudinal analyses

Traditional CLPMs[58] were calculated for all six PPSs using baseline and two-year follow-up

measures. Random-intercept CLPMs[59,60] were applied where PPSs could be calculated for

baseline, one-year and two-year follow-up (i.e., personality/psychopathology and eating

behavior). The methodological advantage of Random-intercept CLPMs lays in the estimation

of latent variables to account for stable interindividual trait differences[59]. However, this

method requires at least three observation waves. Due to varying availability of measures per

study wave, PPSs for CLPM may not be fully congruent with PPSs for DoC (see

Supplementary Table S3).

Data availability

ABCD study data is available at https://abcdstudy.org/.

Code availability

A commented analysis script is available at osf.io/zgda4 for detailed data analysis procedure

insight and replication. Analysis software information is noted in Supplementary Table S9.

Acknowledgments

Uku Vainik has been funded by Estonian Research Council's personal research funding start-

up grant PSG759.

References

1.   Wang, Y. *et al.* Has the prevalence of overweight, obesity and central obesity levelled

off in the United States? Trends, patterns, disparities, and future projections for the

obesity epidemic. *Int. J. Epidemiol.* **49**, 810–823 (2020).

2.   Llewellyn, A., Simmonds, M., Owen, C. G. & Woolacott, N. Childhood obesity as a

predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obes. Rev.*

17, 56–67 (2016).

3.   Hamilton, D., Dee, A. & Perry, I. J. The lifetime costs of overweight and obesity in

childhood and adolescence: a systematic review. *Obes. Rev.* **19**, 452–463 (2018).

4.   Blundell, J. E. & Cooling, J. Routes to obesity: phenotypes, food choices and activity.

Br. J. Nutr. **83 Suppl 1**, S33-8 (2000).

5.   Sutaria, S., Devakumar, D., Yasuda, S. S., Das, S. & Saxena, S. Is obesity associated

with depression in children? Systematic review and meta-analysis. *Arch. Dis. Child.*

104, 64–74 (2019).

6.   Kininmonth, A. *et al.* The association between childhood adiposity and appetite

assessed using the Child Eating Behavior Questionnaire and Baby Eating Behavior

Questionnaire: A systematic review and meta-analysis. *Obes. Rev.* **22**, (2021).

7.   Thamotharan, S., Lange, K., Zale, E. L., Huffhines, L. & Fields, S. The role of

impulsivity in pediatric obesity and weight status: A meta-analytic review. *Clin.*

Psychol. Rev. **33**, 253–262 (2013).

8.   Pearce, A. L., Leonhardt, C. A. & Vaidya, C. J. Executive and Reward-Related

Function in Pediatric Obesity: A Meta-Analysis. *Child. Obes.* **14**, 265–279 (2018).

9.   Saute, R. L., Soder, R. B., Alves Filho, J. O., Baldisserotto, M. & Franco, A. R.

Increased brain cortical thickness associated with visceral fat in adolescents. *Pediatr.*

Obes. **13**, 74–77 (2018).

10.   Prats-Soteras, X. *et al.* Inflammatory agents partially explain associations between

cortical thickness, surface area, and body mass in adolescents and young adulthood.

Int. J. Obes. **44**, 1487–1496 (2020).

11.   Kim, M. S. *et al.* Prefrontal Cortex and Amygdala Subregion Morphology Are

Associated With Obesity and Dietary Self-control in Children and Adolescents. *Front.*

Hum. Neurosci. **14**, 563415 (2020).

12.   Rajan, L. *et al.* Effects of severe obesity and sleeve gastrectomy on cortical thickness in

adolescents. *Obesity* **29**, 1516–1525 (2021).

13.   Thapaliya, G. *et al.* Familial Obesity Risk and Current Excess Weight Influence Brain

Structure in Adolescents. *Obesity (Silver Spring)*. **29**, 184–193 (2021).

14.   Ross, N., Yau, P. L. & Convit, A. Obesity, fitness, and brain integrity in adolescence.
Appetite **93**, 44–50 (2015).

15.   Steegers, C. *et al.* The association between body mass index and brain morphology in
children: a population-based study. *Brain Struct. Funct.* **226**, 787–800 (2021).

16.   Kennedy, J. T., Collins, P. F. & Luciana, M. Higher Adolescent Body Mass Index Is
Associated with Lower Regional Gray and White Matter Volumes and Lower Levels of
Positive Emotionality. *Front. Neurosci.* **10**, (2016).

17.   Hashimoto, T. *et al.* Increased posterior hippocampal volumes in children with lower
increase in body mass index: a 3-year longitudinal MRI study. *Dev. Neurosci.* **37**, 153–
60 (2015).

18.   Alosco, M. L. *et al.* Body mass index and brain structure in healthy children and
adolescents. *Int. J. Neurosci.* **124**, 49–55 (2014).

19.   Laurent, J. S. *et al.* Associations Among Body Mass Index, Cortical Thickness, and
Executive Function in Children. *JAMA Pediatr.* **174**, 170–177 (2020).

20.   Dennis, E., Manza, P. & Volkow, N. D. Socioeconomic status, BMI, and brain
development in children. *Transl. Psychiatry* **12**, 33 (2022).

21.   Ronan, L., Alexander-Bloch, A. & Fletcher, P. C. Childhood Obesity, Cortical
Structure, and Executive Function in Healthy Children. *Cereb. Cortex* **30**, 2519–2528
(2020).

22.   Bohon, C. & Welch, H. Quadratic relations of BMI with depression and brain volume
in children: Analysis of data from the ABCD study. *J. Psychiatr. Res.* **136**, 421–427
(2021).

23.   Hall, P. A., Best, J., Beaton, E. A., Sakib, M. N. & Danckert, J. Morphology of the
Prefrontal Cortex Predicts Body Composition in Early Adolescence: Cognitive
Mediators and Environmental Moderators in the ABCD Study. *Soc. Cogn. Affect.*
Neurosci. (2021) doi:10.1093/scan/nsab104.

24.   Adise, S. *et al.* Multimodal brain predictors of current weight and weight gain in
children enrolled in the ABCD study ®. *Dev. Cogn. Neurosci.* **49**, (2021).

25.   Gray, J. C., Schvey, N. A. & Tanofsky-Kraff, M. Demographic, psychological,
behavioral, and cognitive correlates of BMI in youth: Findings from the Adolescent
Brain Cognitive Development (ABCD) study. *Psychol. Med.* **50**, 1539–1547 (2020).

26.   Vainik, U. *et al.* Neurobehavioral correlates of obesity are largely heritable. *Proc. Natl.*
Acad. Sci. **115**, 9312–9317 (2018).

27. Arumäe, K. *et al.* Two genetic analyses to elucidate causality between body mass index
and personality. *Int. J. Obes. (Lond).* **45**, 2244–2251 (2021).

28. Sahoo, K. *et al.* Childhood obesity: causes and consequences. *J. Fam. Med. Prim. care*
4, 187–92 (2015).

29. Anzman-Frasca, S., Francis, L. A. & Birch, L. L. Inhibitory control is associated with
psychosocial, cognitive, and weight outcomes in a longitudinal sample of girls. *Transl.*
Issues Psychol. Sci. **1**, 203–216 (2015).

30. De Decker, A. *et al.* Fat and lean tissue accretion in relation to reward motivation in
children. *Appetite* **108**, 317–325 (2017).

31. Christensen, K. G. *et al.* Child behaviour and subsequent changes in body weight,
composition and shape. *PLoS One* **14**, e0226003 (2019).

32. Blaine, B. Does Depression Cause Obesity? *J. Health Psychol.* **13**, 1190–1197 (2008).

33. Korczak, D. J., Lipman, E., Morrison, K. & Szatmari, P. Are children and adolescents
with psychiatric illness at risk for increased future body weight? A systematic review.
Dev. Med. Child Neurol. **55**, 980–987 (2013).

34. Derks, I. P. M. *et al.* Testing Bidirectional Associations Between Childhood
Aggression and BMI: Results from Three Cohorts. *Obesity* oby.22419 (2019)
doi:10.1002/oby.22419.

35. Camfferman, R. *et al.* The association between overweight and internalizing and
externalizing behavior in early childhood. *Soc. Sci. Med.* **168**, 35–42 (2016).

36. Bradley, R. H. *et al.* The Relationship between Body Mass Index and Behavior in
Children. *J. Pediatr.* **153**, 629-634.e3 (2008).

37. Shriver, L. H. *et al.* Longitudinal Associations between Emotion Regulation and
Adiposity in Late Adolescence: Indirect Effects through Eating Behaviors. *Nutrients*
11, (2019).

38. Antoniou, E. E. *et al.* Picky eating and child weight status development: a longitudinal
study. *J. Hum. Nutr. Diet.* **29**, 298–307 (2016).

39. Booth, J. N. *et al.* Obesity impairs academic attainment in adolescence: findings from
ALSPAC, a UK cohort. *Int. J. Obes. (Lond).* **38**, 1335–42 (2014).

40. Kase, B. E. *et al.* Longitudinal Associations Between Symptoms of ADHD and BMI
From Late Childhood to Early Adulthood. *Pediatrics* **147**, (2021).

41. Park, B.-Y., Chung, C.-S., Lee, M. J. & Park, H. Accurate neuroimaging biomarkers to
predict body mass index in adolescents: a longitudinal study. *Brain Imaging Behav.* **14**,
1682–1695 (2020).

42.   Hübel, C. *et al.* Genetic correlations of psychiatric traits with body composition and
443   glycemic traits are sex- and age-dependent. *Nat. Commun.* **10**, 5765 (2019).

444   43.   Herle, M. *et al.* The genomics of childhood eating behaviours. *Nat. Hum. Behav.* **5**,
445   625–630 (2021).

446   44.   Lugo-Candelas, C. *et al.* Differences in brain structure and function in children with the
447   FTO obesity-risk allele. *Obes. Sci. Pract.* **6**, 409–424 (2020).

448   45.   Reed, Z. E., Micali, N., Bulik, C. M., Davey Smith, G. & Wade, K. H. Assessing the
449   causal role of adiposity on disordered eating in childhood, adolescence, and adulthood:
450   a Mendelian randomization analysis. *Am. J. Clin. Nutr.* **106**, 764–772 (2017).

451   46.   Munafò, M. R. & Davey Smith, G. Robust research needs many lines of evidence.
452   *Nature* **553**, 399–401 (2018).

453   47.   Volkow, N. D. *et al.* The conception of the ABCD study: From substance use to a
454   broad NIH collaboration. *Dev. Cogn. Neurosci.* **32**, 4–7 (2018).

455   48.   Freedman, D. S. *et al.* BMI z -Scores are a poor indicator of adiposity among 2- to 19-
456   year-olds with very high BMIs, NHANES 1999-2000 to 2013-2014. *Obesity* **25**, 739–
457   746 (2017).

458   49.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
459   Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

460   50.   Carver, C. S. & White, T. L. Behavioral inhibition, behavioral activation, and affective
461   responses to impending reward and punishment: The BIS/BAS Scales. *J. Pers. Soc.*
462   *Psychol.* **67**, 319–333 (1994).

463   51.   Achenbach, T. M. & Edelbrock, C. S. *Manual for the Child Behavior Checklist and*
464   *Revised Child Behavior Profile*. (University Associates in Psychiatry, 1983).

465   52.   Cyders, M. A., Littlefield, A. K., Coffey, S. & Karyadi, K. A. Examination of a short
466   English version of the UPPS-P Impulsive Behavior Scale. *Addict. Behav.* **39**, 1372–
467   1376 (2014).

468   53.   Ratcliff, G. Spatial thought, mental rotation and the right cerebral hemisphere.
469   *Neuropsychologia* **17**, 49–54 (1979).

470   54.   Wechsler, D. *Wechsler Intelligence Scale for Children*. (Psychological Corporation,
471   1949).

472   55.   Steinsbekk, S., Belsky, D., Guzey, I. C., Wardle, J. & Wichstrøm, L. Polygenic Risk,
473   Appetite Traits, and Weight Gain in Middle Childhood. *JAMA Pediatr.* **170**, e154472
474   (2016).

56.   Heath, A. C. *et al.* Testing hypotheses about direction of causation using cross-

sectional family data. *Behav. Genet.* **23**, 29–50 (1993).

57.   Neale, M. C. & Cardon, L. R. *Methodology for genetic studies of twins and families.*
(Kluwer Academic Publishers, 1992).

58.   Bentler, P. M. & Speckart, G. Attitudes "cause" behaviors: A structural equation
analysis. *J. Pers. Soc. Psychol.* **40**, 226–238 (1981).

59.   Hamaker, E. L., Kuiper, R. M. & Grasman, R. P. P. P. A critique of the cross-lagged
panel model. *Psychol. Methods* **20**, 102–116 (2015).

60.   Mund, M. & Nestler, S. Beyond the Cross-Lagged Panel Model: Next-generation
statistical tools for analyzing interdependencies across the life course. *Adv. Life Course*
Res. **41**, 100249 (2019).

61.   Usami, S., Murayama, K. & Hamaker, E. L. A unified framework of longitudinal
models to examine reciprocal relations. *Psychol. Methods* **24**, 637–657 (2019).

62.   Spyridaki, E. C., Avgoustinaki, P. D. & Margioris, A. N. Obesity, inflammation and
cognition. *Curr. Opin. Behav. Sci.* **9**, 169–175 (2016).

63.   Adelantado-Renau, M. *et al.* Inflammatory biomarkers and brain health indicators in
children with overweight and obesity: The ActiveBrains project. *Brain. Behav. Immun.*
81, 588–597 (2019).

64.   Ronan, L. *et al.* Obesity associated with increased brain age from midlife. *Neurobiol.*
Aging **47**, 63–70 (2016).

65.   Wraw, C., Deary, I. J., Der, G. & Gale, C. R. Maternal and offspring intelligence in
relation to BMI across childhood and adolescence. *Int. J. Obes. (Lond).* **42**, 1610–1620
(2018).

66.   Zhou, Z., SooHoo, M., Zhou, Q., Perez, M. & Liew, J. Temperament as Risk and
Protective Factors in Obesogenic Eating: Relations Among Parent Temperament, Child
Temperament, and Child Food Preference and Eating. *J. Genet. Psychol.* **180**, 75–79
(2019).

67.   van den Berg, L. *et al.* Association between impulsivity, reward responsiveness and
body mass index in children. *Int. J. Obes.* **35**, 1301–1307 (2011).

68.   Hill, D. C., Moss, R. H., Sykes-Muskett, B., Conner, M. & O'Connor, D. B. Stress and
eating behaviors in children and adolescents: Systematic review and meta-analysis.
Appetite **123**, 14–22 (2018).

69.   Blundell, J. Behaviour, energy balance, obesity and capitalism. *Eur. J. Clin. Nutr.* **72**,
1305–1309 (2018).

70.   Vainik, U. *et al.* Diet misreporting can be corrected: confirmation of the association

between energy intake and fat-free mass in adolescents. *Br. J. Nutr.* **116**, 1425–1436
(2016).

71. Arumäe, K., Mõttus, R. & Vainik, U. Beyond BMI: Personality traits' associations
with adiposity and metabolic rate. *Physiol. Behav.* **246**, 113703 (2022).

72. Nakamura, B. J., Ebesutani, C., Bernstein, A. & Chorpita, B. F. A Psychometric
Analysis of the Child Behavior Checklist DSM-Oriented Scales. *J. Psychopathol.*
Behav. Assess. **31**, 178–189 (2009).

73. Taylor, B. K. *et al.* Reliability of the NIH toolbox cognitive battery in children and
adolescents: a 3-year longitudinal examination. *Psychol. Med.* 1–10 (2020)
doi:10.1017/S0033291720003487.

 Figures

Fig. 1 | Childhood obesity-behavior and childhood obesity-cognition associations. A, Associations between the percentage of the sex- and age-specific 95th BMI percentile and personality scales, psychopathology scales, as well as eating-specific items of the Child Behavior Checklist. Impulsivity, motivation and prosociality were self-reported, psychopathology (including eating items) parent-reported. **B**, Associations between the percentage of the sex- and age-specific 95th BMI percentile and cognitive test scores. Error bars represent 95% confidence intervals. EB, eating behavior; EF, executive functioning; FDR, false discovery rate; Lang, Language; Perc, perception; PS, prosociality; WM, working memory.

Fig. 2. | Childhood obesity-brain morphology associations. A-C, Associations between

percentage of the sex- and age-specific 95th BMI percentile and cortical thickness (A), cortical

surface area (B), and cortical volume (C). Error bars represent 95% confidence intervals.

FDR, false discovery rate; I, insula.

Fig. 3. | Mapped childhood obesity-brain morphology associations. A-C, Associations between percentage of the sex- and age-specific 95th BMI percentile and (A) cortical thickness, (B) cortical surface area, (C) and cortical volume mapped to a brain model.

Fig. 4. | Causal and longitudinal path coefficients. Estimates for Direction of Causation (DoC), traditional cross-lagged panel models (CLPM) and random-intercept cross-lagged panel models (RI-CLPM). Only best fitting model plotted for DoC. Error bars represent 95% confidence intervals. PPS, poly-phenotype score.