

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

## Triangulating Causality Between Childhood Obesity and Neurobehavior: Twin and Longitudinal Evidence

Leonard Konstantin Kulisch<sup>a,b</sup>, Kadri Arumäe<sup>a</sup>, D. A. Briley<sup>c</sup>, Uku Vainik<sup>a,d,e</sup>

<sup>a</sup> Institute of Psychology, University of Tartu, 50409 Tartu, Estonia

<sup>b</sup> Leipzig University, 04109 Leipzig, Germany

<sup>c</sup> Department of Psychology, University of Illinois at Urbana-Champaign, Champaign, IL 61820, USA

<sup>d</sup> Institute of Genomics, University of Tartu, 50409 Tartu, Estonia

<sup>e</sup> Montreal Neurological Institute, McGill University, Montreal, QC H3A 2B4, Canada

### Author Note

Leonard Konstantin Kulisch  <https://orcid.org/0000-0002-9754-0411>

Kadri Arumäe  <https://orcid.org/0000-0001-9950-904X>

D. A. Briley  <https://orcid.org/0000-0001-6344-597X>

Uku Vainik  <https://orcid.org/0000-0002-9375-9520>

The authors declare no conflict of interest.

Uku Vainik has been funded by Estonian Research Council's personal research funding start-up grant PSG759.

Analysis script <https://osf.io/zgda4/>

Preprint <https://www.medrxiv.org/content/10.1101/2022.04.12.22273769v1>

Correspondence concerning this article should be addressed to Uku Vainik, Institute of Psychology, Faculty of Social Sciences, University of Tartu, Estonia, Näituse 2-216, 50409, Tartu, Estonia

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Email: [uku.vainik@ut.ee](mailto:uku.vainik@ut.ee)

27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

## Abstract

**Objective:** Childhood obesity is a serious health concern that is not yet fully understood. Previous research has linked obesity with neurobehavioral factors such as behavior, cognition, and brain morphology. The causal directions of these relationships remain mostly untested.

**Methods:** We filled this gap by using the Adolescent Brain Cognitive Development study cohort comprising 11,875 children aged 9–10. First, correlations between body mass percentile and neurobehavioral measures were cross-sectionally analyzed. Effects were then aggregated by neurobehavioral domain for causal analyses. Direction of Causation twin modeling was used to test the direction of each relationship. Findings were validated by longitudinal cross-lagged panel modeling. **Results:** Body mass percentile correlated with measures of impulsivity, motivation, psychopathology, eating behavior, and cognitive tests (executive functioning, language, memory, perception, working memory). Higher obesity was also associated with reduced cortical thickness in areas of the frontal and temporal lobe but with increased thickness in parietal and occipital brain areas. Similar although weaker patterns emerged for cortical surface area and volume. Twin modeling suggested causal effects of childhood obesity on eating behavior ( $\beta=.26$ ), cognition ( $\beta=.05$ ), cortical thickness ( $\beta=.15$ ), and cortical surface area ( $\beta=.07$ ). Personality/psychopathology ( $\beta=.09$ ) and eating behavior ( $\beta=.16$ ) appeared to causally influence childhood obesity. Longitudinal evidence broadly supported these findings. Results regarding cortical volume were inconsistent. **Conclusions:** Results supported causal effects of obesity on brain functioning and morphology, consistent with effects of obesity-related brain inflammation on cognition. The present study highlights the importance of physical health for brain development during childhood and may inform interventions aimed at preventing or reducing pediatric obesity.

**Keywords:** childhood obesity, causality, twin modeling, cross-lagged panel modeling, neurobehavior

## 53 **Introduction**

54 Childhood obesity is a growing health issue with every fifth child in the U.S. having obesity  
55 and these numbers expected to rise (Wang et al., 2020). High body weight as indexed by body  
56 mass index (BMI) tends to persist into adulthood and is associated with individual health risks  
57 and increased financial burden to the health and labor system (Hamilton et al., 2018; Llewellyn  
58 et al., 2016). Since all food intake happens through behaviors, such as choosing and eating it  
59 (Blundell & Finlayson, 2004), it is hoped that examining the association between obesity and  
60 neurobehavioral factors like behavior, cognition, and brain structure could aid in designing  
61 interventions that prevent or reduce childhood obesity.

62 Many studies have found links between such neurobehavioral factors and obesity.  
63 Regarding self- and parent-reported traits, correlations have been found between pediatric  
64 weight status and mental health issues like depression (Sutaria et al., 2019) as well as various  
65 eating behaviors (Kininmonth et al., 2021). Child and adolescent obesity have also been found  
66 to correlate with behavioral and self-reported measures of impulsivity (Thamotharan et al.,  
67 2013), motivation, and cognitive functioning (Pearce et al., 2018). Regarding brain  
68 morphology, associations between childhood obesity and cortical thickness have been found in  
69 diverse brain regions. Many studies have reported association to areas within the frontal lobe  
70 (Kim et al., 2020; Prats-Soteras et al., 2020; Rajan et al., 2021; Ross et al., 2015; Saute et al.,  
71 2018; Thapaliya et al., 2021), but other brain regions may also play a role (Steegers et al., 2021).  
72 In participants with higher obesity, cortical surface area was observed to be increased in the left  
73 rostral middle frontal gyrus and the right superior frontal gyrus (Prats-Soteras et al., 2020) while  
74 others found cortical volume to be altered in frontal and temporal brain regions (Alosco et al.,  
75 2014; Hashimoto et al., 2015; Kennedy et al., 2016).

76 However, it is unclear if obesity may have causal associations with these  
77 neurobehavioral domains and aspects of brain morphology and, if so, what the direction of these  
78 associations may be. A recent study in adults showed that neurobehavioral correlates of obesity

79 are largely heritable (Vainik et al., 2018) and follow-up genetic analyses with two different  
80 methods suggest that obesity may cause variance in personality traits (Arumäe et al., 2021). In  
81 childhood, obesity is assumed to be causally influenced by eating, lifestyle, and socio-cultural  
82 behaviors and to have causal effects on cognitive abilities (Sahoo et al., 2015) and brain  
83 morphology (Alosco et al., 2014; Prats-Soteras et al., 2020). Psychopathology is conceptualized  
84 to have a reciprocal relationship with child weight status (Sahoo et al., 2015). However, few  
85 studies have tested these causal assumptions.

86 Genetically informed analyses revealed directional effects from polygenic scores for  
87 BMI to ADHD (Hübel et al., 2019) and overeating (Herle et al., 2021), as well as cortical  
88 volume (Lugo-Candelas et al., 2020). Anorexia nervosa genetic propensity was identified to  
89 predict childhood obesity (Hübel et al., 2019). Another genetically informed study found a  
90 reciprocal causal relationship between BMI and disordered eating (Reed et al., 2017).

91 Longitudinal evidence suggests causal links from BMI to internalizing problems  
92 (Bradley et al., 2008) and depression (Korczak et al., 2013). Additionally, inhibitory control  
93 (Anzman-Frasca et al., 2015), reward responsiveness (De Decker et al., 2017), prosociality  
94 (Christensen et al., 2019), depression (Blaine, 2008; Korczak et al., 2013), aggression (Derks  
95 et al., 2019), psychopathology in general (Camfferman et al., 2016), emotional eating (Shriver  
96 et al., 2019), picky eating (Antoniou et al., 2016), and cortical surface area (Park et al., 2020)  
97 were found to longitudinally influence childhood obesity. Therefore, a reciprocal relationship  
98 between childhood obesity and depressive symptoms seems plausible. Longitudinal studies  
99 found no evidence for any causal relations between BMI and intelligence (Belsky et al., 2013).

100 In this study, we sought to triangulate causality between obesity and its well-known  
101 correlates to gain further understanding about potential causes and consequences of childhood  
102 obesity. First, we aimed to validate previous cross-sectional obesity–neurobehavior  
103 associations using a large population-based study dataset. Childhood obesity was expected to  
104 correlate with impulsivity, motivation, prosociality, psychopathology, eating behavior,

105 cognition, cortical thickness, cortical surface area, and cortical volume. The present  
106 investigation then aimed to shed light on the causal relationships driving these correlations.  
107 Twin and longitudinal designs were chosen for that task. Two causal inference methods with  
108 different assumptions on separate subsets of the data can aid in establishing robust causal effects  
109 (Munafò & Davey Smith, 2018). Based on previous findings, childhood obesity was expected  
110 to influence cortical volume while cortical surface area was expected to influence obesity.  
111 Literature suggested a reciprocal relationship between childhood obesity and behavioral  
112 measures such as personality/psychopathology and eating behavior. We had no expectations  
113 regarding the direction of the association with cognition and cortical thickness as causal  
114 associations have not been previously reported between them. The novelty of the present study  
115 lies in the usage of a large population-based dataset and the simultaneous consideration of a  
116 broad range of brain functioning and structure measures. No previous study has attempted to  
117 apply two causal methods to produce strong causal evidence for the relationships between  
118 childhood obesity and its neurobehavioral correlates.

119

120

## Methods

### *Participants*

121 We used data from the baseline, one-year and two-year follow-ups of the Adolescent Brain  
122 Cognitive Development (ABCD) study (Volkow et al., 2018). The study cohort includes 11875  
123 children aged 9–10 years at baseline. After applying typical exclusion criteria to the available  
124 cases, the final sample comprised 7016 singletons and 2511 siblings (for exclusion criteria see  
125 Supplementary Table S1). Singletons were participants with unique family-IDs. Siblings were  
126 participants who had the same family-ID as one other participant within study waves and had  
127 information about their respective relationship available. Among singletons, the average age at  
128 baseline was 118.6 months (SD=7.3). 52.2% of singletons were boys, 47.7% were girls, and  
129 another gender was reported for 0.1%. Within the sibling subset, the average age at baseline  
130

131 was 119.8 months (SD=7.9). 51.3% of siblings were boys and 48.7% were girls (for  
132 demographics see Supplementary Table S2). Singleton data were used for cross-sectional  
133 analyses of associations between childhood obesity and neurobehavioral indicators as well as  
134 longitudinal analyses of those relationships. Sibling data were the basis for twin modeling.

135 The ABCD study dataset is a popular resource for scientist interested in behavioral  
136 neuroscience. Due to its open accessibility, some of the obesity–neurobehavior correlations  
137 being tested here have already been replicated within the dataset (Adise et al., 2021; Bohon &  
138 Welch, 2021; Dennis et al., 2022; Gray et al., 2020; Hall et al., 2021; Laurent et al., 2020;  
139 Ronan et al., 2020).

140

#### 141 *Measures*

##### 142 *Obesity*

143 The percentage of the sex- and age-specific 95<sup>th</sup> BMI percentile was calculated for each  
144 participant following guidelines for childhood obesity research (Freedman et al., 2017).  
145 Reference data was obtained from the US Centers for Disease Control and Prevention  
146 (<http://www.cdc.gov/growthcharts/>). All analyses used this continuous measure of obesity.  
147 Participants were considered having obesity when the percentage of the sex- and age-specific  
148 95<sup>th</sup> BMI percentile was 100 or greater. BMI is often used in research on obesity in adulthood,  
149 but the dependency of its distribution on sex and age during childhood renders it impractical  
150 for childhood obesity studies.

151

##### 152 *Neurobehavior*

153 To group different neurobehavioral aspects, each variable of interest recorded in the ABCD  
154 study (called *feature*) was assigned to one of six neurobehavioral factors. These were  
155 personality/psychopathology, eating behavior, cognition, cortical thickness, cortical surface  
156 area, and cortical volume. The personality/psychopathology factor included 159 self- and

157 parent-reported questionnaire items measuring impulsivity (Cyders et al., 2014), motivation  
158 (Carver & White, 1994), prosociality (Goodman, 1997), and psychopathology (Achenbach &  
159 Edelbrock, 1983). The eating behavior factor comprised two parent-reported eating items  
160 (Achenbach & Edelbrock, 1983). Thirteen cognitive test scores built the cognition factor  
161 (Lezak, 1976; Ratcliff, 1979; Wechsler, 1949; Weintraub et al., 2013). Cortical thickness,  
162 surface area, and volume included the respective measure in 68 regions of interest obtained  
163 from MRI scans. Brain morphology measures needed to successfully pass internal ABCD  
164 quality control. See Supplementary Table S3 and <https://nda.nih.gov/general-query.html> for  
165 detailed information on the measures.

166

#### 167 *Statistical analyses*

168 All analyses controlled for gender at birth, age, race, Hispanic ethnicity, handedness, puberty  
169 status and study site. Additionally, in analyses with brain morphology measures the individual  
170 mean or total of the respective measure was controlled for. This procedure was chosen to  
171 maintain consistency regarding controls for brain measures. Multiple control variables were  
172 correlated to childhood obesity (see Supplementary Figure S1). False discovery rate correction  
173 (Benjamini & Hochberg, 1995) was applied when screening for features within a  
174 neurobehavioral factor.

175

#### 176 *Correlational analyses within neurobehavioral factors*

177 To validate previously found obesity–neurobehavior associations, cross-sectional correlations  
178 between childhood obesity and each neurobehavioral feature were calculated within the  
179 singleton subset. For an overview, the features within the personality/psychopathology factor  
180 (i.e., items of personality and psychopathology questionnaires) were summarized into their  
181 respective sum scales according to test manuals.

182

183 *Poly-phenotype scores*

184 Poly-phenotype scores (PPS; Vainik et al., 2018) were calculated to summarize the total  
185 association of each neurobehavioral factor with obesity. The PPS method is adapted from  
186 polygenic scores and helps to reduce measurement noise as well as multiple testing of redundant  
187 measures (e.g., Steinsbekk et al., 2016). PPSs are calculated by summing the products of each  
188 feature's measure (i.e., personality/psychopathology/eating behavior item scores, cognitive test  
189 scores, and thickness in mm/surface area in mm<sup>2</sup>/volume in mm<sup>3</sup> of cortical regions of interest)  
190 with its correlation with childhood obesity within a factor. To avoid overfitting, correlations  
191 were obtained from the singleton subset and applied to measures within the subset of siblings  
192 to generate six PPSs for each participant in the sibling subset. These PPSs informed preliminary  
193 correlation and twin modeling analyses. Ten-fold cross-validation within the singleton sample  
194 was later used to generate six corresponding PPSs in an independent sample for longitudinal  
195 analysis (see Supplementary Table S3).

196

197 *Twin modeling*

198 Direction of Causation (DoC) twin modeling was conducted to investigate the causal paths  
199 building the obesity–PPS associations. In DoC, the relationships of heritability variance  
200 components of two phenotypes are structurally modeled (Heath et al., 1993). This builds on the  
201 ACE heritability model which decomposes observed traits into additive genetic (A), common  
202 environment (C), and unique environment (E) variance components by comparing mono- and  
203 dizygotic twin pairs (non-twin siblings were treated equal to dizygotic twins; Neale & Cardon,  
204 1992). If one trait has a causal influence another trait, then the A, C, and E variance components  
205 of this first trait should be proportionally represented in the variance of the second. In this way,  
206 DoC models can be specified for obesity causally influencing PPS, or PPS causally influencing  
207 obesity. A reciprocal model can be defined by allowing both causal paths. Removing any causal  
208 paths leads to the Cholesky model where observed correlations are driven by a third variable

209 (Heath et al., 1993). For each obesity–PPS association, the best-fitting model was chosen by  
210 comparing model fit with regards to model complexity using Chi-squared and Akaike  
211 information criterion. For both, lower values indicate better model fit.

212

### 213 *Longitudinal analyses*

214 To validate DoC results, longitudinal analyses using cross-lagged panel models (CLPM) were  
215 performed. CLPMs estimate autoregressive paths which reflect stability of each measured trait  
216 over time (i.e., *a* paths) by structural modeling. Moreover, paths from previous measures of one  
217 trait to later measures of another trait are modeled. CLPMs refer to these as causal paths or *c*  
218 paths. However, they may not truly reflect causality: CLPMs test the ability of one trait’s past  
219 levels to predict later levels of another trait, but such relationships can be driven by unobserved  
220 third variables as well as causal influences (Usami et al., 2019).

221 Traditional CLPMs (Bentler & Speckart, 1981) were calculated for all six PPSs using  
222 baseline and two-year follow-up measures. Random-intercept CLPMs (Hamaker et al., 2015;  
223 Mund & Nestler, 2019) were applied where PPSs could be calculated for baseline, one-year and  
224 two-year follow-up (i.e., personality/psychopathology and eating behavior factors). The  
225 methodological advantage of Random-intercept CLPMs lays in the estimation of latent  
226 variables to account for stable interindividual trait differences (Hamaker et al., 2015). However,  
227 this method requires at least three observation waves. Due to varying availability of measures  
228 per study wave, PPSs for CLPM may not be fully congruent with PPSs for DoC (see  
229 Supplementary Table S3).

230

### 231 *Data and code availability*

232 ABCD study data is available at <https://abcdstudy.org/>. A commented analysis script is  
233 available at [osf.io/zgda4](https://osf.io/zgda4) for detailed data analysis procedure insight and replication. Analysis  
234 software information is noted in Supplementary Table S9.

235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260

## Results

At baseline, the average percentage of the sex- and age-specific 95th BMI percentile within the singleton subset was 85.0 (SD=17.9) with 17.7% having obesity. The average percentage of the sex- and age-specific 95th BMI percentile within the sibling subset was 82.0 (SD=16.4) with 13.3% having obesity. The results of the cross-sectional correlation analyses within six neurobehavioral factors, obesity–PPS correlation analyses, causal DoC twin modeling, and validating CLPM longitudinal analyses are reported in the following sections.

### *Correlational analyses within neurobehavioral factors*

#### *Personality/psychopathology and eating behavior factors*

In the subset of singletons, obesity was positively correlated to drive, reward responsiveness, social problems, somatic complaint, aggressive behavior, and being depressed. A negative correlation was found with sensation seeking. The strongest associations were between obesity and measures of psychopathology (Figure 1).

Correlation analyses were conducted between childhood obesity and the eating-specific items “overeating” and “doesn’t eat well” of the Child Behavior Checklist which formed the eating behavior factor. Both eating-related and parent-report items had considerable correlations with obesity. A stronger positive correlation was found with “overeating” and a weaker negative correlation with “doesn’t eat well” (Figure 1). The feature “doesn’t eat well” was also represented in the somatic complaints score within the personality/psychopathology factor for correlational analysis. In following analyses, eating-specific items were excluded from the personality/psychopathology factor.

#### *Cognition*

Cognitive test performance showed mainly negative correlations with obesity in the singleton subset. Exceptions were inhibitory control, perception speed, and pattern comparison where no

261 significant associations were found. The obesity–cognition association was strongest for the  
262 Wechsler matrix reasoning task (Figure 1).

263

264 --- Figure 1 here ---

265

### 266 *Brain morphology factors*

267 Analyses of singletons revealed positive associations of posterior cortical thickness in the  
268 occipital and parietal lobe. Moreover, a positive correlation was also found with the precentral  
269 region of interest within the frontal lobe. Negative associations between cortical thickness and  
270 obesity were found for most regions of interest within the frontal lobe as well as superior  
271 temporal and temporal pole regions within the temporal lobe. Higher correlations for the right  
272 compared to the left hemisphere were discovered in some frontal and parietal brain regions  
273 (Figure 2).

274 Fewer meaningful associations were found for cortical surface area and volume. Surface  
275 area–obesity associations were present in the isthmus cingulate, lateral orbitofrontal, medial  
276 orbitofrontal, rostral middle frontal, insula, and temporal pole regions of interest (Figure 2)  
277 while volume–obesity associations were found with the isthmus cingulate, rostral middle  
278 frontal, frontal pole, insula, posterior cingulate, superior temporal, and temporal pole regions  
279 of interest (Figure 2). See Supplementary Figure S2 for mapped visualization of obesity–  
280 morphology correlations.

281

282 --- Figure 2 here ---

283

### 284 *Poly-phenotype scores*

285 Obesity–PPS correlations in the twin sample were  $r=.09$ ,  $p<.001$ ,  $n=2469$  for  
286 personality/psychopathology,  $r=.36$ ,  $p<.001$ ,  $n=2475$  for eating behavior,  $r=.04$ ,  $p=.02$ ,  $n=2296$

287 for cognition,  $r=.14$ ,  $p<.001$ ,  $n=2391$  for cortical thickness,  $r=.06$ ,  $p<.01$ ,  $n=2391$  for cortical  
288 surface area, and  $r=.07$ ,  $p<.001$ ,  $n=2391$  for cortical volume (for full correlation matrix see  
289 Supplementary Table S4).

290

### 291 *Direction of Causation*

292 The heritability of obesity was 89.5% with the remaining 10.5% being due to unique  
293 experiences of a co-sibling. PPS heritability varied between 6.6% (eating behavior) and 57.5%  
294 (cortical thickness, see Supplementary Tables S5–6).

295 Based on model fit, data were consistent with obesity having unidirectional causal  
296 effects on cognition ( $\beta=.05$ ,  $CI=[.00, .09]$ ,  $n_{\text{Pairs}}=1056$ ), cortical thickness ( $\beta=.15$ ,  $CI=[.11, .20]$ ,  
297  $n_{\text{Pairs}}=1138$ ), and cortical surface area ( $\beta=.07$ ,  $CI=[.03, .11]$ ,  $n_{\text{Pairs}}=1138$ ). Contrarily, the  
298 personality/psychopathology ( $\beta=.09$ ,  $CI=[.05, .12]$ ,  $n_{\text{Pairs}}=1213$ ) and cortical volume ( $\beta=.07$ ,  
299  $CI=[.04, .11]$ ,  $n_{\text{Pairs}}=1138$ ) PPS appeared to have unidirectional causal effects on obesity. The  
300 reciprocal causation model had the best fit for eating behavior ( $\beta_{\text{Obesity-to-PPS}}=.26$ ,  $CI_{\text{Obesity-to-PPS}}$   
301  $=[.20, .32]$ ,  $\beta_{\text{PPS-to-Obesity}}=.16$ ,  $CI_{\text{PPS-to-Obesity}}=[.11, .20]$ ,  $n_{\text{Pairs}}=1218$ ). Detailed DoC modeling  
302 statistics are reported in Supplementary Table S7.

303

### 304 *Cross-lagged panel models*

305 Using the preferred random-intercept CLPM, meaningful  $c$  paths were found from obesity to  
306 eating behavior ( $\beta=.11$ ,  $CI=[.03, .19]$ ,  $n=3497$ ) and from personality/psychopathology to  
307 obesity ( $\beta=.04$ ,  $CI=[.02, .07]$ ,  $n=3482$ ). Traditional CLPMs found  $c$  paths from obesity to all  
308 neurobehavioral factors: personality/psychopathology ( $\beta=.07$ ,  $CI=[.04, .10]$ ,  $n=3416$ ), eating  
309 behavior ( $\beta=.22$ ,  $CI=[.18, .26]$ ,  $n=3442$ ), cognition ( $\beta=-.04$ ,  $CI=[-.08, -.01]$ ,  $n=2952$ ), cortical  
310 thickness ( $\beta=.08$ ,  $CI=[.05, .11]$ ,  $n=3215$ ), cortical surface area ( $\beta=.02$ ,  $CI=[.01, .03]$ ,  $n=3215$ ),  
311 and cortical volume ( $\beta=.02$ ,  $CI=[.00, .03]$ ,  $n=3215$ ).  $C$  paths also emerged from  
312 personality/psychopathology ( $\beta=.03$ ,  $CI=[.01, .05]$ ,  $n=3416$ ) and eating behavior ( $\beta=.05$ ,

313 CI=[.02, .08], n=3442) to obesity. Complete CLPM path estimates are shown in Supplementary  
314 Table S8.

315 The directional relationships found in DoC twin modeling were fully replicated except  
316 for the causal effects of cortical volume and eating behavior (only traditional CLPM) on  
317 childhood obesity. Additional longitudinal paths emerged from childhood obesity to  
318 personality/psychopathology (only traditional CLPM) and cortical volume (Figure 3). The  
319 negative longitudinal childhood obesity–cognition effect implies that higher obesity at age nine  
320 to ten is associated with higher cognitive test scores at age eleven to twelve.

321

322 --- Figure 3 here ---

323

## 324 Discussion

325 The purpose of the present study was to apply multiple causal inference methods to advance  
326 the understanding of associations between childhood obesity and neurobehavioral factors.  
327 Replicating previous cross-sectional findings, childhood obesity at age nine to ten was found to  
328 be associated to measures of personality/psychopathology, eating behavior, cognition, cortical  
329 thickness, cortical surface area, and cortical volume. The correlations between obesity and  
330 aggregated PPSs were similar to meta-analytic estimates (Kininmonth et al., 2021; Sutaria et  
331 al., 2019) and previous studies using the PPS method (Vainik et al., 2018) with the exception  
332 of a lower obesity–cognition estimate than previously reported (Pearce et al., 2018).

333 Regarding the direction of these relationships, the data supported causal influences of  
334 obesity on eating behavior, cognition, cortical thickness, and cortical surface area across  
335 analysis methods. Personality/psychopathology showed consistent causal and longitudinal  
336 effects on obesity. A reciprocal relationship may drive the associations between obesity and  
337 eating behavior. However, a random-intercept CLPM found only longitudinal effects from  
338 obesity to eating behavior. Both reciprocal obesity–eating behavior models (i.e., DoC and

339 traditional CLPM) implied the latter to be the stronger path, too. For the association between  
340 obesity and personality/psychopathology, only the methodologically weaker traditional CLPM  
341 suggested a reciprocal relationship. Conflicting findings leave the relationship between obesity  
342 and cortical volume unclear. Together, the results of this multi-method study in a large sample  
343 highlight the importance of obesity as a driving force behind individual differences in brain  
344 functioning and morphology during childhood.

345 Results are in line with previous longitudinal evidence for a causal influence of  
346 personality and psychopathology measures on childhood obesity (Anzman-Frasca et al., 2015;  
347 Blaine, 2008; Camfferman et al., 2016; Christensen et al., 2019; De Decker et al., 2017; Derks  
348 et al., 2019; Korczak et al., 2013) as well as studies suggesting obesity and eating behavior to  
349 have a bidirectional relationship (Antoniou et al., 2016; Herle et al., 2021; Reed et al., 2017).  
350 Some studies suggested that childhood obesity may also influence psychopathology (Bradley  
351 et al., 2008; Hübel et al., 2019; Korczak et al., 2013). Evidence for this direction was only  
352 generated by traditional CLPM modeling. A longitudinal influence of cortical surface area on  
353 body mass (Park et al., 2020) was not replicated. Novel relationships emerged between  
354 childhood obesity and cognition as well as cortical thickness.

355 In recent studies, deficits in cognitive abilities of obese adults have been attributed to an  
356 obesity-associated inflammation in the brain (Spyridaki et al., 2016). Even though less  
357 pronounced, this effect can be partly observed in children, too (Adelantado-Renau et al., 2019).  
358 The beginning inflammation in the brains of obese children may explain the small effect from  
359 obesity to cognition in children aged 9–10. Brain inflammation may cause accelerated brain  
360 aging leading to cortical thinning. Albeit the effect being most pronounced in adulthood, similar  
361 degeneration processes may drive the effect of obesity on cortical thickness and surface area in  
362 the present study (Ronan et al., 2016). Reduced cortical thickness or surface area in frontal brain  
363 areas associated with executive functioning could mediate effects of obesity on performance in  
364 several cognitive tests.

365           However, a longitudinal correlation from increased obesity to *better* cognitive  
366 functioning was observed using a traditional CLPM. The previously negative association  
367 disappeared at age 11–12. Changes in the association between obesity and intelligence across  
368 childhood have been previously described, implying that negative effects of obesity can  
369 potentially be compensated in certain developmental periods (Wraw et al., 2018). However, the  
370 positive effect could not be replicated across methods. Further research is necessary to map this  
371 effect.

372           On the other hand, potential self- and parent-reported contributors to childhood obesity  
373 (i.e., eating behavior, psychopathology, impulsivity, and motivation) were identified. Of them,  
374 strongest effects could be summarized as increased behavioral approach motivation (drive,  
375 reward responsiveness), which may lead to unhealthy food choices that maximize short term  
376 satisfaction (Zhou et al., 2019). Besides food quality, food quantity could also be affected. The  
377 reduced ability to stop eating when hunger is satisfied may lead to overeating (van den Berg et  
378 al., 2011). The other domain seems to be stress-related psychopathology. Children with  
379 increased mental health problems may attempt to regulate adverse emotions by stress eating,  
380 which may contribute to weight gain (Hill et al., 2018). Current evidence favors focusing on  
381 interventions that aid reducing stress and/or try to manage increased sensation seeking.

382           Although the results suggest that eating behavior could contribute to obesity, its effect  
383 seemed trivial compared to the reverse effect of obesity on eating behavior. While rarely  
384 considered, such association is consistent with obesity increasing fat-free mass which in turn  
385 increases food intake, which then maintains obesity (Blundell, 2018). Fat-free mass and food  
386 intake have been linked in adolescents (Vainik et al., 2016), and genetic obesity predicts future  
387 childhood eating behaviors (Herle et al., 2021). Therefore, studies intervening on eating  
388 behaviors should also consider reverse causality.

389           The results are limited using the percentage of the sex- and age-specific 95th BMI  
390 percentile as a proxy for obesity and the assumption of obesity being continuous while most

391 participants did not have obesity. Relative fat mass was explored as an alternative approach to  
392 obesity (Arumäe et al., 2022). However, correlations were highly similar and relative fat mass  
393 was dropped. The PPS method limits the results by ignoring causal relationships of individual  
394 features within a factor, but this was offset by gain in statistical power. DoC results are  
395 influenced by measurement reliability in a way that DoC tends to model the more reliable  
396 variable to have causal effects on the less reliable variable. Reliability could be lower for  
397 behavioral (Nakamura et al., 2009) and cognitive factors (Taylor et al., 2020) which could  
398 contribute to the finding that obesity had causal effects on eating behavior and cognition.  
399 Personality/psychopathology and cognitive test results can be compromised by intrapersonal  
400 components like response bias or motivation which are irrelevant for the directly observed  
401 measures of childhood obesity and brain morphology. At the same time, previous analyses  
402 showed that explicitly modeling reliability did not change DoC conclusions (Arumäe et al.,  
403 2021). Future studies using different causal inference methods should build complementary  
404 evidence (Arumäe et al., 2021; Munafò & Davey Smith, 2018).

405 In summary, the present study provides evidence that childhood obesity may influence  
406 rather than be influenced by the various aspects of brain functioning and morphology that have  
407 been found to differ between children with and without obesity. Therefore, obesity may play an  
408 important role in brain development during childhood. The present findings are hoped to inspire  
409 further investigation of the underlying mechanisms behind these associations and development  
410 of more effective obesity interventions.

411

412

## References

- 413
- 414 Achenbach, T. M., & Edelbrock, C. S. (1983). *Manual for the Child Behavior Checklist and*
- 415 *Revised Child Behavior Profile*. University Associates in Psychiatry.
- 416 Adelantado-Renau, M., Esteban-Cornejo, I., Rodriguez-Ayllon, M., Cadenas-Sanchez, C.,
- 417 Gil-Cosano, J. J., Mora-Gonzalez, J., Solis-Urra, P., Verdejo-Román, J., Aguilera, C. M.,
- 418 Escolano-Margarit, M. V., Verdejo-Garcia, A., Catena, A., Moliner-Urdiales, D., &
- 419 Ortega, F. B. (2019). Inflammatory biomarkers and brain health indicators in children
- 420 with overweight and obesity: The ActiveBrains project. *Brain, Behavior, and Immunity*,
- 421 *81*, 588–597. <https://doi.org/10.1016/j.bbi.2019.07.020>
- 422 Adise, S., Allgaier, N., Laurent, J., Hahn, S., Charani, B., Owens, M., Yuan, D., Nyugen, P.,
- 423 Mackey, S., Potter, A., & Garavan, H. P. (2021). Multimodal brain predictors of current
- 424 weight and weight gain in children enrolled in the ABCD study ®. *Developmental*
- 425 *Cognitive Neuroscience*, *49*(100948). <https://doi.org/10.1016/j.dcn.2021.100948>
- 426 Alosco, M. L., Stanek, K. M., Galioto, R., Korgaonkar, M. S., Grieve, S. M., Brickman, A.
- 427 M., Spitznagel, M. B., & Gunstad, J. (2014). Body mass index and brain structure in
- 428 healthy children and adolescents. *International Journal of Neuroscience*, *124*(1), 49–55.
- 429 <https://doi.org/10.3109/00207454.2013.817408>
- 430 Antoniou, E. E., Roefs, A., Kremers, S. P. J., Jansen, A., Gubbels, J. S., Sleddens, E. F. C., &
- 431 Thijs, C. (2016). Picky eating and child weight status development: a longitudinal study.
- 432 *Journal of Human Nutrition and Dietetics*, *29*(3), 298–307.
- 433 <https://doi.org/10.1111/jhn.12322>
- 434 Anzman-Frasca, S., Francis, L. A., & Birch, L. L. (2015). Inhibitory control is associated with
- 435 psychosocial, cognitive, and weight outcomes in a longitudinal sample of girls.
- 436 *Translational Issues in Psychological Science*, *1*(3), 203–216.
- 437 <https://doi.org/10.1037/tps0000028>
- 438 Arumäe, K., Briley, D., Colodro-Conde, L., Mortensen, E. L., Jang, K., Ando, J., Kandler, C.,

- 439 Sørensen, T. I. A., Dagher, A., Möttus, R., & Vainik, U. (2021). Two genetic analyses to  
440 elucidate causality between body mass index and personality. *International Journal of*  
441 *Obesity (2005)*, *45*(10), 2244–2251. <https://doi.org/10.1038/s41366-021-00885-4>
- 442 Arumäe, K., Möttus, R., & Vainik, U. (2022). Beyond BMI: Personality traits' associations  
443 with adiposity and metabolic rate. *Physiology & Behavior*, *246*, 113703.  
444 <https://doi.org/10.1016/j.physbeh.2022.113703>
- 445 Belsky, D. W., Caspi, A., Goldman-Mellor, S., Meier, M. H., Ramrakha, S., Poulton, R., &  
446 Moffitt, T. E. (2013). Is obesity associated with a decline in intelligence quotient during  
447 the first half of the life course? *American Journal of Epidemiology*, *178*(9), 1461–1468.  
448 <https://doi.org/10.1093/aje/kwt135>
- 449 Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and  
450 Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*  
451 *(Methodological)*, *57*(1), 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- 452 Bentler, P. M., & Speckart, G. (1981). Attitudes “cause” behaviors: A structural equation  
453 analysis. *Journal of Personality and Social Psychology*, *40*(2), 226–238.  
454 <https://doi.org/10.1037/0022-3514.40.2.226>
- 455 Blaine, B. (2008). Does Depression Cause Obesity? *Journal of Health Psychology*, *13*(8),  
456 1190–1197. <https://doi.org/10.1177/1359105308095977>
- 457 Blundell, J. E. (2018). Behaviour, energy balance, obesity and capitalism. *European Journal*  
458 *of Clinical Nutrition*, *72*(9), 1305–1309. <https://doi.org/10.1038/s41430-018-0231-x>
- 459 Blundell, J. E., & Finlayson, G. (2004). Is susceptibility to weight gain characterized by  
460 homeostatic or hedonic risk factors for overconsumption? *Physiology & Behavior*, *82*(1),  
461 21–25. <https://doi.org/10.1016/j.physbeh.2004.04.021>
- 462 Bohon, C., & Welch, H. (2021). Quadratic relations of BMI with depression and brain volume  
463 in children: Analysis of data from the ABCD study. *Journal of Psychiatric Research*,  
464 *136*, 421–427. <https://doi.org/10.1016/j.jpsychires.2021.02.038>

- 465 Bradley, R. H., Houts, R., Nader, P. R., O'Brien, M., Belsky, J., & Crosnoe, R. (2008). The  
466 Relationship between Body Mass Index and Behavior in Children. *The Journal of*  
467 *Pediatrics*, 153(5), 629-634.e3. <https://doi.org/10.1016/j.jpeds.2008.05.026>
- 468 Camfferman, R., Jansen, P. W., Rippe, R. C. A., Mesman, J., Derks, I. P. M., Tiemeier, H.,  
469 Jaddoe, V., & van der Veek, S. M. C. (2016). The association between overweight and  
470 internalizing and externalizing behavior in early childhood. *Social Science & Medicine*  
471 (1982), 168, 35–42. <https://doi.org/10.1016/j.socscimed.2016.09.001>
- 472 Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and  
473 affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal*  
474 *of Personality and Social Psychology*, 67(2), 319–333.
- 475 Christensen, K. G., Nielsen, S. G., Olsen, N. J., Dalgård, C., Heitmann, B. L., & Larsen, S. C.  
476 (2019). Child behaviour and subsequent changes in body weight, composition and shape.  
477 *PloS One*, 14(12), e0226003. <https://doi.org/10.1371/journal.pone.0226003>
- 478 Cyders, M. A., Littlefield, A. K., Coffey, S., & Karyadi, K. A. (2014). Examination of a short  
479 English version of the UPPS-P Impulsive Behavior Scale. *Addictive Behaviors*, 39(9),  
480 1372–1376. <https://doi.org/10.1016/j.addbeh.2014.02.013>
- 481 De Decker, A., De Clercq, B., Verbeken, S., Wells, J. C. K., Braet, C., Michels, N., De  
482 Henauw, S., & Sioen, I. (2017). Fat and lean tissue accretion in relation to reward  
483 motivation in children. *Appetite*, 108, 317–325.  
484 <https://doi.org/10.1016/j.appet.2016.10.017>
- 485 Dennis, E., Manza, P., & Volkow, N. D. (2022). Socioeconomic status, BMI, and brain  
486 development in children. *Translational Psychiatry*, 12(1), 33.  
487 <https://doi.org/10.1038/s41398-022-01779-3>
- 488 Derks, I. P. M., Bolhuis, K., Yalcin, Z., Gaillard, R., Hillegers, M. H. J., Larsson, H.,  
489 Lundström, S., Lichtenstein, P., van Beijsterveldt, C. E. M., Bartels, M., Boomsma, D. I.,  
490 Tiemeier, H., & Jansen, P. W. (2019). Testing Bidirectional Associations Between

- 491 Childhood Aggression and BMI: Results from Three Cohorts. *Obesity*, oby.22419.
- 492 <https://doi.org/10.1002/oby.22419>
- 493 Freedman, D. S., Butte, N. F., Taveras, E. M., Lundeen, E. A., Blanck, H. M., Goodman, A.
- 494 B., & Ogden, C. L. (2017). BMI z -Scores are a poor indicator of adiposity among 2- to
- 495 19-year-olds with very high BMIs, NHANES 1999-2000 to 2013-2014. *Obesity*, 25(4),
- 496 739–746. <https://doi.org/10.1002/oby.21782>
- 497 Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A Research Note. *Journal*
- 498 *of Child Psychology and Psychiatry*, 38(5), 581–586. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-7610.1997.tb01545.x)
- 499 [7610.1997.tb01545.x](https://doi.org/10.1111/j.1469-7610.1997.tb01545.x)
- 500 Gray, J. C., Schvey, N. A., & Tanofsky-Kraff, M. (2020). Demographic, psychological,
- 501 behavioral, and cognitive correlates of BMI in youth: Findings from the Adolescent
- 502 Brain Cognitive Development (ABCD) study. *Psychological Medicine*, 50(9), 1539–
- 503 1547. <https://doi.org/10.1017/S0033291719001545>
- 504 Hall, P. A., Best, J., Beaton, E. A., Sakib, M. N., & Danckert, J. (2021). Morphology of the
- 505 Prefrontal Cortex Predicts Body Composition in Early Adolescence: Cognitive
- 506 Mediators and Environmental Moderators in the ABCD Study. *Social Cognitive and*
- 507 *Affective Neuroscience*. <https://doi.org/10.1093/scan/nsab104>
- 508 Hamaker, E. L., Kuiper, R. M., & Grasman, R. P. P. P. (2015). A critique of the cross-lagged
- 509 panel model. *Psychological Methods*, 20(1), 102–116. <https://doi.org/10.1037/a0038889>
- 510 Hamilton, D., Dee, A., & Perry, I. J. (2018). The lifetime costs of overweight and obesity in
- 511 childhood and adolescence: a systematic review. *Obesity Reviews*, 19(4), 452–463.
- 512 <https://doi.org/10.1111/obr.12649>
- 513 Hashimoto, T., Takeuchi, H., Taki, Y., Yokota, S., Hashizume, H., Asano, K., Asano, M.,
- 514 Sassa, Y., Nouchi, R., & Kawashima, R. (2015). Increased posterior hippocampal
- 515 volumes in children with lower increase in body mass index: a 3-year longitudinal MRI
- 516 study. *Developmental Neuroscience*, 37(2), 153–160. <https://doi.org/10.1159/000370064>

- 517 Heath, A. C., Kessler, R. C., Neale, M. C., Hewitt, J. K., Eaves, L. J., & Kendler, K. S.  
518 (1993). Testing hypotheses about direction of causation using cross-sectional family  
519 data. *Behavior Genetics*, 23(1), 29–50. <https://doi.org/10.1007/BF01067552>
- 520 Herle, M., Abdulkadir, M., Hübel, C., Ferreira, D. S., Bryant-Waugh, R., Loos, R. J. F.,  
521 Bulik, C. M., De Stavola, B., & Micali, N. (2021). The genomics of childhood eating  
522 behaviours. *Nature Human Behaviour*, 5(5), 625–630. [https://doi.org/10.1038/s41562-](https://doi.org/10.1038/s41562-020-01019-y)  
523 [020-01019-y](https://doi.org/10.1038/s41562-020-01019-y)
- 524 Hill, D. C., Moss, R. H., Sykes-Muskett, B., Conner, M., & O'Connor, D. B. (2018). Stress  
525 and eating behaviors in children and adolescents: Systematic review and meta-analysis.  
526 *Appetite*, 123, 14–22. <https://doi.org/10.1016/j.appet.2017.11.109>
- 527 Hübel, C., Gaspar, H. A., Coleman, J. R. I., Hanscombe, K. B., Purves, K., Prokopenko, I.,  
528 Graff, M., Ngwa, J. S., Workalemahu, T., ADHD Working Group of the Psychiatric  
529 Genomics Consortium, Meta-Analyses of Glucose and Insulin-related traits consortium  
530 (MAGIC), Autism Working Group of the Psychiatric Genomics Consortium, Bipolar  
531 Disorder Working Group of the Psychiatric Genomics Consortium, Eating Disorders  
532 Working Group of the Psychiatric Genomics Consortium, Major Depressive Disorder  
533 Working Group of the Psychiatric Genomics Consortium, OCD & Tourette Syndrome  
534 Working Group of the Psychiatric Genomics Consortium, PTSD Working Group of the  
535 Psychiatric Genomics Consortium, Schizophrenia Working Group of the Psychiatric  
536 Genomics Consortium, Sex Differences Cross Disorder Working Group of the  
537 Psychiatric Genomics Consortium, ... Breen, G. (2019). Genetic correlations of  
538 psychiatric traits with body composition and glycemic traits are sex- and age-dependent.  
539 *Nature Communications*, 10(1), 5765. <https://doi.org/10.1038/s41467-019-13544-0>
- 540 Kennedy, J. T., Collins, P. F., & Luciana, M. (2016). Higher Adolescent Body Mass Index Is  
541 Associated with Lower Regional Gray and White Matter Volumes and Lower Levels of  
542 Positive Emotionality. *Frontiers in Neuroscience*, 10.

- 543 <https://doi.org/10.3389/fnins.2016.00413>
- 544 Kim, M. S., Luo, S., Azad, A., Campbell, C. E., Felix, K., Cabeen, R. P., Belcher, B. R., Kim,  
545 R., Serrano-Gonzalez, M., & Herting, M. M. (2020). Prefrontal Cortex and Amygdala  
546 Subregion Morphology Are Associated With Obesity and Dietary Self-control in  
547 Children and Adolescents. *Frontiers in Human Neuroscience*, *14*, 563415.  
548 <https://doi.org/10.3389/fnhum.2020.563415>
- 549 Kininmonth, A., Smith, A., Carnell, S., Steinsbekk, S., Fildes, A., & Llewellyn, C. (2021).  
550 The association between childhood adiposity and appetite assessed using the Child  
551 Eating Behavior Questionnaire and Baby Eating Behavior Questionnaire: A systematic  
552 review and meta-analysis. *Obesity Reviews*, *22*(5). <https://doi.org/10.1111/obr.13169>
- 553 Korczak, D. J., Lipman, E., Morrison, K., & Szatmari, P. (2013). Are children and adolescents  
554 with psychiatric illness at risk for increased future body weight? A systematic review.  
555 *Developmental Medicine & Child Neurology*, *55*(11), 980–987.  
556 <https://doi.org/10.1111/dmcn.12168>
- 557 Laurent, J. S., Watts, R., Adise, S., Allgaier, N., Chaarani, B., Garavan, H., Potter, A., &  
558 Mackey, S. (2020). Associations Among Body Mass Index, Cortical Thickness, and  
559 Executive Function in Children. *JAMA Pediatrics*, *174*(2), 170–177.  
560 <https://doi.org/10.1001/jamapediatrics.2019.4708>
- 561 Lezak, M. D. (1976). *Neuropsychological assessment*. Oxford University Press.
- 562 Llewellyn, A., Simmonds, M., Owen, C. G., & Woolacott, N. (2016). Childhood obesity as a  
563 predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obesity  
564 Reviews*, *17*(1), 56–67. <https://doi.org/10.1111/obr.12316>
- 565 Lugo-Candelas, C., Pang, Y., Lee, S., Cha, J., Hong, S., Ranzenhofer, L., Korn, R., Davis, H.,  
566 McNerny, H., Schebendach, J., Chung, W. K., Leibel, R. L., Walsh, B. T., Posner, J.,  
567 Rosenbaum, M., & Mayer, L. (2020). Differences in brain structure and function in  
568 children with the FTO obesity-risk allele. *Obesity Science & Practice*, *6*(4), 409–424.

- 569 <https://doi.org/10.1002/osp4.417>
- 570 Munafò, M. R., & Davey Smith, G. (2018). Robust research needs many lines of evidence.
- 571 *Nature*, 553(7689), 399–401. <https://doi.org/10.1038/d41586-018-01023-3>
- 572 Mund, M., & Nestler, S. (2019). Beyond the Cross-Lagged Panel Model: Next-generation
- 573 statistical tools for analyzing interdependencies across the life course. *Advances in Life*
- 574 *Course Research*, 41, 100249. <https://doi.org/10.1016/j.alcr.2018.10.002>
- 575 Nakamura, B. J., Ebesutani, C., Bernstein, A., & Chorpita, B. F. (2009). A Psychometric
- 576 Analysis of the Child Behavior Checklist DSM-Oriented Scales. *Journal of*
- 577 *Psychopathology and Behavioral Assessment*, 31(3), 178–189.
- 578 <https://doi.org/10.1007/s10862-008-9119-8>
- 579 Neale, M. C., & Cardon, L. R. (1992). *Methodology for genetic studies of twins and families*.
- 580 Kluwer Academic Publishers.
- 581 Park, B.-Y., Chung, C.-S., Lee, M. J., & Park, H. (2020). Accurate neuroimaging biomarkers
- 582 to predict body mass index in adolescents: a longitudinal study. *Brain Imaging and*
- 583 *Behavior*, 14(5), 1682–1695. <https://doi.org/10.1007/s11682-019-00101-y>
- 584 Pearce, A. L., Leonhardt, C. A., & Vaidya, C. J. (2018). Executive and Reward-Related
- 585 Function in Pediatric Obesity: A Meta-Analysis. *Childhood Obesity*, 14(5), 265–279.
- 586 <https://doi.org/10.1089/chi.2017.0351>
- 587 Prats-Soteras, X., Jurado, M. A., Ottino-González, J., García-García, I., Segura, B., Caldú, X.,
- 588 Sánchez-Garre, C., Miró, N., Tor, C., Sender-Palacios, M., & Garolera, M. (2020).
- 589 Inflammatory agents partially explain associations between cortical thickness, surface
- 590 area, and body mass in adolescents and young adulthood. *International Journal of*
- 591 *Obesity*, 44(7), 1487–1496. <https://doi.org/10.1038/s41366-020-0582-y>
- 592 Rajan, L., McKay, C. C., Santos Malavé, G., Pearce, A. L., Cherry, J. B. C., Mackey, E.,
- 593 Nadler, E. P., & Vaidya, C. J. (2021). Effects of severe obesity and sleeve gastrectomy
- 594 on cortical thickness in adolescents. *Obesity*, 29(9), 1516–1525.

- 595 <https://doi.org/10.1002/oby.23206>
- 596 Ratcliff, G. (1979). Spatial thought, mental rotation and the right cerebral hemisphere.
- 597 *Neuropsychologia*, 17(1), 49–54. [https://doi.org/10.1016/0028-3932\(79\)90021-6](https://doi.org/10.1016/0028-3932(79)90021-6)
- 598 Reed, Z. E., Micali, N., Bulik, C. M., Davey Smith, G., & Wade, K. H. (2017). Assessing the
- 599 causal role of adiposity on disordered eating in childhood, adolescence, and adulthood: a
- 600 Mendelian randomization analysis. *The American Journal of Clinical Nutrition*, 106(3),
- 601 764–772. <https://doi.org/10.3945/ajcn.117.154104>
- 602 Ronan, L., Alexander-Bloch, A. F., Wagstyl, K., Farooqi, S., Brayne, C., Tyler, L. K., &
- 603 Fletcher, P. C. (2016). Obesity associated with increased brain age from midlife.
- 604 *Neurobiology of Aging*, 47, 63–70. <https://doi.org/10.1016/j.neurobiolaging.2016.07.010>
- 605 Ronan, L., Alexander-Bloch, A., & Fletcher, P. C. (2020). Childhood Obesity, Cortical
- 606 Structure, and Executive Function in Healthy Children. *Cerebral Cortex (New York,*
- 607 *N.Y. : 1991)*, 30(4), 2519–2528. <https://doi.org/10.1093/cercor/bhz257>
- 608 Ross, N., Yau, P. L., & Convit, A. (2015). Obesity, fitness, and brain integrity in adolescence.
- 609 *Appetite*, 93, 44–50. <https://doi.org/10.1016/j.appet.2015.03.033>
- 610 Sahoo, K., Sahoo, B., Choudhury, A. K., Sofi, N. Y., Kumar, R., & Bhadoria, A. S. (2015).
- 611 Childhood obesity: causes and consequences. *Journal of Family Medicine and Primary*
- 612 *Care*, 4(2), 187–192. <https://doi.org/10.4103/2249-4863.154628>
- 613 Saute, R. L., Soder, R. B., Alves Filho, J. O., Baldisserotto, M., & Franco, A. R. (2018).
- 614 Increased brain cortical thickness associated with visceral fat in adolescents. *Pediatric*
- 615 *Obesity*, 13(1), 74–77. <https://doi.org/10.1111/ijpo.12190>
- 616 Shriver, L. H., Dollar, J. M., Lawless, M., Calkins, S. D., Keane, S. P., Shanahan, L., &
- 617 Wideman, L. (2019). Longitudinal Associations between Emotion Regulation and
- 618 Adiposity in Late Adolescence: Indirect Effects through Eating Behaviors. *Nutrients*,
- 619 11(3). <https://doi.org/10.3390/nu11030517>
- 620 Spyridaki, E. C., Avgoustinaki, P. D., & Margioris, A. N. (2016). Obesity, inflammation and

- 621 cognition. *Current Opinion in Behavioral Sciences*, 9, 169–175.
- 622 <https://doi.org/10.1016/j.cobeha.2016.05.004>
- 623 Steegers, C., Blok, E., Lamballais, S., Jaddoe, V., Bernardoni, F., Vernooij, M., van der Ende,  
624 J., Hillegers, M., Micali, N., Ehrlich, S., Jansen, P., Dieleman, G., & White, T. (2021).  
625 The association between body mass index and brain morphology in children: a  
626 population-based study. *Brain Structure & Function*, 226(3), 787–800.  
627 <https://doi.org/10.1007/s00429-020-02209-0>
- 628 Steinsbekk, S., Belsky, D., Guzey, I. C., Wardle, J., & Wichstrøm, L. (2016). Polygenic Risk,  
629 Appetite Traits, and Weight Gain in Middle Childhood. *JAMA Pediatrics*, 170(2),  
630 e154472. <https://doi.org/10.1001/jamapediatrics.2015.4472>
- 631 Sutaria, S., Devakumar, D., Yasuda, S. S., Das, S., & Saxena, S. (2019). Is obesity associated  
632 with depression in children? Systematic review and meta-analysis. *Archives of Disease  
633 in Childhood*, 104(1), 64–74. <https://doi.org/10.1136/archdischild-2017-314608>
- 634 Taylor, B. K., Frenzel, M. R., Eastman, J. A., Wiesman, A. I., Wang, Y.-P., Calhoun, V. D.,  
635 Stephen, J. M., & Wilson, T. W. (2020). Reliability of the NIH toolbox cognitive battery  
636 in children and adolescents: a 3-year longitudinal examination. *Psychological Medicine*,  
637 1–10. <https://doi.org/10.1017/S0033291720003487>
- 638 Thamotharan, S., Lange, K., Zale, E. L., Huffhines, L., & Fields, S. (2013). The role of  
639 impulsivity in pediatric obesity and weight status: A meta-analytic review. *Clinical  
640 Psychology Review*, 33(2), 253–262. <https://doi.org/10.1016/j.cpr.2012.12.001>
- 641 Thapaliya, G., Chen, L., Jansen, E., Smith, K. R., Sadler, J. R., Benson, L., Papantoni, A., &  
642 Carnell, S. (2021). Familial Obesity Risk and Current Excess Weight Influence Brain  
643 Structure in Adolescents. *Obesity (Silver Spring, Md.)*, 29(1), 184–193.  
644 <https://doi.org/10.1002/oby.23042>
- 645 Usami, S., Murayama, K., & Hamaker, E. L. (2019). A unified framework of longitudinal  
646 models to examine reciprocal relations. *Psychological Methods*, 24(5), 637–657.


- 647 <https://doi.org/10.1037/met0000210>
- 648 Vainik, U., Baker, T. E., Dadar, M., Zeighami, Y., Michaud, A., Zhang, Y., García Alanis, J.  
649 C., Misic, B., Collins, D. L., & Dagher, A. (2018). Neurobehavioral correlates of obesity  
650 are largely heritable. *Proceedings of the National Academy of Sciences*, *115*(37), 9312–  
651 9317. <https://doi.org/10.1073/pnas.1718206115>
- 652 Vainik, U., Konstabel, K., Lätt, E., Mäestu, J., Purge, P., & Jürimäe, J. (2016). Diet  
653 misreporting can be corrected: confirmation of the association between energy intake and  
654 fat-free mass in adolescents. *British Journal of Nutrition*, *116*(8), 1425–1436.  
655 <https://doi.org/10.1017/S0007114516003317>
- 656 van den Berg, L., Pieterse, K., Malik, J. A., Luman, M., Willems van Dijk, K., Oosterlaan, J.,  
657 & Delemarre-van de Waal, H. A. (2011). Association between impulsivity, reward  
658 responsiveness and body mass index in children. *International Journal of Obesity*,  
659 *35*(10), 1301–1307. <https://doi.org/10.1038/ijo.2011.116>
- 660 Volkow, N. D., Koob, G. F., Croyle, R. T., Bianchi, D. W., Gordon, J. A., Koroshetz, W. J.,  
661 Pérez-Stable, E. J., Riley, W. T., Bloch, M. H., Conway, K., Deeds, B. G., Dowling, G.  
662 J., Grant, S., Howlett, K. D., Matochik, J. A., Morgan, G. D., Murray, M. M., Noronha,  
663 A., Spong, C. Y., ... Weiss, S. R. B. (2018). The conception of the ABCD study: From  
664 substance use to a broad NIH collaboration. *Developmental Cognitive Neuroscience*, *32*,  
665 4–7. <https://doi.org/10.1016/j.dcn.2017.10.002>
- 666 Wang, Y., Beydoun, M. A., Min, J., Xue, H., Kaminsky, L. A., & Cheskin, L. J. (2020). Has  
667 the prevalence of overweight, obesity and central obesity levelled off in the United  
668 States? Trends, patterns, disparities, and future projections for the obesity epidemic.  
669 *International Journal of Epidemiology*, *49*(3), 810–823.  
670 <https://doi.org/10.1093/ije/dyz273>
- 671 Wechsler, D. (1949). *Wechsler Intelligence Scale for Children*. Psychological Corporation.
- 672 Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J.,

673 Carlozzi, N. E., Slotkin, J., Blitz, D., Wallner-Allen, K., Fox, N. A., Beaumont, J. L.,  
674 Mungas, D., Nowinski, C. J., Richler, J., Deocampo, J. A., Anderson, J. E., Manly, J. J.,  
675 Borosh, B., ... Gershon, R. C. (2013). Cognition assessment using the NIH Toolbox.  
676 *Neurology*, 80(11 Suppl 3), S54-64. <https://doi.org/10.1212/WNL.0b013e3182872ded>  
677 Wraw, C., Deary, I. J., Der, G., & Gale, C. R. (2018). Maternal and offspring intelligence in  
678 relation to BMI across childhood and adolescence. *International Journal of Obesity*  
679 (2005), 42(9), 1610–1620. <https://doi.org/10.1038/s41366-018-0009-1>  
680 Zhou, Z., SooHoo, M., Zhou, Q., Perez, M., & Liew, J. (2019). Temperament as Risk and  
681 Protective Factors in Obesogenic Eating: Relations Among Parent Temperament, Child  
682 Temperament, and Child Food Preference and Eating. *The Journal of Genetic*  
683 *Psychology*, 180(1), 75–79. <https://doi.org/10.1080/00221325.2019.1575180>  
684

685 **Figures**

686 **Figure 1**

687 *Associations between the percentage of the sex- and age-specific 95<sup>th</sup> BMI percentile and*  
 688 *personality/psychopathology scales, eating behavior items (A), as well as cognitive test scores*  
 689 *(B)*



690

691 *Note.* Impulsivity, motivation and prosociality were self-reported. Psychopathology (including


692 eating items) was parent-reported. Error bars represent 95% confidence intervals. EB, eating

693 behavior; EF, executive functioning; FDR, false discovery rate; Lang, Language; Perc,

694 perception; PS, prosociality; WM, working memory.

695 **Figure 2**

696 *Associations between the percentage of the sex- and age-specific 95<sup>th</sup> BMI percentile and region*  
 697 *of interest cortical thickness (A), cortical surface area (B) as well as cortical volume (C)*



698


699 *Note.* Error bars represent 95% confidence intervals. FDR, false discovery rate; I, insula.

700

701 **Figure 3**

702 *Causal and c path estimates for Direction of Causation (DoC), traditional cross-lagged panel*

703 *(CLPM) and random-intercept cross-lagged panel (RI-CLPM) models*



704

705 *Note. Only best fitting model plotted for Direction of Causation modeling. Error bars represent*

706 *95% confidence intervals. PPS, poly-phenotype score.*