**A statistical genomics framework to trace bacterial genomic predictors**

**of clinical outcomes in *Staphylococcus aureus* bacteraemia**

Stefano G. Giulieri[a,b,c,d], Romain Guérillot[a], Natasha E. Holmes[d], Sarah L. Baines[a,e], Abderrahman Hachani[a], Diane S. Daniel[a,e], Torsten Seemann[e], Joshua S. Davis[f,g], Sebastiaan Van Hal[h,i], Steve Y. C. Tong[b,c], Timothy P. Stinear[a]*, Benjamin P. Howden[a,d,e]*

[a] *Department of Microbiology and Immunology, The University of Melbourne at the Doherty Institute for Infection and Immunity, Melbourne, Australia*

[b] *Victorian Infectious Disease Service, The Royal Melbourne Hospital, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia*

[c] *Department of Infectious Diseases, The University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia*

[d] *Department of Infectious Diseases, Austin Health, Heidelberg, Australia*

[e] *Microbiological Diagnostic Unit Public Health Laboratory, The University of Melbourne at the Doherty Institute for Infection and Immunity, Melbourne, Australia*

[f] *Department of Infectious Diseases, John Hunter Hospital, Newcastle, New South Wales, Australia*

[g] *Menzies School of Health Research, Charles Darwin University, Casuarina, Northern Territory, Australia*

[h] *Department of Infectious Diseases and Microbiology, Royal Prince Alfred Hospital,*

*Sydney, New South Wales, Australia*

*[i] Central Clinical School, University of Sydney, Sydney, New South Wales, Australia*

*Joint Senior author

#Address correspondence to Benjamin P. Howden, <u>bhowden@unimelb.edu.au</u>

Keywords: bacterial GWAS, machine learning, *Staphylococcus aureus*, bacteraemia,

vancomycin resistance

**ABSTRACT**

Outcomes for patients with severe bacterial infections are determined by the interplay between host, pathogen, and treatments. Most notably, patient age and antibiotic resistance contributes significantly to poor outcomes. While human genomics studies have provided insights into the host genetic factors impacting outcomes of *Staphylococcus aureus* infections, comparatively little is known about *S. aureus* genotypes and disease severity. Building on the idea that bacterial pathoadaptation is a key driver of clinical outcomes, we develop a new genome-wide association study (GWAS) framework to identify adaptive bacterial mutations associated with clinical treatment failure and mortality in three large and independent *S. aureus* bacteraemia cohorts, comprising 1358 episodes. We discovered *S. aureus* loci with previously undescribed convergent mutations linked to both poorer infection outcomes and reduced susceptibility to vancomycin. Our research highlights the potential of vancomycin-selected mutations and vancomycin MIC as key explanatory variables to predict SAB severity. The contribution of bacterial variation was much lower for clinical outcomes (heritability < 5%), however, GWAS allowed us to identify additional, MIC-independent candidate pathogenesis loci. Using supervised machine-learning, we were able to quantify the predictive potential of these adaptive *S. aureus* signatures, along with host determinants of bacteraemia outcomes. The statistical genomics framework we have developed is a powerful means to capture adaptive mutations and find bacterial factors that influence and predict severe infections. Our findings underscore the importance of systematically collected, rich clinical and microbiological data to understand bacterial mechanisms promoting treatment failure.

## INTRODUCTION

The course and outcome of severe bacterial infections such as bacteraemia and sepsis are determined by the interaction of factors related to the host (e.g. genetics, age, chronic conditions), the pathogen and the treatment (Cohen, Vincent et al. 2015). A deeper understanding of these outcome determinants would help design more effective treatments and inform precision medicine strategies. However, in contrast with progress in the immunogenetics of infections (Kwok, Mentzer et al. 2021), it has been challenging to identify bacterial genetic factors influencing infection outcomes (Giulieri, Tong et al. 2020, Denamur, Condamine et al. 2022). This is due to the considerable genetic diversity of bacterial pathogens and the strong clonal structure of bacterial populations that complicate genome-wide association studies (GWAS) (Power, Parkhill et al. 2017).

With up to 20% mortality (Bai, Lo et al. , Battle, Shuping et al. 2021) and increasing incidence (Imam, Tempone et al. 2019), *Staphylococcus aureus* bloodstream infections are a key challenge to human health (Tong, Davis et al. 2015). The clinical determinants of *Staphylococcus aureus* bacteraemia (SAB) outcomes have been described extensively, in particular mortality has been the subject of numerous studies (reviewed here (van Hal, Jensen et al. 2012)). Small studies have shown the impact of host genetic factors of SAB risk and outcomes, for example it was shown that a polymorphism in the DNA methyltransferase *DNMT3A* protects from persistent SAB (Mba Medie, Sharma-Kuinkel et al. 2019). Studies linking specific clones or bacterial characteristics (vancomycin minimum inhibitory concentration [MIC]) with SAB mortality (Holmes, Turnidge et al. 2011) provide indirect evidence for the role of bacterial genetic factors in SAB mortality, however, it has been difficult to identify compelling associations between distinctive genes or mutations and clinical

outcomes (Giulieri, Holmes et al. 2016). In a landmark study Recker *et al*, used a

combination of GWAS and machine learning to explore factors associated with SAB

mortality. Their work showed that both bacterial genetic and phenotypic information

improved the performance of the predictive model. Feature selection from the model

indicated several potential pathogenic mutations, that could represent the basis for

functional validation (Recker, Laabei et al. 2017).

A key concept underlying bacterial phenotype-genotype studies is pathoadaptation,

where bacteria acquire mutations that enhance their ability to survive in the host.

Upon infection, colonising *S. aureus* strains are repeatedly subjected to similar

selective pressures; thus, mutations promoting survival in the host during infection

are expected to be preferentially maintained and become evident when examining *S.*

*aureus* microevolution. The most compelling evidence for the role of pathoadaptation

in SAB comes from within-host evolution studies, which have shown enrichment of

mutations in a small number of genes including *agr* and antibiotic resistance genes

(indicating convergent evolution), both upon transition from colonisation to invasion

and during persistent infection (Young, Wu et al. 2017, Giulieri, Baines et al. 2018).

Bacterial GWAS leverage the concept of adaptation either directly by focusing on

homoplastic mutations (phylogeny-based methods) (Farhat, Shapiro et al. 2013,

Brynildsrud, Bohlin et al. 2016, Collins and Didelot 2018, Saund and Snitkin 2020,

Chen and Shapiro 2021) or indirectly by correcting the analysis for population

structure (Purcell, Neale et al. 2007, Zhou and Stephens 2012, Earle, Wu et al.

2016, Lees, Galardini et al. 2018). For example, Farhat *et al* identified multiple

resistance-associated mutations and genes by applying a phylogenetic convergence

testing approach to a collection of *M. tuberculosis* isolates (Farhat, Shapiro et al.

2013). Convergent evolution was also used in GWAS of *M. tuberculosis* tolerance

(Hicks, Yang et al. 2018), and vancomycin-intermediate *S. aureus* (VISA) phenotype

(Alam, Petit et al. 2014). Approaches based on population structure (allele-counting

methods) don't account for convergent evolution explicitly, but the most robust

associations are expected to occur with homoplastic mutations or genes under

positive selection. For example, in a GWAS of drug resistance in *M. tuberculosis*,

50% of gene targets identified using regression and correction for population

structure were shown to be homoplastic using a convergence-based approach

(Farhat, Freschi et al. 2019).

The examples described above involve antibiotic resistance, an adaptive phenotype

with high heritability and strong selective pressure, where convergent evolution is

expected to be apparent. However, it is unclear whether clinical outcomes of

bacterial infections are dependent on bacterial genotypes, as shown by bacterial

GWAS of clinical outcomes, that have been either negative or have not been

reproduced across independent cohorts. Bacterial adaptation can be frequently

detected in clinical infections (Culyba and Van Tyne 2021), leading to antibiotic

resistance, antibiotic tolerance, or immune evasion, and thus potentially driving

treatment failure (Balaban, Helaine et al. 2019). Here, we propose a statistical

genomics framework that incorporates convergent evolution explicitly and tracks

molecular signatures of adaptation across three independent SAB patient cohorts to

build a model of bacterial genotype-phenotype associations for treatment outcomes.

Given the complexity of identifying genetic correlates of clinical outcomes, we

progress our analysis from outcomes with high expected role of adaptation

(vancomycin MIC) to intermediate outcomes (duration of bacteraemia) to final

endpoints with strong multi-factorial pathogenesis (treatment success, mortality).

## RESULTS

### *S. aureus* bacteraemia cohorts

To identify bacterial genetic correlates of clinical outcomes of bacteraemia, we analysed 1,358 *S. aureus* sequences from 3 cohorts of SAB with extensive clinical metadata. The cohorts included an observational study of vancomycin efficacy in SAB (cohort A, 843 isolates, 738 individual episodes), an observational study of the VISA/heterogenous VISA phenotype in sequence type (ST) 239 methicillin resistant *S. aureus* (MRSA) (cohort B, 296 isolates) and a randomised clinical trial of vancomycin-β-lactam combination therapy in MRSA bacteraemia (cohort C, 324 isolates). The microbiological and clinical characteristics of the isolates are described in Table 1 and Figure 1. The cohorts were different in terms of proportion of MRSA, distribution of sequence-types, mortality outcomes and treatment choice. Therefore, rather than combining all sequences in a single dataset, we analysed the three collections separately, with the aim of finding overlapping hits between the independent samples that would be provide robust evidence for clinically relevant bacterial adaptation.

### Lineage effects and bacterial adaptation may impact clinical outcomes of *S. aureus* bacteraemia

We first assessed the relative impact of clinical factors on SAB outcomes using cohort A because it included all three outcomes of interest for this study (vancomycin MIC, duration of bacteraemia, mortality) (543 episodes) and then fitted a classification model of mortality. In our initial model we included predictors of SAB mortality that are generally available in the clinic including host factors (age, gender,

comorbidities, site of acquisition), anti-staphylococcal treatment, phenotypic bacterial

characteristics (methicillin-resistance, vancomycin MIC) and duration of bacteraemia.

We used a random forest machine learning algorithm that was trained using stratified

cross-validation on 80% of cohort A episodes. To increase the accuracy of the

predictive parameters estimates the model was trained on 100 replicate training-

testing data splits and produced a mean area under the receiver-operating curve

(AUROC) of 0.803 and mean area under the precision-recall curve of 0.395

(AUPRC) (Figure 2A). Thus, traditional predictors of mortality in SAB had an

acceptable predictive performance (Lemeshow, Sturdivant et al. 2013), however,

20% of mortality predictions were incorrect, raising the possibility that the addition of

non-clinical variables (including bacterial genomic features) could increase prediction

accuracy.

To estimate the relative role of these predictors of mortality, we calculate variable

importance (impurity index) based on 100 replicate analyses. While age and

comorbidities were the most important features in the final model (as expected), we

were surprised that vancomycin MIC was as important as comorbidities and more

important than methicillin resistance (Figure 2C). This highlights the role of bacterial

factors in SAB mortality and suggests that adaptive antibiotic resistance is more

important than acquired antibiotic resistance genes.

To explore the relative role of bacterial genetic factors in SAB outcomes, we first

considered whether there were "high-risk" clones or lineages that were associated

with either of the three GWAS outcomes. To assess lineage effects in cohort A, we

first performed a multi-dimensional scaling analysis (MDS) of a distance matrix

generated by Mash, an approach that compares genetic similarity based on a

subsample of 1,000 kmers (Ondov, Treangen et al. 2016). This allowed to include

the whole genome content rather than just the core genome SNP matrix. We then

extracted the top 10 components (85% of genetic variance explained) and assessed

their correlation both with mortality and bacterial covariates. This correlation analysis

showed robust lineage effects for vancomycin MIC, in particular PC2 and PC3 (p <

$10^{-15}$, Figure 2D and Table S2), while no clear lineage effects were found for the

clinical outcomes. As expected, there was a strong correlation between MDS axes

and clonal complexes (CC) (Figure 1B and Figures S1 and S2). For example, the

most prevalent clonal complex (CC 239) was positively correlated with PC2 and

negatively correlated with PC1, while CC 22 was CC93 was strongly correlated with

PC3.

We then calculated the narrow-sense heritability ($h^2$) of clinical outcomes and

bacterial phenotypes to determine what proportion of these parameters was a result

of genetic factors. As expected, vancomycin MIC had a high heritability, while

duration of bacteraemia and mortality had substantially lower values (Figure 2E and

Table S1). These values are consistent bacterial GWAS of *S. aureus* infection vs.

colonisation (Young, Wu et al. 2021) and *E. coli* bacteraemia mortality (Denamur,

Condamine et al. 2022). We also sought to estimate the contribution of lineages to

the phenotypic variance by extracting from the linear mixed model the variance

attributed to the first 10 PC. We calculated that 87% of the vancomycin MIC variance

could be explained by lineage effects, while lineage effects accounted for 12% of the

mortality variance. This is consistent with the analysis of lineage-specific

associations with vancomycin, with highly significant results for first three

components (Figure 2D).

**Identification of loci associated with increased vancomycin MIC**

We first analysed loci associated with vancomycin MIC (as a continuous variable) in two cohorts where vancomycin was determined using E-test. As E-test provides a more granular estimate of vancomycin MIC than broth microdilution (Holmes, Johnson et al. 2012), we expected to have an increased power to detect associations using a linear regression model. The MIC distribution was slightly shifted between the two cohorts with higher median MIC in cohort A than cohort B (1.5 mg/l vs 1mg/l, table 1, figure 1C), however, the shape of the MIC distribution was similar, suggesting that it would be possible to compare the results from the analyses performed in both cohorts separately.

Consistent with the heritability estimate, we identified several mutations that were significantly associated with vancomycin MIC, however the analysis was strongly biased by the population structure, as shown by the quantile-quantile plot (Figure S3). While this indicates inflation (despite correction for population structure in linear mixed model) and confirms the critical role of lineage effects in antibiotic resistance in *S. aureus*, close inspection of variants reaching genome-wide significance showed mutations in genes *mprF* (E269D, *mprF* G637S, *pbp2* A246T) known to be associated with antibiotic resistance in *S. aureus* (Machado, Seif et al. 2021), confirming the ability of this approach to identify resistance-associated loci (Table 2). In addition, we found an intergenic mutation 345 bp upstream of *mprF*, further supporting the importance of *mprF* in our datasets.

We also considered significant mutations outside known vancomycin resistance loci to explore whether our GWAS had the ability to identify new targets of vancomycin resistance. The most significant association in cohort A was found with the mutation V141I in the iron-sulfur cluster synthesis gene *sufU* (Roberts, Al-Tameemi et al.

2017) and with an intergenic mutation at position 1,293,856 of the reference genome

*S. aureus* BPH2947 that was located within a promoter upstream of the predicted

operon that contains the serine-threonine phosphatase *stp1*, which has been

previously associated with vancomycin resistance (Cameron, Ward et al. 2012,

Passalacqua, Satola et al. 2012). Only one locus reached genome-wide significance

in cohort B: an intergenic mutation at position 2,959,009 within a predicted promoter

upstream of the aureolysin gene *aur*.

Inflation of p values and incomplete correction for population structure indicated that

GWAS at variant level was not sufficiently accurate to identify new targets.

Moreover, it is known from within-host adaptive studies that convergence at

mutational level is weak (Young, Wu et al. 2017). Therefore, we aggregated

mutations at the gene level, an approach that has been used by others (Baines, Holt

et al. 2015, Lees, Galardini et al. 2018, Saund and Snitkin 2020) and that may

implicitly consider convergent evolution since different mutations are expected to

arise independently across the phylogeny. Because the effect of non-synonymous

substitutions could be both a gain or loss of functions, we analysed protein-modifying

(all mutations excluding synonymous substitutions) and truncations separately and

aggregated only rare mutations (allele fraction below 1%).

This analysis revealed 8 genes that reached genome-wide significance. Again, some

of these genes (such as *walK and vraG* in cohort B) are known to be associated with

vancomycin resistance but were not identified in the mutations analysis, highlighting

the potential of the gene-burden test to increase the power of GWAS. A strong

association was found with the SAUSA300_0956 locus, a GNAT-family

acetyltransferase (Majorek, Osinski et al. 2017, Whaley, Radka et al. 2021).  The

function of the protein in *S. aureus* is not well understood, however, some indirect

evidence may point to a role in cell-wall synthesis or turn-over. First, it is located

between the autolysin *atl* and the methicillin resistance factor *fmtA (Qamar and*

*Golemi-Kotra 2012).* Second, an *E. coli* homolog of SAUSA300_0956 was linked to

membrane biosynthesis (Whaley, Radka et al. 2021). Another interest association

involved the SAUSA300_1220 locus. This gene (also known as *desR*) was recently

found to be potentially associated with regulation of β-lactam resistance in MRSA

(El-Halfawy, Czarny et al. 2020).

**Gene-burden GWAS identifies novel potential genetic signatures of**

**persistence in *S. aureus* bacteraemia**

We next assessed genetic correlates of duration of bacteraemia in cohort A and C.

As expected, the distribution of this phenotype was skewed with 59 % of strains

being associated with a duration of bacteraemia of 1 day or less (Figure 1D). Using

an automated normalisation optimisation (based on Pearson P statistics), we found

that none of the most common available strategies could improve the distribution of

the variable (Figure S4).

No single site or mutation achieved genome-wide significance in accordance with a

lower heritability and an expected polygenic signal associated with this complex trait

(Dataset S3). However, we identified loci that were associated with duration of

bacteraemia on the truncation burden test (gene-based and direction of change

analysis). The strongest association in cohort A was found with protein truncations in

*murQ*, an esterase that is part of the peptidoglycan recycling machinery (Borisova,

Gaupp et al. 2016). Mapping of *murQ* truncations showed that they were

independently acquired multiples times across the phylogeny, suggesting convergent

evolution (Figure 4B).

We found two significant associations in cohort C. These involved truncations in

*metN2*, a methionine transporter, and protein-modifying mutations in *plsX*. The latter

gene is a phosphate acyltransferase and is part of the phospholipid synthesis

pathway (Whaley, Radka et al. 2021). It interacts with the fatty-acid kinase *fakA*

which has been linked to virulence factors expression (Parsons, Broussard et al.

2014) and resistance to antimicrobial peptides (Li, Rigby et al. 2009).

**Bacterial genomic predictors don't improve mortality prediction of *S. aureus***

**bacteraemia**

We assessed whether there were any mutations or loci associated with SAB

mortality in the three cohorts and the combined dataset. Only in cohort C could we

detect a genome-wide significant association with rare mutations in four putative

metabolic genes (SAUSA300_1088, *hxlA*, *glcU*, *ctaB*) (Figure S5).

This suggested that single locus effects don't have a strong impact on mortality, or

that this impact couldn't be detected with datasets of our size. To investigate whether

combination of mutations in multiple genetic loci could influence SAB mortality, we

assessed the predictive power of bacterial genetic variation in our classification

model of mortality. Since dimensionality can affect the performance of machine

learning algorithms, we didn't assess the single variants, but we rather built a

genotype matrix with all mutated core genes with a score based on the predicted

impact of the mutation (lowest score for no mutation, highest score for predicted

protein truncations). The performance of this unfiltered gene-based approach was

only slightly higher than chance alone (mean AUROC 0.56, mean AUPRC 0.17), and

did not improve when restricting to rare mutations, using the same approach as with

the gene-burden GWAS (mean AUROC 0.52, mean AUPRC 0.17). We also

considered a model that used only the first ten principal components with similar

predictive performance (mean AUROC 0.58, mean AUPRC 0.21). Furthermore, the

performance of the model based on clinical predictors was not improved by adding

genomic predictors (mean AUROC 0.77-0.78, mean AUPRC 0.31-0.35).

**DISCUSSION**

We used a multifaceted strategy including bacterial GWAS and supervised machine

learning to assess the role of bacterial genetic factors in clinical outcomes of *S.*

*aureus* bacteraemia. We found that the bacterial genotype has limited impact on

these outcomes, in particular mortality, as demonstrated by low heritability, poor

predictive performance in our classification model of SAB and limited number of

significant findings in the bacterial GWAS. Despite the weak genotype-outcomes

association, we were able to identify a small number of candidate patho-adaptive loci

associated with duration of bacteraemia, an important intermediate outcome of SAB

that can reflect treatment response. In addition, our GWAS approach expanded the

current picture of the the polygenic background of vancomycin resistance in clinical

*S. aureus* with several new genetic loci implicated.

Identifying causative genomic correlates of bacterial phenotypes has been difficult

even when dealing with phenotypes with expected strong heritability (like antibiotic

resistance) (Chen and Shapiro 2021) but it is particularly challenging for clinical

phenotypes such as invasiveness, treatment response or infection-related mortality

(Lapp, Han et al. 2021, Young, Wu et al. 2021). These outcomes are expected to be

multifactorial and the relative contribution of bacterial genetics can be extremely

variable. Moreover, because the size of the datasets are usually small, negative

findings might reflect an insufficient sample size, not an absence of contributing

bacterial effects.

Among these multifactorial, clinical outcomes, invasiveness is thought to have a

detectable bacterial molecular signature (Lees, Ferwerda et al. 2019). This is

because even if bacterial invasion was exclusively determined by host and

environmental factors (or by chance), some form of adaptive evolution would be

expected as the result of the transition from the colonising niche to blood or human

tissues (Culyba and Van Tyne 2021), as confirmed by within-host evolution studies

where a proportion of invasive isolates carried molecular signatures of adaptation

that were non present in colonising isolates (Lees, Kremer et al. 2017, Young, Wu et

al. 2017). In accordance with this premise, two adequately sized bacterial GWAS of

*S. pneumoniae* meningitis have identified point mutations in both pathogenesis and

antibiotic resistance genes that were significantly associated with bacterial meningitis

(Lees, Ferwerda et al. 2019, Li, Metcalf et al. 2019). Strikingly, in one study the

association was replicated in an independent cohort (Li, Metcalf et al. 2019). In their

study, Lees *et al* estimated the heritability of bacterial meningitis was estimated at a

staggering 70% (Lees, Ferwerda et al. 2019). By contrast, heritability of *S. aureus*

invasiveness was much lower (4%) in the GWAS study by Young et al (Young, Wu

et al. 2021), where they identified a highly-significant association between mutations

in the dihydrofolate reductase gene *dfrB* (associated with antibiotic resistance). This

result is consistent with the association of penicillin-binding protein mutations with

invasive *S. pneumoniae* infections and suggests an intriguing association between

antibiotic resistance and invasiveness in bacteria, possibly due to the pleomorphic

effects of antibiotic resistance and its impact on immune evasion and persistence

(Gao, Cameron et al. 2013, Guerillot, Kostoulias et al. 2019).

The attempt of applying bacterial GWAS to infection outcomes (such as mortality)

has been less successful than antibiotic resistance mechanisms. While Recker *et al*,

were able to identify some molecular correlates of mortality from SAB (Recker,

Laabei et al. 2017), a Danish study found no association between bacterial genetic

determinants and occurrence of infectious endocarditis during SAB (Lilje,

Rasmussen et al. 2017). Similarly, Lees *et al* found a stark contrast between the high

heritability of the invasiveness phenotype and the null heritability of the meningitis

outcome phenotype (Lees, Ferwerda et al. 2019). In a bacterial GWAS of a cohort of

912 patients with *Escherichia coli* bacteraemia, no bacterial genetic variant was

associated with mortality, septic shock or intensive care unit admission. This

correlated with very low heritability estimates for these outcomes (Denamur,

Condamine et al. 2022).

Our heritability estimates and machine learning model confirmed the weak

association between bacterial genotype and clinical outcomes. However, we

hypothesised that it was possible to detect a role for bacterial variation based on the

assumption that niche adaptation of invasive *S. aureus* promotes antibiotic

resistance or tolerance. Building on this framework we performed a multi-cohort

GWAS with a series of phenotypes with expected decreasing importance of bacterial

factors (vancomycin MIC, duration of infection, treatment response and mortality).

Specifically, we focused on the vancomycin MIC, because this phenotype was an

important feature in our random forest model of mortality from SAB, in agreement

with previous analyses of the cohort A (Holmes, Turnidge et al. 2011). In addition,

vancomycin MIC is an example of adaptative phenotype that is relatively easy to

measure and that is available for hundreds of clinical isolates. We then progressed

to duration of bacteraemia, a key intermediary outcome of SAB (Kuehl, Morata et al.

2020) that is related to antibiotic efficacy and is often integrated into clinical

management.

We first identified a key role of lineage effects, in particular the vancomycin MIC

phenotype. Identifying lineage effects is important because they provide an estimate

of the risk of false positive GWAS finding related to the confounding from the

population structure (Earle, Wu et al. 2016). On the other hand, the presence of

lineage effects might signify that there are high-risk clones for resistance or virulence

phenotype, a feature that is important for pathogen surveillance and possibly for

outcome prediction (Lam, Wick et al. 2021).

Previous GWAS of vancomycin MIC have shown the role of known determinants of

vancomycin resistance including *walKR* (Baines, Holt et al. 2015), *rpoB* (Alam, Petit

et al. 2014). Here, in a large GWAS of vancomycin MIC, we confirm the role of some

of these established determinants of vancomycin resistance (*walK, mprF*), and

provide evidence for new loci, the most promising being *desR*, a response regulator

(Kim, Kim et al. 2016) that was recently singled out in a screening for targets of an

antivirulence molecule that is able to reverse beta-lactam resistance in MRSA strains

(El-Halfawy, Czarny et al. 2020). Our results confirm the potential for bacterial

GWAS to assist the discovery of non-canonical antibiotic resistance loci (Farhat,

Freschi et al. 2019).

Overall, only a few molecular markers achieve genome-wide significance in our

GWAS of SAB mortality, indicating that larger sample sizes including thousands of

sequences will be necessary to draw firm conclusion on the impact the bacterial

genotype on mortality. Therefore, we considered duration of bacteraemia, an

intermediate outcome that has strong association with mortality (Kuehl, Morata et al.

2020). We found that truncations in *murQ*, a gene degrades MurNAc-6p, an

intermediate metabolite in peptidoglycan recycling (Borisova, Gaupp et al. 2016),

were significantly associated with persistent SAB, potentially suggesting a role in

antibiotic resistance or tolerance. Two candidate loci of persistent bacteraemia were

the methionine biosynthesis locus *metN2* and the phospholipid biosynthesis gene

*plsX*. MetN2 is an ABC-transporter; a potential clue to its role in bacteraemia comes

from a study where expression of *metN2* was strongly reduced in *S. aureus* mutants

that were resistant to antimicrobial peptides (Kubicek-Sutherland, Lofton et al. 2017).

Interestingly, *plsX* has also a potential association with antimicrobial peptide

resistance: it is closely connected with the fatty acid kinase *fakA* (Parsons,

Broussard et al. 2014), which was found to be linked to resistance to the

antimicrobial peptide dermcidin in a transposon mutant screen (Li, Rigby et al. 2009).

There is ongoing uncertainty around the role of bacterial genetic factors in human

infections. Here we show that despite a likely preponderant role for host and

treatment factors in shaping infection outcomes, it is also possible to disentangle the

multifactorial networks underlying these infections and detect specific bacterial

genetic loci that help drive particular clinical outcomes. While these single loci likely

have a low global predictive value, they might be more relevant in specific, high-risk

settings (e.g. persistent or recurrent infection) and might act in concert with other

bacterial variants, host genetic variants and treatment factors. In addition, the

genetic loci that were enriched in our analysis might represent new clinically relevant

molecular targets of antibiotic resistance and tolerance and important candidates for

functional validation.

**METHODS**

**Study cohorts**

Cohort A was assembled from two multicentre observational studies of SAB that

were performed to assess the impact of vancomycin MIC on clinical outcomes

(Giulieri, Baines et al. 2018). The vancomycin substudy of the Australian and New

Zealand Cooperative on Outcome in Staphylococcal Sepsis (ANZCOSS) study

included 532 cases of SAB from a larger retrospective study of SAB between 2007-

2008 (Turnidge, Kotsanas et al. 2009). The selection of episodes for the substudy

has been described previously (Holmes, Turnidge et al. 2011). Briefly, to ensure a

balanced proportion of cases, each SAB case treated with vancomycin was matched

with the next available case treated with flucloxacillin at each participating centre.

The Vancomycin Efficacy in Staphylococcal Sepsis in Australasia (VANESSA) study

included prospectively 226 patients with SAB between 2012-2013 (Holmes,

Robinson et al. 2018). Data collected within cohort A included patient demographics,

site of acquisition, comorbidities, clinical syndrome, antibiotic treatment, 30-day

mortality and duration of bacteraemia. Baseline isolates were collected for all

patients, while supplementary isolates (colonising isolates from nasal swabs and

serial blood isolates) were obtained when available. Cohort A had

Cohort B included 296 episodes of MRSA bacteraemia due to sequence type 239

that were collected at a single centre in Australia between 1996-2008 to assess the

impact of vancomycin MIC and vancomycin heteroresistance on 30 day mortality

(van Hal, Jones et al. 2011).

Cohort C included data and sequences from the CAMERA2 trial (Combination

Antibiotics for Methicillin Resistant *Staphylococcus aureus*) that was performed

between 2015 and 2018 in Australia, New Zealand, Singapore and Israel, and

randomised participants with MRSA bacteraemia to either monotherapy with

vancomycin or daptomycin or combination therapy with vancomycin or daptomycin

plus an antistaphylococcal beta-lactam (flucloxacillin, cloxacillin or cefazolin) (Tong,

Lye et al. 2020). The primary outcome was a composite endpoint ascertained 90

days after randomisation and included mortality, persistent bacteraemia after 5 days,

microbiological relapse (positive blood culture after a preceding negative culture),

and microbiological treatment failure (positive sterile site culture 14 or more days

after randomisation). In addition, the trial collected information on the day of death,

duration of bacteraemia and day of relapse. Relevant clinical data assessed were

comorbidites, severity scores, primary source and secondary foci, source control.

Details of the antibiotic treatment were collected up to day 90.

**Phenotypic testing**

Bacterial isolates were stored in glycerol broth at -80C, after confirmation of *S.*

*aureus* using coagulase and DNase tests or Matrix-assisted laser desorption-

ionisation time-of-flight (MALDI-TOF). Vancomycin minimum inhibitory concentration

(MIC) was determined by E-test (bioMerieux) following manufacturer's instructions.

An inoculum of 0.5 McFarland was used. Vancomycin E-test was only performed for

cohort A and B strains.

**Machine learning model to select predictors of mortality/treatment response**

Using data from cohort A, we fitted a classification model of 30-day mortality of SAB to assess the predictive performance of non-genomic features including demographic and clinical data (age, Pitt bacteraemia score, Charlson comorbidity index, site of acquisition, gender), microbiological phenotypes (methicillin resistance, vancomycin MIC) and treatment response variables (duration of bacteraemia). The R package tidymodels was used to build a random forest model of SAB mortality. The model was trained on 80% of the data using stratified 10-fold cross-validation and hyperparameter tuning to optimise the area under the receiver operator characteristics curve (AUROC). Tuning of the random forest hyperparameters "mtry" (number of randomly selected predictors) and "min_n" (minimal node size) was done using a 5x5 grid, with range of hyperparameters determined empirically based on initial tuning using 20 automatically selected candidate combinations. Categorical predictors were converted into binary variables an all numeric predictors were centered and scaled. Zero variance predictors were removed. Downsampling was used to compensate for class inbalance (mortality rate 15%). The model was fitted on 100 replicated data-splits to generate estimates of the performance and variable importance as in (Lapp, Han et al. 2021).

**Whole-genome sequencing**

Bacterial strains were subcultured twice from -80C glycerol stock. The Janus® automated workstation (PerkinElmer) or manual extraction kits (Invitrogen PureLink genomic DNA kit or the Sigma GenElute kit) were used to extract genomic DNA from single colonies. DNA concentrations were normalised to 0.2 ng/ul for library preparation using Nextera® XT DNA (Illumina). Whole-genome sequencing was performed on the Illumina MiSeq and NextSeq platforms. Reads were assembled

using Shovill v1.1.0 (https://github.com/tseemann/shovill), a pipeline that streamlines

and optimises Spades (58). To assess the quality of the sequences, mean read

depth, assembly metrics and the percentage of *S. aureus* reads using Kraken 2

v2.1.1 (60), were calculated. Reads were discarded if read depth was < 20, the

fraction of *S. aureus* reads was < 80% or the size of the assembly was > 3.5 Mb or <

2.5 Mb. We used Mash v2.3 to generate a distance matrix from the Shovill

assemblies using 1,000 sketches.

**Variant calling**

Reads were mapped to reference genome *S. aureus* BPH2947 (accession

GCF_900620245.1) using Snippy v4.6.0 (https://github.com/tseemann/snippy ). For

each dataset, a core genome alignment including all positions with at least 90% non-

gaps was constructed using Snippy-core v4.6.0, Goalign v0.3.4 (Lemoine and

Gascuel 2021) and SNP-sites v2.5.1 (Page, Taylor et al. 2016). The alignment was

then used to infer a maximum-likelihood phylogenetic tree with IQ-TREE v2.0.3

(Minh, Schmidt et al. 2020) under a GTR+G4 model. Support was calculated using

the ultrafast support feature and SH-aLRT test. For each variant at positions

included in the core genome alignment, we computed the number of independent

acquisitions across the phylogeny using HomoplasyFinder (Crispell, Balaz et al.

2019).

**Preparation of the bacterial genotype files**

For each dataset, variants at core genome positions (minimum 90% of sequences

with a coverage of at least 10 reads) were merged in a single variant call format

(VCF) file using BCFtools v 1.10.2 (Danecek, Bonfield et al. 2021), without merging

multiallelic records. A catalogue of mutations with predicted impact on protein

function was generated by masking synonymous variants were masked using the

command bcftools view -e 'ANN[0] ~ "[^&]synonymous"'. Mutations with predicted

loss of function (LOF) were extracted using the command bcftools view -i 'ANN[0] ~

"frameshift_variant" | ANN[0] ~ "stop_gained" | ANN[0] ~ "stop_lost" | ANN[0] ~

"start_lost"'.

**Variant annotation**

In addition to the variant annotation based on BPH2947 generated by Snippy, we

used Blast v 2.10.1 to determine homologs in reference genome FPR3757 that were

linked to the AureoWiki (Fuchs, Mehlan et al. 2018) and MicrobesOnline (Dehal,

Joachimiak et al. 2010) databases. Blast hits were kept if alignment coverage was

above 50 % and identity was above 90 %. When multiple hits were available, the hits

with highest identity were selected.

**Bacterial GWAS design**

We ran bacterial GWAS to test the association between mutations with predicted

impact on protein function (filtered as described above) with the phenotypes of

interest, as continuous (vancomycin MIC, duration of bacteraemia) or binary traits

(30 day mortality). For clinical outcomes, we analysed one sample per episode. For

vancomycin MIC GWAS, we included more than one sample per episode when they

were available. For duration of bacteraemia, potential transformations to achieve a

normal distribution were assessed using the bestNormalize package in R (Peterson

and Cavanaugh 2020).

**Bacterial GWAS pipeline**

We used the factored spectrally transformed linear mixed models (FaST-LMM) as

implemented in Pyseer v 1.1.2 (Lees, Galardini et al. 2018), which allows to correct

for population structure by adding a kinship matrix as a random effect. The kinship

matrix was computed from a VCF file including all core-genome variants using the

Gemma, version 0.98.1 (Zhou and Stephens 2012). We assessed phenotype

associations both with individual mutations and with protein-modifying and loss-of-

function aggregated in genes, operons and gene ontology categories as

implemented with the '--burden' option in Pyseer. For the mutation GWAS, we

excluded synonymous substitutions and mutations that were acquired only once

across the phylogeny (based on homoplasy counts obtained using HomoplasyFinder

(Crispell, Balaz et al. 2019)). For the gene-burden GWAS, only rare variants (i.e.

those with a minim allele fraction below 1%) were considered.

**Lineage effects and heritability.**

Heritability (SNP-heritability, $h^2$) - the proportion of phenotypic variance that is

correlated with genotypic variance - was estimated by Pyseer from the linear mixed

model using the kinship matrix computed in Gemma (see above). Lineages of cohort

A isolates were inferred from the Mash distance matrix using the 'cmdscale' function

in R. The association between the first 10 components and the bacterial GWAS

outcomes was computed by running Pyseer with the '—lineage' option. Correlation

between the 10 first components and internal nodes of the tree or major clonal

complexes was calculated as performed in the R package bugwas

([https://github.com/sgearle/bugwas/blob/master/bugwas/R/BUGWAS_functions.R#L](https://github.com/sgearle/bugwas/blob/master/bugwas/R/BUGWAS_functions.R#L)

[962](https://github.com/sgearle/bugwas/blob/master/bugwas/R/BUGWAS_functions.R#L962)) (Earle, Wu et al. 2016).

**Ethics approval**

Human research committee approval was obtained at each site for the three cohorts.

Written informed consent was collected for the prospective component of cohort A

(VANESSA) and for cohort C (CAMERA2).

**Code availability**

The code for the preparation of bacterial genotype files and for running Pyseer is

available at https://github.com/stefanogg/CLOGEN.

**Data availability**

The raw reads of the strains included in this analysis are available in the European

Nucleotide Archive under Bioproject accession numbers PRJEB27932 (cohort A),

PRJEB50796 (cohort B) and PRJEB2561 (cohort C).

**TABLES**

Table 1. Characteristics of *S. aureus* isolates included in the bacteraemia cohorts

| | Cohort A (n = 738) | Cohort B (n = 296) | Cohort C (n = 324) |
|---|---|---|---|
| Sample period | | | |
| 1990-1999 | 0 | 32 (11) | 0 |
| 2000-2009 | 517 (70) | 264 (89) | 0 |
| 2010-2019 | 221 (30) | 0 | 324 (100) |
| Country | | | |
| Australia/NZ | 738 (100) | 296 (100) | 239 (74) |
| Singapore | 0 | 0 | 53 (16) |
| Israel | 0 | 0 | 32 (10) |
| Patient age (years), median (IQR) | 61 (46-74) | 64 (50-73) | 64 (49-76) |
| MRSA | 252 (34) | 296 | 324 |
| Vancomycin MIC (mg/L), median (IQR) | 1.5 (1-2) | 1 (0.75-1.5) | N/A |
| Main antibiotic | | | |
| Flucloxacillin | 384 (52) | 0 | 0 |
| Vancomycin | 301 (41) | 252 (85) | 158 (49) |
| Daptomycin | 0 | 0 | 3 (1) |
| Vancomycin +β-lactam | 0 | 0 | 159 (49) |
| Daptomycin+ β-lactam | 0 | 0 | 4 (1) |
| Other | 53 (7) | 11 (4) | 0 |
| No treatment | 0 | 28 (9) | 0 |
| Duration of bacteraemia (days), median (IQR) | 1 (1-2) | N/A | 1 (1-3) |
| Mortality at 30 days | 109 (15) | 88 (30) | 35 (11) |

580 Values are counts (percent) unless otherwise specified.

581 MRSA: methicillin-resistant *S. aureus*. N/A: not available.

583     Table 2. Selection of loci significantly associated with vancomycin MIC. Loci are

584     displayed if they have the lowest p value within the analysis or if they are located

585     near or in known vancomycin resistance loci

| | | | | P value (effect size) | | |
|---|---|---|---|---|---|---|
| Reference position | Gene | Product | Mutation | Cohort A | Cohort B | Combined analysis |
| 976503 | *sufU* | NifU family SUF system FeS assembly protein | V141I | $3.2 \times 10^{-12}$ (0.886) | | $7.5 \times 10^{-9}$ (0.877) |
| 1293856 | intergenic | Upstream of *def2-fmt-sun-rlmN-stp1-pknB* operon | | | | $6.51 \times 10^{-12}$ (0.779) |
| 2959009 | intergenic | Upstream of *aur* gene | | | $1.12 \times 10^{-4}$ (0.737) | |
| 1458535 | intergenic | Upstream of mprF | | $1.3 \times 10^{-11}$ (0.837) | | $5.52 \times 10^{-8}$ (0.827) |
| 1459663 | *mprF* | oxacillin resistance-related FmtC protein | E269D | $1.04 \times 10^{-7}$ (0.74) | | |
| 1460765 | *mprF* | oxacillin resistance-related FmtC protein | G637S | | | $7.12 \times 10^{-7}$ (-0.479) |
| 1583057 | *pbp2* | penicillin binding protein 2 | A246T | | | $7.12 \times 10^{-7}$ (-0.479) |

**FIGURES**

**Figure 1.** Genomic and phenotypic characterisation of 3 *S. aureus* bacteraemia cohorts. **(**A) Maximum likelihood phylogenetic trees of 1,358 bloodstream *S. aureus* strains included in the analysis. (B) First two principal components of multi-dimensional scaling (MDS) of the Mash distance matrix. Dots represent sequences and are coloured according to the 7 most prevalent clonal complexes (C) Distribution of vancomycin MIC (in mg/L). (D) Distribution of duration of bacteraemia (in days).

**Figure 2.** (A, B) Performance of the random forest model of 30-day mortality when

using clinical and microbiological phenotypic data only. The model was trained and

tested using cohort A data. The performance metrics (area under the curve [AUC]) in

the test dataset were calculated from 100 replicate analyses. Panel A: receiver

operator characteristic curve (ROC). Panel B: precision-recall curve (C) Variable

importance plot (Gini impurity index) based on 100 replicate analyses. (D) Lineage

effects for the vancomycin MIC, duration of bacteraemia and the SAB mortality

outcomes for cohort A. The bars show the significance of the association between

the ten first principal components (based on multidimensional scaling) and the

outcome. (E) Narrow-sense heritability (SNP-heritability) of the GWAS outcomes of

interest: vancomycin MIC, duration of bacteraemia (persistence) and SAB mortality.

**Figure 3.** Gene-burden GWAS of vancomycin MIC in cohorts A, B and A and B combined. (no vancomycin E-test data were available for cohort C). The association was computed using a linear mixed model with vancomycin MIC as continuous outcome variable. All mutations with predicted impact on the protein sequence were aggregated by coding regions (circles: all protein-modifying, triangles: protein-truncating only). The size of the points represents the fractions of strains with a mutated genes and the colour indicated the effect size (impact on vancomycin MIC) in the linear regression.

**Figure 4.** Gene-burden GWAS of duration of bacteraemia in cohorts A and C and A and C combined (duration of bacteraemia was not recorded for cohort B). (A) Manhattan plot showing the strength of the significance of mutated genes. The association was computed using a linear mixed model with duration of bacteraemia as continuous outcome variable. All mutations with predicted impact on the protein sequence were aggregated by coding regions (circles: all protein-modifying, triangles: protein-truncating only). The size of the points represents the fractions of strains with a mutated genes and the colour indicated the effect size (impact on duration of bacteraemia) in the linear regression. (B-D) Positions of *murQ* (B), *plsX* (C) and *metN2* (D) mutations on the phylogenetic tree.

Figure 5. Predictive performance of seven random forest models of SAB mortality in cohort A. The data shown are the summary of 100 replicated random data-splits. Error bars represent mean and standard deviation. Panel A: area under the receiver operating characteristic curve (AUROC). Panel B: area under the precision-recall curve (AUPRC).

**REFERENCES**

Alam, M. T., R. A. Petit, 3rd, E. K. Crispell, T. A. Thornton, K. N. Conneely, Y. Jiang, S. W. Satola and T. D. Read (2014). "Dissecting vancomycin-intermediate resistance in staphylococcus aureus using genome-wide association." Genome Biol Evol **6**(5): 1174-1185.

Bai, A. D., C. K. L. Lo, A. S. Komorowski, M. Suresh, K. Guo, A. Garg, P. Tandon, J. Senecal, O. Del Corpo, I. Stefanova, C. Fogarty, G. Butler-Laporte, E. G. McDonald, M. P. Cheng, A. M. Morris, M. Loeb and T. C. Lee "<em>Staphylococcus aureus</em> bacteremia mortality: A systematic review and meta-analysis." Clinical Microbiology and Infection.

Baines, S. L., K. E. Holt, M. B. Schultz, T. Seemann, B. O. Howden, S. O. Jensen, S. J. van Hal, G. W. Coombs, N. Firth, D. R. Powell, T. P. Stinear and B. P. Howden (2015). "Convergent adaptation in the dominant global hospital clone ST239 of methicillin-resistant Staphylococcus aureus." MBio **6**(2): e00080.

Balaban, N. Q., S. Helaine, K. Lewis, M. Ackermann, B. Aldridge, D. I. Andersson, M. P. Brynildsen, D. Bumann, A. Camilli, J. J. Collins, C. Dehio, S. Fortune, J. M. Ghigo, W. D. Hardt, A. Harms, M. Heinemann, D. T. Hung, U. Jenal, B. R. Levin, J. Michiels, G. Storz, M. W. Tan, T. Tenson, L. Van Melderen and A. Zinkernagel (2019). "Definitions and guidelines for research on antibiotic persistence." Nat Rev Microbiol.

Battle, S. E., M. Shuping, S. Withers, J. A. Justo, P. B. Bookstaver and M. N. Al-Hasan (2021). "Prediction of Mortality in Staphylococcus aureus Bloodstream Infection using Quick Pitt Bacteremia Score." J Infect.

Borisova, M., R. Gaupp, A. Duckworth, A. Schneider, D. Dalugge, M. Muhleck, D.

Deubel, S. Unsleber, W. Yu, G. Muth, M. Bischoff, F. Gotz and C. Mayer (2016).

"Peptidoglycan Recycling in Gram-Positive Bacteria Is Crucial for Survival in

Stationary Phase." mBio **7**(5).

Brynildsrud, O., J. Bohlin, L. Scheffer and V. Eldholm (2016). "Rapid scoring of

genes in microbial pan-genome-wide association studies with Scoary." Genome

Biology **17**(1): 238.

Cameron, D. R., D. V. Ward, X. Kostoulias, B. P. Howden, R. C. Moellering, Jr., G.

M. Eliopoulos and A. Y. Peleg (2012). "Serine/threonine phosphatase Stp1

contributes to reduced susceptibility to vancomycin and virulence in Staphylococcus

aureus." J Infect Dis **205**(11): 1677-1687.

Chen, P. E. and B. J. Shapiro (2021). "Classic genome-wide association methods

are unlikely to identify causal variants in strongly clonal microbial populations."

bioRxiv: 2021.2006.2030.450606.

Cohen, J., J. L. Vincent, N. K. Adhikari, F. R. Machado, D. C. Angus, T. Calandra, K.

Jaton, S. Giulieri, J. Delaloye, S. Opal, K. Tracey, T. van der Poll and E. Pelfrene

(2015). "Sepsis: a roadmap for future research." Lancet Infect Dis **15**(5): 581-614.

Collins, C. and X. Didelot (2018). "A phylogenetic method to perform genome-wide

association studies in microbes that accounts for population structure and

recombination." PLOS Computational Biology **14**(2): e1005958.

Crispell, J., D. Balaz and S. V. Gordon (2019). "HomoplasyFinder: a simple tool to

identify homoplasies on a phylogeny." Microbial Genomics **5**(1).

Culyba, M. J. and D. Van Tyne (2021). "Bacterial evolution during human infection:

Adapt and live or adapt and die." PLOS Pathogens **17**(9): e1009872.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A.

Whitwham, T. Keane, S. A. McCarthy, R. M. Davies and H. Li (2021). "Twelve years

of SAMtools and BCFtools." GigaScience **10**(2).

Dehal, P. S., M. P. Joachimiak, M. N. Price, J. T. Bates, J. K. Baumohl, D. Chivian,

G. D. Friedland, K. H. Huang, K. Keller, P. S. Novichkov, I. L. Dubchak, E. J. Alm

and A. P. Arkin (2010). "MicrobesOnline: an integrated portal for comparative and

functional genomics." Nucleic Acids Res **38**(Database issue): D396-400.

Denamur, E., B. Condamine, M. Esposito-Farèse, G. Royer, O. Clermont, C.

Laouenan, A. Lefort, V. de Lastours, M. Galardini, C. the and S. groups (2022).

"Genome wide association study of Escherichia coli bloodstream infection isolates

identifies genetic determinants for the portal of entry but not fatal outcome." PLOS

Genetics **18**(3): e1010112.

Earle, S. G., C.-H. Wu, J. Charlesworth, N. Stoesser, N. C. Gordon, T. M. Walker, C.

C. A. Spencer, Z. Iqbal, D. A. Clifton, K. L. Hopkins, N. Woodford, E. G. Smith, N.

Ismail, M. J. Llewelyn, T. E. Peto, D. W. Crook, G. McVean, A. S. Walker and D. J.

Wilson (2016). "Identifying lineage effects when controlling for population structure

improves power in bacterial association studies." Nature microbiology **1**: 16041-

16041.

El-Halfawy, O. M., T. L. Czarny, R. S. Flannagan, J. Day, J. C. Bozelli, R. C. Kuiack,

A. Salim, P. Eckert, R. M. Epand, M. J. McGavin, M. G. Organ, D. E. Heinrichs and

E. D. Brown (2020). "Discovery of an antivirulence compound that reverses β-lactam

resistance in MRSA." Nature Chemical Biology **16**(2): 143-149.

Farhat, M. R., L. Freschi, R. Calderon, T. Ioerger, M. Snyder, C. J. Meehan, B. de

Jong, L. Rigouts, A. Sloutsky, D. Kaur, S. Sunyaev, D. van Soolingen, J. Shendure,

J. Sacchettini and M. Murray (2019). "GWAS for quantitative resistance phenotypes

in Mycobacterium tuberculosis reveals resistance genes and regulatory regions."

<u>Nature Communications</u> **10**(1): 2128.

Farhat, M. R., B. J. Shapiro, K. J. Kieser, R. Sultana, K. R. Jacobson, T. C. Victor, R.

M. Warren, E. M. Streicher, A. Calver, A. Sloutsky, D. Kaur, J. E. Posey, B. Plikaytis,

M. R. Oggioni, J. L. Gardy, J. C. Johnston, M. Rodrigues, P. K. C. Tang, M. Kato-

Maeda, M. L. Borowsky, B. Muddukrishna, B. N. Kreiswirth, N. Kurepina, J. Galagan,

S. Gagneux, B. Birren, E. J. Rubin, E. S. Lander, P. C. Sabeti and M. Murray (2013).

"Genomic analysis identifies targets of convergent positive selection in drug-resistant

Mycobacterium tuberculosis." <u>Nature Genetics</u> **45**(10): 1183-1189.

Fuchs, S., H. Mehlan, J. Bernhardt, A. Hennig, S. Michalik, K. Surmann, J. Pané-

Farré, A. Giese, S. Weiss, L. Backert, A. Herbig, K. Nieselt, M. Hecker, U. Völker

and U. Mäder (2018). "AureoWiki ◻ The repository of the Staphylococcus aureus

research and annotation community." <u>International Journal of Medical Microbiology</u>

**308**(6): 558-568.

Gao, W., D. R. Cameron, J. K. Davies, X. Kostoulias, J. Stepnell, K. L. Tuck, M. R.

Yeaman, A. Y. Peleg, T. P. Stinear and B. P. Howden (2013). "The RpoB

H(4)(8)(1)Y rifampicin resistance mutation and an active stringent response reduce

virulence and increase resistance to innate immune responses in Staphylococcus

aureus." <u>J Infect Dis</u> **207**(6): 929-939.

Giulieri, S. G., S. L. Baines, R. Guerillot, T. Seemann, A. Gonçalves da Silva, M.

Schultz, R. C. Massey, N. E. Holmes, T. P. Stinear and B. P. Howden (2018).

"Genomic exploration of sequential clinical isolates reveals a distinctive molecular

signature of persistent Staphylococcus aureus bacteraemia." <u>Genome Medicine</u> **10**(1): 65.

Giulieri, S. G., N. E. Holmes, T. P. Stinear and B. P. Howden (2016). "Use of bacterial whole-genome sequencing to understand and improve the management of invasive Staphylococcus aureus infections." <u>Expert Rev Anti Infect Ther</u> **14**(11): 1023-1036.

Giulieri, S. G., S. Y. C. Tong and D. A. Williamson (2020). "Using genomics to understand meticillin- and vancomycin-resistant Staphylococcus aureus infections." <u>Microb Genom</u> **6**(1).

Guerillot, R., X. Kostoulias, L. Donovan, L. Li, G. P. Carter, A. Hachani, K. Vandelannoote, S. Giulieri, I. R. Monk, M. Kunimoto, L. Starrs, G. Burgio, T. Seemann, A. Y. Peleg, T. P. Stinear and B. P. Howden (2019). "Unstable chromosome rearrangements in Staphylococcus aureus cause phenotype switching associated with persistent infections." <u>Proc Natl Acad Sci U S A</u> **116**(40): 20135-20140.

Hicks, N. D., J. Yang, X. Zhang, B. Zhao, Y. H. Grad, L. Liu, X. Ou, Z. Chang, H. Xia, Y. Zhou, S. Wang, J. Dong, L. Sun, Y. Zhu, Y. Zhao, Q. Jin and S. M. Fortune (2018). "Clinically prevalent mutations in Mycobacterium tuberculosis alter propionate metabolism and mediate multidrug tolerance." <u>Nature Microbiology</u> **3**(9): 1032-1042.

Holmes, N. E., P. D. R. Johnson and B. P. Howden (2012). "Relationship between Vancomycin-Resistant Staphylococcus aureus, Vancomycin-Intermediate S. aureus, High Vancomycin MIC, and Outcome in Serious S. aureus Infections." <u>Journal of Clinical Microbiology</u> **50**(8): 2548-2552.

Holmes, N. E., J. O. Robinson, S. J. van Hal, W. J. Munckhof, E. Athan, T. M. Korman, A. C. Cheng, J. D. Turnidge, P. D. R. Johnson, B. P. Howden and o. b. o. t. A. S. f. I. D. C. R. N. Vanessa study group (2018). "Morbidity from in-hospital complications is greater than treatment failure in patients with Staphylococcus aureus bacteraemia." BMC Infect Dis **18**(1): 107.

Holmes, N. E., J. D. Turnidge, W. J. Munckhof, J. O. Robinson, T. M. Korman, M. V. O'Sullivan, T. L. Anderson, S. A. Roberts, W. Gao, K. J. Christiansen, G. W. Coombs, P. D. Johnson and B. P. Howden (2011). "Antibiotic choice may not explain poorer outcomes in patients with Staphylococcus aureus bacteremia and high vancomycin minimum inhibitory concentrations." J Infect Dis **204**(3): 340-347.

Imam, N., S. Tempone, P. K. Armstrong, R. McCann, S. Johnson, L. J. Worth and M. J. Richards (2019). "Increased incidence of community-associated Staphylococcus aureus bloodstream infections in Victoria and Western Australia, 2011–2016." Medical Journal of Australia **210**(2): 87-88.

Kim, J. W., H.-K. Kim, G. S. Kang, I.-H. Kim, H. S. Kim, Y. S. Lee and J. I. Yoo (2016). "The SAV1322 gene from Staphylococcus aureus: genomic and proteomic approaches to identification and characterization of gene function." BMC microbiology **16**(1): 206-206.

Kubicek-Sutherland, J. Z., H. Lofton, M. Vestergaard, K. Hjort, H. Ingmer and D. I. Andersson (2017). "Antimicrobial peptide exposure selects for Staphylococcus aureus resistance to human defence peptides." The Journal of antimicrobial chemotherapy **72**(1): 115-127.

Kuehl, R., L. Morata, C. Boeing, I. Subirana, H. Seifert, S. Rieg, W. V. Kern, H. B. Kim, E. S. Kim, C. H. Liao, R. Tilley, L. E. Lopez-Cortés, M. J. Llewelyn, V. G.

Fowler, G. Thwaites, J. M. Cisneros, M. Scarborough, E. Nsutebu, M. Gurgui Ferrer,

J. L. Pérez, G. Barlow, S. Hopkins, H. G. Ternavasio-de la Vega, M. E. Török, P.

Wilson, A. J. Kaasch and A. Soriano (2020). "Defining persistent Staphylococcus

aureus bacteraemia: secondary analysis of a prospective cohort study." Lancet

Infect Dis **20**(12): 1409-1417.

Kwok, A. J., A. Mentzer and J. C. Knight (2021). "Host genetics and infectious

disease: new tools, insights and translational opportunities." Nature Reviews

Genetics **22**(3): 137-153.

Lam, M. M. C., R. R. Wick, S. C. Watts, L. T. Cerdeira, K. L. Wyres and K. E. Holt

(2021). "A genomic surveillance framework and genotyping tool for Klebsiella

pneumoniae and its related species complex." Nat Commun **12**(1): 4188.

Lapp, Z., J. H. Han, J. Wiens, E. J. C. Goldstein, E. Lautenbach and E. S. Snitkin

(2021). "Patient and Microbial Genomic Factors Associated with Carbapenem-

Resistant Klebsiella pneumoniae Extraintestinal Colonization and Infection."

mSystems **6**(2): e00177-00121.

Lees, J. A., B. Ferwerda, P. H. C. Kremer, N. E. Wheeler, M. V. Seron, N. J.

Croucher, R. A. Gladstone, H. J. Bootsma, N. Y. Rots, A. J. Wijmega-Monsuur, E. A.

M. Sanders, K. Trzcinski, A. L. Wyllie, A. H. Zwinderman, L. H. van den Berg, W. van

Rheenen, J. H. Veldink, Z. B. Harboe, L. F. Lundbo, L. de Groot, N. M. van Schoor,

N. van der Velde, L. H. Angquist, T. I. A. Sorensen, E. A. Nohr, A. J. Mentzer, T. C.

Mills, J. C. Knight, M. du Plessis, S. Nzenze, J. N. Weiser, J. Parkhill, S. Madhi, T.

Benfield, A. von Gottberg, A. van der Ende, M. C. Brouwer, J. C. Barrett, S. D.

Bentley and D. van de Beek (2019). "Joint sequencing of human and pathogen

genomes reveals the genetics of pneumococcal meningitis." <u>Nat Commun</u> **10**(1): 2176.

Lees, J. A., M. Galardini, S. D. Bentley, J. N. Weiser and J. Corander (2018). "pyseer: a comprehensive tool for microbial pangenome-wide association studies." <u>Bioinformatics</u> **34**(24): 4310-4312.

Lees, J. A., P. H. C. Kremer, A. S. Manso, N. J. Croucher, B. Ferwerda, M. V. Serón, M. R. Oggioni, J. Parkhill, M. C. Brouwer, A. van der Ende, D. van de Beek and S. D. Bentley (2017). "Large scale genomic analysis shows no evidence for pathogen adaptation between the blood and cerebrospinal fluid niches during bacterial meningitis." <u>Microb Genom</u> **3**(1): e000103.

Lemeshow, S., R. X. Sturdivant, D. W. Hosmer, Jr. and D. W. Hosmer, Jr. (2013). <u>Applied Logistic Regression</u>. New York, UNITED STATES, John Wiley & Sons, Incorporated.

Lemoine, F. and O. Gascuel (2021). "Gotree/Goalign : Toolkit and Go API to facilitate the development of phylogenetic workflows." <u>bioRxiv</u>: 2021.2006.2009.447704.

Li, M., K. Rigby, Y. Lai, V. Nair, A. Peschel, B. Schittek and M. Otto (2009). "*Staphylococcus aureus* Mutant Screen Reveals Interaction of the Human Antimicrobial Peptide Dermcidin with Membrane Phospholipids." <u>Antimicrobial Agents and Chemotherapy</u> **53**(10): 4200-4210.

Li, Y., B. J. Metcalf, S. Chochua, Z. Li, H. Walker, T. Tran, P. A. Hawkins, R. Gierke, T. Pilishvili, L. McGee and B. W. Beall (2019). "Genome-wide association analyses of invasive pneumococcal isolates identify a missense bacterial mutation associated with meningitis." <u>Nature Communications</u> **10**(1): 178.

Lilje, B., R. V. Rasmussen, A. Dahl, M. Stegger, R. L. Skov, V. G. Fowler, K. L. Ng,

K. Kiil, A. R. Larsen, A. Petersen, H. K. Johansen, H. C. Schonheyder, M. Arpi, F. S.

Rosenvinge, E. Korup, U. Host, C. Hassager, S. U. A. Gill, T. F. Hansen, T. B.

Johannesen, J. Smit, P. Sogaard, P. Skytt Andersen and N. Eske-Bruun (2017).

"Whole-genome sequencing of bloodstream Staphylococcus aureus isolates does

not distinguish bacteraemia from endocarditis." Microb Genom **3**(11).

Machado, H., Y. Seif, G. Sakoulas, C. A. Olson, Y. Hefner, A. Anand, Y. Z. Jones, R.

Szubin, B. O. Palsson, V. Nizet and A. M. Feist (2021). "Environmental conditions

dictate differential evolution of vancomycin resistance in Staphylococcus aureus."

Communications Biology **4**(1): 793.

Majorek, K. A., T. Osinski, D. T. Tran, A. Revilla, W. F. Anderson, W. Minor and M.

L. Kuhn (2017). "Insight into the 3D structure and substrate specificity of previously

uncharacterized GNAT superfamily acetyltransferases from pathogenic bacteria."

Biochimica et biophysica acta. Proteins and proteomics **1865**(1): 55-64.

Mba Medie, F., B. K. Sharma-Kuinkel, F. Ruffin, L. C. Chan, M. Rossetti, Y. L.

Chang, L. P. Park, A. S. Bayer, S. G. Filler, R. Ahn, E. F. Reed, D. Gjertson, M. R.

Yeaman and V. G. Fowler, Jr. (2019). "Genetic variation of DNA methyltransferase-

3A contributes to protection against persistent MRSA bacteremia in patients." Proc

Natl Acad Sci U S A **116**(40): 20087-20096.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von

Haeseler and R. Lanfear (2020). "IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era." Molecular Biology and Evolution **37**(5):

1530-1534.

Ondov, B. D., T. J. Treangen, P. Melsted, A. B. Mallonee, N. H. Bergman, S. Koren

and A. M. Phillippy (2016). "Mash: fast genome and metagenome distance

estimation using MinHash." <u>Genome Biology</u> **17**(1): 132.

Page, A. J., B. Taylor, A. J. Delaney, J. Soares, T. Seemann, J. A. Keane and S. R.

Harris (2016). "SNP-sites: rapid efficient extraction of SNPs from multi-FASTA

alignments." <u>Microbial Genomics</u> **2**(4).

Parsons, J. B., T. C. Broussard, J. L. Bose, J. W. Rosch, P. Jackson, C.

Subramanian and C. O. Rock (2014). "Identification of a two-component fatty acid

kinase responsible for host fatty acid incorporation by *Staphylococcus aureus*."

<u>Proceedings of the National Academy of Sciences</u> **111**(29): 10532-10537.

Passalacqua, K. D., S. W. Satola, E. K. Crispell and T. D. Read (2012). "A mutation

in the PP2C phosphatase gene in a Staphylococcus aureus USA300 clinical isolate

with reduced susceptibility to vancomycin and daptomycin." <u>Antimicrob Agents</u>

<u>Chemother</u> **56**(10): 5212-5223.

Peterson, R. A. and J. E. Cavanaugh (2020). "Ordered quantile normalization: a

semiparametric transformation built for the cross-validation era." <u>Journal of Applied</u>

<u>Statistics</u> **47**(13-15): 2312-2327.

Power, R. A., J. Parkhill and T. de Oliveira (2017). "Microbial genome-wide

association studies: lessons from human GWAS." <u>Nat Rev Genet</u> **18**(1): 41-50.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J.

Maller, P. Sklar, P. I. W. de Bakker, M. J. Daly and P. C. Sham (2007). "PLINK: a

tool set for whole-genome association and population-based linkage analyses."

<u>American journal of human genetics</u> **81**(3): 559-575.

Qamar, A. and D. Golemi-Kotra (2012). "Dual roles of FmtA in Staphylococcus

aureus cell wall biosynthesis and autolysis." <u>Antimicrobial agents and chemotherapy</u>

**56**(7): 3797-3805.

Recker, M., M. Laabei, M. S. Toleman, S. Reuter, R. B. Saunderson, B. Blane, M. E.

Torok, K. Ouadi, E. Stevens, M. Yokoyama, J. Steventon, L. Thompson, G. Milne, S.

Bayliss, L. Bacon, S. J. Peacock and R. C. Massey (2017). "Clonal differences in

Staphylococcus aureus bacteraemia-associated mortality." <u>Nat Microbiol</u> **2**(10):

1381-1388.

Roberts, C. A., H. M. Al-Tameemi, A. A. Mashruwala, Z. Rosario-Cruz, U. Chauhan,

W. E. Sause, V. J. Torres, W. J. Belden and J. M. Boyd (2017). "The Suf Iron-Sulfur

Cluster Biosynthetic System Is Essential in Staphylococcus aureus, and Decreased

Suf Function Results in Global Metabolic Defects and Reduced Survival in Human

Neutrophils." <u>Infect Immun</u> **85**(6).

Saund, K. and E. S. Snitkin (2020). "Hogwash: three methods for genome-wide

association studies in bacteria." <u>Microbial genomics</u> **6**(11): mgen000469.

Tong, S. Y., J. S. Davis, E. Eichenberger, T. L. Holland and V. G. Fowler, Jr. (2015).

"Staphylococcus aureus infections: epidemiology, pathophysiology, clinical

manifestations, and management." <u>Clin Microbiol Rev</u> **28**(3): 603-661.

Tong, S. Y. C., D. C. Lye, D. Yahav, A. Sud, J. O. Robinson, J. Nelson, S. Archuleta,

M. A. Roberts, A. Cass, D. L. Paterson, H. Foo, M. Paul, S. D. Guy, A. R.

Tramontana, G. B. Walls, S. McBride, N. Bak, N. Ghosh, B. A. Rogers, A. P. Ralph,

J. Davies, P. E. Ferguson, R. Dotel, G. L. McKew, T. J. Gray, N. E. Holmes, S.

Smith, M. S. Warner, S. Kalimuddin, B. E. Young, N. Runnegar, D. N. Andresen, N.

A. Anagnostou, S. A. Johnson, M. D. Chatfield, A. C. Cheng, V. G. Fowler, Jr, B. P.

Howden, N. Meagher, D. J. Price, S. J. van Hal, M. V. N. O'Sullivan, J. S. Davis and

f. t. A. S. f. I. D. C. R. Network (2020). "Effect of Vancomycin or Daptomycin With vs

Without an Antistaphylococcal β-Lactam on Mortality, Bacteremia, Relapse, or

Treatment Failure in Patients With MRSA Bacteremia: A Randomized Clinical Trial."

JAMA **323**(6): 527-537.

Turnidge, J. D., D. Kotsanas, W. Munckhof, S. Roberts, C. M. Bennett, G. R. Nimmo,

G. W. Coombs, R. J. Murray, B. Howden, P. D. Johnson, K. Dowling and S. Australia

New Zealand Cooperative on Outcomes in Staphylococcal (2009). "Staphylococcus

aureus bacteraemia: a major cause of mortality in Australia and New Zealand." Med

J Aust **191**(7): 368-373.

van Hal, S. J., S. O. Jensen, V. L. Vaska, B. A. Espedido, D. L. Paterson and I. B.

Gosbell (2012). "Predictors of mortality in Staphylococcus aureus Bacteremia." Clin

Microbiol Rev **25**(2): 362-386.

van Hal, S. J., M. Jones, I. B. Gosbell and D. L. Paterson (2011). "Vancomycin

Heteroresistance Is Associated with Reduced Mortality in ST239 Methicillin-

Resistant Staphylococcus aureus Blood Stream Infections." PLOS ONE **6**(6):

e21217.

Whaley, S. G., C. D. Radka, C. Subramanian, M. W. Frank and C. O. Rock (2021).

"Malonyl-acyl carrier protein decarboxylase activity promotes fatty acid and cell

envelope biosynthesis in Proteobacteria." The Journal of biological chemistry **297**(6):

101434-101434.

Young, B. C., C.-H. Wu, J. Charlesworth, S. Earle, J. R. Price, N. C. Gordon, K.

Cole, L. Dunn, E. Liu, S. Oakley, H. Godwin, R. Fung, R. Miller, K. Knox, A.

Votintseva, T. P. Quan, R. Tilley, M. Scarborough, D. W. Crook, T. E. Peto, A. S.

Walker, M. J. Llewelyn and D. J. Wilson (2021). "Antimicrobial resistance

determinants are associated with <em>Staphylococcus aureus</em> bacteraemia

and adaptation to the hospital environment: a bacterial genome-wide association

study." medRxiv: 2021.2001.2013.21249734.

Young, B. C., C.-H. Wu, J. Charlesworth, S. Earle, J. R. Price, N. C. Gordon, K.

Cole, L. Dunn, E. Liu, S. Oakley, H. Godwin, R. Fung, R. Miller, K. Knox, A.

Votintseva, T. P. Quan, R. Tilley, M. Scarborough, D. W. Crook, T. E. Peto, A. S.

Walker, M. J. Llewelyn and D. J. Wilson (2021). "Antimicrobial resistance

determinants are associated with Staphylococcus aureus bacteraemia and

adaptation to the healthcare environment: a bacterial genome-wide association

study." Microbial Genomics **7**(11).

Young, B. C., C.-H. Wu, N. C. Gordon, K. Cole, J. R. Price, E. Liu, A. E. Sheppard,

S. Perera, J. Charlesworth, T. Golubchik, Z. Iqbal, R. Bowden, R. C. Massey, J.

Paul, D. W. Crook, T. E. Peto, A. S. Walker, M. J. Llewelyn, D. H. Wyllie and D. J.

Wilson (2017). "Severe infections emerge from commensal bacteria by adaptive

evolution." eLife **6**: e30637.

Zhou, X. and M. Stephens (2012). "Genome-wide efficient mixed-model analysis for

association studies." Nature Genetics **44**(7): 821-824.