Epidemiological investigation of the COVID-19 outbreak in Vellore district in South

India using Geographic Information Surveillance (GIS)

Malathi Murugesan[1], Padmanaban Venkatesan[2], Senthil Kumar[3], Premkumar Thangavelu[4],

Winsley Rose[5], Jacob John[3], Marx Castro[6], Manivannan T[6], Venkata Raghava Mohan[3*]and

Priscilla Rupali[4*]

[1]Department of Clinical Microbiology & Hospital Infection Control Committee, Christian
Medical College, Vellore, Tamil Nadu, India
10
11   [2]Department of Biochemistry, Christian Medical College, Vellore, Tamil Nadu, India
12
13   [3]Department of Community Health, Christian Medical College, Vellore, Tamil Nadu, India
14
15   [4] Department of Infectious diseases, Christian Medical College, Vellore, Tamil Nadu, India
16
17   [5]Department of Paediatrics, Christian Medical College, Vellore, Tamil Nadu, India
18
19   [6]Deputy Director of Health Services, Vellore, Tamil Nadu, India
20

*Shared corresponding authors

Corresponding author 1

Dr. Venkata Raghava Mohan
Professor, Department of Community Health
Christian Medical College, Vellore
Tamil Nadu, India – 632 004
28

29   **Corresponding author 2**

30   Dr. Priscilla Rupali
31   Professor, Department of Infectious Diseases
32   Christian Medical College, Vellore
33   Tamil Nadu, India – 632 004
34

Abstract

Objectives: Geographical Information Surveillance (GIS) is an advanced digital technology tool that maps location-based data and helps in epidemiological modeling. We applied GIS to analyze patterns of spread and hotspots of COVID-19 cases in Vellore district in South India.

Methods: Laboratory-confirmed COVID-19 cases from the Vellore district and neighboring taluks from March 2020 to June 2021 were geo-coded and spatial maps were generated. Time trends exploring urban-rural burden with an age-sex distribution of cases and other variables were correlated with outcomes.

Results: A total of 45,401 cases of COVID-19 were detected with 20730 cases during the first wave and 24671 cases during the second wave. The overall incidence rates of COVID-19 were 462.8 and 588.6 per 100,000 populations during the first and second waves respectively. The pattern of spread revealed epicenters in densely populated urban areas with radial spread sparing rural areas in the first wave. The case fatality rate was 1.89% and 1.6% during the first and second waves that increased with advancing age.

Conclusions: Modern surveillance systems like GIS can accurately predict the trends and pattern of spread during future pandemics. A real-time mapping can help design risk mitigation strategies thereby preventing the spread to rural areas.

Keywords: COVID-19; GIS; Spatial analysis; mapping

Background

Various epidemiological mapping systems have been employed in the past to track the spatial and temporal patterns of infectious diseases like cholera, influenza, and even plague (Sarfo and Karuppannan, 2020). During the COVID-19 pandemic, integrated and innovative surveillance methods using new technologies with accurate and speed reporting of the disease have been crucial due to the dynamic and evolving nature of the infection (Ibrahim, 2020). Geospatial technology helps alleviate this by using its different methods namely geographical information systems (GIS), global positioning systems (GPS), and other satellite mediated technology systems. GIS is a technology that is used for geographic data capture based on location which helps in mapping and epidemiological modelling using computer-assisted programs (Musa et al., 2013). In 1854, John Snow has plotted the location of Cholera patients in a hand-drawn map which linked the cases to the source (broad street pump) confirming the fact that cholera was likely waterborne (Gilbert, 1958). Over the years, there has been tremendous advancement in GIS which have provided accurate information in a real-time manner.

Geographic information surveillance (GIS) system is an advanced tool for infectious diseases and could play a vital role in analyzing spatial and temporal patterns of spread of infectious diseases. GIS mainly helps identify clusters thereby linking the social and geographical factors of disease (Musa et al., 2013). The information generated by mapping will help understand risk factors, predict the susceptible population that is going to get infected in near future, and can estimate the incidence rates of the disease. In addition, patterns of spread with the identification of high-risk populations can help formulate risk mitigation strategies by creating new healthcare infrastructure and

improving existing facilities. Since there were many uncertainties regarding transmission

in the initial stages of the COVID-19 pandemic (World Health Organization, 2020) with

the possibility of transmission from pre-asymptomatic and asymptomatic individuals (Zou

et al., 2020) we felt that it would be prudent to track the epidemiology and patterns of

spread in Vellore district. In this study, we present a spatial mapping of COVID-19

infected cases in Vellore district in Tamil Nadu, South India, and adjacent taluks to

identify epidemic clusters or hotspots in the community, track the pattern of spread and

study the demographic characteristics of study population.

Methodology

Study area: In this observational cohort study, mapping the burden of disease and further

analyses have been performed using the earlier undivided Vellore district region. Study

regions depicting different taluks, rural administrative areas, urban town panchayats, and

city corporation limits are shown in Supplementary figure 1. The undivided Vellore district

(12°55′13″N 79°08′00″E) spanned over 392.62 sq. km and had a population of 906,745 as

per the census 2011 data by the Government of India (Vellore City Population Census

2011-2022 | Tamil Nadu, n.d.). In the year 2020, the Vellore district was divided into three

districts namely Vellore, Tirupattur, and Ranipet.

Study population: Documented laboratory-confirmed COVID-19 patients from the

Vellore district for the period 28 March 2020 to 30 June 2021 were included in the study

analyses.

Data collection: The data used for the analysis in this study was obtained from the Vellore district surveillance portal, after obtaining necessary permission from the district authorities. De identified data of all the COVID-19 laboratory-confirmed positive patients were collected from the district portal with patient address, unique epidemiological identification number, symptoms, comorbidities, complications during hospitalization and outcome.

Spatial mapping of COVID-19 cases: The documented addresses from the district portal were geocoded to the closest neighborhoods using Google Earth Pro (Earth Versions, n.d.) by trained GIS technicians at Christian Medical College (CMC), Vellore. Spatial data on geocoordinates was linked to the available attribute data of COVID-19 cases and personal identifiers were removed from the database used for final analyses. Village and taluk level vector layers of the study region including population-level data sourced from Survey of India, which is the National Survey and Mapping Organization under the Government of India were used as base layers for spatial mapping. Using the local neighborhood level spatial information on cases, all cases were mapped onto the base layer using the software ArcMap v 10.8.1 (ArcGIS Desktop quick start guide—ArcMap | Documentation, n.d.).

Statistical analysis: Descriptive analysis was performed to study the characteristics of the study population for age, gender, and locality (urban or rural), presentation of COVID-19 symptoms, complications and survival rate was calculated as a percentage among the study population. We analyzed the age-sex distribution of COVID-19 cases along with positivity rates by age groups, gender, and area of residence (rural/urban) with 95% CI. In addition, time trends exploring urban-rural burden and other variables were correlated with outcome

data and compared between the first wave (28 March 2020 to 31 March 2021) and the

second wave (01 April 2021 to 30 June 2021).

Spatial analysis: Spatial maps depicting the distribution of COVID-19 case counts by

villages and urban wards in the Vellore district were generated using the geocoded

addresses and layered as points on the base map to illustrate the distribution of all COVID-

19 cases from March 2020 to June 2021. A total of 42,921 cases of COVID-19 were

available for spatial mapping based on the completeness of documented addresses in the

study area. There was a total of 938 villages, urban localities, and reserved forest areas

available as individual polygons for the Vellore district. The total number of cases falling

under each of these individual area polygons was estimated by spatially joining the case

layer with the polygon layer and sub-block level maps depicting caseloads for the overall

period and waves 1 and 2 were generated using ArcMap. Densities of COVID-19 cases

around each documented case were estimated using the point density tool within the Spatial

Analyst toolbox in ArcMap and quarterly heat maps were generated to demonstrate the

spread of the disease for the period March 2020 to June 2021. The earlier undivided

Vellore district was subdivided into eight taluks which included Arakkonam, Wallajah,

Arcot, Katpadi, Vellore, Gudiyattam, Vaniyambadi and Tirupattur spanning from west to

east (Revenue Divisions | Vellore District, Government of Tamil Nadu | India, n.d.). Taluk

level population counts were obtained from the recently published figures on the taluk

websites and were used as denominators to estimate and map incidence rates of COVID-19

disease during the two waves.

Results

A total of 45,401 cases of COVID-19 were detected between 28 March 2020 to 31 June

2021 with 20730 cases during the first wave (28 March 2020 to 31 March 2021) and 24671

cases during the second wave (1 April 2021 to 30 June 2021). Data was available in 42921

cases for spatial mapping based on the completeness of documented addresses during both

waves of which 19,021 (44.3%) and 23,900 (55.7%) had occurred during the first and

second waves respectively. A higher number of cases resided in Vellore, Katpadi and

Gudiyattam taluks (Supplementary Figure 2). We mapped weekly and monthly trends in

the epidemiological spread of infection and were able to identify geographical clusters,

hotspots and spatial-temporal trends using GIS.

Of the 938 smaller geographical areas within the study region, 188 were classified as

reserved forest areas and the rest comprised villages, urban localities, town panchayat and a

city corporation located in Vellore taluk. Throughout both waves, COVID-19 cases were

documented from 636 of the inhabited villages and urban areas in the Vellore district. The

mean (SD) number of cases in these smaller units was 67 (278) with the minimum being

one case documented in one of the villages and the maximum documented caseload was

3323 in Konavattam urban ward in Vellore city. Ten urban areas had recorded over 1000

cases each; of which one, four, and five of them were Gudiyattam, Katpadi and Vellore

taluks in the region. Village and urban ward-wise caseloads during both the waves and

overall study period are presented in Figure 1 and a similar trend was observed with the

same urban and peri-urban neighborhoods experiencing higher burden during both the

waves and with relative sparing of rural areas.

Heatmaps generated estimating the point density around geolocations of cases in the initial

stages of the pandemic from April to June 2020 revealed that the epicenters were in the

middle of Vellore city, a densely populated city corporation and the town panchayat area of

Gudiyattam taluk. Between July and September 2020 when the first wave was at its peak, it

was observed that more cases occurred in the urban corporation and town panchayat areas

and spread to neighboring smaller urban areas sparing the rural areas in Vellore,

Gudiyattam and Katpadi taluks. Even though the other taluks (Arakkonam, Wallajah, Arcot

and Tirupattur) had documented COVID-19 cases significant spatial hotspots were not

detected. As the first wave started to ebb, between December 2020 and March 2021, very

few neighborhoods, especially in the Vellore corporation area still had cases but at a lower

density. During the second wave predominantly due to the Delta variant, which was

officially documented in late March 2021, the region experienced many more COVID-19

cases as compared to the first wave. Although the pattern of spread was similar with the

epicenters in urban geographies spreading radially and sparing rural areas, heat maps

revealed much higher densities at these epicenters when compared to the first wave and

with more peri-urban and rural area involvement (Figure 2).

The overall incidence rates of COVID-19 across the study region were 462.8 per 100,000

and 588.6 per 100,000 populations during the first and second waves respectively. Of the

eight taluks, five of them (Tirupattur, Arakkonam, Arcot, Wallajah and Vaniyambadi)

consistently had incidence rates less than 100 per 100,000 population during both the

waves. During the first wave, the burden of disease in Vellore and Katpadi taluks were

1427.8 and 1344.4 per 100,000 respectively while during the second wave, the Katpadi

region experienced a higher rate (2098.2 per 100,000) as compared to the Vellore region

(1731.4 per 100,000 population). Incidence rates across the eight taluks during both the

waves and overall study duration are presented in Figure 3.

The age distribution among the COVID-19 cases during both the waves is given in table 1.

Most of them belonged to the age group of 21-60 years. Among the study population, 3.1%

in the first wave and 1.99% in the second wave were children less than 10 years. Elderly

(>60 years) were 16% in the first wave and 17.3% in the second wave. Overall, COVID-19

incidence was higher in males (first wave 58.71%; second wave 56.86%) when compared

to females (first wave 41.29%; second wave 42.90%). Figure 4 shows the overall time

trends of the outbreak in this location. The first case documented to be COVID positive

was on 28 March 2020. The outcome details were documented for 20730 cases in the first

wave and 21672 cases in the second wave. Case fatality rate was 1.89% during the first

wave and 1.6% during the second wave and steadily increased with advancing age, i.e.,

7.38% were aged more than 60 years in the first wave and 5.02% in the second wave (Table

2). Case fatality rates were higher in men; first wave 2.40%, and 1.76% in the second wave

as compared to women first wave 1.16%, second wave 1.38%. When correlating

comorbidities with mortality, data from the first wave revealed that the fatality rates were

highest among those who had >2 comorbidities (9.52%).

The data on risk factors, comorbidities and clinical symptoms, and outcomes were available

only for 17748 out of 19092 cases during the first wave of the pandemic but were not

available for the second wave as recording this was not mandatory during the second wave.

On analyzing risk factors for acquiring COVID-19 infection, 4% of them had a recent

history of travel outside India and 2.1% reported a history of travel within India but 62.9%

had a history of contact with a COVID-19 infected individual. The comorbidity profile data

available for 17748 cases showed that 14.4% of the patients had at least one co-morbidity

but 7% of the patients had more than one co-morbidity. Diabetes, hypertension, and asthma

were the most seen co-morbidities in these patients (Supplementary Table 1). Among the

61.93% of patients who were symptomatic, fever (39.7%), cough (24.8%) and sore throat

(15.9%) were the most common symptoms. Breathlessness was reported in 9% of all cases

(Supplementary Table 2). 99 out of 17748 (0.6%) were pregnant among the study

population during the first wave of the pandemic.

Discussion

In our spatial-temporal analysis, we noted that the first wave of the pandemic started with a

returned traveler from the UK in March after which a peak was noted in travelers who had

attended a religious meet in another part of the country (Tablighi Jamaat, 2020).

Subsequently there was a steady rise in cases over five months (June to October 2020) with

continuing sporadic cases reported thereafter till March 2021. Contrary to the first wave,

the epidemic noted a sharp spike in April with a rapid decline over two months (April to

June 2021) during the second wave of the pandemic. This was presumed to be mainly due

to the delta variant B.1.617.2 which had a high transmissibility and immune evasion as

compared to the original strain (Hwang et al., 2022; Mlcochova et al., 2021). A similar

pattern was observed, 100 years ago (1917-1918) when there were a successive occurrence

of three waves caused by Spanish flu in one year (Taubenberger and Morens, 2006); a

smaller spring wave, a larger fall wave and again a smaller winter wave. The R_0 of Spanish

flu in 1918 was 1.5 – 3.8 (Chowell et al., 2006) and the subsequent H1N1 influenza

pandemic in 2009 was 1.33 (White et al., 2013). However, R_0 for COVID-19 based on the

incidence data of European countries was in fact 2.2 (95% CI 1.9-2.6) for the ancestral

strain (Locatelli et al., 2021) , however China reported a higher R_0 of 3.32 (95% CI 2.81 to

3.82) (Alimohamadi et al., 2020). Subsequent waves caused by the Delta wave revealed a

R_0 of 5.08 (3.2-8) (Liu and Rocklöv, 2021) and the Omicron 9.5 (5.5-20) (Liu and Rocklöv,

2022). This suggested that R_0 of SARS-Co-2 was second only to measles in being highly

transmissible warranting strict quarantine and containment strategies for effective control.

The rapid transmission left many countries and states unprepared. Hence, the usage of

digital methods to map the cluster areas, predict hotspots would have enabled contact

tracing in real-time helping to cordon off affected households to prevent ongoing

transmission.

The geospatial mapping in our study population showed that the pattern of spread of the

cases is mainly in highly populated urban areas followed by less populated semi-urban

areas and then rural areas. In both the first and the second wave, the initial cluster of cases

is seen in the urban population. But in the second wave, the peri-urban and rural areas were

also affected. Since 60% of population in India is rural it is imperative to curb spread from

urban to rural areas. Similar trend has been shown worldwide where the metropolitan cities

and urban areas are affected earlier due to a higher population density, indoor crowding,

and high usage of crowded public transport systems (Lee et al., 2020). A study conducted

by Gangwar *et al.,* analyzed COVID-19 cases in India using geospatial technology from

statistics available in the public domain and showed similar results suggesting that 80% of

the confirmed cases during the first wave occurred in high population areas as cities

underwent gradual unlocking after a stringent lockdown phase (Gangwar and Ray, 2021).

The case fatality rate in our study population was 1.89% and 1.6% during the first wave

and second wave respectively. A study published by Shah *et al.,* found that the CFR for

SAARC countries were lower than the developed countries and the estimated crude and

adjusted CFR for India was 1.542% and 1.601% (95% CI 1.548 - 1.684) during the first

wave (Shah et al., 2021). Another study by Bogam *et al.,* estimated that the fatality rate was

47% lower in the delta variant driven second wave than in the first wave (adjusted CFR

0.43; 95% CI 0.41 – 0.45) (Bogam et al., 2021) despite the large numbers. This suggested

that although, the transmission was higher during the second wave, the CFR is probably

lower due to a predominantly younger population being affected, pre-existing immunity

obtained from either previous COVID-19 infection or other coronaviruses in circulation,

COVID-19 vaccination or easy access to health care services as all hospitals were directed

to accept COVID-19 infected patients under the epidemic act of 1897 (The epidemic

disease act, 2020, n.d.). Even though the country managed well during the first wave of the

pandemic, the rapid hit of the second wave decimated the health care system (National

Health Profile, 2019).

The COVID-19 pandemic has also caused an unprecedented global economic crisis due to

lockdown, travel restrictions, border closures and loss of employment. The UNICEF has

estimated that 42 to 66 million children could fall into extreme poverty because of the

COVID-19 crisis in addition to the already existing poor population of 386 million children

(2019) (UNSDG | Policy Brief, n.d.). Thus, an accurate mapping of the diseased population

which could predict the spread of future respiratory pandemics would help adapt future

strategies of containment and management including testing, physical distancing, contact

tracing, training, and public awareness based on the resources and capacity of each health

setting. Educational awareness and targeted training of public and health care sectors can

be done in a phased manner in clusters/hot spots followed by other areas. The less affected

areas with susceptible populations can be targeted for vaccination by doorstep delivery of

vaccines through mobile camps. Even though the Epidemic diseases act 1897 was helpful

in the containment of the disease spread nationwide during the pandemic, digitalized

methods and real-time mapping of the clusters were not employed and thus interventions

which could have helped during the second wave could not be implemented effectively.

Previous respiratory pandemics have led to enormous mortality and morbidity around the

world with measles epidemic in 1960s infecting 3-4 million people in the United States

leading to 500 deaths (Scary measles history has been forgotten by many - The Washington

Post, n.d.) and plague in India in 1900s causing over 2-3 million deaths (Plague, n.d.). This

calls for appropriate measures which can accurately predict mortality, morbidity, and the

course of the outbreak during future respiratory pandemics and help with real-time

mapping of cases [latitude and longitude coding] on an individual case basis thereby

efficiently implementing risk mitigation strategies.

As new diseases emerge and old ones re-emerge, the routinely used surveillance methods in

public health include active, passive, categorical, integrated, syndromic, case-based,

sentinel and serological surveillance etc. (Ibrahim, 2020; Nsubuga et al., 2006) are helpful.

However, due to rapid spread of respiratory pandemics with a high R_0 in the past, it makes

sense to employ digital surveillance technology as it can save time, manpower and

resources. During the pandemic, many organizations including WHO created various

geospatial dashboards like the WHO Coronavirus (COVID-19) dashboard (WHO

Coronavirus (COVID-19) Dashboard, n.d.), John Hopkins University COVID-19

dashboard (Coronavirus COVID-19 (2019-nCoV), n.d.), COVID-19 India (Coronavirus in

India, n.d.) and many others regionally. This was extremely useful for epidemiologists and

local health policymakers to implement isolation and quarantine measures based on the

geographic spread. Mobile-based digital apps like the COVID symptom study tracker (ZOE

COVID Study, n.d.) and Aarogya Setu (Aarogya Setu, n.d.) helped in collecting

demographic characteristics of the diseased individuals but lacked real-time mapping. In

India, due to the usage of the same mobile numbers in a family, network accessibility and

loss of privacy due to location access limited the usage of mobile-based apps. Based on the

lessons learned during this pandemic, futuristic plans, and budget allocation to more

advanced information technology systems using satellite and geomapping would help in

outbreak management.

Limitations

As the analysis is based on the data available in the public domain and conducted

retrospectively, geographic location of the study population (5.46%) and outcome details

(6.60%) were not available for the entire study population. The clinical features and

comorbidity profile was able to be retrieved only for 85.62% of the first wave population.

This limited our comparison of the risk factors and clinical features during the first wave

and second wave. The data captured is from the reported cases from the district portal and

not from the state wise data where there may be some patients from this district who got

admitted in other places. But due to lockdown and other restrictions, we believe that almost

all of them were admitted within the district.

Conclusion

Respiratory pandemics with a high R_0 seem to start in the urban areas and then spread to

rural areas. Therefore, GIS surveillance as a contact tracing digital tool would help in rapid

implementation of risk mitigation and containment strategies, thus preventing spread to

rural areas where access to appropriate health care services is limited. The nation must

build a catastrophe resilient public health care system with well-equipped facilities under

the existing National Rural Health Mission (NRHM) to manage pandemics in urban and

rural areas.

Disclosures

None

Conflict of Interest

None to declare

Funding Source

This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Ethical Approval Statement

This study conducted jointly by Christian Medical College, Vellore, and the district health

authorities which is approved by the institutional review board, Christian Medical College,

Vellore located in South India (IRB no:13261 dated 26.08.2020).

Author contributions

Malathi Murugesan - Conceptualization, Methodology, Resources, Validation, Formal

analysis, Data Curation, Writing- Original draft & Editing

Padmanaban Venkatesan - Methodology, Resources, Formal analysis, Data Curation,

Writing- Original draft & Editing

Senthil Kumar - Methodology, Resources, Validation, Data Curation

Premkumar Thangavelu - Methodology, Resources, Validation, Data Curation

Winsley Rose - Validation, Formal analysis, Data Curation, Writing - Review & Editing

Jacob John - Validation, Formal analysis, Data Curation, Writing - Review & Editing

Venkata Raghava Mohan - Conceptualization, Methodology, Resources, Validation,

Formal analysis, Data Curation, Writing - Review & Editing

Priscilla Rupali - Conceptualization, Methodology, Resources, Validation, Formal analysis,

Data Curation, Writing - Review & Editing

Acknowledgement

We wish to thank Mr. Varunkumar, Department of Community Health who assisted in

geocoding and mapping.

391   **References**

393   Aarogya Setu. n.d. https://www.aarogyasetu.gov.in/ (accessed March 29, 2022).

394
395   Alimohamadi Y, Taghdir M, Sepandi M. Estimate of the Basic Reproduction Number for

396   COVID-19: A Systematic Review and Meta-analysis. J Prev Med Public Health

397   2020;53:151–7. https://doi.org/10.3961/jpmph.20.076.

398
399   ArcGIS Desktop quick start guide—ArcMap | Documentation. n.d.

400   https://desktop.arcgis.com/en/arcmap/latest/get-started/setup/arcgis-desktop-quick-start-

401   guide.htm (accessed March 28, 2022).

402
403   Bogam P, Joshi A, Nagarkar S, Jain D, Gupte N, Shashidhara LS, et al. Burden of COVID-

404   19 and Case Fatality Rate in Pune India: An Analysis of First and Second Wave of the

405   Pandemic 2021:2021.06.21.21259225. https://doi.org/10.1101/2021.06.21.21259225.

406
Chowell G, Ammon CE, Hengartner NW, Hyman JM. Estimation of the reproductive

number of the Spanish flu epidemic in Geneva, Switzerland. Vaccine 2006;24:6747–50.

https://doi.org/10.1016/j.vaccine.2006.05.055.

Coronavirus COVID-19 (2019-nCoV).

https://gisanddata.maps.arcgis.com/apps/dashboards/bda7594740fd40299423467b48e9ecf6

(accessed March 29, 2022).

Coronavirus in India: Latest Map and Case Count. n.d. https://www.covid19india.org

(accessed March 29, 2022).

Earth Versions. Google Earth n.d. https://www.google.com/intl/en_in/earth/versions/

(accessed March 28, 2022).

Gangwar HS, Ray PKC. Geographic information system-based analysis of COVID-19

cases in India during pre-lockdown, lockdown, and unlock phases. International Journal of

Infectious Diseases 2021;105:424. https://doi.org/10.1016/j.ijid.2021.02.070.

Gilbert EW. Pioneer Maps of Health and Disease in England. The Geographical Journal

1958;124:172–83. https://doi.org/10.2307/1790244.

Hwang H, Lim J-S, Song S-A, Achangwa C, Sim W, Kim G, et al. Transmission Dynamics

of the Delta Variant of SARS-CoV-2 Infections in South Korea. J Infect Dis

2022;225:793–9. https://doi.org/10.1093/infdis/jiab586.

Ibrahim NK. Epidemiologic surveillance for controlling Covid-19 pandemic: types,

challenges and implications. Journal of Infection and Public Health 2020;13:1630–8.

https://doi.org/10.1016/j.jiph.2020.07.019.

Lee VJ, Ho M, Kai CW, Aguilera X, Heymann D, Wilder-Smith A. Epidemic preparedness

in urban settings: new challenges and opportunities. Lancet Infect Dis 2020;20:527–9.

https://doi.org/10.1016/S1473-3099(20)30249-8.

Liu Y, Rocklöv J. The effective reproductive number of the Omicron variant of SARS-

CoV-2 is several times relative to Delta. Journal of Travel Medicine 2022:taac037.

https://doi.org/10.1093/jtm/taac037.

Liu Y, Rocklöv J. The reproductive number of the Delta variant of SARS-CoV-2 is far

higher compared to the ancestral SARS-CoV-2 virus. Journal of Travel Medicine

2021;28:taab124. https://doi.org/10.1093/jtm/taab124.

Locatelli I, Trächsel B, Rousson V. Estimating the basic reproduction number for COVID-

19 in Western Europe. PLOS ONE 2021;16:e0248731.

https://doi.org/10.1371/journal.pone.0248731.

Mlcochova P, Kemp SA, Dhar MS, Papa G, Meng B, Ferreira IATM, et al. SARS-CoV-2

B.1.617.2 Delta variant replication and immune evasion. Nature 2021;599:114–9.

https://doi.org/10.1038/s41586-021-03944-y.

Musa GJ, Chiang P-H, Sylk T, Bavley R, Keating W, Lakew B, et al. Use of GIS Mapping

as a Public Health Tool—From Cholera to Cancer. Health Serv Insights 2013;6:111–6.

https://doi.org/10.4137/HSI.S10471.

National Health Profile, 2019, Government of India,

http://www.cbhidghs.nic.in/WriteReadData/l892s/8603321691572511495.pdf (accessed

April 5,2022)

Nsubuga P, White ME, Thacker SB, Anderson MA, Blount SB, Broome CV, et al. Public

Health Surveillance: A Tool for Targeting and Monitoring Interventions. In: Jamison DT,

Breman JG, Measham AR, Alleyne G, Claeson M, Evans DB, et al., editors. Disease

Control Priorities in Developing Countries. 2nd ed., Washington (DC): World Bank; 2006.

Plague: deaths in India by region 1896-1904. Statista n.d.

https://www.statista.com/statistics/1115019/plague-deaths-india-circa-1900/ (accessed

April 9, 2022).

471
472   Revenue Divisions | Vellore District, Government of Tamil Nadu | India. n.d.

473   https://vellore.nic.in/revenue-divisions/ (accessed March 28, 2022).

474   Sarfo AK, Karuppannan S. Application of Geospatial Technologies in the COVID-19 Fight

475   of Ghana. Trans Indian Natl Acad Eng 2020;5:193–204. https://doi.org/10.1007/s41403-

476   020-00145-3.

477
478   Scary measles history has been forgotten by many - The Washington Post. n.d.

479   https://www.washingtonpost.com/national/health-science/measles-why-its-so-deadly-and-

480   why-vaccination-is-so-vital/2019/02/15/a452e5c4-2fd0-11e9-8ad3-

481   9a5b113ecd3c_story.html (accessed April 9, 2022).

482
483   Shah MRT, Ahammed T, Anjum A, Chowdhury AA, Suchana AJ. Finding the real

484   COVID-19 case-fatality rates for SAARC countries. Biosafety and Health 2021;3:164–71.

485   https://doi.org/10.1016/j.bsheal.2021.03.002.

486
487   Tablighi Jamaat: The group blamed for new Covid-19 outbreak in India. BBC News 2020

488   (accessed April 5, 2022).

489
490   Taubenberger JK, Morens DM. 1918 Influenza: the Mother of All Pandemics. Emerg Infect

491   Dis 2006;12:15–22. https://doi.org/10.3201/eid1201.050979.

492
The application of geographic information system (GIS) in the field of public health. n.d.

https://ieeexplore.ieee.org/document/5603111/ (accessed March 29, 2022).

The epidemic diseases act, 2020, Government of India,

https://egazette.nic.in/WriteReadData/2020/222125.pdf (accessed April 9,2022)

UNSDG | Policy Brief: The Impact of COVID-19 on children. n.d.

https://unsdg.un.org/resources/policy-brief-impact-covid-19-children,

https://unsdg.un.org/resources/policy-brief-impact-covid-19-children (accessed April 5,

2022).

Vellore City Population Census 2011-2022 | Tamil Nadu. n.d.

https://www.census2011.co.in/census/city/472-vellore.html (accessed March 28, 2022).

White LF, Archer B, Pagano M. Estimating the reproductive number in the presence of

spatial heterogeneity of transmission patterns. Int J Health Geogr 2013;12:35.

https://doi.org/10.1186/1476-072X-12-35.

507
508   WHO Coronavirus (COVID-19) Dashboard. n.d. https://covid19.who.int (accessed March

509   29, 2022).

510
511   World Health Organization. Modes of transmission of virus causing COVID-19:

512   implications for IPC precaution recommendations: scientific brief, 29 March 2020. World

513   Health Organization; 2020.

514
515   ZOE COVID Study. n.d. https://covid.joinzoe.com (accessed March 29, 2022).

516
517   Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in

518   Upper Respiratory Specimens of Infected Patients. N Engl J Med 2020;382:1177–9.

519   https://doi.org/10.1056/NEJMc2001737.

520
521

Legends for figures

Figure 1: Distribution of COVID-19 cases across different geographical subunits in the

study area

Figure 2: Heatmaps depicting spread of COVID-19 cases during the first and second waves

in the study area

Figure 3: Subdistrict level incidence of COVID-19 during the first and second waves

Figure 4: Epidemic curve of the COVID-19 pandemic during the first and the second waves

Supplementary figure 1: Map of undivided Vellore district showing rural and urban

settlements

Supplementary figure 2: Spatial mapping of COVID-19 cases in the study region for the

period March 2020 to June 2021.

Legends for Tables

Table 1: Age and gender of COVID-19 patients during the first and second wave

Table 2: Outcome by age during first wave (N=20730) and second wave (N=21672) of the

COVID-19 pandemic

Supplementary table 1: Co-morbidities among the COVID-19 affected patients in the first

wave (N = 17748)

Supplementary table 2: Clinical features of COVID-19 infection in the first wave (N =

17748)

Figure 1: Distribution of COVID-19 cases across different geographical subunits in

the study area

Figure 2: Heatmaps depicting spread of COVID-19 cases during the first and second

waves in the study area

A: April - June 2020

B: July - September 2020

C: October - December 2020

D: January - March 2021

E: April - June 2021

F: April 2020 to June 2021

Point density maps depicting quarterly distribution of Covid -19 cases

Figure 3: Subdistrict level incidence of COVID-19 during the first and second waves

Figure 4: Epidemic curve of the COVID-19 pandemic during the first and the second

waves

Table 1: Age and gender of COVID-19 patients during the first and second wave

	First wave			Second wave			
Age (years)	**Female n (%)**	**Male n (%)**	**Total n (%)**	**Female n (%)**	**Male n (%)**	**No data n %**	**Total n (%)**
0-10	295 (3.4)	349 (2.9)	644 (3.1)	222 (2.10)	270 (1.92)	0	492 (1.99)
11-20	624 (7.3)	806 (6.6)	1430 (6.9)	629 (5.94)	703 (5.01)	3 (5.00)	1335 (5.41)
21-30	1849 (21.5)	2100 (17.3)	3949 (19)	2114 (19.97)	2397 (17.09)	6 (10.00)	4517 (18.31)
31-40	1660 (19.3)	2297 (18.9)	3957 (19.1)	2128 (20.10)	3115 (22.21)	18 (30.00)	5261 (21.32)
41-50	1591 (18.5)	2271(18.7)	3862 (18.6)	2017 (19.06)	2543 (18.13)	13 (21.67)	4573 (18.54)
51-60	1390 (16.2)	2207 (18.2)	3597 (17.4)	1861 (17.58)	2353 (16.78)	10 (16.67)	4224 (17.12)
61-70	845 (9.8)	1406 (11.6)	2251 (10.9)	1092 (10.32)	1721 (12.27)	8 (13.33)	2821 (11.43)
71-80	276 (3.2)	570 (4.7)	846 (4.1)	421 (3.98)	718 (5.12)	2 (3.33)	1141 (4.62)
81-90	50 (0.6)	127 (1)	177 (0.9)	94 (0.89)	192 (1.37)	0	286 (1.16)
91-100	6 (0.1)	11 (0.1)	17 (0.1)	7 (0.07)	14 (0.10)	0	21 (0.09)
Total	8586	12144	20730	10585	14026	60	24671

Table 2: Outcome by age during first wave (N=20730) and second wave (N=21672) of

COVID-19 pandemic

	First wave		Second wave			
Age (years)	Death	Discharged	No data	Death	Discharged	In Admission
0-14	1 (0.10)	1010 (99.90)	13 (1.82)	0 (0.00)	545 (76.33)	156 (21.85)
15-44	37 (0.36)	10266 (99.64)	161 (1.47)	38 (0.35)	8762 (79.87)	2010 (18.32)
45-59	111 (1.81)	6014 (98.19)	55 (0.89)	118 (1.90)	4773 (76.92)	1259 (20.29)
60 and above	243 (7.38)	3048 (92.62)	12 (0.32)	190 (5.02)	2810 (74.30)	770 (20.36)

Supplementary tables:

Supplementary table 1: Co-morbidities among the COVID-19 affected patients in the

first wave (N = 17748)

Comorbidities	Count (n)	Percentage (%)
Diabetes	1427	8.0
Hypertension	1258	7.1
Asthma	848	4.8
Liver Disease	330	1.9
Malignancy	119	0.7
Pregnancy	99	0.6
Chronic Renal Disease	92	0.5
Heart disease	90	0.5
Psychiatric disorders	74	0.4
Hypothyroidism	72	0.4
COPD	57	0.3
Post-partum (<6 weeks)	12	0.1
Chronic neurological or neuromuscular disease	11	0.1
AKI	10	0.1
Immunocompromised condition including HIV	9	0.1
Immunocompromised condition including TB	1	0.0

	15198	86.7
None		

640 **Supplementary table 2: Clinical features of COVID-19 infection in the first wave (N =**

641 **17748)**

Clinical features	Count (n)	Percentage (%)
Asymptomatic	6757	38.07
Symptomatic	10991	61.93
Fever	7042	39.7
Cough	4396	24.8
Sore throat	2828	15.9
Breathlessness	1615	9.1
General Weakness	1515	8.5
Diarrhoea	1468	8.3
Headache	637	3.6
Nausea/Vomiting	213	1.2
Irritability/Confusion	50	0.3
Others	3088	17.3

Supplementary figure 1: Map of undivided Vellore district showing rural and urban

settlements

Supplementary figure 2: Spatial mapping of COVID-19 cases in the study region for

the period March 2020 to June 2021.