The impact of sexual violence in mid-adolescence on girls' mental health: evidence from a longitudinal population-based study

Francesca Bentivegna, MSc [1] & Praveetha Patalay, PhD[2,3]

[1] UCL Institute of Education, Department of Psychology and Human Development, University College London, London, UK

[2] Centre for Longitudinal Studies, Social Research Institute, University College London, London, UK

[3] MRC Unit for Lifelong Health and Ageing at UCL, University College London, London, UK

Correspondence: Praveetha Patalay – p.patalay@ucl.ac.uk

Funding: This study was supported by grants MR/N013867/1 from the Medical Research Council. The Millennium Cohort Study is supported by the Economic and Social Research Council and a consortium of UK government departments. The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of this report.

Acknowledgements: The authors are grateful for the cooperation of the Millennium Cohort Study families who voluntarily participate in the study. They would also like to thank a large number of stakeholders from academic, policy-maker and funder communities and colleagues at the Centre for Longitudinal Studies involved in data collection and management. We are also grateful to feedback provided on an earlier draft of the manuscript by Drs. Francesca Solmi and Gemma Lewis, Division of Psychiatry, UCL.

Conflicts of Interest: None.

Data availability: The Millennium Cohort Study data is available to use by researchers via the UK Data Service (https://ukdataservice.ac.uk/).

Abstract

Background: A large gender gap appears in internalising mental health during adolescence. There is little high-quality longitudinal population-based research investigating the role of sexual violence experiences, which are disproportionately experienced by females. This study estimates the impact of sexual violence experiences in adolescent girls on mental ill-health.

Methods: Longitudinal data from girls in the UK Millennium Cohort Study are used. The impact of sexual violence experiences on psychological distress, self-harm and attempted suicide at age 17 is examined using both multivariable confounder adjusted and propensity matching approaches.

Findings: The analytic samples for multivariable and matched analyses were, respectively, 5,134 and 4,355 participants. In the final adjusted model, sexual violence was associated with greater mean psychological distress (mean difference 2.23 [1.67 – 2.79])), and higher risk of high distress (RR 1.74 [1.44 – 2.09]), self-harming (RR 2.04 [1.70 – 2.45]), and attempting suicide (RR 2.26 [1.64 – 3.11]) at age 17 years. Similar results were found for the matched analyses for all mental health outcomes: psychological distress (mean difference 1.97 [1.58 – 2.36]; RR 1.60 [1.41 – 1.81]), self-harm (RR 1.64 [1.48 – 1.82]), and attempted suicide (RR 1.81 [1.47 – 2.22]). Estimates of population attributable fractions suggested that, in a scenario with no sexual violence, we could expect 25-37% fewer incident mental health problems at this age.

Interpretation: Our findings highlight the substantial role of sexual violence experiences in the poorer mental health outcomes experienced by adolescent girls and the substantial changes needed at societal and policy levels to prevent sexual violence and its impacts.

Funding: Medical Research Council.

Keywords: depression, self-harm; suicide; gender-based violence; sexual harassment.

Introduction

There is a well-established and substantial gender gap in common mental illnesses, whereby women experience higher rates of depression, anxiety and self-harm from adolescence onwards (1–4). For instance, recent prevalence estimates in the UK indicate that, in late adolescence, 24% of females have an emotional disorder compared to 10% of males (5). There are many hypothesized explanations for this gender gap, including differential social risks, biological development, and specific vulnerability factors in interaction with stressors that are particularly salient in adolescence (6–8); and it is likely that several factors contribute to the larger burden of these mental health problems faced by women. One potential risk factor that is disproportionately experienced by females from early adolescence is sexual violence, including sexual abuse, assault, and harassment (9–11). Teenage girls experience over five times the levels of sexual assault compared to their male peers (12). Estimating the impact of sexual violence in mid-adolescence on subsequent mental health is, therefore, critical (13).

The available evidence on the relationship between sexual violence in mid-adolescence and mental health is scarce, as much attention pre-adulthood has been given to adverse childhood experiences, including sexual abuse earlier in childhood (14–17), experiences of sexual violence in the university and college environment (18,19), and intimate partner violence in older adolescent age groups (20). Several existing studies we identified that focused on mid-adolescence, an age at which many girls begin to experience increasing levels of harassment and assault after puberty, are limited in their ability to make assumptions on the population level impacts of sexual violence due to either the use of selective samples, such as those recruited from sexual assault referral centres (21,22), the examination of cross-sectional associations (23–26), or the use of retrospective recollections of both violence experienced and its perceived subsequent impact on mental health (13,23). As for the studies with any longitudinal examination, these were limited to convenience samples in some North American school settings (16,17) and do not include relevant wider and earlier life factors that might be important confounders of this relationship (27,28).

This paper aims to estimate the impact of sexual violence experienced in mid-adolescence on depressive symptoms, self-harm and attempted suicide in a prospective longitudinal cohort study. Although in many health contexts randomised control trials represent the gold standard for establishing the causal impact of an experience X on health outcomes, the ethical considerations for avoiding a design where sexual violence experiences are experimentally manipulated are clear. However, there are methods that allow the creation of a pseudo-experimental design, including propensity score matching (PSM (29,30)). In this matching approach, each individual from the treatment group is matched to an individual in the control group based on similarity on pre-specified criteria. The nature of sexual violence at this age, which is not an experience that an individual selects into, but rather is done to them, also increases the methodological scope for inferring causality using such pseudo-experimental approaches (in contrast to smoking or physical activity, for instance, which often involve self-selection into these behaviours). Due to the greatly gendered nature of these sexual violence experiences and different salience of these experiences in males (17), we focus on adolescent females. We hypothesise that adolescent girls who experienced sexual violence will have worse subsequent mental health and that effects will be robust to rich confounder adjustment and under the conditions of a pseudo-experimental design. Lastly, we estimate the population attributable fraction (PAF) to predict the reduction in adverse mental health outcomes if

sexual violence against adolescent girls was successfully limited via necessary policy and societal changes.

Methods

Design

We used a longitudinal study with rich data on early life factors and confounders to estimate the impacts of sexual violence on mental health using two main approaches: 1) multivariable regression with adjustment for a wide range of relevant confounders; 2) pseudo-experimental design with a group that experienced sexual violence between ages 14 and 17 and a matched control group. Moreover, as a further robustness check, we combined these two methods by adjusting for a wider set of confounders in the context of the pseudo-experimental design.

Participants

Data from a national birth cohort study, the Millennium Cohort Study, are used. The main analysis focuses on data from two consecutive sweeps at age 14 (2015-16; 11,726 families) and age 17 (2018-19; 10,625 families) (31) with confounders being from earlier sweeps at ages 3, 5, 7, 11 years as relevant.

In line with the aims of the study, we restricted the possible analytical sample to participants who were born female or identified as female. For the multivariable analyses, participants had to have information available on both experience of sexual violence prior to age 17 and one of the outcomes, i.e. psychological distress, self-harm and/or attempted suicide at age 17. For the pseudo-experimental design with matched sexual violence-control group, the included participants were also required to have data on all the key matching variables, and we excluded 199 (4.9%) of girls who had reported experiences of sexual violence prior to age 14 years.

Measures

Sexual violence

Sexual violence was assessed using three relevant questions from the self-reported questionnaire at age 17. Specifically, participants were asked about their experiences as victims in the previous 12 months with regards to sexual assault, unwelcomed sexual approach, and harassment via mobile phone or email. A dichotomous answer (yes/no) was given for each item. We created a binary overall sexual violence variable by combining information from these three items, and also analysed each of the three separately in additional analyses.

Mental health outcomes

Psychological distress at age 17 was assessed using the Kessler-6 (K-6 (32)), a six-item, self-administered inventory measuring distress over the previous 30 days. Items refer to both depressive and anxiety symptoms and rated on a 5-point Likert scale ('all of the time' to 'none of the time'). A total score was obtained from summing the items; higher scores indicating greater distress; scores of 13 or above indicating high distress (33).

Self-harm was assessed by asking participants whether (yes/no responses) they hurt themselves on purpose in the previous 12 months, in one or more ways, from a list of six specific situations, including cutting, burning, bruising, taking an overdose, pulling out hair, and hurting themselves in

other ways. A binary variable was created indicating self-harming in the previous 12 months if participants responded affirmatively to any of these items.

Attempted Suicide was assessed via responses (yes/no) to the question 'Have you ever hurt yourself on purpose in an attempt to end your life?'.

Confounding variables

We chose the following matching and confounding factors in line with previous literature indicating the following as relevant for either or both the main exposure of sexual violence and the mental health outcomes: ethnicity (34); pubertal status (35); parental education (36); sexual identity (Coulter et al., 2018); psychological distress at age 14 (36); self-harm at age 14 (24); body mass index (BMI (37)); family income (36); number of siblings (38,39); single parent household (36,40); age of menarche and early sexual activity (41); relationship status (42); missing school without parents' permission (43); poor peer relationships (41); having been bullied (44,45); risky behaviours (37,46); life satisfaction (47); disability/chronic illness/special needs (36,48). Detailed information regarding confounding and matching variables used can be found in the **Supplementary material**.

Statistical analyses

All analyses were conducted using Stata version 16.0 (49).

Multivariable analyses

We carried out multiple imputations using chained equations including the exposure, outcomes, and confounder variables (10 imputed datasets). We conducted linear regressions for the continuous outcome (psychological distress) and Poisson regressions for the binary outcomes (high psychological distress, self-harm and attempted suicide).

For each outcome we began with an unadjusted model, and this was followed by models with increasing levels of adjustment: first we included prior mental health problems (psychological distress and self-harm at age 14); then socio-demographic characteristics (ethnicity, parental education, family income, number of siblings, number of parents/caretakers, sexual orientation); puberty-related variables (age of menarche, pubertal status, early sexual activity, sexual violence at age 14); interpersonal relationships (relationship status, peer relationships, bullying); and lastly, in the fully adjusted model, we added health-related variables (BMI, risky behaviours, disability, life satisfaction). The complex survey design of the cohort was accounted for and sample and attrition weights were used in all analyses.

Matched analyses

We used PSM to generate a control group that was matched to the sexual violence group on key variables (see **Supplementary material** for variables), using *psmatch2* in STATA (50). The quality of the matching was assessed using covariate imbalance testing. Once matched, simple linear and Poisson regressions were estimated to investigate the effect of group membership on outcomes. As a sensitivity analysis, matched control groups involving different matching variables were created to examine whether findings were robust to differences in the matching variables used.

In a further examination of the robustness of our results, we combined both approaches whereby we estimated the impact of sexual violence on mental health in the PSM matched groups with a larger set of confounder adjustment (beyond the matching variables).

As an exploratory analysis, we run multivariate analyses separately for harassment online, unwelcomed sexual approach, and sexual assault, in order to evaluate whether the observed effect sizes were similar for these three exposures considered separately.

Lastly, we estimated PAFs to provide an interpretable estimate of effect size of overall mental health problems that would not occur if no adolescent girls experienced sexual violence (51). This was calculated using *punaf* in STATA (52).

Results

As seen in **Figure S1**, the analysed imputed sample for the multivariable regression included 5,134 participants, whereas the analytic sample for the matched analysis comprised of 4,355 participants (due to missing information on matching variables and exclusion of 199 girls with sexual violence experienced before age 14), with the 1,265 participants who experienced sexual violence 1:1 matched to 1,265 controls for analyses.

Descriptive analyses
Table 1 shows the weighted numbers and percentages, and means and standard deviations, for the full analytic sample (sexual violence group vs non-sexual violence group) and for the PSM sample (sexual violence group vs matched control group). **Figure 1** shows the descriptive prevalence for the matched groups at ages 14 and 17 years.

As shown in **Table S1** and **Table S2,** the percentages and means for exposure, outcomes, and each of the key matching variables, were very similar in the full sample and in the PSM sample (for a comparison of the matching variables before and after matching see **Table S3**). **Figure 2** shows both multivariable regression analyses and PSM analyses for all the mental health outcomes for the sexual violence group compared to the non-sexual violence group.

Multivariable analysis
Experiencing sexual violence prior to age 17 years was associated with worse mental health-outcomes at age 17 (**Table 2**). Specifically, in the final adjusted model, having experienced sexual violence was associated with a higher mean psychological distress (mean difference 2.23 [1.67 – 2.79]), and with a higher risk of experiencing high psychological distress (RR 1.74 [1.44 – 2.09]), self-harming (RR 2.04 [1.70 – 2.45]), and attempting suicide (RR 2.26 [1.64 – 3.11]). Unweighted estimates are presented in **Table S4** (for a comparison between the weighted and non-weighted final adjustment see **Table S5**).

Matched sample analyses
The quality of the matching was assessed through evaluation of the percentage of bias for the exposure, outcomes, and matching variables (see **Table S6**). Such bias was found to be low (<5%) for all the variables, with a slightly larger bias for sexual identity (% bias = 6.5).

The sexual violence group, on average, scored higher on psychological distress at 17 years compared to the matched control group (mean difference 1.97 [1.58 – 2.36]), and had greater risk of high distress (RR 1.60 [1.41 – 1.81]), self-harm (RR 1.64 [1.48 – 1.82]) and attempted suicide (RR 1.81 [1.47 – 2.22]).

As sensitivity analyses for the PSM, we used different matching variables to evaluate the best match as well as the difference in their estimates. Slightly different effect sizes were estimated based on the variables included in the matching, with often higher estimates when the matching was conducted with fewer variables (see **Table 2**).

To further assess robustness, we merged the matched PSM groups with the full sample in order to adjust the analysis in the matched sample for all the additional confounders not accounted for by matching (**Table 2**). We found effect sizes slightly attenuated (psychological distress (continuous): 1.84 [1.47 – 2.21]; psychological distress (binary): 1.56 [1.37 – 1.76]; self-harm: 1.59 [1.44 – 1.76]; attempted suicide: 1.70 [1.39 – 2.07].

As a final exploratory analysis, we run multivariate analyses for harassment online, unwelcomed sexual approach, and sexual assault, in order to evaluate whether the effect sizes are similar for these three exposures considered separately. As seen in **Table 3**, all these exposures were associated with worse mental health outcomes. Effect sizes were similar for harassment online and unwelcomed sexual approach, while sexual assault was associated with higher effect sizes for all outcomes.

Finally, we estimated PAFs based on our findings. The PAF defines the fraction of all cases that is attributable to a specific exposure and can be used to estimate the percentage of a certain outcome in a hypothetical scenario where the exposure is null. In other words, we can estimate the percentage of overall mental health problems that would occur if no adolescent girls experienced sexual violence. We calculated such percentage based on the fully adjusted multivariable, the matched and the combined matched-multivariable adjusted models (**Table S7**). In a scenario where girls did not experience sexual violence, there would approximately 25-37% fewer incident mental health problems at age 17 with up to 37% fewer cases for attempted suicide – translating to 9% prevalence (compared to 13.6% observed currently).

Discussion

In our sample, girls who were victims of sexual violence prior to age 17 years reported greater levels of psychological distress, self-harm, and attempted suicide at age 17 years compared to girls without these experiences. These effects persisted even after accounting for previous psychological distress and self-harm and a wide range of relevant confounders, and were robust to multiple methodological approaches and sensitivity checks.

Because this study was based on data from the general population, the reported effect sizes can be considered in a public health context as they apply to the population. Due to the sizes of effect being large for all the analysed mental health outcomes (RRs > 1.50), these findings hold potentially important real-world implications. Based on these estimates, in a world where the examined sexual violence against girls was absent, we could expect a 25-37% reduction in the observed incidences at age 17, thus also substantially narrowing the gender gap reported at this age (53).

In other words, sexual violence at this age contributes non-trivially to the large prevalence of mental ill-health observed in adolescent girls, and actively tackling sexual violence could entail significant improvements in adolescent mental health. In a recent report in the UK, estimates from school

populations showed high levels of sexual violence, with 79% of the surveyed girls reporting sexual assault of any kind (54). The same report highlighted the perceived inadequacy of school-based sex education. However, peer-on-peer in schools could be tackled with appropriate school-based intervention and education (55). This is even more relevant when considering the high prevalence of girls with a history of sexual violence and the fact that many of these girls do not report such experiences due to the stigma around these problems (56). This highlights the importance of formulating more comprehensive research questions about sexual violence experiences as well as the need to gather more data on such experiences in adolescence. Not to mention, sexual violence is known to be widespread in adulthood as well (57), thus suggesting a continuation and potential worsening of mental health sequalae of these experiences. Because sexual violence is a global phenomenon, it is imperative to increase efforts to understand its role in the mental health of adolescent girls while also promoting the development of effective preventative strategies, both to prevent sexual violence and to mitigate their subsequent impacts when it does occur.

Many factors, from biological, social to familial and ascertainment biases, have been attributed as causes for the emergent gender gap in depression and other mental health difficulties in adolescence (1,6). For instance, greater pressures around body image and academic achievements are often cited as important gendered risk factors. Our findings clearly highlight the role of sexual violence, that is disproportionally experienced by girls, as a key driver of the gender gap in adolescent mental health. While it is recommended that future studies also investigate mental health outcomes in males with experiences of sexual violence, it is undeniable that these experiences are much more commonly experienced by females (12,58). Their risk factors, mechanisms and pathways are likely different in males and these differences are important to further understand (17).

The strengths of this study include a population-based cohort, longitudinal data and adjustment for a rich set of important factors that might confound this association. These strengths of the data, accompanied with methods that aim to improve causal inference, help provide robust estimates of the mental health impacts of sexual violence in adolescence. However, there were also some limitations. First, information on mental health and prior sexual violence were collected at the same time in the age 17 assessment, and there is some evidence that mental health can be a predictor in the recall and reporting of these adverse experiences (59). In addition, reported prevalence of harassment, albeit high, is lower than suggested by some other data sources (9,10). Hence it is possible that the association we observe is overestimated as girls without mental health difficulties might not have reported their sexual harassment experiences; on the other hand, underreporting might have resulted in downward bias and an underestimation of effects. Future studies with more longitudinal follow-up data will allow further unpacking of these findings, however, the underreporting of sexual violence experiences is likely to remain an issue. Therefore, better ways of collecting these data in this age group need to be developed and used in large population-based studies. Second, there are limitations of the actual questions asked. For instance, participants were assessed about sexual violence experiences in the previous 12 months, meaning that experiences before this period might not have been reported. The questions about sexual violence were dichotomous, hence it was not possible to investigate important aspects such as severity, frequency and perpetrator. Lower severity experiences such as harassment but experienced constantly might have substantial impacts, as potentially suggested by the findings related to harassment from our disaggregated analyses. The lack of data on frequency, severity and perpetrator is a limitation and

more studies need to collect this information from teenagers. Third, despite PSM being a very useful technique to be used in observational studies where RCTs are not possible, it has been argued that, while matching can improve efficiency and precision, it is limited in reducing confounding by unobserved variables (60). However, the inclusion of a large set of relevant confounders reduces the risk of bias due to unobserved confounding. Future research should aim to triangulate these findings using alternative approaches to assess the impact of sexual violence on mental health.

The implications of our findings are clear and multifold. Given that a reduction in sexual violence is likely to lead to a decrease in mental health problems, this can in turn reduce the societal costs not only for the women affected, but also for their families and wider society (61). Starting early, interventions aimed at educating young people in schools with regards to sexual violence are one actionable avenue (62). Moreover, more attention should be paid at the societal level by policy makers with a focus on reducing the general societal tolerance and permissiveness currently shown for sexual violence (63). The low conviction rates for perpetrators (64) have severe impacts on victims' mental health (65), and erode women's trust in the law enforcement systems that are meant to protect victims. Legal and law enforcement changes might reduce risks, for example by tackling the existing policies regarding prosecutions and more sexual violence deterrents. The effect sizes observed also highlight the urgent need for better tailored and targeted support for victims, to try and mitigate the long-term mental health impacts.

Conclusion

This study aimed at investigating the impact of sexual violence on adolescent girls' mental health using a robust methodology in a longitudinal population-based cohort study. We estimated the extent of the impacts of sexual violence experienced in mid-adolescence on subsequent mental health difficulties. The findings highlight the large burden of sexual violence experienced at this age and stress the importance of reducing these experiences for young women. Based on our findings, if sexual violence experiences in adolescence could be eliminated, 25-37% fewer teenage girls might engage in self-harming and attempted suicide in mid-adolescence. Addressing this risk factor would also contribute substantially to closing the observed gender gap in these mental health outcomes that emerges in adolescence. The findings of this study highlight the need to consider this common and important, yet often gendered and understudied, risk factor more seriously and urgently in adolescent mental health research, clinical practice, and policymaking.

References

1.   Van de Velde S, Bracke P, Levecque K. Gender differences in depression in 23 European countries. Cross-national variation in the gender gap in depression. Soc Sci Med [Internet]. 2010;71(2):305–13. Available from: https://www.sciencedirect.com/science/article/pii/S0277953610002844

2.   Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. Arch Gen Psychiatry. 1994;51(1):8–19.

3.   McManus S, Hassiotis A, Jenkins R, Dennis M, Aznar C, Appleby L, et al. Suicidal thoughts, suicide attempts and self-harm. Ment Heal Wellbeing Engl Adult Psychiatr Morb Surv 2014. 2014;

4.   Salk RH, Hyde JS, Abramson LY. Gender differences in depression in representative national samples: Meta-analyses of diagnoses and symptoms. Psychol Bull. 2017;143(8):783.

5.   Sadler. Mental Health of Children and Young People in England, 2017. Sadler. 2018;(November 2018):233–44.

6.   Piccinelli M, Wilkinson G. Gender differences in depression: Critical review. Br J Psychiatry. 2000;177(6):486–92.

7.   Hyde JS, Mezulis AH, Abramson LY. The ABCs of depression: integrating affective, biological, and cognitive models to explain the emergence of the gender difference in depression. Psychol Rev. 2008;115(2):291.

8.   Schmitz A, Brandt M. Gendered patterns of depression and its determinants in older Europeans. Arch Gerontol Geriatr. 2019;82:207–16.

9.   Eom E, Restaino S, Perkins AM, Neveln N, Harrington JW. Sexual harassment in middle and high school children and effects on physical and mental health. Clin Pediatr (Phila). 2015;54(5):430–8.

10.   Schnoll JS, Connolly J, Josephson WJ, Pepler D, Simkins-Strong E. Same- and Cross-Gender Sexual Harassment Victimization in Middle School: A Developmental-Contextual Perspective. J Sch Violence [Internet]. 2015;14(2):196–216. Available from: http://dx.doi.org/10.1080/15388220.2014.906311

11.   Landstedt E, Gillander Gådin K. Experiences of violence among adolescents: Gender patterns in types, perpetrators and associated psychological distress. Int J Public Health. 2011;56(4):419–27.

12.   Finkelhor D, Shattuck A, Turner HA, Hamby SL. The lifetime prevalence of child sexual abuse and sexual assault assessed in late adolescence. J Adolesc Heal. 2014;55(3):329–33.

13.   Elkin M. The lasting impact of violence against women and girls. Office for National Statistics. 2021.

14. Karatekin C. Adverse childhood experiences (ACEs), stress and mental health in college students. Stress Heal. 2018;34(1):36–45.

15. Schilling EA, Aseltine RH, Gore S. Adverse childhood experiences and mental health in young adults: a longitudinal survey. BMC Public Health. 2007;7(1):1–10.

16. Rinehart SJ, Espelage DL, Bub KL. Longitudinal Effects of Gendered Harassment Perpetration and Victimization on Mental Health Outcomes in Adolescence. J Interpers Violence [Internet]. 2017 Aug 21;35(23–24):5997–6016. Available from: https://doi.org/10.1177/0886260517723746

17. Chiodo D, Wolfe DA, Crooks C, Hughes R, Jaffe P. Impact of Sexual Harassment Victimization by Peers on Subsequent Adolescent Victimization and Adjustment: A Longitudinal Study. J Adolesc Heal [Internet]. 2009 Sep 1;45(3):246–52. Available from: https://doi.org/10.1016/j.jadohealth.2009.01.006

18. Carey KB, Norris AL, Durney SE, Shepardson RL, Carey MP. Mental health consequences of sexual assault among first-year college women. J Am Coll Heal. 2018;66(6):480–6.

19. Rothman K, Georgia Salivar E, Roddy MK, Hatch SG, Doss BD. Sexual assault among women in college: Immediate and long-term associations with mental health, psychosocial functioning, and romantic relationships. J Interpers Violence. 2021;36(19–20):9600–22.

20. Herbert A, Heron J, Barter C, Szilassy E, Barnes M, Howe LD, et al. Risk factors for intimate partner violence and abuse among adolescents and young adults: findings from a UK population-based cohort. Wellcome open Res. 2020;5.

21. Khadr S, Clarke V, Wellings K, Villalta L, Goddard A, Welch J, et al. Mental and sexual health outcomes following sexual assault in adolescents: a prospective cohort study. Lancet Child Adolesc Heal. 2018;2(9):654–65.

22. Valencia-Agudo F, Kramer T, Clarke V, Goddard A, Khadr S. Correlates and predictors of self-harm in a prospective sample of sexually assaulted adolescents. Clin Child Psychol Psychiatry. 2020;25(4):847–59.

23. Bendixen M, Daveronis J, Kennair LEO. The effects of non-physical peer sexual harassment on high school students' psychological well-being in Norway: consistent and stable findings across studies. Int J Public Health [Internet]. 2018;63(1):3–11. Available from: https://doi.org/10.1007/s00038-017-1049-3

24. Landstedt E, Gillander Gådin K. Deliberate self-harm and associated factors in 17-year-old Swedish students. Scand J Public Health. 2011;39(1):17–25.

25. Baiden P, Panisch LS, Kim YJ, LaBrenz CA, Kim Y, Onyeaka HK. Association between first sexual intercourse and sexual violence victimization, symptoms of depression, and suicidal behaviors among adolescents in the United States: findings from 2017 and 2019 National Youth Risk Behavior Survey. Int J Environ Res Public Health. 2021;18(15):7922.

26. Ståhl S, Dennhag I. Online and offline sexual harassment associations of anxiety and depression in an adolescent sample. Nord J Psychiatry. 2021;75(5):330–5.

27. Dahlqvist HZ, Landstedt E, Young R, Gådin KG. Dimensions of Peer Sexual Harassment Victimization and Depressive Symptoms in Adolescence: A Longitudinal Cross-Lagged Study in a Swedish Sample. J Youth Adolesc. 2016;45(5):858–73.

28. Marshall SK, Faaborg-Andersen P, Tilton-Weaver LC, Stattin H. Peer sexual harassment and deliberate self-injury: Longitudinal cross-lag investigations in Canada and Sweden. J Adolesc Heal [Internet]. 2013;53(6):717–22. Available from: http://dx.doi.org/10.1016/j.jadohealth.2013.06.009

29. Lanza ST, Moore JE, Butera NM. Drawing causal inferences using propensity scores: A practical guide for community psychologists. Am J Community Psychol. 2013;52(3–4):380–92.

30. Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. Biometrika. 1983;70(1):41–55.

31. Joshi H, Fitzsimons E. The Millennium Cohort Study: the making of a multi-purpose resource for social science and policy. Longit Life Course Stud. 2016;7(4):409–30.

32. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand S-L, et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. Psychol Med. 2002;32(6):959–76.

33. Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, et al. Screening for serious mental illness in the general population. Arch Gen Psychiatry. 2003;60(2):184–9.

34. Coulter RWS, Mair C, Miller E, Blosnich JR, Matthews DD, McCauley HL. Prevalence of Past-Year Sexual Assault Victimization Among Undergraduate Students: Exploring Differences by and Intersections of Gender Identity, Sexual Identity, and Race/Ethnicity. Prev Sci [Internet]. 2017 Aug;18(6):726–36. Available from: https://pubmed.ncbi.nlm.nih.gov/28210919

35. Mendle J, Ryan RM, Mckone KM. Early Childhood Maltreatment and Pubertal Development: Replication in a Population-Based Sample. 2015;

36. Butler AC. Child sexual assault: Risk factors for girls. Child Abuse Negl. 2013;37(9):643–52.

37. Aucott L. Mental Well-Being Related To Lifestyle and Risky Behaviours in 18-25 Year Old: Evidence from North-East Scotland. Int J Public Heal Res [Internet]. 2014 Mar 1;4(1 SE-Public Health Research Articles). Available from: http://spaj.ukm.my/ijphr/index.php/ijphr/article/view/207

38. Finkelhor D, Ormrod RK, Turner HA. Re-victimization patterns in a national longitudinal sample of children and youth. Child Abus Negl. 2007;31(5):479–502.

39. Lawson DW, Mace R. Siblings and childhood mental health: Evidence for a later-born advantage. Soc Sci Med. 2010;

40. Barrett AE, Turner RJ. Family structure and mental health: The mediating effects of socioeconomic status, family process, and social stress. J Health Soc Behav. 2005;46(2):156–69.

41. Vicary JR, Klingaman LR, Harkness WL. Risk factors associated with date rape and sexual assault of adolescent girls. J Adolesc. 1995;18(3):289–306.

42. Organization WH. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. World Health Organization; 2013.

43. RUNTZ M, BRIERE J. Adolescent "Acting-Out" and Childhood History of Sexual Abuse. J Interpers Violence [Internet]. 1986 Sep 1;1(3):326–34. Available from: https://doi.org/10.1177/088626086001003005

44. Coggan C, Bennett S, Hooper R, Dickinson P. International Journal of Mental Health Promotion Association between Bullying and Mental Health Status in New Zealand Adolescents E E Association between Bullying and Mental Health Status in New Zealand Adolescents. Int J Ment Heal Promot Vol [Internet]. 2003 [cited 2021 May 6];5(1):16–22. Available from: https://www.tandfonline.com/action/journalInformation?journalCode=rijm20

45. DUNCAN RD. Peer and Sibling Aggression:: An Investigation of Intra- and Extra-Familial Bullying. J Interpers Violence [Internet]. 1999 Aug 1;14(8):871–86. Available from: https://doi.org/10.1177/088626099014008005

46. Charak R, Koot HM, Dvorak RD, Elklit A, Elhai JD. Unique versus cumulative effects of physical and sexual assault on patterns of adolescent substance use. Psychiatry Res. 2015 Mar 5;230(3):763–9.

47. Schou-Bredal I, Bonsaksen T, Ekeberg Ø, Skogstad L, Grimholt TK, Lerdal A, et al. Sexual Assault and the Association With Health, Quality of Life, and Self-Efficacy in the General Norwegian Population. 2020;

48. Alriksson-Schmidt AI, Armour BS, Thibadeau JK. Are adolescent girls with a physical disability at increased risk for sexual violence? J Sch Health. 2010;80(7):361–7.

49. StataCorp. Stata Statistical Software: Release 16. Coll Station TX. 2019;StataCorp.

50. Leuven E, Sianesi B. PSMATCH2: Stata module to perform full Mahalanobis and propensity score matching, common support graphing, and covariate imbalance testing. 2003;

51. Mansournia MA, Altman DG. Population attributable fraction. BMJ [Internet]. 2018;360(February):2–3. Available from: http://dx.doi.org/doi:10.1136/bmj.k757

52. Newson R. PUNAF: Stata module to compute population attributable fractions for cohort studies. 2010;

53. Patalay P, Fitzsimons E. Psychological distress, self-harm and attempted suicide in UK 17-year olds: prevalence and sociodemographic inequalities. Br J Psychiatry. 2021;1–3.

54. Ofsted. Review of sexual abuse in schools and colleges. 2021;

55. Miller E. Prevention of and Interventions for Dating and Sexual Violence in Adolescence. Pediatr Clin North Am [Internet]. 2017;64(2):423–34. Available from: https://www.sciencedirect.com/science/article/pii/S0031395516411521

56. Kennedy AC, Prock KA. "I still feel like I am not normal": A review of the role of stigma and stigmatization among female survivors of child sexual abuse, sexual assault, and intimate

partner violence. Trauma, Violence, Abus. 2018;19(5):512–27.

57.   de Toledo Blake M, Drezett J, Vertamatti MA, Adami F, Valenti VE, Paiva AC, et al. Characteristics of sexual violence against adolescent girls and adult women. BMC Womens Health. 2014;14(1):1–7.

58.   Barth J, Bermetz L, Heim E, Trelle S, Tonia T. The current prevalence of child sexual abuse worldwide: a systematic review and meta-analysis. Int J Public Health. 2013;58(3):469–83.

59.   Langeland W, Smit JH, Merckelbach H, de Vries G, Hoogendoorn AW, Draijer N. Inconsistent retrospective self-reports of childhood sexual abuse and their correlates in the general population. Soc Psychiatry Psychiatr Epidemiol. 2015;50(4):603–12.

60.   Pearce N. Analysis of matched case-control studies. bmj. 2016;352.

61.   Trautmann S, Rehm J, Wittchen H-U. The economic costs of mental disorders: Do our societies react appropriately to the burden of mental disorders? EMBO Rep [Internet]. 2016/08/04. 2016 Sep;17(9):1245–9. Available from: https://pubmed.ncbi.nlm.nih.gov/27491723

62.   Letourneau EJ, Schaeffer CM, Bradshaw CP, Feder KA. Preventing the Onset of Child Sexual Abuse by Targeting Young Adolescents With Universal Prevention Programming. Child Maltreat [Internet]. 2017/01/01. 2017 May;22(2):100–11. Available from: https://pubmed.ncbi.nlm.nih.gov/28413921

63.   Brown CS, Biefeld SD, Elpers N. A Bioecological Theory of Sexual Harassment of Girls: Research Synthesis and Proposed Model. Rev Gen Psychol [Internet]. 2020 Sep 10;24(4):299–320. Available from: https://doi.org/10.1177/1089268020954363

64.   Temkin J, Krahé B. Sexual assault and the justice gap: A question of attitude. Vol. 5. Bloomsbury Publishing; 2008.

65.   Bohner G, Eyssel F, Pina A, Siebler F, Viki GT. Rape myth acceptance: Cognitive, affective and behavioural effects of beliefs that blame the victim and exonerate the perpetrator. In: Rape. Willan; 2013. p. 40–68.

66.   Messer SC, Angold A, Costello EJ, Loeber R, al et. Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents: Factor composition and structure across development. Int J Methods Psychiatr Res. 1995;5(4):251–62.

67.   Lundervold AJ, Breivik K, Posserud M-B, Stormark KM, Hysing M. Symptoms of depression as reported by Norwegian adolescents on the Short Mood and Feelings Questionnaire. Front Psychol. 2013;4:613.

Tables and Figures

Table 1. Descriptive analysis of mental health outcomes in different analytic samples

	Full analytic sample (N = 5,134)	Full analytic sample: Sexual violence group (N = 1,550)	Full analytic sample: Non-sexual violence group (N = 3,572)	PSM sample: Sexual violence group (N = 1,265)	PSM sample: Matched control group (N = 1,265)
	N (%) or M (SE) – [95% CI]	N (%) or M (SE) – [95% CI]	N (%) or M (SE) – [95% CI]	N (%) or M (SE) – [95% CI]	N (%) or M (SE) – [95% CI]
Mental health at age 17					
Psychological distress (continuous)					
	8.39 (0.18) – [8.02 – 8.75]	10.76 (0.20) – [10.36 – 11.15]	7.34 (0.25) – [6.85 – 7.84]	10.43 (0.20) – [10.05 – 10.82]	8.37 (0.20) – [7.98 – 8.76]
Psychological distress (binary)					
< 13 tot score	3,895 (77.8) – [75.23 – 80.16]	941 (62.3) – [58.27 – 66.07]	2,949 (84.2) – [80.90 – 87.10]	796 (65.3) – [61.77 – 68.65]	972 (78.6) – [75.30 – 81.48]
≥ 13 tot score	1,097 (22.2) – [19.84 – 24.77]	550 (37.7) – [33.93 – 41.73]	544 (15.8) – [12.90 – 19.10]	438 (34.7) – [31.35 – 38.23]	275 (21.4) – [18.52 – 24.70]
Self-harm					
No	3,524 (71.7) – [69.35 – 73.91]	748 (50.9) – [45.12 – 53.15]	2,774 (80.9) – [78.08 – 83.41]	635 (48.3) – [44.18 – 52.50]	878 (72.9) – [69.62 – 75.95]
Yes	1,439 (28.3) – [26.09 – 30.65]	733 (49.1) – [46.88 – 54.88]	704 (19.1) – [16.59 – 21.92]	594 (51.7) – [47.50 – 55.82]	367 (27.1) – [24.05 – 30.38]
Attempted suicide					
No	4,442 (89.2) – [87.72 – 90.60]	1,187 (78.4) – [74.63 – 81.80]	3,252 (93.8) – [92.40 – 95.01]	1,012 (80.4) – [75.93 – 84.14]	1,121 (91.0) – [88.96 – 92.71]
Yes	512 (10.8) – [9.40 – 12.28]	289 (21.6) – [18.20 – 25.37]	220 (6.2) – [4.99 – 7.60]	216 (19.6) – [15.86 – 24.07]	121 (9.0) – [7.29 – 11.04]
Mental health at age 14					
Psychological distress (continuous)					
	7.12 (0.15) – [6.83 – 7.41]	9.40 (0.30) – [8.80 – 9.99]	6.05 (0.15) – [5.75 – 6.35]	8.85 (0.32) – [8.22 – 9.47]	8.79 (0.28) – [8.24 – 9.34]
Psychological distress (binary)					
< 12 tot score	3,550 (77.4) [75.71 – 78.99]	913 (66.1) [62.31 – 69.61]	2,634 (82.7) [80.85 – 84.44]	840 (68.7) – [64.70 – 72.49]	856 (69.4) – [65.81 – 72.68]
≥ 12 tot score	1,023 (22.6) [21.01 – 24.29]	475 (33.9) [30.4 – 37.7]	547 (17.3) [15.56 – 19.15]	395 (31.3) – [27.51 – 35.30]	393 (30.7) – [27.32 – 34.19]
Self-harm					
No	3,601 (77.8) – [76.00 – 79.54]	907 (64.8) – [60.75 – 68.58]	2,692 (83.9) – [81.88 – 85.67]	841 (67.7) – [63.67 – 71.47]	872 (69.7) – [65.97 – 73.18]
Yes	976 (22.2) – [20.46 – 24.00]	478 (35.2) – [31.42 – 39.25]	496 (16.1) – [14.33 – 18.12]	394 (32.3) – [28.53 – 36.33]	377 (30.3) – [26.82 – 34.03]

N numbers; % percentage; M mean; SD standard deviation; CI confidence interval.

Table 2. Regression coefficients and risk ratios for sexual violence predicting mental health outcomes from a range of modelling approaches

	Psychological distress (continuous)	High psychological distress	Self-harm	Attempted Suicide
	Coefficient [95% CI]	RR [95% CI]	RR [95% CI]	RR [95% CI]
Multivariable regressions with increasing levels of confounder adjustment				
Unadjusted	3.44 [2.78 – 4.10]	2.39 [1.91 – 3.00]	2.68 [2.25 – 3.20]	3.55 [2.61 – 4.82]
Adjustment 1	2.32 [1.64 – 3.00]	1.78 [1.42 – 2.23]	2.17 [1.78 – 2.65]	2.37 [1.65 – 3.40]
Adjustment 2	2.19 [1.57 – 2.82]	1.72 [1.40 – 2.11]	2.04 (1.70 – 2.45]	2.23 [1.58 – 3.14]
Adjustment 3	2.26 [1.65 – 2.87]	1.75 [1.42 – 2.15]	2.06 [1.72 – 2.47]	2.22 [1.58 – 3.14]
Adjustment 4	2.20 [1.59 – 2.81]	1.72 [1.40 – 2.11]	2.06 (1.71 – 2.47]	2.26 [1.62 – 3.15]
Fully adjusted model	2.23 [1.67 – 2.79]	1.74 [1.44 – 2.09]	2.04 [1.70 – 2.45]	2.26 [1.64 – 3.11]
Regressions in Propensity score matched (PSM) samples				
PSM (all matching variables)	1.97 [1.58 – 2.36]	1.60 [1.41 – 1.81]	1.64 [1.48 – 1.82]	1.81 [1.47 – 2.22]
PSM (all minus sexual identity, pubertal status, parental education)	2.44 [2.05 – 2.83]	1.80 [1.57 – 2.05]	1.77 [1.59 – 1.97]	1.84 [1.49 – 2.27]
PSM (all minus psychological distress and self-harm at age 14)	2.97 [2.58 – 3.36]	2.04 [1.77 – 2.35]	1.96 [1.75 – 2.20]	2.65 [2.08 – 3.37]
Combined PSM sample with full confounder adjustment on non-matching confounder variables				
Adjusted regression analysis	1.84 [1.47 – 2.21]	1.56 [1.37 – 1.76]	1.59 [1.44 – 1.76]	1.70 [1.39 – 2.07]

RR risk ratio; CI confidence interval. All *p* < 0.0001.

Table 3. Regression coefficients and risk ratios for mental health outcomes by sexual violence type from a multivariable model including all listed confounders

Full-sample and PSM sample merged	Psychological distress (continuous)	Psychological distress (binary)	Self-harm	Attempted Suicide
	Coefficient [95% CI]	RR [95% CI]	RR [95% CI]	RR [95% CI]
Adjusted regression analysis				
Harassment online	2.65 [2.31 – 2.99]	1.73 [1.55 – 1.92]	1.78 [1.64 – 1.95]	2.26 [1.91 – 2.67]
Unwelcomed sexual approach	2.26 [1.93 – 2.59]	1.72 [1.55 – 1.91]	1.81 [1.66 – 1.97]	1.82 [1.54 – 2.16]
Sexual assault	3.09 [2.51 – 3.67]	1.80 [1.57 – 2.07]	1.94 [1.74 – 2.16]	2.34 [1.89 – 2.90]

Number of observations: 5,134.

Figure 1. Psychological distress, self-harm, and attempted suicide prevalence at age 14 and at age 17 in the group that experienced sexual violence between these ages compared to the group that did not experience sexual violence.

Note. This graph is based on prevalence in propensity score matched groups. N in each group = 1,265. Prevalence of high depression/distress are not comparable across ages 14 and 17. Depressive symptoms at 14 assessed using the Short Moods and Feelings Questionnaire and distress at age 17 using the Kessler K-6 measure of psychological distress.

Figure 2. Multivariable regression analyses and PSM analyses for psychological distress (as both a continuous and binary measure), self-harm, and attempted suicide at age 17 in the sexual violence group compared to the non-sexual violence group.

Data sharing statement: The Millennium Cohort Study data is available to use by researchers via the UK Data Service (https://ukdataservice.ac.uk/).