

**Title:**

**Global burden of mental health problems among children and adolescents during COVID-19 pandemic: A systematic umbrella review**

**Authors:**

**M. Mahbub Hossain, Fazilatun Nesa, Jyoti Das, Roaa Aggad, Samia Tasnim, Mohan Bairwa, Ping Ma, Gilbert Ramirez**

**Affiliations:**

- a. School of Public Health, Texas A&M University, TX 77840.
- b. Research Initiative for Health Equity, Khulna, Bangladesh.
- c. North South University, Dhaka, Bangladesh.
- d. All Indian Institute of Medical Sciences, New Delhi, India.
- e. Robert Stempel College of Public Health and Social Work, Florida International University, FL 33174.

**\*Corresponding author:** M. Mahbub Hossain, School of Public Health, Texas A&M University, TX 77840. Email: mahbub321@gmail.com

## Abstract

Mental health problems among children and adolescents are increasingly reported amidst the coronavirus disease (COVID-19) pandemic. In this umbrella review, we aimed to synthesize global evidence on the epidemiologic burden and correlates of child and adolescent mental health (CAMH) problems during this pandemic from existing systematic reviews and meta-analyses. We evaluated 422 citations and identified 17 eligible reviews on the epidemiology of CAMH in the context of COVID-19. Most of the reviews reported a high prevalence of anxiety, depression, sleep disorders, suicidal behavior, stress-related disorders, attention-deficit/hyperactivity disorder, and other mental health problems. Also, factors associated with CAMH such as age, gender, place of residence, educational attainment, household income, sedentary lifestyle, social media and internet use, comorbidities, family relationships, parents' psychosocial conditions, COVID-19 related experiences, closure of schools, online learning, and social support were reported across reviews. As most studies were cross-sectional and used nonrepresentative samples, future research on representative samples adopting longitudinal and intervention designs is needed. Lastly, multipronged psychosocial care services, policies, and programs are needed to alleviate the burden of CAMH problems during and after this pandemic.

**Keywords:** Mental Health; Psychiatry; Psychology; COVID-19; Pandemic; Child Health; Adolescent Health; Systematic Review; Meta-analysis; Umbrella Review

## Introduction

The world has become acquainted with the coronavirus disease 2019 (COVID-19) pandemic that has triggered a wide range of biological, psychological, and socio-economic consequences globally [1–3]. A growing body of empirical research informs a high burden of mental health problems in recent times that is increasingly recognized as a psychiatric epidemic synergistically occurring alongside this pandemic [4, 5]. Several reviews suggested a heavy psychosocial impact of COVID-19 pandemic resulting in mental health problems such as anxiety, depression, posttraumatic stress disorders (PTSD), and sleep disorders [4, 6–9]. An umbrella review of eight systematic reviews and meta-analyses found that preventive measures such as quarantine and isolation for infection prevention may have immediate and prolonged mental health impacts [10]. This evidence suggests ongoing and future mental health crises associated with this pandemic that should be investigated and addressed across various population groups globally.

Mental health problems in childhood and adolescence are unique as they have versatile correlates and may have long-term health biopsychosocial consequences [11–13]. The global prevalence of mental health disorders in children and adolescents has been inconsistent yet high across previous syntheses of research [14, 15]. Specifically, several childhood mental health problems, including depression and anxiety, have become highly prevalent over the past years [8, 11, 14, 15]. For example, a recent study found that the incidence of depression or anxiety in ages 6 to 17 has risen from 5.4% in 2003 to 8.4% in 2011-2012 in the US [16]. Another study that was published in 2010 reported that 1 in 5 US children experience a serious or debilitating mental health problem in their lifetime [17]. Similar studies from Canada [18, 19], Europe [20, 21], and Australia [22, 23] suggest comparative high burden of child and adolescent mental health (CAMH) problems. Moreover, countries with limited resources, mostly those in the global south, often lack high-quality evidence on mental health epidemiology. Yet, emerging literature from many low- and middle-income countries suggests a high burden of CAMH problems in those contexts that is a global mental health concern [24–26]. These numbers are expected to increase amidst and after the COVID-19 pandemic.

The impacts of the COVID-19 pandemic on the mental health outcomes of children have drawn considerable attention amongst global researchers. Children and adolescents constitute the

most vulnerable population affected by the preventive measures for COVID-19, such as the closure of schools, leading to reduced interaction with peers, and lessened opportunities for physical activity and exploration [27, 28]. These effects are expected to have adverse impacts on the welfare and mental health of children through sleeping disorders, depression, stress and anxiety [5, 28]. Studies conducted across the globe provide evidence of the detrimental impact of the pandemic on the mental health of adolescents [29]. During the COVID-19 pandemic lockdown, the percentage of Bangladeshi children who suffered from mental health disturbances ranged from 43% for subthreshold disturbances to 7.2% for severe disturbances [30]. In Turkey, children of health workers had significantly higher anxiety levels measured using the State-Trait Anxiety Inventory for Children (STAI-C) than non-health workers' children [31]. In Ireland, a qualitative study showed that adolescents and children experienced increased rates of social isolation, maladaptive behavior, anxiety, and depression during the pandemic [32]. Moreover, the pandemic resulted in a significant psychosocial impact on Chinese adolescents and children in terms of emotional distress, with a prevalence of 24.9% for anxiety, 19.7% for depression, and 15.2% for stress [33]. Another systematic review of Chinese studies showed that the prevalence of COVID-19 related posttraumatic stress, sleep disorders, anxiety, and depression were 48%, 44%, 26%, and 29%, respectively [34].

An increasing body of primary studies has facilitated the development of many such reviews that should be synthesized to provide an extensive overview of all CAMH problems and their associated factors. Such systematic overview of reviews, also known as umbrella reviews, have become increasingly useful in understanding the evidence landscape in a given domain [35]. The findings of an umbrella review can offer a “bird’s eye” view of a scientific field that may help evidence-based decision-making, synthesizing future evidence, and conducting primary research on least-explored areas of knowledge and practice [35–37]. This umbrella review aimed to summarize the global evidence on the epidemiological burden and correlates of mental health problems among children and adolescents during the COVID-19 pandemic from existing systematic reviews and meta-analyses. The results of this review can help understand and address mental health problems among children and adolescents in this pandemic and future public health emergencies.

## **Methodology**

### **Guidelines, data sources, and search strategy**

In this umbrella review, we have followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guideline [38] and the Joanna Briggs Institute (JBI) recommendations [39]. The conduct of an umbrella review aims to address the broader scope of a scholarly topic and related issues presenting a wide picture of existing evidence. Umbrella reviews identify and examine evidence from existing systematically conducted reviews through a rigorous process. In this umbrella review, we conducted an extensive literature search in PubMed/MEDLINE, APA PsycInfo, Academic Search Ultimate, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Health Source- Nursing/Academic Edition, Health Policy Reference Center, and Web of Science using a set of keywords as outlined in Table 1.

### **Inclusion criteria**

#### ***Participants***

This review focused on children and adolescents irrespective of the age range adopted in systematic reviews and the primary studies included in the same. For studies with mixed participants such as adolescents with young adults or children as a part of other population groups, a decision was made prior to conducting the review about including a review if at least 70% of its studies focused on children or adolescents.

#### ***Phenomenon of interest***

This review studied a wide range of mental health problems, including but not limited to psychiatric disorders enlisted in the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) and the International Classification of Diseases (ICD-10) Classification of Mental and Behavioral Disorders. To make the evidence inclusive of any problem that affects mental health and wellbeing, this review considered research that emphasized psychosocial problems alongside psychiatric conditions.

**Table 1: Keywords and search strategy**

<b>Search query</b>	<b>Search topic</b>	<b>Search keywords (titles, abstracts, general keywords, and subject headings)</b>
1	Condition/ outcome of interest	“mental health” OR “mental wellbeing” OR “mental well-being” OR “mental disorder*” OR “mental illness” OR “psychiatr*” OR “psychological health” OR “psychological distress” OR “psychological impact” OR “psychological outcomes” OR “psychological consequence*” OR “psychological comorbid*” OR “psychosocial problem*” OR “behavioral problem*” OR “behavioral disorder*” OR “cognitive disorder*” OR “cognitive impairment*” OR “emotional distress” OR “depression” OR “depressive disorder*” OR “anxiety” OR “PTSD” OR “PTSS” OR “posttraumatic” OR “posttraumatic” OR “addiction” OR “substance use disorders” OR “mood disorder*” OR “affective disorder*” OR “DSM*” OR “psychosis” OR “psychotic” OR “oppositional defiant disorder” OR “hyperactiv*” OR “conduct disorder” OR “obsess*” OR “phobi*” OR “schizophren*” OR “bipolar disorder” OR “anorexia” OR “bulimi*” OR “challenging behav*”
2	Population	“child*” OR “adolescen*” OR “young people” OR “school students” OR “youth” OR “teen*”
3	Exposure/ Context	“Coronavirus” OR “COVID-19” OR “COVID19” OR “SARS-CoV-2” OR “2019 novel coronavirus” OR “2019-nCoV”
4	Phenomenon	“prevalence” OR “incidence” OR “rate” OR “diagno*” OR “disease burden” OR “epidemiolog*” OR “frequen*” OR “determinant*” OR “predictor*” OR “correlate*” OR “risk factors” OR “protective factors” OR “associated factors”
5	Study types	“systematic review” OR “systematic literature review” OR “meta-analy*” OR “metaanaly*” OR “quantitative review” OR “qualitative review” OR “meta-synthesis” OR “pooled prevalence” OR “pooled estimate*” OR “scoping review” OR “critical review” OR “integrative review” OR “evidence-based review” OR “synthesized evidence”
<b>Final search query</b>	Intersection of four topics	1 AND 2 AND 3 AND 4 AND 5

## **Context**

The context of this review was the COVID-19 pandemic. Therefore, studies conducted during and in the context of this pandemic were considered eligible for this review. If a review used data collected beyond the scope of this pandemic, it was considered ineligible for inclusion. Moreover, studies with mixed data from COVID-19 and previous infectious outbreaks were reviewed carefully. In such cases, a study was considered eligible if at least 70% of the respondents on included studies informed evidence in the context of COVID-19.

## **Types of studies**

Any review article with a systematic approach to retrieve and synthesize evidence was considered eligible for inclusion in this umbrella review. This decision was made considering methodological differences within systematically conducted review articles. Therefore, systematic literature reviews, scoping reviews, and meta-analyses were included, whereas primary studies or narrative reviews without a clear review method were excluded. Moreover, articles were included in the full text was available in the English language. Lastly, studies published from January 1, 2020, to December 31, 2021, were primarily included in this review, and the search was last updated on March 12, 2022.

## **Study selection**

We collated and uploaded all identified citations were to the cloud-based screening tool Rayyan. Two reviewers independently reviewed those citations using the above-mentioned criteria. All conflicts were addressed through a discussion, and the opinion of a third reviewer was taken whenever it was necessary. Articles that appeared to be qualified at this stage were considered for full-text review, and a final set of articles meeting all eligibility criteria was retained for data extraction and synthesis.

## **Data extraction and synthesis**

We used a standardized data extraction form that included key variables of interest such as the bibliographic information of each article, the type of the reviews, number of primary studies, databases accessed in respective reviews, characteristics of the study samples and populations, epidemiologic measures used in those studies, and finally, the burden of mental health problems

and factors associated with the same. Two reviewers (FN and AD) conducted the data extraction independently in the supervision of a third reviewer (ST). The extracted data were further reviewed for accuracy and quality by another reviewer (MMH). As this review included a wide range of systematically conducted reviews with and without quantitative synthesis, there was marked heterogeneity in the methods and outcomes of the included reviews. For this reason, a narrative synthesis of overall evidence on the burden of and factors associated with mental health problems in children and adolescents during the COVID-19 pandemic was presented in this review.




Figure 1: Flow diagram of the umbrella review process

### Assessment of methodological quality

In this umbrella review, we used the JBI checklist for the critical appraisal of systematic reviews and research syntheses to assess the methodological quality of studies included in this umbrella review [35, 39]. This checklist has ten items evaluating different methodological aspects of a

systematically conducted review, including the appropriateness of the search strategies, the approach to synthesizing evidence, potential sources of biases, and prospects for future research and policymaking. We used this ten-items checklist and allocated one point for each item. So, the overall score of a study can range from zero to ten. In this review, studies receiving zero to four, five to seven, and eight to ten points were categorized as the low, medium, and high-quality studies, respectively.

## **Results**

### **Overview of the included studies**

We found a total of 422 citations using the search strategy described earlier (See Figure 1 for the review process). After eliminating 181 duplicate citations, we retained 241 unique citations and reviewed their titles and abstracts. A total of 24 citations met our preliminary criteria and were retained for full-text review. After full-text evaluation, 17 systematic reviews [29, 34, 48–54, 40–47], including 6 meta-analytic reviews [34, 42, 43, 46, 49, 51] met all criteria of this umbrella review and were selected for further analysis (see Table 2).

### **Characteristics of the included studies**

Seventeen reviews included in this umbrella review used different databases for retrieving primary studies. The number of databases ranged from 2 to 11 with a median number of 5 databases, whereas MEDLINE and WHO COVID-19 databases are the leading sources of information across multiple reviews.

The number of primary studies included in different reviews ranged from 5 to 102. Moreover, the sample size across studies ranged from 12 to 1,199,320 in systematic reviews, whereas a meta-analytic review used a pooled sample of 80,879 participants [49]. The majority of the primary studies were cross-sectional, and online data collection measures were widely used across studies. China was the most common source of primary studies, followed by the North American and European countries. Limited primary studies were reported from LMICs such as India, Brazil, Bangladesh, and countries from Sub-Saharan Africa [41, 46, 48–50, 52].

The methodological quality of the reviews was assessed using the JBI critical appraisal checklist (See Supplementary Material). Most of the reviews (n = 13) included in this umbrella review were of high quality [29, 34, 51, 53, 54, 40–44, 48–50], whereas the remaining (n = 4) reviews had medium quality [45–47, 52]. None of the reviews were identified to have low methodological quality as per the JBI checklist.

## **Epidemiological burden of CAMH during COVID-19**

### ***Anxiety***

Thirteen reviews reported a varying burden of anxiety among children and adolescents during this pandemic [29, 34, 51, 53, 54, 40, 43, 44, 46–50]. The prevalence of anxiety ranged from 1.85% to 74% across primary studies included in the reviews. For example, studies reported in the review by Panchal et al. (2021) found the prevalence of anxiety ranging from 1.85% to 49.5%. Another review by Samji et al. (2021) reported the prevalence of anxiety between 8% and 74%. A meta-analytic review by Racine et al. (2021) found that 20.5% (95% CI: 17% - 24%) participants had anxiety disorders during this pandemic.

### ***Depression***

Among the included studies, thirteen reviews and meta-analyses evaluated a high prevalence of depression among children and adolescents ranging from 2% to 71.5% [29, 34, 52–54, 41, 43, 44, 46–50]. For example, Marchi et al. (2021) found that the prevalence of depression was between 2% and 4% across the reviewed literature. Another review by Oliveira et al. (2022) reported the prevalence of depression between 6.3% and 71.5%. A meta-analytic review by Ma et al. (2021) reported the pooled prevalence of depression as 29% (95% CI: 17% - 40%) among children and adolescents.

### ***Sleep disorders***

Seven reviews reported varying magnitude of sleep disorders among children and adolescents during this pandemic [34, 41, 42, 44, 48, 51, 54]. For example, Panchal et al. (2021) reported that the prevalence of sleep disorders was 20% and 55.9% among children and adolescents, respectively. Another review by Sharma et al. (2021) reported a pooled prevalence of any sleep disorders as 54%

**Table 2: Overview of evidence from existing systematic reviews and meta-analyses on CAMH problems during COVID-19**

Author and year of publication	Name(s) and timeframe of searching Databases	Number of primary studies; type of review (meta-analysis or systematic review)	Sample sizes in the included primary studies (total or range as reported in respective reviews)	Characteristics of the study populations (demographics, recruitment strategy, and location)	Burden of mental health (prevalence, rate, correlates or any quantitative measures of epidemiologic burden)	Factors associated with children and adolescent mental health in respective studies
<b>Nearchou et al. (2020)</b>	APA PsycInfo, MEDLINE, CINAHL, Scopus, PubMed, Embase, ERIC, and the WHO Global Health database till June 2020	12; Systemic Review	Sample size range from 17 to 8072.	The study focused on human participant ≤18 years old; applied quantitative cross-sectional design; data collected through online and open access forum; conducted in China, Italy, Poland, Turkey, and United State	The prevalence of depression and anxiety in young people ranged from 22.6% to 43.7%. and 18.9% to 37.4% respectively. Moreover, among children and adolescent 22%-62.2% were found to have Covid-19 related emotional reaction as well as 2.39%-22% in terms of having somatic syndrome	Factors like fear of affecting with Covid-19, social distancing, Covid-19 circumstances, and emergencies were found to be associated with mental health outcome. However, some positive domains were found such as belongingness, relationship with peer, hope, meaning of life, life satisfaction.
<b>Panda et al. (2021)</b>	MEDLINE, Embase, Web of Science, CENTRAL, medRxiv and bioRxiv till August 2020	15; Systemic Review	Sample size range from 50 to 8079	The study included children up to 18 year and their caregiver; participants are recruited through direct and online interview; studies were cross-sectional design from France Italy, China, India	Studies had found 79.4% children were found to have psychological state due to the pandemic while 22.5%, 35.2%, 21.3% suffered from fear of covid-19 infection, boredom, and sleep disturbance respectively. Overall, 34.5%, 41.7%,	Studies found multiple associated factors like female, urban citizen, clinical depression level, family member infected with covid-19, non-medical occupation, poor academic records, addicted to internet or smartphone. On the other hand, higher exercise,

				Hong Kong, Brazil, Turkey, Bangladesh and Korea were reported	42.3% and 30.8% of children were found to be suffering from anxiety, depression, irritability, and inattention. Similarly, 52.3% and 27.4% of caregivers developed anxiety and depression, respectively.	knowledge on covid-19, prevention, and control measures, having siblings found to be protective factors.
<b>Bussières et al. [42]</b>	MEDLINE, ERIC, APA PsycInfo, and Google Scholar till June 2021	28; Meta-Analysis	Sample size ranges from 12 to 8124; total of 14209 participants.	Children aged 5-13 years old (including pre-existing conditions like ASD, epilepsy, obesity) were recruited; The quantitative synthesis of cross sectional, longitudinal and retrospective studies done; most study were from European countries followed by middle east then America	Studies found that children demonstrated longer sleeping hours ( $g = 0.324$ ; 95% CI [0.10, 0.55]; $p = 0.004$ ; $k = 9$ ) and higher levels of both internalized ( $g = 0.215$ ; 95% CI [0.06, 0.37]; $p < 0.001$ ; $k = 19$ ) and externalized ( $g = 0.141$ (95% CI [0.08, 0.21]; $p < 0.01$ ; $k = 15$ ) symptoms during lockdown. effect sizes in studies where the child was the informant ( $g = 0.57$ ; 95% CI [-0.36, 1.50]; $p = 0.23$ ; $k = 3$ )	
<b>Chai et al. [43]</b>	PubMed, Web of Science, APA PsycInfo, Google Scholar, and the China National Knowledge Infrastructure (CNKI) till March 2021	12; Meta-Analysis	Total sample size 34,276. Sample size ranges from 396 to 8079.	The studies recruited Chinese children and adolescents; all data collected through online measures; Studies were conducted in China	The pooled prevalence of mental problems was 28% (95% confidence interval, CI: 0.22–0.34) while for depression it was 22% (95% CI: 0.16–0.30) and for anxiety it was 25% (95% CI: 0.20–0.32)	The percentage of boy participants was found to be an associated moderator factor.

<b>Chawla et al. [44]</b>	PubMed, Cochrane, WHO global health research till March 2021	102; Systemic review	Sample size ranges from 61 to 10,000.	The children/ adolescents with psychological effects, communication issues, stress or sleep disturbances were recruited; Age of participants ranges from pre-schoolers to 24; Online survey and telephone interview were conducted; Most of the studies were cross sectional or retrospective design and recruited participants from China	Studies found mostly 10%-30% participants suffered from different level of depression and anxiety.	Being women and older aged found to be aggravating factor for anxiety and depression. Furthermore, factors like reduced physical activity, increased internet use, screen time and social media use, sedentary lifestyle were associated with severe anxiety. According to most of the studies, reduced sleep latency, increased sleep time as well as reduced sleep quality was correlated with the condition.
<b>Cunning et al. [45]</b>	PubMed, MEDLINE, APA PsycInfo, AMED, BNI, CINAHL, Embase, EMCare and HMIL till January 2021	6; Systemic review	Sample size ranges from 101-600 participants	The children and adolescent under age 21 having symptoms of OCD were included in this study; recruitment strategies were not specified, the studies were conducted in Denmark, Israel, the USA, Turkey, and Iran	The prevalence of OCD among adolescents ranged from 44.6% to 73% causing exaggerating symptoms	Being female seemed to be an exaggerating factor with slightly higher prevalence (72.1%) than men. Studies found some more associated factors like family history of ADHD, financial strain, or new unemployment, increased age, family accommodation.
<b>Sanja et al. (2021)</b>	Ovid MEDLINE, Embase, Global Health and APA PsycInfo till November 2020	27; Meta-Analysis	Total sample size 52,797.	The study recruited children and adolescents from 6 to 19 years old (54.12% were female); recruitment strategies	After pandemic, the prevalence of anxiety is 49.5% and severe anxiety is 20.5% (p <0.0001), for depression it is 16% among adolescents	Factors to increase anxiety and depression level are female gender (32.2%, M = 2.85, SE = .06), smartphone and internet addiction, low family income, perceived

				were not specified; conducted in Asia (20%), Europe (50%), America (20%), Africa (5%), Oceania (5%)		threats, higher grades at school, loneliness, social distancing, poor academic records, social stigma, friends or family conflict, emotion-focused coping.
<b>Jones et al.</b> [29]	EBSCOhost, MEDLINE, APA PsycInfo, APA Psych articles, Socindex, and CINAHL till January 2021	16; Systemic Review	Sample sizes varied from 102 to 9554.	13-17 years old adolescents along with mental illness were included in this study; 75% studies conducted through online measures, and 25% not clearly reported; recruited in China, UK, the USA, Canada, Germany, Japan, Denmark and Philippine	The pandemic increased the likelihood of anxiety (OR=2.29, 95% CI: 1.94-2.48 for low support, OR=3.18, 95% CI: 2.54-3.98 for moderate support) and 31% of the adolescents reported of depression. Also 16.3% adolescents suffered from psychological impairment and experienced higher rates of posttraumatic stress disorder (PTSD) (effect size beta = 0.16~0.27) and higher rates of anxiety (effect size beta = 0.32~0.47)	Factors which worsened the condition by social media (OR=1.8, 95% CI: 1.29-2.8) and internet addiction (OR=3.1 95% CI: 1.2-7.7). Female gender, social distancing, negative coping skills, adolescents with diagnosed OCD (44.6%) found to worsen the condition.
<b>Ma et al.</b> [34]	PubMed, APA PsycInfo, and Web of Science, and two Chinese academic databases: China National Knowledge Infrastructure (CNKI) and	23; Systemic review & Analysis.	Sample size vary from 46 to 9554; total sample size 57,927.	The studies focused on the children and adolescents age up to 18 years, with suicidal behavior, suicidal ideation and attempt; computer-based simulation studies with no human participants, most	The prevalence of depression ranged from 10-71% with a pooled prevalence of 29% (95%CI: 17%, 40%). The prevalence of anxiety ranged from 7%-55% with a pooled prevalence of 26% (95%CI: 16%, 35%). Prevalence of sleep	Females gender with age between 13-18 year were found to be higher risk factors for both anxiety s (29.1%, 95%CI: 17.1%, 41.1%, p<0.001) and depression (33.9%, 95%CI: 24.6%, 43.1%, p<0.001).

	Wanfang September 2020			study were from China 2 study from turkey	disorder was between 32%-56%, with a pooled prevalence of 44% (95%CI: 21%, 68%). The pooled prevalence of PTSD was 48% (95%CI: -0.25, 1.21, p=0.200)	
<b>Marchi et al. (2021)</b>	PubMed, APA PsycInfo, Web of Science, CINHAL, and Social Science Premium Collection till December 2020	59; Systematic scoping review		The study focused on child and adolescent below 18 years old; Study was identified in an iterative way; most of the study were took in China and United states	The prevalence of depression and anxiety among children and adolescents ranges from 2%-44%. And these symptoms seemed to reach peak around April- May with the decline by May-June, 2020	The protective factors found to be associated with the symptoms were strong resilience, positive emotion regulation, physical activity, parental self-efficacy, family functioning, social support. On the other hand, negative factors to impact were emotional reactivity, experiential avoidance, exposure to excessive information, internet use, closure of schools, social isolation, decline support from friend, family's crisis related hardships (such as job loss, income loss, caregiving burden, and illness).
<b>Panchal et al. (2021)</b>	Embase, Ovid, Global Health, APA	61; Systemic review	Sample size ranges from 15-7772.	The children and adolescent $\leq 19$ years old (female 49.7%)	The prevalence of anxiety and depression ranges from 1.85 to 49.5% and	Risk factors to increase anxiety and depression are lack of routine (p<0.001),

	PsycInfo, Web of Science, and pre-print databases till April 1, 2021			exposed to covid 19 lockdown were included in this study; most study involved participant through self-report; studies were conducted across 5 continents, Europe (n=35), Asia (n=22), Australia(n=1), North America (n=1), South America (n=1).	2.2% to 63.8% respectively, Irritability ranged from 16.7%-73.2% and anger from 30%-51.1%. Symptoms of ADHD ranges from 55.9% to 76.6% (p<0.001); sleeping disorder observed among 55.6% adolescents and 20% children. In addition, The prevalence of non-suicidal self-injury (OR=1.35, (p<0.001), suicide ideation (OR=1.32, p=0.008), suicide planning (OR=1.71, p<0.001), suicide attempts (OR=1.74, p<0.001)	female sex (p<0.001), adolescence (p=0.005), excessive COVID-19 information (p<0.05), media exposure (OR=2.4), previous psychiatric treatment (OR=4.4), increased social media usage s (OR=1.83, p=0.001), suffering from Autism Spectrum Disorder, high amount of COVID-19 cases in the area (OR=2.3, p<0.001), first-line job responsibilities related to COVID-19 (p<0.05). The protective was found to be all types of routine, family communication, social support, appropriate play, leisure and physical activity.
<b>Racine et al. [49]</b>	PsycInfo, Cochrane Central Register of Controlled Trials (CENTRAL), Embase, and MEDLINE till February 2021.	29; Meta-Analysis	Total sample size 80879; sample size ranges from 168 to 8079.	Participants up to 18 years old (mean age 13 years) were included in this study; recruitment strategies were not specified; Studies were performed in East Asia (n=16), Europe (n=4), North America (n=6), South America (n=2) and middle east (n=1)	The pooled prevalence rate for depression and anxiety are 25.2% (95% CI, 0.21-0.30) and 20.5% (95% CI, 0.17-0.24) respectively. The rate in European countries is higher (k = 4; rate = 0.34; 95% CI, 0.23-0.46; P = .01) in comparison to East Asian countries (k = 14; rate = 0.17; 95% CI, 0.13-0.21; P < .001)	For depression, the rate is increased with the increased months of year (b = 0.26; 95% CI, 0.06-0.46), increased child age (b = 0.08; 95% CI, 0.01-0.15), for female (b = 0.03; 95% CI, 0.01-0.05). Again, For anxiety, the rate is increased with the increased months of year (b = 0.27; 95% CI, 0.10-0.44), for female (b = 0.04; 95% CI, 0.01-0.07).

<b>Samji et al. (2021)</b>	MEDLINE, PsycInfo, PubMed, Embase, CINAHL, Web of Science, medRxiv, PsyArxiv, and Scopus till February 2021.	116; Systemic review	Total sample size 127,923.	The studies include youth up to 18 years of age, one third of the studied focused on neurodiverse conditions such as ASD, ADH and OCD; studies were performed in East Asia, north America, south Asia, Australia, West Asia, South America, Southeast Asia, Sub-Saharan Africa and North Africa	The prevalence of depression and anxiety ranged from 13%-64% and 8%-74% respectively. Moreover, prevalence of suicidal ideation ranged from 6%-37% and non-suicidal self-injury increased from 32% to 42%	Older age and female gender were positively associated the symptoms. Studies found some more aggravating factors like more screen time, internet use, social media, physical and social distancing, pandemic control measures, increased time at home, lack of daily routine, online learning of children, adverse childhood experiences (family abuse, neglect, and household dysfunction), covid-19 news, suffering from ASD, ADHD. However, some protective factors may improve the symptoms like engaging in hobbies, listening to music, praying, maintaining a routine, physical activity, Increasing knowledge and awareness of COVID-19 prevention and control measures.
<b>Sharma et al. [51]</b>	MEDLINE, Embase, and Web of Science	16; Meta-Analysis	Total sample size 21018; sample sizes ranges from 37 to 7736.	Studies includes children, adolescents and youth age ranging from 6 months to 29 years; all the studies recruited through online surveys; location of the studies were not specified.	The pooled prevalence of any sleep disturbance in children during the COVID-19 pandemic was 54% (95%CI: 50%-57%). The pooled prevalence of children with worsening of sleep quality and sleep duration were 27% (95%	The positive factors to improve sleep quality are harmonious family atmosphere, increased parent-children communication, good parenting practice, younger parent, parents in physical activity, parents

					CI: 12%-49%) and 16% (95%CI: 5%-40%) respectively. The pooled prevalence of children not meeting sleep duration recommendation was 49% (95%CI: 39%-58%)	discouraging screen time, parents encouraging children to sleep. The associated negative factors were poor maternal factors (poor sleep pattern, poor self-control, poor control of emotion, Maternal COVID-19 anxiety), decreased outdoor activities, female gender etc.
<b>Elharake et al. (2022)</b>	PubMed and Collabovid; January 2020- July 2021	5; Systematic review	Sample size ranged from 584 to 4342	Five studies included children and 16 studies included college students with varying ages; studies were from China, India, France, Jordan, Nigeria, Switzerland, and the US	The prevalence of depression (11.78% - 22.6%), anxiety (18.9% - 24.9%), PTST (14%) and stress (15.2%) were reported among children-based samples. Among the college students, 4.18% - 50.3% had depression, 7.7% - 79.66% had anxiety, 24.7% - 71% had stress, and 30.8% - 67.05% had PTSD symptoms.	Risk factors associated with adverse mental health outcomes in children included lower parents educational attainment and having a family member infected with COVID-19. Moreover, among college students, several factors such as family financial challenges, food insecurity (less than three meals per day), living in rural areas, closed on infected with COVID-19 were associated with poor mental health outcomes.
<b>Oliveira et al. [53]</b>	Embase, Epistemonikos, LILACS, APA PsycInfo, PubMed, Scopus, Web of Science, WHO Covid database, and Google	19; Systematic review	More than 35,543 children and adolescents were included	17 cross-sectional and two cohort studies were conducted using internet based approaches; nine countries were represented- most	The prevalence of fear (61.9%), helplessness (66.1%), worry (68.5%), emotional problems (27.4%), anxiety (17.6% - 43.7%), depression (6.3% - 71.5%), stress (7% - 25%), PTSD (85.5%), suicidal	Differences were observed between age groups (6-18 years versus 2-8 years) and responders (parents versus self-reported)

	Scholar; up to February 1, 2021			studies were from China	ideation (29.7% - 31.3%) behavioral problems (5.7% -13.9%) were reported across the included studies	
<b>Viner et al. (2022)</b>	PubMed, PsycInfo, Web of Science Social Citation Index, Australian Education Index, British Education Index, Education Resources Information Centre, WHO Global Research Database on COVID-19, MedRxiv, PsyArXiv, Research Square, and COVID-19 Living Evidence; up to September 1, 2020	36; Systematic review	Studies included 79,781 children and 18,028 parents	Five cohort studies, 9 pre-post studies, one modeling, and 21 cross-sectional studies were included; 13 studies were from LMICs (8 from China), 11 from UK, 4 from the US, 5 from Italy, and one study each from several countries	No significant increase in UK suicide rates; however, COVID-19 related problems were associated with 48% of the reported suicide deaths. The prevalence of anxiety (boys: 44%, girls: 53.3%, overall: 10%-21.8%), depression (boys: 21.9%, girls: 19.4%, overall: 17% - 28.6%), emotional difficulties (37%), suicidal ideation (17.5% in 16-18 years compared to 6% of pre-COVID estimates), conduct problems (43%), hyperactivity/inattention (41%), insomnia (23.2%), low life satisfaction (18% of 2001), low wellbeing (26.9% of 1201), child protection referral fell by 36% (156 per quarter in 2019 to 99 in 2020)56 to 39% (incidence rate ratio for 2020 compared with 2018/2019 was 0.61 [95%CI, 0.43-0.86])	School closure during lockdown was studied in the included studies highlighting the psychosocial impacts of such measures. Most studies found mental and emotional difficulties in this period, whereas a study with preexisting mental health problems in children found a significant reduction in emotional difficulties during lockdown. Child abuse reporting and referral services declines leading to low reporting. Increased social media and internet use, lack of healthy food consumption, and low-level of family education were reported across samples.

(95% CI: 50% - 57%), whereas the pooled prevalence of worsening sleep quality was 27% (95% CI: 12% - 49%) and the pooled estimate of abnormal sleep duration was 16% (95% CI: 5% - 40%).

### ***Trauma and stress-related disorders***

Four reviews examined the burden of posttraumatic stress disorders in different samples of children and adolescents [29, 34, 52, 53]. For example, Ma et al. [34] found a pooled prevalence of PTSD as 48%, whereas another review by Jones et al. [29] found that adolescents experienced a high effect size of PTSD (beta = 0.16 to 0.27). Another review by Elharake et al. [52] found that 30.8% to 67.05% of the study samples had PTSD symptoms among young people amidst the COVID-19 pandemic. Moreover, three reviews have reported a varying burden of psychological distress and stress-related disorders [41, 52, 53]. For example, Oliveira et al. (2022) reported that 7% to 25% of children and adolescents experienced stress-related disorders amidst this pandemic.

### ***Suicidal thoughts, ideations, and behavior***

Four reviews studied the burden of suicidal thoughts, ideations, and self-harm behavior among children and adolescents during this pandemic [48, 50, 53, 54]. For example, Panchal et al. (2021) reported a high prevalence of suicide ideation (OR = 1.32,  $p = 0.008$ ), suicide planning (OR = 1.71,  $p < 0.001$ ), suicide attempts (OR = 1.74,  $p < 0.001$ ), and non-suicidal self-harm (OR = 1.35,  $p < 0.001$ ). Another review by Samji et al. (2021) found that the prevalence of suicidal ideation ranged from 6% to 37%, and non-suicidal self-harm increased from 32% to 42% during this pandemic. Viner et al. (2022) found that the prevalence of suicidal ideation increased to 17.5% among adolescents aged 16-18 years compared to 6% of pre-COVID estimates.

### ***Attention-deficit/hyperactivity disorder***

Several empirical studies reported the burden of attention-deficit/hyperactivity disorder in children and adolescents during this pandemic that was synthesized in three reviews [41, 48, 54]. Viner et al. (2022) found that 41% of the participating children had hyperactivity/inattention, whereas Panchal et al. (2021) reported that the symptoms of ADHD ranged from 55.9% to 76.6% ( $p < 0.001$ ) among children and adolescents across multiple studies. Panda et al. (2021) found that 30.8% of the participating children had attention-deficit symptoms alongside other mental health problems.

### ***Other mental health problems***

Several other mental health problems were reported in multiple reviews. Cunning et al. [45] reported that the prevalence of OCD among adolescents ranged from 44.6% to 73% causing exaggerating symptoms. Moreover, emotional disorders were reported by three reviews that found 22% to 62.2% of children and adolescents experienced emotional difficulties during this pandemic [40, 53, 54]. A review by Nearchou et al. (2020) found that 2.39% to 37% of participants had somatic syndrome. Furthermore, psychological problems such as fear of COVID-19 [41, 53], boredom [41], irritability [41, 48], internalized and externalized problems [42], helplessness [53], conduct problems [54], and low life satisfaction [54] were reported across reviews. In addition, child protection referral fell by 36% (156 per quarter in 2019 to 99 in 2020) to 39% (incidence rate ratio for 2020 compared with 2018/2019 was 0.61 [95%CI, 0.43-0.86]), suggesting a potential burden of child abuse and related mental health problems [54]. Overall mental and psychosocial health problems deteriorated across all reviews that included children, adolescents, and their family caregivers in respective empirical studies.

### **Correlates of mental health problems in children and adolescents during COVID-19**

Multiple factors associated with mental health problems were identified across multiple reviews that affected the mental health of children and adolescents during this pandemic. The age of the children and adolescents was reported to be associated with mental health problems in different studies. For example, Chawla et al. (2021) found that older children and adolescents was an aggravating factor for anxiety and depression. Increased age was also reported as a risk factor by Cunning and Hodes [45], Samji et al. [50], and Ma et al. (2021). Specifically, Racine et al. [49] found that increased child age was associated with a higher burden of depression ( $b = 0.08$ , 95% CI: 0.01 – 0.15). Moreover, differences in mental health outcomes were observed between participants aged 2-8 years and those aged 6-18 years in a review by Oliveira et al. (2022).

Gender was associated with mental health outcomes as reported in several reviews. Chai et al. [43] reported the percentage of boys was a moderator of mental health outcomes across study samples. In contrast, several reviews reported girls had a higher burden of mental health problems during this pandemic. For example, female gender was associated with increased

anxiety and depression level (32.2%,  $M = 2.85$ ,  $SE = .06$ ) as reported by Sanja et al. [46]. Another review by Panchal et al. [48] found that the female gender was significantly associated with anxiety and depression ( $p < .001$ ). Racine et al. [49] reported that female participants had higher rates of depression ( $b = 0.03$ , 95% CI: 0.01 – 0.05).

Urban residence was found to be associated with mental health outcomes in a review by Panda et al. [41], whereas Elharake et al. [52] reported poor mental health outcomes in college students living in rural areas. Moreover, the time of the study appeared to affect the study outcomes. For example, Racine et al. [49] found that later months of the year was associated with higher burden of anxiety ( $b = 0.27$ , 95% CI: 0.10 – 0.44) and depression ( $b = 0.26$ , 95% CI: 0.06 – 0.46). Furthermore, poor academic records were associated with adverse mental health outcomes in one review [41], whereas another review found that higher academic grades were associated with anxiety and depression [46].

Several reviews synthesized evidence on behavioral and lifestyle-related factors associated with mental health problems. For example, Chawla et al. [44] found that a sedentary lifestyle, increased screen time, and lack of exercise were associated with depression and anxiety. Physical activity and regular exercise were associated with improved mental health across multiple reviews [41, 44, 47, 48, 50, 51]. Staying home for increased duration and lack of daily routine affected mental health in many children and adolescents, as reported by Panchal et al. (2021) and Samji et al. [50]. A Review by Viner et al. [54] found that a lack of healthy food consumption was associated with adverse mental health outcomes among children amidst this pandemic. Moreover, better emotional regulation and psychological resilience were associated with improved mental health outcomes. Furthermore, better knowledge about the pandemic was associated with better mental health among children and adolescents [47], whereas excessive information was negatively associated with mental health, as reported by Panchal et al. (2021).

Individual behaviors such as hobbies, praying, and listening to music were reported to be associated with positive mental health [50]. Internet and smartphone addiction appeared to be a common risk factor associated with mental health problems reported in several reviews [29, 41, 44, 46–48, 50, 51, 54]. Moreover, the past history of mental health problems was associated with adverse mental health outcomes, as reported in several reviews [29, 45, 48, 50, 54]. Severe

events related to COVID-19 affected mental health among many children and adolescents. For example, Panda et al. (2021) found that having a family member who was infected with COVID-19 was associated with adverse psychological outcomes. Another review by Panchal et al. [48] reported an increased number of COVID-19 cases within the same geographic area affected mental health (OR = 2.3,  $p < 0.001$ ) in children and adolescents.

Several family issues were reported to be associated with mental health outcomes during this pandemic. Nearchou et al. [40] found that belongingness and better relationships with closed ones were associated with improved mental health. Parental self-efficacy and family functioning were protective factors for mental health, as reported by Marchi et al. [47]. Moreover, adverse childhood and family experiences such as household dysfunction, family neglect, and child abuse were associated with poor mental health outcomes [50]. Moreover, the parents education and responsiveness were associated with the mental health of children and adolescents [51–54]. In addition, family financial crises affected the mental health of children and adolescents in many reviews. For example, unemployment and financial constraints affected families and the mental health of young people [45]. Elharake et al. [52] found that family financial challenges and food insecurity (less than three meals per day) affected mental health in children and college students.

School closure was associated with a wide range of mental health outcomes, as studied in a review by Viner et al. [54] and Marchi et al. [47]. Specifically, Viner et al. [54] found that child abuse referral reports declined during this pandemic due to school closure suggesting interrupted services in those contexts. As many institutions transitioned to online learning systems, Samji et al. [50] found such online learning to be associated with adverse mental health outcomes in children and adolescents.

Social factors such as social isolation and distancing, social stigma, and social support were associated with overall mental health outcomes in many reviews [29, 46–48, 50]. For example, social distancing was associated with adverse mental health outcomes in a review by Jones et al. [29]. Moreover, Samji et al. [50] reported that social distancing was associated with poor mental health outcomes. Both social stigma and social distancing were related to a higher burden of anxiety in a review by Sanja et al. Furthermore, Panchal et al. [48] and Marchi et al. [47] found social support to have a protective role on mental health outcomes in children and adolescents.

## Discussion

This umbrella review provides a systematic overview of global evidence on CAMH during the COVID-19 pandemic from existing systematic reviews and meta-analyses. To the best of our knowledge, this is one of the few umbrella reviews on mental health in COVID-19 [55, 56] and the first one emphasizing CAMH. The findings of this review may facilitate mental and child health policymaking and practice and inform further research strengthening the global knowledge base. In this review, we found a varying burden of several mental health problems such as anxiety, depression, sleep disorders, PTSD, suicidal behavior, psychological distress, and other psychosocial challenges among children and adolescents. Moreover, we identified several critical factors associated with mental health problems in this young population. Individual social demographic factors associated with mental health outcomes included older age of the children and adolescents, female gender, location of the residence, educational attainment, sedentary lifestyle, lack of routine activities, spending more time at home, social media addiction, Internet overuse, knowledge about pandemic or excessive information, emotional regulation, resilience, self-efficacy, physical activity and exercise behavior, recreational activities, spending time on hobbies, and history of mental health problems. Furthermore, several family-related factors associated with mental health were identified, which included belongingness and relationship with family members, parents' education, household income and economic hardship, adverse childhood and family experience, and challenging family relationships. In addition, community and social factors related to mental health included social distancing practices, social support, social stigma, closure of the schools, online learning activities, and COVID-19 related life events and preventive measures. These factors influenced mental health outcomes among children and adolescents in different contexts as identified in existing reviews and respective primary studies included within those reviews.

Most studies reported a high prevalence of CAMH problems across contexts suggesting a heavy psychosocial impact of this pandemic that is consistent with other reviews conducted in different populations [5, 55–58]. Socioecological perspectives may inform the complex epidemiology of these crises emerging at different levels. Children and adolescents experienced an unusual interruption in their everyday lives that included prolonged stay within home, altered relationship dynamics with their caregivers and family members, less-frequent

interactions with peers and relatives, lack of access to outdoor activities, changed nutritional and lifestyle behaviors and other changes at the household level [59, 60]. These problems compounded the effects of community and social changes associated with this pandemic, which included social distancing, closure of schools and institutions, and interruption of health and social services [54, 61, 62]. Lastly, universal challenges such as socioeconomic problems and health hazards of COVID-19 impacted children, adolescents, and their families that also affected mental health and wellbeing [5, 47, 63]. Taken together, a variety of psychosocial stressors emerged during this pandemic, whereas mental health promotion and protective services were mostly disrupted or unavailable. These factors affected biopsychosocial health and wellbeing across populations resulting in a high burden of CAMH problems that require clinical, public health, social, and policy considerations.

The overall evidence on the epidemiologic burden of CAMH and associated factors reflected high heterogeneity that can be attributed to several factors. First, earlier studies on psychosocial impacts were conducted in China [34, 45], whereas a majority of overall studies were from the US, UK, Canada, and several high-income countries [42, 49, 52, 54]. Most reviews informed a scarcity of evidence from the global south reflecting a critical knowledge gap in those contexts. Moreover, online measures of collecting data and non-standardized scales were widely used in cross-sectional studies reported across reviews [40, 44, 45, 47, 52], which may not provide representative estimates on CAMH impacts of COVID-19. Therefore, standardized measurements approaches should be used in representative samples and preferably in longitudinal and interventions studies that more accurately reflect the psychosocial impacts locally and globally. A brief overview of the recommendations for future research on CAMH is provided in Table 3-A.

The current evidence on CAMH problems amidst COVID-19 necessitates multipronged efforts to alleviate the immediate and future health and social consequences of this burden. The policymakers and practitioners need to evaluate the evidence-based guidelines and recommendations that may fit the mental health needs of children and adolescents in different contexts, as listed in Table 3-B. At the individual level, awareness of mental health and pandemic safety should be strengthened, addressing the fear of infection and health hazards [41, 43, 45, 46]. Moreover, positive psychology-based targeted interventions should be promoted alongside

public mental health interventions [40, 52, 64]. Furthermore, parents, teachers, and other caregivers should be engaged in mental health promotion [41, 43, 44, 47]. As many scholars suggest, those formal and informal caregivers may require psychoeducational and behavioral support to stay healthy and contribute to interpersonal and community health efforts [44, 47, 48]. Also, healthcare professionals and social workers should be empowered to offer psychosocial care services to the affected children and adolescents in diverse settings [46, 48]. Institutional care services should be made available and accessible to address the growing burden of mental health epidemics accompanying COVID-19 pandemic [40, 64]. Health systems and services must be strengthened to actualize these efforts at the societal level [45, 46, 64]. Lastly, a stronger policy commitment is needed to improve social determinants of mental health in children and adolescents that may enable them to overcome the current mental health problems and develop resilience to better psychosocial outcomes in the future [48, 49, 52].

## **Limitations**

This umbrella review has several limitations that must be addressed in future evidence syntheses. First, we included major databases but did not include preprint servers and other potential sources that may contain eligible reviews. Second, we identified multiple reviews with different methods and synthesized them narratively. A patient-level data synthesis from individual studies could have provided a more accurate and less overlapping view of the evidence landscape. Third, as systematic reviews and meta-analyses included primary studies published before conducting those reviews, a synthesis of the reviews is likely to miss insights from the most recent primary studies. Lastly, heterogeneity and publication biases were not assessed in this narrative umbrella study. Understanding those and addressing the same should be prioritized in future meta-research.

**Table 3: Evidence-based recommendations for future research, policymaking, and practice**

---

<b>A. Recommendations for future research</b>
<ul style="list-style-type: none"><li>• Conducting research using longitudinal/cohort designs to understand temporal changes and long-term effects on CAMH [5, 34, 44–46, 48, 52, 53]</li><li>• Using representative samples of children and adolescent populations rather than using small, non-representative samples, or sub-groups [40, 52]</li><li>• Examining the biopsychosocial determinants of CAMH during and after COVID-19 pandemic to explore specific risk and protective factors [40, 50, 52]</li><li>• Applying appropriate psychometric tools for CAMH problems with standardized cut-off values [34, 40, 43, 53]</li><li>• Exploring the impacts of COVID-19 on education, work, and living conditions and their associations with CAMH [42]</li><li>• Investigating the impacts of parents’ psychosocial health and family relationships on CAMH during COVID-19 [42]</li><li>• Studying the psychosocial correlates of pandemic-related events and their impacts on CAMH [44, 45]</li><li>• Investigating the persistence of psychosocial symptomology or development of mental disorders in the future [46, 48, 50]</li><li>• Strengthening research capacities of the researchers and institutions on CAMH research in the context of COVID-19 [46]</li><li>• Emphasizing younger children’s mental health research- an area with little evidence but high significance for long-term impacts [34]</li><li>• Examining psychiatric comorbidities alongside identified physical and mental health problems [34]</li><li>• Examining the role of timing and context of studies (e.g., studied undertaken during lockdown vs. after returning to school) on CAMH [50]</li><li>• Exploring pre-pandemic and existing mental health gaps to develop stronger mental health systems in the future [50, 52]</li><li>• Conducting implementation research on mental health services for children and adolescents emphasizing trends, barriers, and facilitators of mental health utilization in different contexts [5, 52, 64]</li></ul>

---

---

## B. Recommendations for policymaking and practice

---

- Improving social determinants of CAMH as a local and global health priority [5, 49, 52, 65–67]
  - Integrating clinical and psychosocial interventions in public health and child health policies and programs with special focus on targeted mental health components whenever possible [5, 40, 64, 68]
  - Establishing and strengthening mental health safety nets for children and adolescents [52]
  - Psychoeducational interventions on pandemic preparedness and mental health promotion for children and adolescents engaging parents, family members, teachers, and other stakeholders [41, 43, 45, 46, 48]
  - Designing and implementing physical activity-based interventions that may improve psychosocial outcomes alongside overall health and wellbeing [29, 41, 44, 47, 48, 53]
  - Addressing food insecurity and nutritional problems that may influence mental health outcomes [41, 52, 69]
  - Providing extensive and integrative care for children with special needs ensuring mental wellbeing [48, 70]
  - Providing social support, social connectedness, peer-networking, and social interventions for improving prosocial coping mechanisms among children and adolescents with poor mental health [41, 48, 50, 63]
  - Educating and engaging parents on positive parenting and mental health promotion [41, 43, 47]
  - Providing family-based interventions for holistic mental health development of children, adolescents, and their family caregivers [41, 43, 44, 47, 48]
  - Engaging academic institutions and providers for delivering nonpharmacological mental health education and services [41]
  - Developing public health messages and programs emphasizing positive psychology and better mental health [45, 48–50]
  - Leveraging evidence-based digital approaches for delivering online education ensuring optimal mental health of the young learners [41]
  - Organizing safe and planned school opening and resuming in-person/routine activities ensuring public health compliance for better psychosocial outcomes [41, 49]
-

- 
- Funding mental health services for children and adolescents ensuring wider access and sustainability [48]
  - Promoting problem solving skills and empowerment programs among children and adolescents [41, 53]
  - Facilitating recreational activities such as reading books, story reading/telling, listening to music, spending time on hobbies [41, 47]
  - Restoring mental health services and increasing access to mental health care [45, 46, 64]
  - Adopting digital mental health resources and services for improving mental health access and utilization in children and adolescents [29, 41, 48–50, 53]
  - Providing special attention to early childhood mental health development interventions [52]
  - Restoring and improving child protection and referral services [54, 71, 72]
  - Empowering health and other care providers for addressing CAMH problems through capacity building exercises, developing evidence-based practice guidelines, and mobilizing care resources [5, 46, 48]
  - Examining and implementing evidence-based guidance on screen time, social media and internet use that may protect mental health and wellbeing [43, 44, 47]
  - Strengthening public mental health services for rural and other marginalized population groups [5, 52]
- 

## Conclusions

Psychosocial stressors associated with the COVID-19 pandemic have resulted in a wide range of mental health problems in children and adolescents. The current evidence suggests a high burden of anxiety, depression, psychological distress, PTSD, sleep disorders, suicidal behavior, addiction disorders, and other psychosocial problems that require evidence-based measures to address the same. Also, many of these problems may persist and develop further mental health crises that should be addressed by adopting multifaceted psychiatric, psychological, and public health efforts. Transdisciplinary mental health research, policymaking, and practice should be emphasized to mitigate the mental health impacts of this pandemic and ensure optimal mental, physical, and social health of children and adolescents locally and globally.

**Funding information:** No funding was received at any stage of conceptualizing or conducting this umbrella review.

## References

- [1] Fauci AS, Lane HC, Redfield RR. Covid-19 — Navigating the Uncharted. *N Engl J Med* 2020; 382: 1268–1269.
- [2] Bashir MF, Ma B, Shahzad L. A brief review of socio-economic and environmental impact of Covid-19. *Air Qual Atmos Heal* 2020; 13: 1403–1409.
- [3] Bialek S, Bowen V, Chow N, et al. Geographic Differences in COVID-19 Cases, Deaths, and Incidence — United States, February 12–April 7, 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69: 465–471.
- [4] Robinson E, Sutin AR, Daly M, et al. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J Affect Disord* 2022; 296: 567–576.
- [5] Hossain M, Tasnim S, Sultana A, et al. Epidemiology of mental health problems in COVID-19: a review [version 1; peer review: 2 approved] report report. Epub ahead of print 2020. DOI: 10.12688/f1000research.24457.1.
- [6] Ren X, Huang W, Pan H, et al. Mental Health During the Covid-19 Outbreak in China: a Meta-Analysis. *Psychiatr Q* 2020; 91: 1033–1045.
- [7] Şimşir Z, Koç H, Seki T, et al. The relationship between fear of COVID-19 and mental health problems: A meta-analysis. <https://doi.org/10.1080/0748118720211889097> 2021; 46: 515–523.
- [8] Wu T, Jia X, Shi H, et al. Prevalence of mental health problems during the COVID-19 pandemic: A systematic review and meta-analysis. *J Affect Disord* 2021; 281: 91–98.
- [9] Hessami K, Romanelli C, Chiurazzi M, et al. COVID-19 pandemic and maternal mental health: a systematic review and meta-analysis. <https://doi.org/10.1080/1476705820201843155>. Epub ahead of print 2020. DOI: 10.1080/14767058.2020.1843155.
- [10] Hossain MM, Sultana A, Purohit N. Mental health outcomes of quarantine and isolation for infection prevention: a systematic umbrella review of the global evidence. *Epidemiol Health*; 42. Epub ahead of print 2020. DOI: 10.4178/EPIH.E2020038.
- [11] Bassuk EL, Richard MK, Tsertsvadze A. The Prevalence of Mental Illness in Homeless Children: A Systematic Review and Meta-Analysis. *J Am Acad Child Adolesc Psychiatry* 2015; 54: 86-96.e2.
- [12] Lund C, Brooke-Sumner C, Baingana F, et al. Social determinants of mental disorders and the Sustainable Development Goals: a systematic review of reviews. *The Lancet Psychiatry* 2018; 5: 357–369.

- [13] Yoshikawa H, Aber JL, Beardslee WR. The effects of poverty on the mental, emotional, and behavioral health of children and youth. *Am Psychol* 2012; 67: 272–284.
- [14] Silva SA, Silva SU, Ronca DB, et al. Common mental disorders prevalence in adolescents: A systematic review and meta-analyses. *PLoS One* 2020; 15: e0232007.
- [15] Polanczyk G V., Salum GA, Sugaya LS, et al. Annual Research Review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *J Child Psychol Psychiatry* 2015; 56: 345–365.
- [16] Bitsko RH, Holbrook JR, Ghandour RM, et al. Epidemiology and impact of health care provider-diagnosed anxiety and depression among US children. *J Dev Behav Pediatr* 2018; 39: 395–403.
- [17] Merikangas KR, He JP, Burstein M, et al. Lifetime Prevalence of Mental Disorders in U.S. Adolescents: Results from the National Comorbidity Survey Replication–Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry* 2010; 49: 980–989.
- [18] Spady DW, Schopflocher DP, Svenson LW, et al. Prevalence of Mental Disorders in Children Living in Alberta, Canada, as Determined From Physician Billing Data. *Arch Pediatr Adolesc Med* 2001; 155: 1153–1159.
- [19] Gadermann AM, Gagné Petteni M, Janus M, et al. Prevalence of Mental Health Disorders Among Immigrant, Refugee, and Nonimmigrant Children and Youth in British Columbia, Canada. *JAMA Netw Open* 2022; 5: e2144934–e2144934.
- [20] Deighton J, Lereya ST, Casey P, et al. Prevalence of mental health problems in schools: poverty and other risk factors among 28 000 adolescents in England. *Br J Psychiatry* 2019; 215: 565–567.
- [21] Kovess-Masfety V, Husky MM, Keyes K, et al. Comparing the prevalence of mental health problems in children 6–11 across Europe. *Soc Psychiatry Psychiatr Epidemiol* 2016; 51: 1093–1103.
- [22] Sawyer MG, Reece CE, Sawyer ACP, et al. Has the Prevalence of Child and Adolescent Mental Disorders in Australia Changed Between 1998 and 2013 to 2014? *J Am Acad Child Adolesc Psychiatry* 2018; 57: 343–350.e5.
- [23] Kamieniecki GW. Prevalence of psychological distress and psychiatric disorders among homeless youth in Australia: A comparative review. *Aust N Z J Psychiatry* 2001; 35: 352–358.
- [24] Patel V. Mental health in low- and middle-income countries. *Br Med Bull* 2007; 81–82: 81–96.
- [25] Patel V, Flisher AJ, Nikapota A, et al. Promoting child and adolescent mental health in low and middle income countries. *J Child Psychol Psychiatry* 2008; 49: 313–334.
- [26] Hossain M, Purohit N. Improving child and adolescent mental health in India: Status, services, policies, and way forward. *Indian J Psychiatry* 2019; 61: 415.
- [27] Rajmil L, Hjern A, Boran P, et al. Original research: Impact of lockdown and school

- closure on children's health and well-being during the first wave of COVID-19: a narrative review. *BMJ Paediatr Open*; 5. Epub ahead of print 25 May 2021. DOI: 10.1136/BMJPO-2021-001043.
- [28] Larsen L, Helland MS, Holt T. The impact of school closure and social isolation on children in vulnerable families during COVID-19: a focus on children's reactions. *Eur Child Adolesc Psychiatry* 2021; 1: 1–11.
- [29] Jones EAK, Mitra AK, Bhuiyan AR. Impact of COVID-19 on Mental Health in Adolescents: A Systematic Review. *Int J Environ Res Public Heal* 2021, Vol 18, Page 2470 2021; 18: 2470.
- [30] Yeasmin S, Banik R, Hossain S, et al. Impact of COVID-19 pandemic on the mental health of children in Bangladesh: A cross-sectional study. *Child Youth Serv Rev* 2020; 117: 105277.
- [31] Almis H, Han Almis B, Bucak IH. Mental health of children of health workers during the COVID-19 pandemic: A cross-sectional study. *Clin Child Psychol Psychiatry* 2022; 27: 104–111.
- [32] O'Sullivan K, Clark S, McGrane A, et al. A Qualitative Study of Child and Adolescent Mental Health during the COVID-19 Pandemic in Ireland. *Int J Environ Res Public Heal* 2021, Vol 18, Page 1062 2021; 18: 1062.
- [33] Tang S, Xiang M, Cheung T, et al. Mental health and its correlates among children and adolescents during COVID-19 school closure: The importance of parent-child discussion. *J Affect Disord* 2021; 279: 353–360.
- [34] Ma L, Mazidi M, Li K, et al. Prevalence of mental health problems among children and adolescents during the COVID-19 pandemic: A systematic review and meta-analysis. *J Affect Disord* 2021; 293: 78–89.
- [35] Aromataris E, Fernandez R, Godfrey CM, et al. Summarizing systematic reviews: methodological development, conduct and reporting of an umbrella review approach. *Int J Evid Based Healthc* 2015; 13: 132–140.
- [36] Hossain MM, Sultana A, Tasnim S, et al. Prevalence of mental disorders among people who are homeless: An umbrella review. *Int J Soc Psychiatry* 2020; 66: 528–541.
- [37] Dang Y, Hou Y. The prognostic value of late gadolinium enhancement in heart diseases: an umbrella review of meta-analyses of observational studies. *Eur Radiol* 2021; 31: 4528–4537.
- [38] Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*; 372. Epub ahead of print 29 March 2021. DOI: 10.1136/BMJ.N71.
- [39] Aromataris E, Fernandez R, Godfrey C, et al. JBI Manual for Evidence Synthesis - JBI Global Wiki. *JBI*, <https://jbi-global-wiki.refined.site/space/MANUAL/3283910768/Chapter+10%3A+Umbrella+reviews> (2020, accessed 19 March 2022).
- [40] Nearchou F, Hennessy E, Flinn C, et al. Exploring the impact of covid-19 on mental health outcomes in children and adolescents: A systematic review. *Int J Environ Res Public*

*Health* 2020; 17: 1–19.

- [41] Panda PK, Gupta J, Chowdhury SR, et al. Psychological and Behavioral Impact of Lockdown and Quarantine Measures for COVID-19 Pandemic on Children, Adolescents and Caregivers: A Systematic Review and Meta-Analysis. *J Trop Pediatr*; 67. Epub ahead of print 2021. DOI: 10.1093/tropej/fmaa122.
- [42] Bussières EL, Malboeuf-Hurtubise C, Meilleur A, et al. Consequences of the COVID-19 Pandemic on Children’s Mental Health: A Meta-Analysis. *Front Psychiatry*; 12. Epub ahead of print 2021. DOI: 10.3389/fpsy.2021.691659.
- [43] Chai J, Xu H, An N, et al. The Prevalence of Mental Problems for Chinese Children and Adolescents During COVID-19 in China: A Systematic Review and Meta-Analysis. *Front Pediatr*; 9. Epub ahead of print 2021. DOI: 10.3389/fped.2021.661796.
- [44] Chawla N, Tom A, Sen MS, et al. Psychological Impact of COVID-19 on Children and Adolescents: A Systematic Review. *Indian J Psychol Med* 2021; 43: 294–299.
- [45] Cunning C, Hodes M. The COVID-19 pandemic and obsessive–compulsive disorder in young people: Systematic review. *Clin Child Psychol Psychiatry* 2022; 27: 18–34.
- [46] Sanja Đ, Ileana CG, Vesna D, et al. Anxiety and depressive symptomatology among children and adolescents exposed to the COVID-19 pandemic - meta-analysis. *Vojnosanit Pregl* 2021; 1–21.
- [47] Marchi J, Johansson N, Sarkadi A, et al. The Impact of the COVID-19 Pandemic and Societal Infection Control Measures on Children and Adolescents’ Mental Health: A Scoping Review. *Front Psychiatry*; 12. Epub ahead of print 2021. DOI: 10.3389/fpsy.2021.711791.
- [48] Panchal U, Salazar de Pablo G, Franco M, et al. The impact of COVID-19 lockdown on child and adolescent mental health: systematic review. *Eur Child Adolesc Psychiatry*. Epub ahead of print 2021. DOI: 10.1007/s00787-021-01856-w.
- [49] Racine N, McArthur BA, Cooke JE, et al. Global Prevalence of Depressive and Anxiety Symptoms in Children and Adolescents during COVID-19: A Meta-analysis. *JAMA Pediatr* 2021; 175: 1142–1150.
- [50] Samji H, Wu J, Ladak A, et al. Review: Mental health impacts of the COVID-19 pandemic on children and youth – a systematic review. *Child Adolesc Ment Health*. Epub ahead of print 2021. DOI: 10.1111/camh.12501.
- [51] Sharma M, Aggarwal S, Madaan P, et al. Impact of COVID-19 pandemic on sleep in children and adolescents: a systematic review and meta-analysis. *Sleep Med* 2021; 84: 259–267.
- [52] Elharake JA, Akbar F, Malik AA, et al. Mental Health Impact of COVID-19 among Children and College Students: A Systematic Review. *Child Psychiatry Hum Dev*. Epub ahead of print 2022. DOI: 10.1007/s10578-021-01297-1.
- [53] Oliveira JMD de, Butini L, Pauletto P, et al. Mental health effects prevalence in children and adolescents during the COVID-19 pandemic: A systematic review. *Worldviews*

*Evidence-Based Nurs* 2022; 1–8.

- [54] Viner R, Russell S, Saulle R, et al. School Closures during Social Lockdown and Mental Health, Health Behaviors, and Well-being among Children and Adolescents during the First COVID-19 Wave: A Systematic Review. *JAMA Pediatr* 2022; 1–10.
- [55] Sahebi A, Nejati-Zarnaqi B, Moayedi S, et al. The prevalence of anxiety and depression among healthcare workers during the COVID-19 pandemic: An umbrella review of meta-analyses. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2021; 107: 110247.
- [56] Fernandez R, Sikhosana N, Green H, et al. Anxiety and depression among healthcare workers during the COVID-19 pandemic: a systematic umbrella review of the global evidence. *BMJ Open* 2021; 11: e054528.
- [57] Bourmistrova NW, Solomon T, Braude P, et al. Long-term effects of COVID-19 on mental health: A systematic review. *J Affect Disord* 2022; 299: 118–125.
- [58] Li K, Tang F, Hanna S, et al. The Impact of COVID-19 Concerns on the Mental Health of Older Adults: A Rapid Review. *Innov Aging* 2021; 5: 727–728.
- [59] Settersten RA, Bernardi L, Härkönen J, et al. Understanding the effects of Covid-19 through a life course lens. *Curr Perspect Aging Life Cycle* 2020; 45: 100360.
- [60] de Figueiredo CS, Sandre PC, Portugal LCL, et al. COVID-19 pandemic impact on children and adolescents' mental health: Biological, environmental, and social factors. *Prog Neuropsychopharmacol Biol Psychiatry* 2021; 106: 110171.
- [61] Barach P, Fisher SD, Adams MJ, et al. Disruption of healthcare: Will the COVID pandemic worsen non-COVID outcomes and disease outbreaks? *Prog Pediatr Cardiol* 2020; 59: 101254.
- [62] Nguyen PH, Kachwaha S, Pant A, et al. COVID-19 Disrupted Provision and Utilization of Health and Nutrition Services in Uttar Pradesh, India: Insights from Service Providers, Household Phone Surveys, and Administrative Data. *J Nutr* 2021; 151: 2305.
- [63] Preston AJ, Rew L. Connectedness, Self-Esteem, and Prosocial Behaviors Protect Adolescent Mental Health Following Social Isolation: A Systematic Review. *Issues Ment Health Nurs* 2022; 43: 32–41.
- [64] Mohd W, Mohd W, Wan A, et al. Registered psychiatric service use , self - harm and suicides of children and young people aged 0 – 24 before and during the COVID - 19 pandemic : a systematic review. 2022; 3: 1–14.
- [65] Abrams EM, Szeffler SJ. COVID-19 and the impact of social determinants of health. *Lancet Respir Med* 2020; 8: 659.
- [66] Ashcroft R, Lam S, Kourgiantakis T, et al. Protocol: Preparing social workers to address health inequities emerging during the COVID-19 pandemic by building capacity for health policy: a scoping review protocol. *BMJ Open* 2021; 11: 53959.
- [67] Smitherman LC, Golden WC, Walton JR. Health Disparities and Their Effects on Children and Their Caregivers During the Coronavirus Disease 2019 Pandemic. *Pediatr Clin North*

*Am* 2021; 68: 1133.

- [68] Boldt K, Coenen M, Movsisyan A, et al. Interventions to Ameliorate the Psychosocial Effects of the COVID-19 Pandemic on Children-A Systematic Review. *Int J Environ Res Public Health* 2021; 18: 1–31.
- [69] Zemrani B, Gehri M, Masserey E, et al. A hidden side of the COVID-19 pandemic in children: the double burden of undernutrition and overnutrition. *Int J Equity Health*; 20. Epub ahead of print 1 December 2021. DOI: 10.1186/S12939-021-01390-W.
- [70] Asbury K, Fox L, Deniz E, et al. How is COVID-19 Affecting the Mental Health of Children with Special Educational Needs and Disabilities and Their Families? *J Autism Dev Disord* 2021; 51: 1772–1780.
- [71] Renov V, Risser L, Berger R, et al. The impact of the COVID-19 pandemic on child protective services caseworkers and administrators. *Child Abuse Negl*. Epub ahead of print 2021. DOI: 10.1016/J.CHIABU.2021.105431.
- [72] Katz C, Fallon B. Protecting children from maltreatment during COVID-19: Struggling to see children and their families through the lockdowns. *Child Abuse Negl* 2021; 116: 105084.

**Supplementary Material 1: Critical appraisal of the methodological quality of the included reviews**

Authors/ Sources	Is the review question clearly and explicitly stated?	Were the inclusion criteria appropriate for the review question?	Was the search strategy appropriate?	Were the sources and resources used to search for studies adequate?	Were the criteria for appraising studies appropriate?	Was critical appraisal conducted by two or more reviewers independently?	Were the methods used to combine studies appropriate?	Was the likelihood of publication bias assessed?	Were recommendations for policy and/or practice supported by the reported data?	Were the specific directives for new research appropriate?	Overall score and quality rating
Nearchou et al., 2020	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	CD	Yes	8 (High)
Panda et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	CD	9 (High)
Panda et al., 2021	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	9 (High)
Bussi�eres et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10 (High)
Chai et al., 2021	Yes	Yes	Yes	Yes	Yes	CD	Yes	Yes	Yes	Yes	9 (High)
Chawla et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	9 (High)
Cunning and Hodes, 2022	Yes	Yes	CD	CD	Yes	CD	CD	No	Yes	Yes	5 (Medium)
Sanja et al., 2021	Yes	Yes	Yes	Yes	Yes	CD	CD	No	Yes	Yes	7 (Medium)
Jones et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	CD	No	Yes	Yes	8 (High)
Ma et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10 (High)
Marchi et al., 2021	Yes	Yes	Yes	Yes	NA	No	Yes	NA	Yes	Yes	7 (Medium)
Panchal et al., 2021	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	9 (High)

Racine et al., 2021	Yes	Yes	Yes	Yes	Yes	CD	Yes	Yes	Yes	Yes	9 (High)
Samji et al., 2021	Yes	Yes	Yes	Yes	Yes	CD	CD	No	Yes	Yes	8 (High)
Sharma et al., 2021	Yes	No	Yes	Yes	10 (High)						
Elharake et al., 2022	Yes	Yes	Yes	Yes	No	No	Yes	No	Yes	Yes	7 (Medium)
Oliveira et al., 2022	Yes	No	Yes	Yes	9 (High)						
Viner et al., 2022	Yes	No	Yes	Yes	9 (High)						

(Abbreviations: CD= could not determine; NA= not applicable)