Pronounced Regional Variation in Esketamine and Ketamine Prescribing to US Medicaid Patients

Alexia G. Aguilar[1], Burke A. Beauregard[1], Christopher P. Conroy[1], Yashoda T. Khatiwoda[1], Shantia M. E. Horsford[1], Stephanie D. Nichols[2], Brian J. Piper[1,3*]

[1]Department of Medical Education, Geisinger Commonwealth School of Medicine, Scranton, PA, United States of America

[2]Department of Pharmacy Practice, University of New England, Portland, ME, United States of America

[3]Center for Pharmacy Innovation & Outcomes, Forty Fort, PA, United States of America

Disclosures: BJP was part of an osteoarthritis research team supported by Pfizer/Eli Lilly from 2019 to 2021.

*psy391@gmail.com; bpiper@som.geisinger.edu

Abstract: 204 words

Manuscript: 1,672 words

Figures: 2, **Tables**: 0

Supplementary Materials: 3 Appendices, 4 Figures

Abstract

Background: Ketamine, and its isomer esketamine are useful for treatment resistant depression (depression not responsive to traditional antidepressants). Compared to other antidepressants, ketamine and esketamine lack a therapeutic delay and do not increase risk of suicidality in adolescents. Esketamine gained FDA approval in March of 2019.

Objective: This study aimed to geographically characterize ketamine and esketamine prescriptions in the United States among Medicaid patients.

Methods: Ketamine and esketamine prescription rate data per state per quarter were obtained from the Medicaid State Drug Utilization Database for 2019-2020. Total prescription rates per year for ketamine and esketamine were obtained for 2009-2020.

Results: Between 2009-2020, ketamine prescribing rates peaked in 2013 followed by a general decline. For ketamine and esketamine prescriptions in 2019, Montana and Indiana showed significantly higher prescription rates, respectively, relative to the national average. A total of 21 states prescribed neither ketamine or esketamine in 2019. Since its approval, esketamine prescriptions have surpassed those of ketamine. There was a 121.3% increase in esketamine prescriptions from 2019 to 2020. Ten states prescribed neither ketamine or esketamine in 2020.

Conclusion: Despite the effectiveness of ketamine and esketamine for treatment resistant depression, their use among Medicaid patients was limited and variable in many areas of the US.

Introduction

Suicide rates continued to increase in the United States marking the presence of a suicide epidemic. Suicide was the second leading cause of death for people aged 10-34 years old in 2019 [1]. Suicide ranks as the 10th leading cause of death with 4.8% of adults having thoughts of suicide and 0.6% (1,969,200) reporting having had attempted suicide in the past year [1]. Since the emergence of COVID-19, serious suicide attempts have increased in adolescent girls but not in boys [2]. The rate of serious suicide attempt ED visits increased by 50.6% in girls in 2021 compared with the same time frame in 2019 [2]. Among boys, the increase in serious suicide attempts was only 3.7% [2]. Unfortunately, traditional antidepressants are often ineffective in preventing suicide linked to depression for two reasons. First, treatment resistant depression, or failure to respond to traditional antidepressants is present in up to two-thirds of patients and second, a time lag of weeks to months delays the onset of therapeutic effects [3]. Second, traditional antidepressants have been found to counterintuitively increase the risk of suicidality in the first month of therapy in patients who are under 25. This effect was not found in adults and a protective effect was observed in geriatric adults, emphasizing age related differences in suicidality and response to medications [4].

Ketamine, a well-known anesthetic agent, has gained attention as an antidepressant using a novel mechanism of action and is very useful for reducing suicidal ideation [5]. Its absence of therapeutic delay and effectiveness for treatment-resistant depression are well-suited for patients at risk of suicide [3]. By being effective in both major unipolar and bipolar depressive episodes, unlike other antidepressants, it is

useful for depression even when the diagnostic picture is unclear. In March of 2019, the FDA issued a conditional approval of esketamine, a stereoisomer of ketamine, as nasal spray which is administered under direct supervision of a medical provider. It is approved for use in patients with treatment resistant depression or with depression and acute suicidality. This is a landmark development as existing antidepressants have a black-box warning for young-adults for increasing suicide.

The objective of this study was to examine patterns in ketamine and esketamine prescription rates throughout the United States among Medicaid patients. Determining geographic variability in prescription could be useful for correlating use of ketamine and esketamine to rates of suicide and beginning to examine corrective efforts on a population level. Given the recency of esketamine to the market and its current high visibility in the field, these trends may be particularly useful.

METHODS

Procedures

Ketamine and esketamine prescription rates were obtained from the Medicaid State Drug Utilization database [6]. Medicaid is a joint federal and state program that provides coverage for 75 million people, approximately 21% of the United States population. All states provide coverage for outpatient prescription drugs [6]. We evaluated the Medicaid State Drug Utilization database for total outpatient ketamine and esketamine prescriptions per state per quarter during 2019-2020 [6]. Prescription rates were reported per 1,000,000 Medicaid enrollees. Formulations were categorized by National Drug Codes (Supplemental Appendix A & B) as ketamine versus esketamine and by route of administration (nasal vs injection). Procedures were approved as exempt by the Geisinger and the University of New England IRBs.

Data analysis

The following analyses were completed: 1) esketamine percent change of prescription rate from 2012 to 2013, and 2) a 95% confidence interval (mean ± 1.96*SD) with states outside this range interpreted as significantly different from the mean. We analyzed the data and constructed figures using SAS, JMP and GraphPad Prism.

RESULTS

Figure 1. National prescription rate of ketamine and esketamine to Medicaid patients from 2009 to 2020.

Figure 1 shows that prescribing rates for ketamine from 2009 to 2020 were in elevated in 2013 with a 91.2% increase compared to 2012. This was followed by a gradual decrease from 2014 to 2018. With the approval of esketamine, the total (i.e. ketamine and esketamine) prescriptions tripled from 2018 (106/million enrollees) to 2020 (292/million).

There were 27,238 prescriptions nationally for ketamine and esketamine in 2019. Of these, 12,668 prescriptions (46.5%) were for esketamine nasal spray and 14,570 (53.5%) for ketamine injections.

Pronounced geographical differences were also identified. For ketamine, Montana (967.3) showed a statistically significant ($p < .05$) higher rate of prescription relative to the national average (Figure 2A). There was a 116-fold higher rate of prescriptions in Montana relative to the lowest state (MA = 8.3). States in the southern US from New Mexico to South Carolina had no ketamine prescriptions (Supplemental Figure 1). For esketamine, Indiana (424.6) showed a significantly higher rate of prescription which was also 61-fold higher than the lowest state (Supplemental Figure 2). Twenty states (AK, AL, AR, DE, GA, HI, IA, ID, IL, KS, ME, MS, ND, NH, OK, RI, SC, VT, WI, WY) and Washington DC prescribed neither ketamine nor esketamine.

In 2020, Montana (1,705.6) was significantly higher than the national average and 245-fold higher than the lowest state (Michigan = 7.0, Figure 2B) for ketamine. States that were located at or below the 35° latitude (Arizona proceeding East to South Carolina) had the lowest prescribing (Supplemental Figure 3). Three-states, North Dakota, North Carolina, and Montana were significantly elevated for esketamine in 2020. There was a 104-fold difference between the highest (North Dakota = 1,423.0) and lowest (Alaska = 13.7) states (Figure 2D, Supplemental Figure 4). Ten states (NY, OK, RI, SC, TX, VI, VT, WV, WI, & WY) prescribed neither ketamine nor esketamine.

Figure 2. Ketamine (A, C) and esketamine (B, D) prescription rate per state to Medicaid patients. States not shown had a value of 0. * *p* < .05 vs the state average.

DISCUSSION

There are two key findings from this report. First was the increasing, but still quite low, quarterly increase in esketamine relative to ketamine. Esketamine prescriptions in 2019 were less than half (44.2%) those of ketamine in 2019, likely due to its recent FDA-approval and introduction to the market. Second, the state-level differences in esketamine prescription rates were notable. For example, one state, Indiana prescribed 25% of the national total in 2019. At the beginning of 2019, quarterly esketamine was prescribed at a rate close to zero, but as the year progressed, a subset of states quickly adapted to use of esketamine. This was perhaps due to the large marketing budget in those states, increasing clinical evidence and wider distribution of trial results, and the rapid establishment of protocols and procedures that meet the FDA Risk Evaluation and Mitigation Strategy program requirements. The pronounced differences in esketamine prescribing at a state-level could largely be attributed to marketing rather than long-standing preset preferences in patients and physicians since this is a new drug to the market. It will likely be a fruitful endeavor to determine how use of these agents change in future years.

The identification of the rapid therapeutic and anti-suicidality effects of ketamine is potentially one of the most important developments in psychopharmacology in the past two decades. However, we found that the use of esketamine and ketamine among Medicaid patients was both modest, relative to the prevalence of depression and treatment resistant depression, and extremely variable [11]. This variability is shown by the absence of ketamine/esketamine prescriptions in two-fifths of US states in an outpatient setting. The role of ketamine in the US as an antidepressant is continues to evolve [12,13].

Regardless of the underlying reasons, it is interesting to note that the rates of ketamine prescribing are nowhere near as high as one might expect from an antidepressant in a class on its own which overcomes the therapeutic lag of selective serotonin reuptake inhibitors and serotonin norepinephrine reuptake inhibitors. Among the top two-hundred most prescribed medications known for their antidepressant

effects written for Medicaid patients in 2019 [6], fluoxetine was the most prescribed. The ratio of fluoxetine to ketamine prescription was nearly an eight-hundred-fold difference [6]. While it is not yet conclusive that a "miracle drug" has been found to treat depression, the rates of prescribing suggest peripheral challenges like the logistics of proper administration and/or social inertia on the use of this psychoactive and controlled substance that may be causing delays. The COVID-19 pandemic likely also contributed to the limited utilization of these agents. However, there have also been fourteen ketamine/esketamine manuscripts which were subsequently retracted (e.g. use of ketamine in the ER) which may also be contributing to some ambiguity [14, Supplemental Appendix C]. There are also serious concerns with misuse, the development of tolerance [15] and tremendous costs of esketamine and monitoring relative to more traditional treatments [16-18].

Depression leads to significant long term negative effects for both social and personal health indicators, leading to a decreased effectiveness in the workforce and for the individual's self-actualization [19]. Suicide was the leading cause of death in 2019 among individuals who belonged to the age of group 10-35 years old. Depression prevalence has been evaluated in various studies to be as low as 5% and as high as 22.3% within the adult Medicaid population, correlating to the format of the Patient Health Questionnaire used in each individual study [11]. Nearly 58% of respondents in a 2005/06 survey on National Health and Drug use stated that there was either a gap in their major depressive disorder treatment: receiving a lack of actual treatment, or an unmet need while receiving treatment, or both [19]. While rates of bipolar disorder are much lower than those of major depressive disorder, rates of death by suicide in this population approach 20% and are 20 – 30 times that of the general population making this a key area of intervention [17]. Ketamine and esketamine may play a pivotal role in helping to bridge this care gap. Regardless, continual tracking of ketamine prescription rate helps one better understand the standard of care for depression and among treatment resistant patients at heightened risk of suicide and identify policy opportunities aimed at optimizing care in this vulnerable population.

In conclusion, this study found that low and extremely regionally disparate, rates of ketamine and esketamine prescribing to Medicaid patients in 2019 and 2020. It will likely be a fruitful endeavor to continue to monitor how use of these agents change in future years.

Acknowledgements

BJP was supported by the Health Resources Services Administration ((D34HP31025). Software used in this effort was provided by the NIEHS (T32- ES007060-31A1). Iris Johnston provided technical support.

References

1. U.S. Department of Health and Human Services. (n.d.). *Suicide*. National Institute of Mental Health. Retrieved February 11, 2022, from https://www.nimh.nih.gov/health/statistics/suicide

2. Centers for Disease Control and Prevention. (2021, June 17). *Emergency department visits for suspected suicide attempts among persons aged 12–25 years before and during the COVID-19 pandemic - United States, January 2019–May 2021*. Centers for Disease Control and Prevention. Retrieved February 11, 2022, from https://www.cdc.gov/mmwr/volumes/70/wr/mm7024e1.htm

3. Duman RS. Ketamine and rapid-acting antidepressants: a new era in the battle against depression and suicide. *F1000Res*. 2018;7:F1000 Faculty Rev-659. doi:10.12688/f1000research.14344.1

4. Nischal, A, Tripathi A, Nischal A, Trivedi J K. Suicide and antidepressants: What current evidence indicates. *Mens sana monographs* 2012; Retrieved February 11, 2022, from https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3353604

5. Dadiomov D, Lee K. The effects of ketamine on suicidality across various formulations and study settings. *Mental Health Clin* 2019; 9(1):48-60. doi: 10.9740/mhc.2019.01.048.

6. *State Drug Utilization Data*. Medicaid. (n.d.). Retrieved February 11, 2022, from https://www.medicaid.gov/medicaid/prescription-drugs/state-drug-utilization-data/index.html

7. Murrough JW, Iosifescu DV, Chang LC, Al Jurdi RK, Green CE, Perez AM, et al. Antidepressant efficacy of ketamine in treatment-resistant major depression: a two-site randomized controlled trial. *Am J Psychiatry* 2013; 170(10), 1134–1142.https://doi.org/10.1176/appi.ajp.2013.13030392

8. Aan Het Rot, M, Zarate CA, Jr, Charney DS, Mathew SJ. Ketamine for depression: Where do we go from here?. *Biol Psychiatry* 2012; 72(7), 537–547. https://doi.org/10.1016/j.biopsych.2012.05.003

9. Zarate CA, Jr, Brutsche NE, Ibrahim L, Franco-Chaves J, Diazgranados N, Cravchik A., et al. Replication of ketamine's antidepressant efficacy in bipolar depression: a randomized controlled

add-on trial. *Biol Psychiatry* 2012; 71(11):939–946.

https://doi.org/10.1016/j.biopsych.2011.12.010

10. Chapel JM, Ritchey MD, Zhang D, Wang G. Prevalence and medical costs of chronic diseases among adult Medicaid beneficiaries. *Am J Prev Med* 2017; 53(6). https://doi.org/10.1016/j.amepre.2017.07.019

11. McIntyre RS, Rosenblat JD, Nemeroff CB, et al. Synthesizing the evidence for ketamine and esketamine in treatment-resistant depression: an international expert opinion on the available evidence and implementation. *Am J Psychiatry* 2021; 178:383–399

12. Mattes J. Ketamine after two antidepressants?. *Am J Psychiatry*, 178(12), p. 1129

13. The Retraction Watch Database [Internet]. New York: The Center for Scientific Integrity. 2018. [Cited (2/1/2022)]. Available from: http://retractiondatabase.org/.

14. Mojtabi R. Unmet need for treatment of major depression in the US. *Psychiatric Services* 2003; 60(3). P. 297-305

15. Lengvenyte A, Strumila R, Oliéa E, Courtet P. Ketamine and esketamine for crisis management in patients with depression: Why, whom, and how? *Eur Neuropsychopharm* 2022; 57:88-104.

16. Agboola F, Atlas SJ, Touchette DR, Fazioli K, Pearson SD. The effectiveness and value of esketamine for the management of treatment-resistant Depression: A summary from the Institute for Clinical and Economic Review's Midwest Comparative Effectiveness Public Advisory Council. *J Manag Care Spec Pharm*. 2020; 26(1):16-20.

17. Maldi KD, Asellus P, Myléus A, Norström F. Cost-utility analysis of esketamine and electroconvulsive therapy in adults with treatment-resistant depression. *BMC Psychiatry* 2021; 21:610. https://doi.org/10.1186/s12888-021-03601-8.

18. Ross EL, Soeteman DL. Cost-effectiveness of esketamine nasal spray for patients with treatment-resistant depression in the United States. *Psychiatr Serv*. 2020; 71(10): 988–997. doi:10.1176/appi.ps.201900625.

19. Miller JN, Black DW. Bipolar disorder and suicide: A review. *Current Psychiatry Rep* 2020; 22(2):6. doi: 10.1007/s11920-020-1130-0.

Supplemental Figure 1. Heat map showing prescriptions of ketamine per million Medicaid enrollees in 2019.

Supplemental Figure 2. Heat map showing prescriptions of esketamine per million Medicaid enrollees in 2019.

Supplemental Figure 3. Heat map showing prescriptions of ketamine per million Medicaid enrollees in 2020.

Supplemental Figure 4. Heat map showing prescriptions of esketamine per million Medicaid enrollees in 2020.

Supplementary Appendix A. Ketamine product name, National Drug Codes (NDC), and number of Medicaid prescriptions.

Product Name	NDC	Number of prescriptions
Ketamine 5	67457000110	2,139
	409205105	851
	143950810	659
	409205310	468
	143950801	374
	67457000100	362
	143950910	137
	143950901	97
Ketamine H	67457000110	1,099
	409205105	456
	143950810	375
	409205310	306
	67457018100	293
	67457018120	235
	67457000100	67
	67457010800	54
	143950801	36
	143950910	34
	143950901	14
	42023013810	14
	42023013710	0
	42023013910	0
	67457010810	0
Ketamine 50	42023011410	1065
	42023011510	801
Ketalar 20	42023011310	1530
Ketalar C-	42023011310	898
	42023011410	408
	42023011510	223
Ketalar	42023011310	204
	42023011510	180
	42023011410	56
Ketamine 2	67457018120	740
	67457018100	222
Ketamine 1	67457010810	81
	409205105	80
	67457010800	12

Supplemental Appendix B. Eskatamine product name and National Drug Code (NDC), and Medicaid prescriptions.

Product Name	NDC	Number of Prescriptions
Nasal spray	45802041059	4,007
	49348002827	1,136
	24385006710	812
	46122016510	446
	113030410	336
	36800006510	220
	36800030410	160
	49348023127	149
	36800081710	132
	49348023027	71
	49348013027	67
	113006510	32
	36800038810	25
	24385030411	0
	24385049810	0
	36800031213	0
	36800031221	0
	36800064810	0
	46122014903	0
	46122016511	0
	49348027627	0
Spravato N	50458002803	974
	50458002802	511
Spravato 8	50458002803	1,818
Spravato 5	50458002802	1,091
Spravato	50458002803	454
	50458002802	227

Supplementary Appendix C. Retracted ketamine articles reported to the RetractionWatch database. Rows in <mark>green</mark> highlights are especially relevant to depression or bipolar treatment.

Title	Journal	Outcome
Nardostachys jatamansi targets BDNF-TrkB to alleviate ketamine-induced schizophrenia-like symptoms in rats	*Neuropsychobiology*	Retracted Sept 13, 2021
The sustained antidepressant effects of ketamine are independent of the lateral habenula	*Journal of Neuroscience*	Retracted May 5, 2021
Ketamine reduces LPS-induced HMGB1 through HO-1 and Nrf2/p38 MAPK	*Journal of Surgical Research*	Retracted November, 2020
<mark>Negative trial of scopolamine in major depressive disorder does not demonstrate neurophysiological changes seen with the antidepressant response of ketamine</mark>	<mark>*Biological Psychiatry*</mark>	<mark>Retracted May 1, 2018</mark>
<mark>Efficacy of ketamine in bipolar depression: Systematic review and meta-analysis</mark>	<mark>*Journal of Psychiatric Practice*</mark>	<mark>Retracted July, 2018</mark>
The influence of ketamine on the locus coeruleus and the noradreneg system- a randomized fMRI study	*Clinical Neurophysiology*	Retracted Sept 4, 2018
<mark>A preliminary naturalistic study of low-dose ketamine for depression and suicide ideation in the emergency department</mark>	<mark>*International Journal of Neuropsychopharmacology*</mark>	<mark>Retracted July 1, 2017</mark>
Postoperative analgosedation with S(+) ketamine decreased the incidences of postanesthetic shivering and nausea and vomiting after cardiac surgery	*Medical Science Monitor*	Retracted Dec 1, 2008
Perioperative ketamine for acute	*Cochrane Database of*	Retracted (withdrawn) July 2,

postoperative pain	*Systematic Reviews*	2015
A randomized, double-blind, placebo controlled study assessing the anti-inflammatory effects of ketamine in cardiac surgical patients	*Journal of Cardiothoracic and Vascular Anesthesia*	Retracted Oct, 2014
Efficacy of lignocaine plus ketamine at different doses in the prevention of pain due to propofol injection	*Clinical Drug Investigation*	Retracted May, 2014
Sevoflurane-emergence agitation: Effect of supplementary low-dose oral ketamine premedication in preschool children undergoing dental surgery	*European Journal of Anaesthesiology*	Retracted April, 2011
Assessment of recovery, dreaming, hemodynamics, and satisfaction in post cardiac surgery patients receiving supplementary propofol sedation with S(+)- ketamine	*Minerva Anestesiologica*	Retracted May, 2011