1   **Using Explainable-AI to Find Geospatial Environmental and Sociodemographic**
2   **Predictors of Suicide Attempts**
3
4

Mirko Pavicic [1][†], Angelica M. Walker [2][†], Kyle A. Sullivan [1], John Lagergren [1], Ashley Cliff [2],

Jonathon Romero [2], Jared Streich [1], Michael R. Garvin [1], MVP Suicide Exemplar Workgroup, the

Million Veteran Program, John Pestian [1,3], Benjamin McMahon [4], David W. Oslin [5,6], Jean C.

Beckham [7,8,9], Nathan A. Kimbrel [*7,8,10,11], Daniel A. Jacobson [*1]

Affiliations:

[1]Oak Ridge National Laboratory, Computational and Predictive Biology, Oak Ridge, TN.
[2]The Bredesen Center for Interdisciplinary Research and Graduate Education, University of Tennessee Knoxville,
Knoxville, TN.
[3]Cincinnati Children's Hospital Medical Center, University of Cincinnati, Cincinnati, OH.
[4]Theoretical Biology and Biophysics, Los Alamos National Laboratory, Los Alamos, NM.
[5]VISN 4 Mental Illness Research, Education, and Clinical Center, Center of Excellence, Corporal Michael J. Crescenz
VA Medical Center, Philadelphia, PA, USA.
[6]Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, PA, USA.
[7]Durham Veterans Affairs Health Care System, Durham, NC.
[8]VA Mid-Atlantic Mental Illness, Research, Education, and Clinical Center.
[9]Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC, USA.
[10]Duke University School of Medicine, Duke University, Durham, NC.
[11]VA Health Services Research and Development Center of Innovation to Accelerate Discovery and Practice
Transformation, Durham, NC.
24
25
26 *Correspondence: nathan.kimbrel@va.gov, jacobsonda@ornl.gov
27 [†] Contributed equally
28

Abstract

Despite a global decrease in suicide rates in recent years, death by suicide has increased in the
United States. It is therefore imperative to identify the risk factors associated with suicide attempts
in order to combat this growing epidemic. In this study, we use an explainable artificial intelligence
(X-AI) method, iterative Random Forest (iRF), to predict suicide attempts using data from the
Million Veteran Program (MVP). Our predictive model incorporates multiple environmental
variables (e.g., elevation, light wavelength absorbance, temperature, humidity, etc) at ZIP code-
level geospatial resolution. We additionally consider demographic variables from the American
Community Survey as well as the number of firearms and alcohol vendors per 10,000 people in
order to assess the contributions of proximal environment, access to means, and restraint

decreasement to suicide attempts. Our results show that geographic areas with higher
concentrations of married males living with spouses are predictive of lower rates of suicide
attempts, whereas geographic areas where males are more likely to live alone and to rent housing
are predictive of higher rates of suicide attempts. We also identified climatic features that were
associated with suicide attempt risk by age group. Additionally, we observed that firearms and
alcohol vendors were associated with increased risk for suicide attempts irrespective of the age
group examined, but that their effects were small in comparison to the top features. Taken
together, our findings highlight the importance of social determinants and contextual factors in
understanding suicide risk among veterans.

48 Introduction

Suicide rates in the U.S. have been increasing in recent years despite these rates declining
globally[1]. According to the biopsychosocial model of suicide risks, there are distal, developmental,
and proximal factors that can affect the probability of suicide attempt[2]. Distal factors are those
related to familial and genetic predisposition and early-life adversity. Developmental factors
include personality traits associated with suicidal behavior, cognitive deficits, and chronic
substance misuse. Proximal, or precipitating factors include but are not limited to psychiatric,
psychological, socioeconomic, and environmental factors. Even though socioeconomic and
environmental factors are classified as proximal factors, they can simultaneously influence distal
and developmental factors such as early-life adversity and individual development. Several
studies have found associations between a number of demographic factors and suicide such as
age, ethnicity, socioeconomic status, marital status, religion, etc.[3]. Furthermore, climatic variables
such as cumulative solar radiation has been negatively associated with suicide risk, while
temperature and diurnal temperature range have been positively associated with suicide risk[4–6].
However, to thoroughly analyze the effect of environmental and socioeconomic factors on an
individual-level risk of attempting suicide, a study at high geospatial resolution is needed to
account for within-county variation[7].
65
Other relevant proximal factors include access to means and substance misuse[8,9]. For example,
occupations with access to lethal means are associated with increased risk of death by suicide
and method of choice[10,11]. Moreover, controlling access to lethal means is an effective strategy
for decreasing suicide risk[12]. In the U.S., death by suicide is the leading cause of violent deaths,
and firearms are responsible for approximately half of these deaths[13]. Substance misuse also
plays an important role in suicide prevention because acute substance intoxication can increase
individual's disinhibition. For example, a prior study showed that suicide decedents have an
increased risk of alcohol ingestion and intoxication before their death relative to controls[14].
74
The objective of the present research was to conduct an analysis of climatic and demographic
features that are associated with increased risk for suicide attempts among U.S. veterans using
an explainable artificial intelligence (X-AI) model. We were particularly interested in the
relationship of the number of firearms and alcohol vendors per 10,000 people as proxies for
access to means and decreased restraint, which were also included in the final model. Together,

we identify a number of novel factors at zip code-level resolution that impact individual-level risk
for attempting suicide.

Methods

Data and Data Pre-processing

Patient Data

The cohort used in this study was the same as previously described[15], with some modifications.
The patients in this study were enrolled in the U.S. Department of Veterans Affairs' (VA) Million
Veteran Program (MVP). All subjects provided informed consent and the research presented here
were approved by the VA Central Institutional Review Board. Cases were defined as having a
history of one or more suicide attempts (including both fatal and non-fatal), whereas controls were
defined has having no history of suicide attempts or ideation. This vector was created from
electronic health records (EHR) from the VA corporate data warehouse (CDW), using
International Classification of Diseases (ICD) diagnostic codes, survey data from the mental
health domain, and the Suicide Prevention Application Network (SPAN) data set[16]. We encoded
these data as a binary vector, where controls were represented by 0 and suicide attempts were
set to 1, regardless of whether the attempt was or was not fatal. In total, this resulted in 405,540
patients with 391,409 controls and 14,131 attempts. Each attempt and control were associated
with climatic and demographic features according to the most recent patient ZIP code. Sex
distribution was 8% female and 92% male. This cohort was predominantly white (73%), 18% were
black or African American, 1% American Indian or Alaska Native, 1% Asian, and 7% mixed race,
missing, or unknown. The mean age was 62.4 years for the whole cohort, 63.4 for males and 50.8
years for females (**Supplementary Figure 1A**). Notably, this cohort was unbalanced in terms of
cases and controls where only 3% attempted suicide versus 97% who did not, which is reflective
of the low occurrence of suicide attempts in the general population. Most of the suicide attempts
in this cohort were concentrated around 60 years old, likely since this age group is
overrepresented among these patients (**Supplementary Figure 1A and B**). However, the
proportion of attempts grouped by age decreased steadily after age 60 (**Supplementary Figure
1C**). Due to this steady decrease in attempts after age 60, we explored the demographic and
climatic features that were associated with suicide attempts in patients greater than 60 years of
age separately from those younger than 60. The split with patients above or equal to 60 contained
a total of 267,447 individuals with 4,231 attempts and 263,216 controls. The split with patients
below age 60 was composed of 138,093 individuals with 9,900 suicide attempts and 128,193
controls.

Climatic Features

The climatic features included two groups: static measurements and monthly measurements.
Monthly measurements included longitudinal features such as monthly average precipitation and
maximum temperature. There were 12 distinct measurement types recorded each month, totaling

144 features. The 30 static features included features such as elevation and percent urban cover.
This led to a total of 174 climatic and weather-related features, mapped to 33,144 ZIP codes
across the U.S.[17–23] (**Supplementary Table 1**).

Demographic Features

The demographic features were collected from the 2019 American Community Survey, produced
by the United States Census Bureau[24]. These features included demographic, socioeconomic,
and housing features, such as the percentage of the population that were white females between
the age of 30 and 34 living in the ZIP code, percentage of rented housing units in the ZIP code
that have adequate plumbing facilities, and the Gini Index of Income Inequality. These 1606
features were captured using the tidycensus software package in R using 5 years estimate for
2019[25]. These features were normalized to represent a percentage of the total population or age
bracket within each ZIP code. We also included two additional features: population density
(people per square mile) and the ratio of water to land area, which led to a total of 1608
demographic features that were mapped to 33,120 ZIP codes across the U.S (**Supplementary
Table 2**).

Alcohol and Firearms Features

133 The Historical Business Database[26] contains information on the businesses that exist in every ZIP
134 code in the U.S. We utilized two important features from this dataset: one that measured the
135 number of firearms vendors per 10,000 residents in each ZIP code, averaged across the years
136 2010 and 2019, and the similarly calculated number of alcohol vendors. This information was
137 included for 31,378 ZIP codes across the U.S (**Supplementary Table 3**).

Correlation Analysis

In this study, we used iRF[27,28], an X-AI algorithm that ranks input features by importance through
iterative feature weighting, to associate proximal environmental features with suicide attempts.
Thus, data pre-processing was necessary to ensure accuracy and objectivity in the results by
reducing the number of highly correlated, redundant features explaining variance in the model.
Large pairwise correlations between input features may negatively impact the iRF model, since
the importance of one feature is then split across the correlated features, thereby decreasing their
overall importance in predicting suicide attempt. To account for this, if one feature was highly
correlated with another, one of the features was selected to be a representative of both features,
while the other feature was removed from consideration. Pearson's correlation coefficients were
calculated between all 1,784 features. Pairs of features with a correlation coefficient greater than
or equal to 0.90 in absolute value were considered to be highly correlated. This analysis found
390 features that were highly correlated with each other. Of these 390 features, 142 of them were
kept as representative features (**Supplementary Table 4**). Thus, 248 features were dropped,
reducing the total number of features to 1,536.

iRF k-Fold Cross Validation and Accuracy Calculation

To obtain accuracy scores for the iRF model, k-fold cross validation was used. For this study, 5-fold cross validation was repeated ten times, for a total of 50 computational runs. For each run, the data was separated into an 80/20 training/testing split, where 80% of the data was used for training the model and the remaining 20% was reserved for testing. To prevent geographical bias, patients within the same ZIP code were moved to either the training or testing dataset. Prediction accuracy was calculated using the average Area Under the Precision Recall curve (AUPRC), where precision was defined as *true positives / true positives + false positives* and recall as *true positives / true positives + false negatives*. Each random forest in the iRF model includes 1000 trees with a leaf node size of 1000 patients. We set the number of iRF iterations to five to rank the importance of each feature in predicting suicide attempt. In addition to ranking input features by importance, we next identified the feature-level explainability of our model by determining whether each feature was predictive for or against suicide attempt. To estimate if a feature predicts suicide attempt or controls, the result of each split was averaged using the given feature and mapping those to a linear effect. This provided both the feature effect in the slope of the line and an R^2 with how closely that related to each of the set of splits. If the slope of the line was positive, then the feature effect size direction was positive, i.e., the value and the feature were positively correlated. If the slope of the line was negative, then the feature effect size direction was negative.

Model selection

Initially we ran iRF models with *k*-fold cross validation using all 1,536 features to capture the most important features predicting suicide attempt in patients above or equal 60 years old and below 60 years old. Then, we used several combinations of the top most important features together with alcohol and firearm vendors per 10,000 residents to run reduced iRF models with *k*-fold cross validation to evaluate if this improved the accuracy of the models. The accuracy of the reduced models was better than using all features in all cases as shown in the area under the precision recall curves (AUPRC) for all 50 data splits used to test the models (**Supplementary Figure 2 and 3**). These results show that by using all features we may introduce noise that may reduce accuracy and explainability of the model. Interestingly, most of the models had some degree of predictability, since most of them had AUPRC above random chance, and this effect was stronger in the age group of patients below 60 years old (red dashed line in **Supplementary Figure 2**). Based on these results, we selected the model including the top 20 features plus firearms and alcohol vendors for model explanation.

iRF-LOOP

To show how features or groups of features interconnect each other, we applied iRF-Leave One Out Prediction (iRF-LOOP), which is an extension of the iRF model[27]. In this framework, iRF was used to compute all-against-all predictions of each vector of data from all other vectors. The results of this analysis were captured as networks, in which nodes (i.e., features) were connected by an edge if the pair of features were predictive of each other, thereby revealing functional

relationships and subgroups within and across data layers. We performed iRF-LOOP using the
pre-processed input matrix which consisted of climate, census, and alcohol and firearm business
data, with a total of 1,536 features and 31,378 samples or ZIP codes. This analysis created an
all-to-all directed feature association network that captured the relationships between data layers.
The resulting network was then thresholded to the top 1% of edges to capture the most important
connections between features.

Results

Features that were most strongly associated with suicide attempts

iRF predictive models can compute the importance of each feature predicting the dependent
variable as well as the direction of the prediction, which in this case was predicting suicide attempt
vs control. We examined iRF models trained with 1,536 features to identify the 20 most important
predictive features in patients below, and above or equal to age 60, plus *firearms and alcohol*
vendors per 10,000 residents (**Supplementary Figure 4 and 5**). The resulting features were
further analyzed to determine directionality (i.e., if they predict suicide attempts or controls) and
can be aggregated in groups related to emotional support, housing, ancestry, commuting and
mobility, access to healthcare, cognitive difficulties, access to means, decrease restraint and
climate (**Figure 1 and 2**).

In the model using patients of above or equal 60 years old, the features *married males* and
household with spouse present between 35 - 64, and 65 years and older were associated with
decreased risk for attempting suicide, whereas *males never married of 75 - 84 years*, and *married*
males of 85 years old with spouse absent were associated with increased risk for attempting
suicide (**Figure 1**). Ancestry also appeared among the top features explaining suicide attempts.
French Canadian, *Northern European*, *Pennsylvania German ancestries* were associated with
increased risk for suicide attempts, whereas *Dutch ancestry* and *speaking French (Haitian or*
Cajun dialects) at home were associated with reduced risk (**Figure 1**). *Females with a work*
commute lasting 10 to 14 minutes, and *number of firearms* and *alcohol vendors per 10,000*
residents were also associated with increased risk (**Figure 2**). Several climate features were also
predictive: *precipitation in April*, *solar radiation in July,* and *wind speed in July* were associated
with decreased risk, whereas *wind speed in April* and *maximum temperature in December* were
associated with increased risk **(Figure 2)**.

For the model using only patients younger than 60, the two most important features associated
with decreased risk were *married males*, and *females with no cognitive difficulties between 35*
and 64 years old (**Figure 1 and 2**). Conversely, the features *living alone* and *living with unmarried*
partner were associated with increased risk (**Figure 1**). Similarly, *commuting between 10 to 14*
mins, regardless of gender, *monthly housing costs between $700-800, cash rent between $600-*
649, *house heating using gas*, *moved within county,* and *having two types of public insurance*
were all associated with increased risk (**Figure 1 and 2**). On the other hand, only *Pennsylvania*
German ancestry was associated with increased risk in the ancestry category (**Figure 1**). Similarly
to the model from patients 60 years of age or older, *number of firearm* and *alcohol vendors per*

10,000 inhabitants were associated with increased risk (**Figure 2**). Regarding climatic features,
precipitation in January and *water vapor in May* were associated with reduced risk (**Figure 2**).
Contrary to the first model, in patients younger than 60 the climatic variables *solar radiation in*
July and *wind speed in July* were predictive of individuals with a history of suicide attempt
(**Figures 1 and 2**). Additionally, *wind speed in April* and *terrain elevation* were also associated
with suicide attempts (**Figure 2**).

239 Feature network by iRF-LOOP

It is well known that alcohol abuse increases suicide risk[2]. Thus, we included the number of
alcohol vendors per 10,000 people in our final model. Interestingly, when used in combination
with the top 20 features, alcohol vendors per 10,000 people ranked 12th and 20th in feature
importance in patients above or equal to 60 and below 60 years old respectively, even though it
ranked 360th (above age of 60) and 74th (below age of 60) in the models using all features
(**Supplementary Figure 4 and 5**). This could mean that other features are competing with
number of alcohol vendors per 10,000 people within the iRF model to classify suicide attempt and
controls. Therefore, we created a feature network using iRF-LOOP to explore how features relate
to each other. **Figure 3** shows the immediate neighbors of the feature measuring the number of
alcohol vendors per 10,000 people, where each node in the subnetwork represents a feature and
the arrows represent the edges (connections) between features. Edges are weighted by their
normalized feature importance value and the arrow direction shows what feature is predicting
another one. Several features related to geographic mobility, population density, low gross rent
with cash, and high home value were observed. Further, the subnetwork shows associations
among features including number of widowed females, several European ancestries, and the
population-adjusted density of alcohol vendors. Regarding climate features, minimum
temperature in July and maximum temperature in December were observed.

257 Discussion

Most prior suicide prevention studies have focused on a relatively small number of features.
Moreover, most have typically relied on individual-level information only (e.g., clinical features) or
aggregated data for a given geographical area with low geospatial resolution. In the present study
we showed that ZIP code-level data can improve prediction in individuals with a history of suicide
attempt greater than random chance, supporting the role of the surrounding environment as
proximal/precipitating factors that influence the propensity for an individual to attempt suicide.
Furthermore, using X-AI we were able to screen more than a thousand demographic and climatic
features to obtain several that appear to potentially protect or increase veterans' propensity for
attempting suicide.
267
An individual's environment can have profound effects on their psychological development and
subsequent risk of suicide. In fact, it has been well documented that socioeconomic and
demographic factors such as poverty, education and population density are related to suicide
risk[29–31]. In our study we used the EHR data from VA CDW, ACS and climate data to build a
predictive model of suicide attempts in U.S. veteran population. It is worth highlighting that ACS

and climate data are not direct information of the individuals under study, but rather a
representation of the environment in which they currently reside. Moreover, in the cohort used
here, there was a rapid decline in the number of suicide attempt after the age of 60
(**Supplementary Figure 1**), in agreement with a surveillance summary from the Center for
Disease Control and Prevention (CDC), where the highest suicide rates were observed in age
group between 35 and 64 years[32]. Thus, we divided the cohort into patients younger or older than
60 for further analysis, since age groups may be affected differently by risk factors[33].

Interestingly, the top features can be classified into nine main groups namely: emotional support,
housing, ancestry, commuting and mobility, climate, decrease restraint, access to means,
cognitive difficulties and access to healthcare (**Figure 1 and 2**). In the emotional support group,
the protective effect of marital status on suicide risk is well-documented[34]. Studies have shown
that married people showed the lowest rate of suicide rates[34–36], whereas divorced or separated
persons are twice as likely to commit suicide than married persons, and this effect is stronger in
divorced males[37]. In our study, for all age groups we observed that living in areas with high
proportions of married males and individuals living with a spouse were protective factors against
suicide attempts for those aged 60 years old or above (similar to the individual-level protective
factor of marital status). Conversely, areas with high proportions of individuals living alone and
unmarried males were associated with higher rates of suicide attempts[38].

Regarding the commuting and mobility group, commuting time has been associated with
depression in a dose responsive manner[39]. Our results showed that living in areas with high
proportions of individuals with commutes between 10 and 14 minutes has been associated with
increased risk for suicide attempts, irrespective of age. The iRF-LOOP network showed that
commuting features are predictive of each other regardless of the commuting time
(**Supplementary figure 6**). Thus, commuting between 10 and 14 minutes may be acting as a
proxy for commuting in general. We also observed housing-related features for both age groups
in relation to suicide attempts. These results are consistent with Lorant *et al.* (2005)[40], who
showed that higher education and home ownership status decreases the risk of suicide. Other
studies have found that living in rented units increased the risk for suicide in middle-aged males
and females[31]. This effect might be especially important for females who live in large urban
areas[41].

In the ancestry group, we observed that living in areas with a higher proportion of Northern
European ancestry was associated with suicide attempts. Although suicide attempts vary widely
in Europe, countries of Finno-Ugrian origin show disproportionally higher suicide rates than the
rest of Europe, suggesting a genetic cause[42]. Voracek (2006)[43], tested this hypothesis in the
U.S.A. using state-level self-reported ancestry from census data. The study found support for this
hypothesis using historical data from 1913-1924 and 1928-1932, but not from 1990-1994. Taken
together, these findings encourage an analysis with higher geospatial resolution of the sample
(e.g. ZIP or county level) and a genetic characterization of the individuals ancestries, since self-
reporting may be inaccurate. We also found that living in a ZIP code with higher proportion of
Native Hawaiian and other pacific islander females between ages 45 to 54 was associated with
suicide attempts. These results are consistent with Ji and colleagues (2020)[44] finding, who

showed that Asian or Pacific Islander ancestry was a risk factor for suicide in healthcare
professionals.
319

Seasonality of suicide attempts is also a well-known phenomenon, and climate has been
suggested to potentially be the main cause of this pattern[45]. However, results have varied
depending on the methods and location used[46,47]. Several studies have shown that solar
radiation, precipitation, and temperature are associated with suicide risk[4–6]. In our results we
observed that precipitation in April, solar radiation in July and wind speed in July were protective,
whereas wind speed in April and maximum temperature in December (with lower temperatures)
were associated with increased risk among patients 60 years of age or older. In patients younger
than 60 years old, precipitation in January and water vapor in May were protective, whereas solar
radiation, wind speed in April and July were associated with increased risk; however, additional
research is still needed in independent samples to confirm these findings, as climatic features of
neighboring months are highly correlated, making it difficult to dissect suicide attempt seasonality
using our method. Furthermore, while climatic variation exists even within zip code-level
resolution, climatic features exhibit the closest overlap between individual-level and zip code-level
features.
334

Another goal of the present work was to explore decreased restraint and access to means for
suicide attempts using the number of alcohol and firearms vendors per 10,000 residents as
proxies. Importantly, regardless of age, we observed that living in areas with more alcohol and
firearm vendors was associated with increased risk for attempting suicide. These results provide
additional evidence regarding the importance of access to means as a risk factor for suicide,
especially in the U.S., where a larger proportion of suicides are committed using firearms in
comparison to other high income countries where only 5% use them[13,48,49]. Our findings also
confirm the importance of including alcohol abuse in models of suicide risk[50,51].
343

Study Limitations and Future Directions

The present study also had a number of limitations that should be considered when interpreting
these findings. First, we utilized a cross-sectional design, and a lifetime suicide attempts variable.
Thus, additional work is still needed to examine the degree to which the features identified in the
present study might be predictive of future suicide attempts. Second, our findings regarding
alcohol and firearms vendors should be interpreted cautiously, as there are several other
variables (**Supplementary figure 7**) that could also potentially explain their association with
suicide attempts which should be considered in future work. Third, while the present study
provides clear evidence that zip code-level geospatial variables can predict suicide attempt risk
better than random chance, the degree to which these variables might predict suicide attempts
above and beyond well-validated individual-level risk factors (e.g., psychiatric diagnosis) remains
unknown. While such work is beyond the scope of the present study, future work is still needed
to ascertain if and how geospatial variables might interact with individual-level data to predict
suicide risk.

Conclusion

In this study we demonstrated the use of X-AI to explore the impact of more than 1,000 demographic and climatic features on suicide attempt risk with high geospatial resolution. This research provides additional evidence for the role of several demographic and climatic features in suicide attempts and demonstrates the utility of using geospatial features within an X-AI framework to improve suicide risk prediction.

Acknowledgements

This research is based on data from the Million Veteran Program, Office of Research and Development (ORD), Veterans Health Administration (VHA), and was supported by award #I01CX001729 from the Clinical Science Research and Development (CSR&D) Service of VHA ORD. This publication does not represent the views of the Department of Veteran Affairs or the United States Government. We also thank and acknowledge MVP and the MVP Suicide Exemplar Workgroup for their contributions to this manuscript. This work was also supported in part by the joint U.S. Department of Veterans Affairs and US Department of Energy MVP CHAMPION program. This manuscript has been co-authored by UT-Battelle, LLC under contract no. DE-AC05-00OR22725 with the U.S. Department of Energy. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The Department of Energy will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (http://energy.gov/downloads/doe-public-access-plan, last accessed September 16, 2020).

Conflict of interest

The authors have no conflict of interest to declare at this time.

References

1. WHO. World Health Statistics data visualizations dashboard. Noncommunicable diseases and mental health. World Health Organization. Published 2019. Accessed December 6, 2021. https://www.who.int/data/gho/data/themes/mental-health/suicide-rates

2. Turecki G, Brent DA, Gunnell D, et al. Suicide and suicide risk. *Nature Reviews Disease Primers*. 2019;5(1). doi:10.1038/s41572-019-0121-0

3. Huang X, Ribeiro JD, Musacchio KM, Franklin JC. Demographics as predictors of suicidal thoughts and behaviors: A meta-analysis. *PLoS One*. 2017;12(7):e0180793.

4. Dixon PG, Kalkstein AJ. Where are weather-suicide associations valid? An examination of nine US counties with varying seasonality. *Int J Biometeorol*. 2018;62(5):685-697.

5. Holopainen J, Helama S, Partonen T. Does diurnal temperature range influence seasonal

suicide mortality? Assessment of daily data of the Helsinki metropolitan area from 1973 to
2010. *Int J Biometeorol*. 2014;58(6):1039-1045.

6.   Ruuhela R, Hiltunen L, Venäläinen A, Pirinen P, Partonen T. Climate impact on suicide
rates in Finland from 1971 to 2003. *Int J Biometeorol*. 2009;53(2):167-175.

7.   Lin CY, Hsu CY, Gunnell D, Chen YY, Chang SS. Spatial patterning, correlates, and
inequality in suicide across 432 neighborhoods in Taipei City, Taiwan. *Soc Sci Med*.
2019;222:20-34.

8.   Pompili M, Serafini G, Innamorati M, et al. Suicidal behavior and alcohol abuse. *Int J*
Environ Res Public Health. 2010;7(4):1392-1431.

9.   Nock MK, Green JG, Hwang I, et al. Prevalence, correlates, and treatment of lifetime
suicidal behavior among adolescents: results from the National Comorbidity Survey
Replication Adolescent Supplement. *JAMA Psychiatry*. 2013;70(3):300-310.

10.   Milner A, Witt K, Maheen H, LaMontagne AD. Access to means of suicide, occupation and
the risk of suicide: a national study over 12 years of coronial data. *BMC Psychiatry*.
2017;17(1):125.

11.   Skegg K, Firth H, Gray A, Cox B. Suicide by occupation: does access to means increase
the risk? *Aust N Z J Psychiatry*. 2010;44(5):429-434.

12.   Sarchiapone M, Mandelli L, Iosue M, Andrisano C, Roy A. Controlling access to suicide
means. *Int J Environ Res Public Health*. 2011;8(12):4550-4562.

13.   Ertl A, Sheats KJ, Petrosky E, Betz CJ, Yuan K, Fowler KA. Surveillance for Violent Deaths
- National Violent Death Reporting System, 32 States, 2016. *MMWR Surveill Summ*.
2019;68(9):1-36.

14.   Kaplan MS, Huguet N, McFarland BH, et al. Use of alcohol before suicide in the United
States. *Annals of Epidemiology*. 2014;24(8):588-592.e2.
doi:10.1016/j.annepidem.2014.05.008

15.   Kimbrel NA, Ashley-Koch AE, Qin XJ, et al. A Genome-Wide Association Study of Suicide
Attempts in the Million Veterans Program Identifies Evidence of Pan-Ancestry and
Ancestry-Specific Risk. *Mol Psychiatry*.

16.   Hoffmire C, Stephens B, Morley S, Thompson C, Kemp J, Bossarte RM. VA Suicide
Prevention Applications Network: A National Health Care System-Based Suicide Event
Tracking System. *Public Health Rep*. 2016;131(6):816-821.

17.   Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global
land areas. *International Journal of Climatology*. 2017;37(12):4302-4315.
doi:10.1002/joc.5086

18.   Fischer G, Nachtergaele F, Prieler S. Global agro-ecological zones assessment for
agriculture (GAEZ 2008). *IIASA, Laxenburg, Austria*. Published online 2008.

19.   Fischer G, Nachtergaele FO, Prieler S. Global Agro-Ecological Zones (GAEZ v3. 0): Model
Documentation. International Institute for Applied systems Analysis (IIASA), Laxenburg.
Published online 2012.

20. Igbp-Dis S. A program for creating global soil-property databases. *IGBP Global Soils Data*
Task, France. Published online 1998.

21. Trabucco A, Zomer RJ. High-Resolution Global Soil-Water Balance explicit for Climate–
Standard Vegetation and Soil Conditions. *CGIAR Consortium for Spatial Information*.
Published online 2010.

22. Zomer, J. R, Bossio, et al. *Trees and Water: Smallholder Agroforestry on Irrigated Lands in*
Northern India. IWMI; 2007.

23. Zomer RJ, Trabucco A, Bossio DA, Verchot LV. Climate change mitigation: A spatial
analysis of global land suitability for clean development mechanism afforestation and
reforestation. *Agriculture, Ecosystems & Environment*. 2008;126(1-2):67-80.
doi:10.1016/j.agee.2008.01.014

24. United States Census Bureau. American Community Survey: Data Profiles.

25. Walker K, Herman M. tidycensus: Load US Census Boundary and Attribute Data as
"tidyverse" and "sf"-Ready Data Frames. Published online 2021. https://CRAN.R-
project.org/package=tidycensus

26. Data Axle. 2010-2019 Historical Business Data.

27. Cliff A, Romero J, Kainer D, Walker A, Furches A, Jacobson D. A High-Performance
Computing Implementation of Iterative Random Forest for the Creation of Predictive
Expression Networks. *Genes* . 2019;10(12). doi:10.3390/genes10120996

28. Basu S, Kumbier K, Brown JB, Yu B. Iterative random forests to discover predictive and
stable high-order interactions. *Proc Natl Acad Sci U S A*. 2018;115(8):1943-1948.

29. Bantjes J, Iemmi V, Coast E, et al. Poverty and suicide research in low- and middle-income
countries: systematic mapping of literature published in English and a proposed research
agenda. *Global Mental Health*. 2016;3. doi:10.1017/gmh.2016.27

30. Vichi M, Vitiello B, Ghirini S, Pompili M. Does population density moderate suicide risk? An
Italian population study over the last 30 years. *Eur Psychiatry*. 2020;63(1):e70.

31. Johansson LM, Sundquist J, Johansson SE, Qvist J, Bergman B. The influence of ethnicity
and social and demographic factors on Swedish suicide rates. *Soc Psychiatry Psychiatr*
Epidemiol. 1997;32(3):165-170.

32. Ivey-Stephenson AZ, Crosby AE, Jack SPD, Haileyesus T, Kresnow-Sedacca MJ. Suicide
Trends Among and Within Urbanization Levels by Sex, Race/Ethnicity, Age Group, and
Mechanism of Death — United States, 2001–2015. *MMWR Surveillance Summaries*.
2017;66(18):1-16. doi:10.15585/mmwr.ss6618a1

33. Suresh Kumar PN, Anish PK, George B. Risk factors for suicide in elderly in comparison to
younger age groups. *Indian J Psychiatry*. 2015;57(3):249-254.

34. Rendall MS, Weden MM, Favreault MM, Waldron H. The protective effect of marriage for
survival: a review and update. *Demography*. 2011;48(2):481-506.

35. Smith JC, Mercy JA, Conn JM. Marital status and the risk of suicide. *Am J Public Health*.

1988;78(1):78-80.

36. Kyung-Sook W, SangSoo S, Sangjin S, Young-Jeon S. Marital status integration and suicide: A meta-analysis and meta-regression. *Soc Sci Med*. 2018;197:116-126.

37. Kposowa AJ. Marital status and suicide in the National Longitudinal Mortality Study. *J Epidemiol Community Health*. 2000;54(4):254-261.

38. Calati R, Ferrari C, Brittner M, et al. Suicidal thoughts and behaviors and social isolation: A narrative review of the literature. *J Affect Disord*. 2019;245:653-667.

39. Wang X, Rodríguez DA, Sarmiento OL, Guaje O. Commute patterns and depression: Evidence from eleven Latin American cities. *J Transp Health*. 2019;14:100607.

40. Lorant V, Kunst AE, Huisman M, Costa G, Mackenbach J, EU Working Group on Socio-Economic Inequalities in Health. Socio-economic inequalities in suicide: a European comparative study. *Br J Psychiatry*. 2005;187:49-54.

41. Johansson LM, Sundquist J, Johansson SE, Bergman B. Ethnicity, social factors, illness and suicide: a follow-up study of a random sample of the Swedish population. *Acta Psychiatr Scand*. 1997;95(2):125-131.

42. Marusic A, Farmer A. Genetic risk factors as possible causes of the variation in European suicide rates. *Br J Psychiatry*. 2001;179:194-196.

43. Voracek M. Ancestry, genes, and suicide: a test of the Finno-Ugrian Suicide Hypothesis in the United States. *Percept Mot Skills*. 2006;103(2):543-550.

44. Ji YD, Robertson FC, Patel NA, Peacock ZS, Resnick CM. Assessment of Risk Factors for Suicide Among US Health Care Professionals. *JAMA Surg*. 2020;155(8):713-721.

45. Lee HC, Lin HC, Tsai SY, Li CY, Chen CC, Huang CC. Suicide rates and the association with climate: a population-based study. *J Affect Disord*. 2006;92(2-3):221-226.

46. Ajdacic-Gross V, Lauber C, Sansossio R, et al. Seasonal associations between weather conditions and suicide--evidence against a classic hypothesis. *Am J Epidemiol*. 2007;165(5):561-569.

47. Volpe FM, Tavares A. Re: "Seasonal associations between weather conditions and suicide--evidence against a classic hypothesis." *Am J Epidemiol*. 2007;166(2):237; author reply 237-238.

48. Yip PSF, Caine E, Yousuf S, Chang SS, Wu KCC, Chen YY. Means restriction for suicide prevention. *Lancet*. 2012;379(9834):2393-2399.

49. Grinshteyn E, Hemenway D. Violent death rates in the US compared to those of the other high-income countries, 2015. *Prev Med*. 2019;123:20-26.

50. Arsenault-Lapierre G, Kim C, Turecki G. Psychiatric diagnoses in 3275 suicides: a meta-analysis. *BMC Psychiatry*. 2004;4:37.

51. Dumais A, Lesage AD, Alda M, et al. Risk factors for suicide completion in major depression: a case-control study of impulsive and aggressive behaviors in men. *Am J*

Figures

510
511   **Figure 1.** Summary of features found by iRF related to emotional support, housing, and ancestry.
512   Cyan: predict control, yellow: predict attempt, blue border: above 60 years of age, red below 60
513   years of age, purple: both age groups.
514
515

Figure 2. Summary of features found by iRF related to commuting and mobility, climate, decreased restraint, access to means, access to healthcare and cognitive difficulties. Cyan: predict control, yellow: predict attempt, blue border: above 60 years of age, red below 60 years of age, purple: both age groups.

523
524
525   **Figure 3**. iRF-LOOP Subnetwork showing first neighbors of alcohol vendors per 10000 people.
526   Only the top 1% of edges weighed by normalized importance are shown. Arrow direction
527   corresponds to one feature predicting another feature.