Interactions between seasonal temperature variation and temporal synchrony drive

increased arbovirus co-infection incidence

Marya L. Poterek[1], Chantal B.F. Vogels[2], Nathan D. Grubaugh[2], Gregory D. Ebel[3], T. Alex

Perkins[1], Sean M. Cavany[1]

1. Department of Biological Sciences, University of Notre Dame

2. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale

University

3. Department of Microbiology, Immunology, and Pathology, Colorado State University

ABSTRACT

Though instances of arthropod-borne (arbo)virus co-infection have been documented clinically,

the overall incidence of arbovirus co-infection and its drivers are not well understood. Now that

dengue, Zika, and chikungunya viruses are all in circulation across tropical and subtropical

regions of the Americas, it is important to understand the environmental and biological

conditions that make co-infections more likely to occur. To understand this, we developed a

mathematical model of cocirculation of two arboviruses, with transmission parameters

approximating dengue, Zika, and/or chikungunya viruses and co-infection possible in both

humans and mosquitoes. We examined the influence of seasonal timing of arbovirus

cocirculation on the extent of co-infection. By undertaking a sensitivity analysis of this model,

we examined how biological factors interact with seasonality to determine arbovirus co-infection

transmission and prevalence. We found that temporal synchrony of the co-infecting viruses and

average temperature were the most influential drivers of co-infection incidence. For seasonal

patterns typical of a tropical region, we observed non-negligible incidence irrespective of arrival

time when two arboviruses arrived simultaneously. Under our default parameter settings, this

corresponded to a maximum co-infection cumulative incidence of 83 per 1,000 individuals and a

minimum cumulative incidence of 32 per 1,000 individuals in the year following arrival. For

seasonal patterns typical of a more temperate region, co-infections only occurred if arrivals took

place near the seasonal peak, and even then, did not reach 0.01 co-infections per 1,000

individuals. Our model highlights the synergistic effect of co-transmission from mosquitoes,

which leads to more than double the number of co-infections than would be expected in a

scenario without co-transmission. Our results show that arbovirus co-infections are unlikely to

occur in appreciable numbers unless epidemics overlap in space and time and in a tropical

region.

COMPETING INTEREST

The authors declare no competing interests

FUNDING

This work was supported by the NIH National Institute of General Medical Sciences R35 MIRA

award (R35GM143029) to TAP. MLP was additionally supported by a Richard and Peggy

Notebaert Premier Fellowship.

INTRODUCTION

The past decade has seen the Americas affected by epidemics of both Zika and chikungunya,

adding to the burden of arthropod-borne (arbo)viral disease in a region where seasonal dengue

epidemics were already a regular occurrence in most countries (1–3). All three of the viruses that

cause these diseases are spread by the same vectors: *Aedes aegypti* and *Aedes albopictus*

mosquitoes. Hence, the diseases' spatiotemporal distribution is largely determined by the same

environmental and climatological drivers (4–7). This has led to overlapping epidemics of two

and three viruses, which in turn has led to many reports of co-infections (8). The rate of co-

infections with multiple arboviruses is magnified by the ability of the vector to be simultaneously

co-infected with two or more viruses and to co-transmit two or more viruses with a single bite

(9).

The phenomenon of arbovirus co-infection is still largely understudied with many unknowns (8).

For instance, while some studies have reported an increased risk of severe outcomes in co-

infections of dengue virus (DENV) and chikungunya virus (CHIKV), other studies have not

observed this (10,11). Similarly, while co-infection involving Zika virus (ZIKV) does not alter

the clinical presentation of uncomplicated infections, it is unclear whether it alters the risk of

severe disease (12). It is also unclear the extent to which prior or recent infection with one virus

can enhance or protect against subsequent infection with another (13–16). When multiple

arboviruses circulate in the same region at the same time, the combination of uncertainty about

cross-protection versus mutual enhancement, differing importation times of each virus, and

strong seasonal climate drivers, leads to potentially complex temporal patterns of single infection

and co-infection (17).

Seasonal climate drivers play an important role in arbovirus infection dynamics, as variations in

temperature determine environmental suitability for mosquito vector survival and virus

transmission (18–21). Arbovirus epidemic size and duration are a product of both mean

temperatures and seasonal variation and are maximized under conditions that promote mosquito

survival (22). Tropical climates are generally more suitable for arbovirus vectors (6) and are

therefore more likely to experience recurring arbovirus epidemics, which leads to the accrual of

immunity in human populations who live there (23). While much remains unknown about the

level of cross-immunity between arboviruses, preexisting immunity in a population is likely to

impact the dynamics and observed patterns of arbovirus co-infections, as well.

Data on the frequency of arbovirus co-infection remain sparse (8) and where data does exist

there are many factors which could lead to variability between studies, such as cross-immunity,

epidemic timing, and seasonality. In this context, mathematical modeling provides a useful way

to synthesize our understanding of arbovirus transmission and explore the conditions which may

most likely give rise to a heightened burden of arbovirus co-infection. To do this, we built a

temperature-dependent mathematical model of arbovirus co-circulation and co-transmission that

permits cross-protection between arboviruses and asynchronous epidemics. We first use the

model to understand the interplay of differing importation times and seasonal transmission in an

immunologically naive population. Next, we describe how the burden of co-infection could

change under differing levels of immunity and cross-protection. Finally, we undertake a global

sensitivity analysis of our model's parameters to provide a holistic view of the conditions which

may lead to the highest frequency of co-infection in humans.

METHODS

Model

We used a deterministic SEIR-SEI model to explore the influence of temperature on arbovirus

co-infection magnitude and timing. This model incorporates two arboviruses, referred to as virus

A and virus B, with identical transmission and human recovery rate parameters. We relied upon

several of the structural assumptions and parameter values reported by Vogels et al. (8),

particularly those governing co-transmission. In our model, transmission from co-infected

humans and mosquitoes occurs with the same probability as transmission from singly-infected

humans and mosquitoes, and there is no mortality effect from infection in either humans or

mosquitoes (9,24). Consistent with Rückert et al. (9), we assume that 60% of mosquitoes become

co-infected following a blood meal on a co-infected human, while 20% become singly-infected

with virus A and 20% singly-infected with virus B. Our model implements an intermediate

transmission scenario, in which 50% of bites from a co-infected mosquito lead to co-infection

and 50% lead to a single infection, the latter split evenly between the two arboviruses. We

assessed model sensitivity to this assumption of intermediate transmission, which has been used

in previous modeling work (8). Viruses A and B have identical, dengue-like parameters (Table

2).

We additionally made several assumptions about the roles that exposed and infected individuals

play in transmission. Co-infection in humans and mosquitoes can occur via co-transmission to

susceptible individuals or sequential transmission involving individuals who are susceptible and

then exposed to the first of two viruses. Following their infectious period, individuals recover,

and if they have only been singly infected by one virus can then be singly infected with the other

virus. Infection while an individual is exposed restarts the incubation period. In our baseline

analysis, we assumed no cross-protection or enhancement from a prior infection but explored this

in later analyses. Structural details of the model are illustrated in Figure 1, with sequential

transmission indicated in red and co-transmission in blue.

We incorporated a seasonal component into the model by using temperature-dependent

parameters, where biologically appropriate (Table 1). We used sinusoidal temperature curves

with a period of one year to drive the values of these parameters, with mean and amplitude

chosen to reflect specific climate regimes (Rio de Janeiro for tropical regions, Beijing for

temperate regions). We did not consider diurnal temperature variation. Temperature-dependent

parameters were those describing *Aedes aegypti* life history: mosquito biting rate, probability

that an infected mosquito transmits to a human during feeding, probability that a mosquito

becomes infected after feeding on an infected human, mosquito mortality rate, and virus extrinsic

incubation rate. Their values were chosen with reference to previous modeling work on fitted

thermal responses for *Aedes aegypti* (22). This approach allowed us to explore the relative

influences of seasonal timing and temporal synchrony on arbovirus co-infection under several

climate scenarios.

Human

Mosquito

Figure 1: Model diagrams for the human and mosquito components of the model. Red arrows indicate sequential transmissions into a co-infected state, while blue arrows indicate co-transmissions into a co-infected state. Superscripts refer to co-transmission (C) or sequential transmission (S); co-transmission can occur to susceptible or exposed individuals. Subscript "12" refers to co-infection, and "1,2" refers to secondary infection with the second virus following recovery from the first. Compartments are colored by state.

Equations

Model equations for humans are as follows, with parameter values and meanings shown in

Tables 1-2. Eqs. 1-3 define the forces of infection for each infection for each virus and for co-

transmitted viruses, which involves input from mosquito states and transmission probabilities.

Eqs. 7-8 and 11-12 distinguish between co-transmitted co-infections, in which individuals

become infected with two arboviruses simultaneously, and sequentially-transmitted co-

infections, in which individuals initially infected with a single arbovirus become infected with a

second. Eqs. 16-19 describe the process of acquiring a second infection for individuals who have

completely recovered from an initial infection, with possible cross-protective immunity.

$$\lambda_1 = mab(I_1^M + p_1 I_{12}^M) \qquad (1)$$
$$\lambda_2 = mab(I_2^M + p_2 I_{12}^M) \qquad (2)$$
$$\lambda_{12} = mabp_{12}I_{12}^M \qquad (3)$$
$$\frac{dS}{dt} = -(\lambda_1 + \lambda_2 + \lambda_{12})S \qquad (4)$$
$$\frac{dE_1}{dt} = \lambda_1 S - ((\lambda_2 + \lambda_{12})(1 - \alpha_1) + \frac{1}{\epsilon^H})E_1 \quad (5)$$
$$\frac{dE_2}{dt} = \lambda_2 S - ((\lambda_1 + \lambda_{12})(1 - \alpha_2) + \frac{1}{\epsilon^H})E_2 \quad (6)$$
$$\frac{dE_{12}^C}{dt} = \lambda_{12} S + \lambda_{12}((1 - \alpha_1)E_1 + (1 - \alpha_2)E_2) - \frac{1}{\epsilon^H}E_{12}^C \qquad (7)$$
$$\frac{dE_{12}^S}{dt} = \lambda_2(1 - \alpha_1)E_1 + \lambda_1(1 - \alpha_2)E_2 - \frac{1}{\epsilon^H}E_{12}^S \qquad (8)$$
$$\frac{dI_1}{dt} = \frac{1}{\epsilon^H}E_1 - \frac{1}{r}I_1 \qquad (9)$$
$$\frac{dI_2}{dt} = \frac{1}{\epsilon^H}E_2 - \frac{1}{r}I_2 \qquad (10)$$
$$\frac{dI_{12}^C}{dt} = \frac{1}{\epsilon^H}E_{12}^C - \frac{1}{r}I_{12}^C \quad (11)$$
$$\frac{dI_{12}^S}{dt} = \frac{1}{\epsilon^H}E_{12}^S - \frac{1}{r}I_{12}^S \quad (12)$$
$$\frac{dR_1}{dt} = \frac{1}{r}I_1 - (\lambda_{12} + \lambda_2)(1 - \alpha_1)R_1 \qquad (13)$$
$$\frac{dR_2}{dt} = \frac{1}{r}I_2 - (\lambda_{12} + \lambda_1)(1 - \alpha_2)R_2 \qquad (14)$$
$$\frac{dR_{12}}{dt} = \frac{1}{r}(I_{1,2} + I_{2,1} + I_{12}^C + I_{12}^S) \qquad (15)$$
$$\frac{dE_{1,2}}{dt} = (\lambda_{12} + \lambda_2)(1 - \alpha_1)R_1 - \frac{1}{\epsilon^H}E_{1,2} \qquad (16)$$
$$\frac{dE_{2,1}}{dt} = (\lambda_{12} + \lambda_1)(1 - \alpha_2)R_2 - \frac{1}{\epsilon^H}E_{2,1} \qquad (17)$$
$$\frac{dI_{1,2}}{dt} = \frac{1}{\epsilon^H}E_{1,2} - \frac{1}{r}I_{1,2} \qquad (18)$$
$$\frac{dI_{2,1}}{dt} = \frac{1}{\epsilon^H}E_{2,1} - \frac{1}{r}I_{2,1} \qquad (19)$$
168

Model equations for mosquitoes are as follows, with parameter values and meanings shown in

Tables 1-2. Eqs. 20-22 define the forces of infection for each infection for each virus and for co-

transmitted viruses, which involves input from human states and transmission probabilities. Eqs.

26 and 29 describe mosquito co-infection, which is driven by temperature-dependent parameters.

Eqs. 30-31 outline how mosquitoes in this model become infected with a second arbovirus after

recovering from their first infection.

175
$$\lambda_1^M = ac(I_1 + I_{2,1} + p_1^M(I_{12}^C + I_{12}^S)) \qquad (20)$$

176
$$\lambda_2^M = ac(I_2 + I_{1,2} + p_2^M(I_{12}^C + I_{12}^S)) \qquad (21)$$

177
$$\lambda_{12}^M = acp_{12}^M(I_{12}^C + I_{12}^S) \qquad (22)$$

178
$$\frac{dS^M}{dt} = g(E_1^M + E_2^M + E_{12}^M + I_1^M + I_2^M + I_{12}^M + I_1E_2^M + I_2E_1^M) - (\lambda_1^M + \lambda_2^M + \lambda_{12}^M)S_M \qquad (23)$$

179
$$\frac{dE_1^M}{dt} = \lambda_1^M S_M - (\lambda_2^M + \lambda_{12}^M + g)E_1^M - \epsilon^M E_1^M \qquad (24)$$

180
$$\frac{dE_2^M}{dt} = \lambda_2^M S_M - (\lambda_1^M + \lambda_{12}^M + g)E_2^M - \epsilon^M E_2^M \qquad (25)$$

181
$$\frac{dE_{12}^M}{dt} = \lambda_{12}^M S_M + (\lambda_2^M + \lambda_{12}^M)E_1^M + (\lambda_1^M + \lambda_{12}^M)E_2^M - gE_{12}^M - \epsilon^M E_{12}^M \qquad (26)$$

182
$$\frac{dI_1^M}{dt} = \epsilon^M E_1^M - \lambda_2^M I_1^M - gI_1^M \qquad (27)$$

183
$$\frac{dI_2^M}{dt} = \epsilon^M E_2^M - \lambda_1^M I_2^M - gI_2^M \qquad (28)$$

184
$$\frac{dI_{12}^M}{dt} = \epsilon^M(E_{12}^M + I_1E_2^M + I_2E_1^M) - gI_{12}^M \qquad (29)$$

185
$$\frac{dI_1E_2^M}{dt} = \lambda_2^M I_1^M - \epsilon^M I_1E_2^M - gI_1E_2^M \qquad (30)$$

186
$$\frac{dI_2E_1^M}{dt} = \lambda_1^M I_2^M - \epsilon^M I_2E_1^M - gI_2E_1^M \qquad (31)$$

Equations to address seasonal fluctuations in temperature and thermal traits across a 365-day

period are as follows, modeled after the approach to seasonal forcing used by Huber et al. (22).

In Eq. 32, T_{max}, T_{min}, and T_{mean} represent the maximum, minimum, and mean temperature for a

region across a calendar year. In Eqs. 33 and 34, c, T_{max}, T_{min}, and T represent the fitted rate

constant, critical temperature maximum, critical temperature minimum, and temperature at a

given time, respectively. As in Mordecai et al., we assumed that values above the critical

maximum and below the critical minimum were zero (25).

196
$$T(t) = \frac{T_{max}-T_{min}}{2}\sin\left(\frac{2\pi}{365}t\right) + T_{mean} \qquad (32)$$

198
$$Q(T) = c(T - T_{min})(T - T_{max}) \qquad (33)$$

199
$$B(T) = cT(T - T_{min})\sqrt{T_{max} - T} \qquad (34)$$

Parameters

Fitted parameters describing *Aedes aegypti* life traits and arbovirus transmission are shown in

Table 1. Temperature dependence of traits was described using quadratic or Briére functions and

fitted to experimental data (22,25). The value of the parameters describing these traits varies in

our model as seasonal temperatures fluctuate.

Table 1: Temperature-dependent parameters describing fitted *Aedes aegypti* life traits and arbovirus transmission (22).

Parameter	Definition	Function	Fitted Parameters		
a	Mosquito biting rate	Briere	c=13.4	$T_{min} = 40.1$	$T_{max} = 2.0 \times 10^{-4}$
b	Probability that an infected mosquito transmits to a human during feeding	Briere	c=17.1	$T_{min} = 35.8$	$T_{max} = 8.5 \times 10^{-4}$
c	Probability that a mosquito becomes infected after feeding on an infected human	Briere	c = 12.2	$T_{min} = 37.5$	$T_{max} = 4.9 \times 10^{4}$
g	Mosquito mortality rate	Quadratic	c = 9.2	$T_{min} = 37.7$	$T_{max} = -1.5 \times 10^{-1}$
ε^M	Virus extrinsic incubation rate	Briere	c = 10.7	$T_{min} = 45.9$	$T_{max} = 6.7 \times 10^{-5}$

Additional population-level parameters were governed by temperature. To ensure that the ratio

of mosquitoes to humans, *m(T)*, remained biologically feasible regardless of climate, we

followed the approach of Siraj et al. (26) and developed a mosquito ratio scaling factor, γ, such

that

$$\gamma = 1.24 g\left(T_{Tropical}\right) \qquad (35)$$

and

$$m = \frac{\gamma}{g(T)} \qquad (36).$$

γ is defined here as the product of the estimated ratio of mosquitos to humans in Rio de Janeiro

in 2012 (1.24) and the temperature-varying fitted mosquito mortality rate at temperatures typical

of that city, such that $T_{Tropical} = 24.3$ (Eq. 35) (22,26,27). Using this value to scale m across

various temperature environments ensured that the ratio of mosquitoes to humans remained

biologically feasible, between 1.18 and 1.25 over the course of each simulation (Eq. 36).

Temperature-independent parameter values and definitions are consistent with those in Vogels et

al. (8), which developed a generic model of arbovirus co-infection that ours is built on.

Transmission parameters pertaining to *Aedes aegypti* mosquitoes follow those used in previous

dengue modeling studies (16,28,29) while transmission parameters pertaining to co-transmission

from co-infected humans to mosquitoes were informed by data from Rückert et al. (9).

228
229   **Table 2: Temperature-independent parameters.**

Parameter	Definition	Value	One-at-a-time sensitivity analysis range	Source(s)
α	Coefficient of cross-protection	0-1, varied	0-1	n/a
r	Average time for a human to recover	5 days	3-10 days	(30–32)
p_{12}	Probability that when a co-infected mosquito transmits, it transmits both viruses to a human	0.5	0-1	(8)
p_1	Probability that when a co-infected mosquito transmits, it transmits only virus X to a human	0.25	$0.5(1- p_{12})$	(8)
p_2	Probability that when a co-	0.25	$0.5(1- p_{12})$	(8)

	infected mosquito transmits, it transmits only virus Y to a human			
p_1^M	Probability that when a mosquito becomes infected after feeding on a co-infected human, it only becomes infected by virus X	0.2	n/a	(8,9)
p_2^M	Probability that when a mosquito becomes infected after feeding on a co-infected human, it only becomes infected by virus Y	0.2	n/a	(8,9)
p_{12}^M	Probability that when a mosquito becomes infected after feeding on a co-infected human, it becomes infected by both viruses	0.6	n/a	(8,9)
ε^H	Incubation period (human)	7 days	4-8 days	(28,29,33)

Analyses

Outcomes of interest

We focused on four model outputs: 1) cumulative incidence of infection with virus A, 2)

cumulative incidence of infection with virus B, 3) cumulative incidence of co-infection, and 4)

proportion of all infections that were co-infections. All quantities were defined as cumulative

values across the course of a year-long simulation, at which time all arbovirus outbreaks had run

their course.

One-at-a-time sensitivity analysis of temperature-independent parameters

While the majority of the parameters governing arbovirus infection and co-infection in our

model were temperature-dependent, three were not: cross-protection (α), recovery time (r), and

human incubation period (ε^H). We considered the individual impact of these parameters on

model outputs in a series of one-at-a-time sensitivity analyses. For each parameter, we varied the

value across a plausible range (Table 2) while holding the remaining two temperature-

independent variables constant and examined the relationship between the varied parameter and

selected model outputs. We then repeated this analysis under several assumptions about

population-level preexisting immunity—25% immunity to virus A, 25% immunity to virus B,

and 25% immunity to both—and considered the aforementioned model outputs' response to

these population scenarios.

Explore differing roles of importation time and seasonality

The seasonal component of the model made it possible to examine the effect of seasonal

temperature variation on the cumulative incidence of co-infection, as well as the effect of

temporal synchrony or asynchrony of the co-infecting viruses. We evaluated this effect under

two temperature regimes, defined by mean temperatures and seasonal amplitudes for a given

region and based upon 2019 monthly mean values obtained from Weather Underground

(wunderground.com). These included an environment with temperatures typical of a tropical

region (mean 25.1 °C, amplitude 3.4 °C; similar to Rio de Janeiro), and an environment with

temperatures typical of a more temperate region (mean 13.8 °C, amplitude 14.7 °C; similar to

Beijing). Using monthly mean temperatures for the two cities, we calculated associated mean,

minimum, and maximum temperatures across a year to determine the average temperature and

seasonality observed in the most recent full calendar year. We systematically considered each

possible combination of virus arrival times within a simulation and compared simulation results

between the two temperature settings.

Global sensitivity analysis

To evaluate the interaction components of our model parameters, we conducted a global,

variance-based sensitivity analysis, also known as a Sobol sensitivity analysis, using the SALib

library in Python (34). This approach is not dependent on the presence of monotonic

relationships between input parameters and outputs. With this analysis, we were able to quantify

the amount of variance in the aforementioned model outputs that could be attributed to individual

input parameters, as well as the amount of variance that could be attributed to pairwise and

higher interactions among these parameters. We varied all temperature-independent parameters

and the mean and amplitude of the yearly temperature curve in this analysis. We used the Saltelli

sampling scheme to generate 1.8 million parameter combinations from a range of plausible

values (Table 3) to ensure that we covered the parameter space of biological interest. Our

sensitivity analysis was conducted on the corresponding 1.8 million model outputs, once for each

of four population immunity scenarios: no existing immunity, 25% existing immunity to virus A,

25% existing immunity to virus B, and 25% existing immunity to both infections.

Table 3: Sampling ranges for global sensitivity analysis.

Parameter	Sampling Range

Mean temperature	0 – 36 °C
Mean amplitude	0 – 20 °C
Date of importation of virus A	0 – 365
Human recovery time (r)	3 – 10 days
Human incubation period (ε^H)	5 – 8 days
Virus introduction interval	0 – 75 days
Coefficient of cross-protection (α)	0 – 1
Probability of human co-transmission (p_{12})	0 – 1

RESULTS

The role of virus importation timing, seasonality, and temperature

To better understand what combinations of epidemic timing and seasonality lead to a high

incidence of co-infections, we explored a range of virus arrival times throughout the year under

tropical and temperate temperature regimes. This scenario is reflective of a situation where two

arboviruses are imported to a location within a year of each other, as happened with Zika and

chikungunya viruses in some South American countries in the 2014-2016 period. When seasonal

patterns resembled those in a tropical region (25.1 °C, amplitude 3.4 °C), simultaneous (same-

day) virus importation resulted in incidence of co-infection that was always greater than 19 per

1,000 individuals, although seasonal differences were observed (Fig. 2A-B). Simultaneous

arrival of viruses A and B resulted in co-infection incidence ranging from 32 to 83 per 1,000

individuals per year, with low incidence being associated with periods of significant negative

temperature change, particularly in late summer (days 100-150) (Fig. 2A). Asynchronous virus

arrival resulted in fewer co-infections than simultaneous arrival did, with larger gaps between

virus arrival dates corresponding to lower incidence of co-infection (Fig. 2B).

In contrast, when seasonal patterns resembled those of a more temperate region (mean 13.8 °C,

amplitude 14.7 °C), both simultaneous virus arrival and seasonal high temperatures were

required to observe non-negligible co-infection, with a maximum incidence of 0.0079 per 1,000

individuals (Fig. 2C-D). Asynchronous arbovirus arrival resulted in seasonal trends in co-

infection consistent with simultaneous arrival but produced negligible co-infections as the time

between arrivals grew (Fig. 2D). Arrivals after the summer (around day 100) generated similarly

negligible co-infection incidence, as temperatures fell below those conducive to virus

transmission by *Aedes aegypti* mosquitoes (Fig. 2C).

Figure 2: Seasonal temperature and co-infection responses under different temperature
regimes. Fill colors indicated total co-infections observed in a year (or the attack rate), rather
than an instantaneous measurement, given each possible combination of virus importation dates.
Temperature curves are shown in blue on each axis of the contour plots, and approximate those
in Rio de Janeiro (A-B, 25.1°C, amplitude 3.4°C), and those in Beijing (C-D, mean 13.8°C,
amplitude 14.7°C). Colored lines on each plot show how the attack rate observed in each year
changes given different intervals between virus importation dates.
315

In addition to being temperature-driven, model outputs are influenced by several biologically

important temperature-independent parameters, including immunological cross-protection,

recovery time, and the human incubation period. The coefficient of cross-protection (α) refers to

the level of protection each infection provides against the other, where 0 indicates no protection,

and 1 indicates complete protection. When values of α increased, the incidence of virus A

decreased slightly, and the proportion of co-infections decreased dramatically (Figs. 3A, D). This

decrease in the incidence of virus A when cross-protection was high was driven by individuals

with virus B who experienced a reduced force of infection of virus A and was limited by the later

arrival of virus B into the population. The proportion of co-infections declined steeply as high

values of α inhibit both sequential and co-transmitted co-infections from occurring. However,

high α favors co-transmitted co-infections over sequentially transmitted co-infections because

co-transmission is limited by cross-protection to the extent that sequential transmission is.

Model outputs are also noticeably influenced by changes in the value of recovery time (r), the

average time in days it takes for a human to recover from either infection. Longer recovery times

correspond to higher virus A incidence and a higher proportion of co-infections, as more time

spent infectious allows for greater exposure to a second infection and increases the reproduction

number of both viruses (Figs. 3B, E). Longer recovery times lead to a lower ratio of co-

transmitted co-infections to sequentially transmitted co-infections for the same reason—time

spent infectious, where another infection cannot be acquired immediately, favors sequential

infection transmission (Fig. 3H).

In contrast with the previous two parameters, model outputs do not appear particularly

susceptible to changing incubation period values. Human incubation period (ε^H) is also measured

in days, and has been approximated in studies of dengue and Zika virus to be 5-8 days (28,35).

Within this range, ε^H does not strongly influence the incidence of either single or co-infections

(Figs. 3C, F). However, the seasonal and non-seasonal models produce noticeably different

model outputs observe noticeable differences in seasonal model outputs depending on the day

virus A is imported (Figs. 3C, F, I). When virus A was imported in fall or winter, the incidence

of virus A and the proportion of infections that were co-infections were relatively low as

compared to spring or summer importation days. While both the seasonal and non-seasonal

models have parameters that approximate values for a tropical-like region, earlier importation

dates within the year appear more suitable for arbovirus infection and co-infection, as rising

temperatures following importation reach values ideal for transmission once incidence has

grown.

351
Figure 3: Univariate sensitivity to temperature-independent parameters under seasonal
and non-seasonal models. Varying initial importation dates of virus A were considered when
the seasonal model was used to explore the range of temperature environments possible within a
year. Vertical lines indicate the baseline value for each parameter, and y-axes differ for each
subplot. Panels D-F show the proportion of all infections that are co-infections. Panels G-J show
the ratio of co-transmitted co-infections to sequentially-transmitted co-infections.
358

Role of preexisting immunity

In many tropical environments, transmission of some arboviruses occurs in semi-regular seasonal

cycles, which will impact the incidence of co-infection as part of the population will have prior

immunity to one or more of the viruses. To consider how this might influence co-infections, we

examined the behavior of the model with respect to four scenarios about initial conditions for

population immunity: 1) none; 2) 50% immune to virus A; 3) 50% immune to virus B; and 4)

50% immune to both viruses (the same 50% of the population immune to virus A was also

immune to virus B) (Fig. 4). Scenario 2, in which virus B is introduced to a population with

some immunity to virus A, reflects patterns in immunity similar to those observed when Zika or

chikungunya viruses have been introduced in dengue-endemic settings. We used the non-

seasonal model for this analysis, and we set the time between importation of the two viruses to

30 days. Results from the baseline scenario were equivalent to those from the non-seasonal

scenario in Figure 3.

When high levels of cross-protection were present, immunity to virus B limited infection by

either virus once virus B became prevalent (Figs. 4A, D). This resulted in a decrease in virus A

incidence driven by the lower proportion of individuals not immune to virus B and, thereby,

partially immune to virus A. In contrast, scenarios with immunity to virus A and immunity to

both produced negligible incidence of virus A infections (Figs. A, D), since both limited the

population susceptible to virus A.

Preexisting immunity had the most noticeable impact on the ratio of co-transmitted co-infections

to sequentially transmitted co-infections, where we observed that co-transmitted co-infections

were more heavily represented under the scenario with immunity to both viruses (black line) than

they were under the no-immunity scenario (red dashed line) (Fig. 4J). When there was no initial

immunity, there was a larger group of individuals susceptible to virus A at the beginning, which

provided more opportunities for sequentially-transmitted co-infections. Since that population of

individuals was much smaller when there was immunity to both viruses at the beginning,

sequential transmission occurred less frequently. More generally, though, immunity of any type

led to a much smaller proportion of infections that were co-infections (Figs. 4D-F). While this

proportion was low even under the no-immunity scenario, immunity to even a single virus

limited the occurrence of both sequential and co-transmitted co-infections.

391
Figure 4: Non-seasonal model output in response to varied temperature-independent
parameters under different initial immunity conditions. In each immunity scenario, 50% of
the population are immune to a given virus or viruses, while the baseline immunity scenario
includes no preexisting immunity. Vertical lines indicate the baseline value for each parameter.
Panels D-F show the proportion of all infections that are co-infections. Panels G-J show the ratio
of co-transmitted co-infections to sequentially-transmitted co-infections.
398

Variance-Based Sensitivity Analysis

To gain a holistic view of the effect of the examined biological and environmental factors and

their interactions on the epidemiology of arbovirus co-infection, we used a variance-based

sensitivity analysis. This approach allowed us to explore the contribution of each parameter to

the variance of each model output. Examining the first-order indices, or those describing the

direct relationship between each parameter and each model output, revealed that interactions

between parameters accounted for more than 50% of the total variance in all four outputs (Fig.

5). However, when total-order indices were considered, which measure all contributions of input

parameters to output variance (including interactions), parameters related to temperature and

timing had the greatest effect on the outputs, especially temperature amplitude, temperature

mean, the importation date of virus A, and the importation date of virus B. Together, these four

parameters accounted for 85% of the total variance in the proportion of co-infection we

observed, implying that the timing of outbreaks is by far the most important determinant of the

level of co-infection. Additionally, the parameter governing the time between arbovirus

introductions (virus B importation date) was influential on co-infection-related model outputs

when there was preexisting immunity to virus B, as larger intervals between virus importation

times could severely limit the potential for any temporal overlap between the viruses (Supp. Figs.

2-3). We also observed that the coefficient of cross-protection had a much smaller effect on

incidence of virus A than it did on the other three outputs, due to interactions with the parameters

responsible for temperature and its timing within the year (Fig. 5, Supp. Figs. 1-3).

Figure 5: Output of variance-based sensitivity analysis for baseline immunity scenario (no preexisting immunity). Both first-order (A) and total-order (B) indices are shown. Fill colors indicate the parameter responsible for a given fraction of the variance in a given output. First-order indices sum to one, while total-order indices additionally account for all variance caused by a parameter's interactions and therefore have no such constraint.

We further explored the interactions between parameters that contribute to variance in the

incidence of co-infection and found these relationships to be consistent with the results from

first- and total-order indices alone (Fig. 6). Interactions between mean temperature and other

429   timing and temperature-related parameters, particularly virus A importation date, explained

430   much more variance in this output than did other parameter combinations.

432
433
Figure 6: Total-order variance-based sensitivity analysis parameter interactions for total
co-infection output under the baseline immunity scenario (no preexisting immunity). Black
boxes on the diagonal represent first-order interactions, which were not considered here.

Additionally, we assessed the role that interactions between the viruses themselves play in

observed co-infections, particularly via the mechanism of co-transmission. As our model allows

for simultaneous arbovirus co-transmission between mosquitoes and humans, a phenomenon

documented frequently in laboratory studies (9), the frequency of arbovirus co-infection was

substantially greater than would be expected if there were no interaction between the viruses.

When we did not allow co-transmission from co-infected mosquitoes in our model, the

cumulative incidence of co-infection was 22/1,000 individuals per year, a substantial reduction

from 46/1,000 individuals per year, the cumulative incidence when co-transmission is allowed.

Further, we found that the prevalence of co-infection in our model was always greater than the

product of the prevalences of the two individual viruses (Fig. 7). This suggests that these viruses

have a synergistic relationship brought on by the presence of co-transmission in the model.

449
450   **Figure 7: Prevalence of virus A, virus B, observed co-infections, and expected co-infections**
451   **throughout a year-long simulation of the nonseasonal model, using baseline parameter**
452   **values.** Expected co-infections are defined as the product of the individual prevalences of virus
453   A and B at each time point.
454

DISCUSSION

By incorporating seasonal temperature variation, differential importation times, and virus co-

transmission, the model implemented here explores the drivers of arbovirus co-infection and

assesses conditions under which increased arbovirus co-infection may be likely. To observe

substantial co-infection incidence, our model suggests a need for both the consistently favorable

temperatures typical of the tropics as well as temporal synchrony between the viruses. In more

temperate regions co-infections of the studied arboviruses are rare, only occurring during

summer months and even then at very low levels. Repeated seasonal arbovirus outbreaks could

result in some degree of cross-protective natural immunity in populations living in tropical

environments (13,36,37), which adds an additional layer of complexity to the processes modeled

here. Our results suggest that such preexisting immunity to one or more arboviruses could inhibit

significant co-infection incidence even when environmental and temporal circumstances are

otherwise ideal. However, regardless of immunity, sensitivity analyses indicate that parameters

related to seasonality and timing were the primary contributors to variance in model outputs.

This study provides a novel exploration of temperature variation in mosquito life traits and co-

infection dynamics within a single modeling framework. Previous modeling studies have

characterized the relationship between temperature and arbovirus transmission and have

emphasized that warm climates and highly variable moderate climates have high epidemic

suitability (28,38). Our results concur with this, and further show that, in some cases, seasonal

temperature variation can drive changes in co-infection incidence far more than temperature-

independent population parameters do (Fig. 3). As climate change and human mobility patterns

lead to expanded arbovirus vector ranges (6), arbovirus co-infection may affect increasing

proportions of the global population. The association between higher *Aedes*-borne disease

incidence and higher poverty levels (7) indicates that this could become the subject of

humanitarian concern, to the extent that co-infections might be associated with more severe

outcomes. Increased clinical testing for multiple arboviruses, even if one positive diagnosis has

already been obtained, is necessary to provide more informative data on this spread in the future.

We made several assumptions to support parsimonious and computationally tractable model

scenarios. First, while temporal synchrony between arriving arboviruses was a crucial

component of this study, it is perhaps more likely that an arbovirus would be introduced to an

environment where another arbovirus is already endemic, as has been noted in studies of Zika

virus and endemic dengue in the Americas (15,39). We explored the dynamics of these scenarios

by imposing preexisting immunity to the viruses in turn, as well as simultaneously, and found

that immunity to one or both viruses reduces the incidence of co-infection, particularly in the

presence of substantial cross-immunity. A similar outcome might be expected in real-world

populations where an arbovirus is endemic, although this reduction in incidence of co-infection

could be limited if cross-protection is incomplete. Conversely, situations where two novel

viruses invade in quick succession, as was observed with Zika and chikungunya viruses in South

America in 2013-14 (40) could increase the incidence of co-infection, as model simulations

showed here.

Second, for the sake of simplicity, we assumed that the population was homogenous and well-

mixed. This could lead to an overestimation of transmission, as high-risk clusters are not always

localized both in time and space (17). Incorporating heterogeneity reduces the herd immunity

threshold for a population and can lower the arbovirus reproduction number (R_0), as well

(41,42). Within our model, this could reduce the attack rate and influence the ratio of co-

transmitted co-infections to sequentially-transmitted co-infections. Finally, model parsimony

also influenced our choice of a single mosquito population scaling factor, $\gamma(T)$, a parameter used

to ensure that mosquito populations remained within a reasonable range. While this neglects

many complexities of mosquito population dynamics, our use of simplifying assumptions more

generally made it possible to isolate the effects of parameters of interest in a straightforward

way. Sensitivity analysis of our model allowed us to further examine all parameter interactions

and explore the full parameter space.

In this study, we built upon existing modeling work (8) to explore a range of possible influences

on arbovirus co-infection through a largely theoretical lens. Expansions of this analysis in the

future could benefit from incorporating the growing body of empirical studies exploring the

biological mechanisms and outcomes of arbovirus co-infection, particularly those investigating

cross-protection and antibody-dependent enhancement (13–16). Data on the frequency of

arbovirus co-infection during overlapping epidemics could be informative to this model as well,

as could varying parameters between the modeled viruses. While arbovirus co-infection remains

a growing area of study, significant work has been done on the interactions and outcomes of a

variety of other co-infections, from HIV and tuberculosis (43) to respiratory viral co-infections

(44,45). As such, understanding the dynamics of co-infecting pathogens and the clinical

consequences of co-infection, especially in the context of global change, is of growing

importance for disease mitigation and human health around the world.

ACKNOWLEDGEMENTS

This work was supported by the NIH National Institute of General Medical Sciences R35 MIRA

program to TAP (R35GM143029) and a Richard and Peggy Notebaert Premier Fellowship from

the University of Notre Dame to MLP.

REFERENCES

1.   Epidemiology of Chikungunya in the Americas | The Journal of Infectious Diseases | Oxford Academic [Internet]. [cited 2022 Mar 30]. Available from: https://academic-oup-com.proxy.library.nd.edu/jid/article/214/suppl_5/S441/2632641?login=true

2.   Model-based projections of Zika virus infections in childbearing women in the Americas | Nature Microbiology [Internet]. [cited 2022 Mar 30]. Available from: https://www-nature-com.proxy.library.nd.edu/articles/nmicrobiol2016126

3.   Epidemiology of dengue: past, present and future prospects - PMC [Internet]. [cited 2022 Mar 30]. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3753061/

4.   Messina JP, Brady OJ, Golding N, Kraemer MUG, Wint GRW, Ray SE, et al. The current and future global distribution and population at risk of dengue. Nat Microbiol. 2019 Sep;4(9):1508–15.

5.   Messina JP, Kraemer MU, Brady OJ, Pigott DM, Shearer FM, Weiss DJ, et al. Mapping global environmental suitability for Zika virus. Jit M, editor. eLife. 2016 Apr 19;5:e15272.

6.   Kraemer MUG, Reiner RC, Brady OJ, Messina JP, Gilbert M, Pigott DM, et al. Past and future spread of the arbovirus vectors Aedes aegypti and Aedes albopictus. Nature Microbiology. 2019 May;4(5):854–63.

7.   Morgan J, Strode C, Salcedo-Sora JE. Climatic and socio-economic factors supporting the co-circulation of dengue, Zika and chikungunya in three different ecosystems in Colombia. PLOS Neglected Tropical Diseases. 2021 Mar 11;15(3):e0009259.

8.   Vogels CBF, Rückert C, Cavany SM, Perkins TA, Ebel GD, Grubaugh ND. Arbovirus coinfection and co-transmission: A neglected public health concern? PLOS Biology. 2019 Jan 22;17(1):e3000130.

9.   Rückert C, Weger-Lucarelli J, Garcia-Luna SM, Young MC, Byas AD, Murrieta RA, et al. Impact of simultaneous exposure to arboviruses on infection and transmission by *Aedes* aegypti mosquitoes. Nature Communications. 2017 May;8:ncomms15412.

10. Mercado-Reyes M, Acosta-Reyes J, Navarro-Lechuga E, Corchuelo S, Rico A, Parra E, et al. Dengue, chikungunya and zika virus coinfection: results of the national surveillance during the zika epidemic in Colombia. Epidemiology and Infection [Internet]. 2019 [cited 2019 Feb 19];147. Available from: https://www.cambridge.org/core/product/identifier/S095026881800359X/type/journal_article

11. Furuya-Kanamori L, Liang S, Milinovich G, Soares Magalhaes RJ, Clements ACA, Hu W, et al. Co-distribution and co-infection of chikungunya and dengue viruses. BMC Infectious Diseases. 2016;16:84.

12.   Lobkowicz L, Ramond A, Clemente NS, Ximenes RA de A, Miranda-Filho D de B,
Montarroyos UR, et al. The frequency and clinical presentation of Zika virus coinfections: a
systematic review. BMJ Global Health. 2020 May 1;5(5):e002350.

13.   Gordon A, Gresh L, Ojeda S, Katzelnick LC, Sanchez N, Mercado JC, et al. Prior dengue
virus infection and risk of Zika: A pediatric cohort in Nicaragua. PLoS Med [Internet].
2019 Jan 22 [cited 2021 Mar 19];16(1). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6342296/

14.   Subramaniam KS, Lant S, Goodwin L, Grifoni A, Weiskopf D, Turtle L. Two Is Better
Than One: Evidence for T-Cell Cross-Protection Between Dengue and Zika and
Implications on Vaccine Design. Front Immunol [Internet]. 2020 [cited 2021 May 7];11.
Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2020.00517/full

15.   Rodriguez-Barraquer I, Costa F, Nascimento EJM, Nery N, Castanha PMS, Sacramento
GA, et al. Impact of preexisting dengue immunity on Zika virus emergence in a dengue
endemic region. Science. 2019 Feb 8;363(6427):607–10.

16.   Andrade DV, Harris E. Recent advances in understanding the adaptive immune response to
Zika virus and the effect of previous flavivirus exposure. Virus Res. 2018 Aug 2;254:27–
33.

17.   Freitas LP, Cruz OG, Lowe R, Sá Carvalho M. Space–time dynamics of a triple epidemic:
dengue, chikungunya and Zika clusters in the city of Rio de Janeiro. Proc Biol Sci
[Internet]. 2019 Oct 9 [cited 2021 Mar 19];286(1912). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6790786/

18.   Couret J, Dotson E, Benedict MQ. Temperature, larval diet, and density effects on
development rate and survival of Aedes aegypti (Diptera: Culicidae). PLoS One.
2014;9(2):e87468.

19.   Kamimura K, Matsuse IT, Takahashi H, Komukai J, Fukuda T, Suzuki K, et al. Effect of
temperature on the development of Aedes aegypti and Aedes albopictus. Medical
Entomology and Zoology. 2002;53(1):53–8.

20.   Marinho RA, Beserra EB, Bezerra-Gusmão MA, Porto V de S, Olinda RA, Dos Santos
CAC. Effects of temperature on the life cycle, expansion, and dispersion of Aedes aegypti
(Diptera: Culicidae) in three cities in Paraiba, Brazil. J Vector Ecol. 2016 Jun;41(1):1–10.

21.   Geoghegan JL, Walker PJ, Duchemin J-B, Jeanne I, Holmes EC. Seasonal drivers of the
epidemiology of arthropod-borne viruses in Australia. PLoS Negl Trop Dis. 2014
Nov;8(11):e3325.

22.   Huber JH, Childs ML, Caldwell JM, Mordecai EA. Seasonal temperature variation
influences climate suitability for dengue, chikungunya, and Zika transmission. PLOS
Neglected Tropical Diseases. 2018 May 10;12(5):e0006451.

23.   Li Z, Wang J, Cheng X, Hu H, Guo C, Huang J, et al. The worldwide seroprevalence of
DENV, CHIKV and ZIKV infection: A systematic review and meta-analysis. PLOS
Neglected Tropical Diseases. 2021 Apr 28;15(4):e0009337.

24.   Göertz GP, Vogels CBF, Geertsema C, Koenraadt CJM, Pijlman GP. Mosquito co-infection
with Zika and chikungunya virus allows simultaneous transmission without affecting vector
competence of Aedes aegypti. PLOS Neglected Tropical Diseases. 2017
Jun;11(6):e0005654.

25.   Mordecai EA, Cohen JM, Evans MV, Gudapati P, Johnson LR, Lippi CA, et al. Detecting
the impact of temperature on transmission of Zika, dengue, and chikungunya using
mechanistic models. PLOS Neglected Tropical Diseases. 2017 Apr 27;11(4):e0005568.

26.   Siraj AS, Oidtman RJ, Huber JH, Kraemer MUG, Brady OJ, Johansson MA, et al.
Temperature modulates dengue virus epidemic growth rates through its effects on
reproduction numbers and generation intervals. Althouse B, editor. PLoS Negl Trop Dis.
2017 Jul 19;11(7):e0005797.

27.   Massad E, Amaku M, Coutinho FAB, Struchiner CJ, Lopez LF, Wilder-Smith A, et al.
Estimating the size of Aedes aegypti populations from dengue incidence data: Implications
for the risk of yellow fever outbreaks. Infect Dis Model. 2017 Dec 8;2(4):441–54.

28.   Fourié T, Grard G, Leparc-Goffart I, Briolant S, Fontaine A. Variability of Zika Virus
Incubation Period in Humans. Open Forum Infect Dis [Internet]. 2018 Nov 1 [cited 2020
Oct 2];5(11). Available from: https://academic.oup.com/ofid/article/5/11/ofy261/5128777

29.   Rudolph KE, Lessler J, Moloney RM, Kmush B, Cummings DAT. Incubation Periods of
Mosquito-Borne Viral Infections: A Systematic Review. Am J Trop Med Hyg. 2014 May
7;90(5):882–91.

30.   Burattini MN, Chen M, Chow A, Coutinho F a. B, Goh KT, Lopez LF, et al. Modelling the
control strategies against dengue in Singapore. Epidemiol Infect. 2008 Mar;136(3):309–19.

31.   Flasche S, Jit M, Rodríguez-Barraquer I, Coudeville L, Recker M, Koelle K, et al. The
Long-Term Safety, Public Health Impact, and Cost-Effectiveness of Routine Vaccination
with a Recombinant, Live-Attenuated Dengue Vaccine (Dengvaxia): A Model Comparison
Study. PLOS Medicine. 2016 Nov 29;13(11):e1002181.

32.   Newton EA, Reiter P. A model of the transmission of dengue fever with an evaluation of
the impact of ultra-low volume (ULV) insecticide applications on dengue epidemics. Am J
Trop Med Hyg. 1992 Dec;47(6):709–20.

33.   Chan M, Johansson MA. The Incubation Periods of Dengue Viruses. PLoS One [Internet].
2012 Nov 30 [cited 2020 Oct 2];7(11). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3511440/

34.   Herman J, Usher W. SALib: An open-source Python library for Sensitivity Analysis. JOSS.
2017 Jan 10;2(9):97.

35.   Siler JF, Hall MW, Hitchens AP. Dengue: Its History, Epidemiology, Mechanism of
Transmission, Etiology, Clinical Manifestations, Immunity, and Prevention. Philipp J Sci
[Internet]. 1926 [cited 2021 Feb 11];29(1–2). Available from:
https://www.cabdirect.org/cabdirect/abstract/19261000360

36.   Katzelnick LC, Bos S, Harris E. Protective and enhancing interactions among dengue
viruses 1-4 and Zika virus. Curr Opin Virol. 2020 Aug;43:59–70.

37.   Ribeiro GS, Kikuti M, Tauro LB, Nascimento LCJ, Cardoso CW, Campos GS, et al. Does
immunity after Zika virus infection cross-protect against dengue? The Lancet Global
Health. 2018 Feb 1;6(2):e140–1.

38.   Burt FJ, Chen W, Miner JJ, Lenschow DJ, Merits A, Schnettler E, et al. Chikungunya virus:
an update on the biology and pathogenesis of this emerging pathogen. The Lancet
Infectious Diseases. 2017 Apr 1;17(4):e107–17.

39.   Borchering RK, Huang AT, Mier-y-Teran-Romero L, Rojas DP, Rodriguez-Barraquer I,
Katzelnick LC, et al. Impacts of Zika emergence in Latin America on endemic dengue
transmission. Nature Communications. 2019 Dec 16;10(1):5730.

40.   WHO | Zika: the origin and spread of a mosquito-borne virus [Internet]. WHO. World
Health Organization; [cited 2021 Mar 19]. Available from:
http://www.who.int/bulletin/online_first/16-171082/en/

41.   Ferrari MJ, Bansal S, Meyers LA, Bjørnstad ON. Network frailty and the geometry of herd
immunity. Proceedings of the Royal Society B: Biological Sciences. 2006 Nov
7;273(1602):2743–8.

42.   Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population
heterogeneity on herd immunity to SARS-CoV-2. Science. 2020 Aug 14;369(6505):846–9.

43.   Pawlowski A, Jansson M, Sköld M, Rottenberg ME, Källenius G. Tuberculosis and HIV
Co-Infection. PLOS Pathogens. 2012 Feb 16;8(2):e1002464.

44.   Scotta MC, Chakr VCBG, de Moura A, Becker RG, de Souza APD, Jones MH, et al.
Respiratory viral coinfection and disease severity in children: A systematic review and
meta-analysis. J Clin Virol. 2016 Jul;80:45–56.

45.   Asner SA, Science ME, Tran D, Smieja M, Merglen A, Mertz D. Clinical Disease Severity
of Respiratory Viral Co-Infection versus Single Viral Infection: A Systematic Review and
Meta-Analysis. PLOS ONE. 2014 Jun 16;9(6):e99392.

 SUPPLEMENTARY MATERIALS

668
669 **Figure S1: Output of variance-based sensitivity analysis for virus A immunity scenario**
670 **(50% preexisting immunity to virus A).** Both first order (A) and total order (B) indices are
671 shown.

672
673 **Figure S2: Output of variance-based sensitivity analysis for virus B immunity scenario**
674 **(50% preexisting immunity to virus B).** Both first order (A) and total order (B) indices are
675 shown.

676
677 **Figure S3: Output of variance-based sensitivity analysis for dual virus immunity scenario**
678 **(50% preexisting immunity to both viruses).** Both first order (A) and total order (B) indices
679 are shown.