**TITLE**

Accurate Screening for Early-Stage Breast Cancer by Detection and Profiling of

Circulating Tumor Cells

**SHORT TITLE**

CTC Based Breast Cancer Screening

**AUTHORS**

Timothy Crook[1], Darshana Patil[2], Dadasaheb Akolkar[2], Anantbhushan Ranade[3],

Sewanti Limaye[4], Raymond Page[5], Vineet Datta[2], Pradip Fulmali[2], Sachin Apurwa[2],

Stefan Schuster[6], Ajay Srinivasan[2*], Rajan Datar[2]

**AFFILIATIONS**

[1]Leaders in Oncology Care, London, UK

[2]Datar Cancer Genetics, Nasik, India

[3]Avinash Cancer Clinic, Pune, India

[4]Sir HN Reliance Foundation Hospital and Research Centre, Mumbai, India

[5]Worcester Polytechnic Institute, Worcester, USA

[6]Datar Cancer Genetics Europe GmbH, Eckersdorf, Germany

**CORRESPONDING AUTHOR**

Dr. Ajay Srinivasan. E-mail: ajays@datarpgx.org, Phone: +91-253-6690803.

**KEYWORDS**

Breast Cancer; Screening; Circulating Tumor Cells; Immunocytochemistry

**ABSTRACT**

BACKGROUND:

Screening of asymptomatic women for early detection of Breast Cancer (BrC) is

associated with improved survival. Presently, mammography is the standard of care

(SoC) for BrC screening but has lower accuracy for invasive cancers as well as in

women with higher breast tissue density.

METHODS:

In this manuscript, we describe an accurate blood-based breast cancer detection test

based on functional enrichment of breast adenocarcinoma associated circulating

tumor cells (BrAD-CTCs) and their identification via multiplexed fluorescence

immunocytochemistry (ICC) profiling for GCDFP15, GATA3, EpCAM, PanCK and

CD45 status.

RESULTS:

The test accurately detects BrAD-CTCs in breast cancers irrespective of age,

ethnicity, disease stage, grade and hormone receptor status. Analytical validation

established the high accuracy and reliability of the test under intended use

conditions. A case-control study with samples from 9,632 healthy women and 548

known BrC cases established 100% specificity and 92.07% overall sensitivity; stage-

wise sensitivities were 70.00% for Stage 0, 89.36% for Stage I, 95.74% for Stage II

and 100% for Stage III and Stage IV cancers. In a prospective clinical study with 141

suspected cases of breast cancer who underwent a biopsy after blood collection, the

test showed 93.1% specificity and 94.64% overall sensitivity in differentiating breast

cancer cases (n = 112) from those with benign breast conditions (n = 29); stage-wise

sensitivities were 87.50% for Stage 0, 95.83% for Stage I and Stage II, 95.00% for

Stage III and 100% for Stage IV cancers.

CONCLUSION:

The findings reported in this manuscript support the clinical potential of this test for

blood based BrC detection.

**BACKGROUND**

Breast Cancer (BrC) is the most common malignancy and a leading cause of cancer related mortality among women globally (1). Although mammography is the standard of BrC screening in asymptomatic females, there is a need for improved BrC detection which addresses the risks and limitations of mammography, such as radiation exposure, lower specificity in differentiating benign conditions from malignancy, lower sensitivity for invasive carcinomas, as well as incompatibility with dense breast tissue. Circulating tumor analytes in peripheral blood have been evaluated for potential application in more accurate, non-radiological and non- / minimally invasive screening for breast cancer. Circulating tumor cells (CTCs) are an ideal analyte for detection of cancers since they are intact malignant cells which harbour the imprint of the parent tumor. CTCs have distinct advantages over nucleic acid fragments or serum antigens since the latter may also be released by non-malignant cells and are associated with lower sensitivity and specificity respectively. There is evidence of sufficient viable CTCs being released into blood even during early stages of carcinogenesis. In breast cancer, it is reported that angiogenesis commences at the DCIS stage itself which can facilitate the dissemination of tumor cells (2–4). Such disseminated tumor cells (DTCs) have been reported in bone marrow of 20% to >50% of patients with DCIS or DCIS with microinvasion, respectively (5–7). Prior studies have also indicated high detection rates of CTCs in blood samples of patients with early-stage breast cancers. Using nanostructured coated slides, Krol et al (8) reported 62.5% CTC detection rate for Stage I and II BrC. Using filtration-based devices, Reduzzi et al (9) showed 76% CTC detection rate in early-stage breast cancer. Similarly, Jin et al (10) used the CytoSorter® CTC capture system and showed 50% and >80% sensitivity in DCIS and Stage I / II BrC. These studies support the biological plausibility of CTC-based cancer screening approaches. Although CTCs have been evaluated for cancer detection, the inability of prior technologies to effectively enrich and harvest sufficient CTCs has hindered meaningful downstream applications. Most prior attempts at evaluating CTCs for cancer screening have been based on epitope capture using the CellSearch platform which, while not approved for CTC detection, finds frequent research use. Several prior studies have highlighted the lower performance of epitope capture arising due to its inability to efficiently harvest or detect CTCs with lower expression of EpCAM and PanCK which are the most routinely employed target markers (11–17). We have

previously described a novel functional enrichment method with high CTC detection sensitivity, which yields sufficient CTCs for downstream applications such as immunocytochemistry (ICC) profiling (18,19). In this manuscript we describe the validation of this technology for use as BrC detection test. We show that the technology has vastly improved CTC detection sensitivity even in stage 0 BrC (DCIS) and addresses several limitations of prior CTC based cancer detection efforts.

**METHODS**

**Study Participants and Samples**

Samples for method development and validation were obtained from participants in two ongoing observational studies of the sponsor, TRUEBLOOD (WHO ICTRP https://trialsearch.who.int/?TrialID=CTRI/2019/03/017918, CTRI http://ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=31879) and RESOLUTE (WHO ICTRP https://trialsearch.who.int/?TrialID=CTRI/2019/01/017219, CTRI http://ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=30733) the design of which were intended to support the identification and characterization of blood-based malignant tumor derived analytes for non- / minimally invasive cancer detection. The TRUEBLOOD study enrolled known cases of cancers and symptomatic individuals suspected of solid organ cancers. The RESOLUTE study enrolled asymptomatic adults with no prior diagnosis of cancer, no current symptoms or findings suspected of cancer and only age associated risk of cancer. Both studies were approved by the Ethics Committees of the sponsor (Datar Cancer Genetics, DCG) as well as the participating institutes and were performed in accordance with the Declaration of Helsinki. Fifteen millilitres of peripheral blood were collected from all enrolled study participants in EDTA vacutainers after obtaining written informed consent. Where possible, tissue samples were also obtained from TRUEBLOOD study participants posted for a biopsy as per standard of care (SoC) procedures (tissue samples were used for method development). In addition, leftover blood samples from suspected or known (recently diagnosed or pre-treated) cancer patients who availed of the study sponsor's commercial services for cancer management as well as healthy (asymptomatic) volunteers at the study sponsor's organization were also obtained after due consent. Blood samples (15 mL) from suspected cases of cancers were collected prior to the patients undergoing an invasive biopsy. All biological samples

were assigned alphanumeric barcodes and stored at 2°C - 8°C during transport to
reach the clinical laboratory within 46 h. Sample blinding avoided systematic
differences between groups due to (un)known baseline variables that could affect the
test findings and also eliminated potential biases that could have otherwise arisen
due to operator's knowledge of the sample. From the originally collected 15 mL
blood samples, a 5 mL aliquot was set aside for processing (CTC enrichment and
ICC profiling) as part of clinical studies. The remaining blood samples were used for
various method development studies. All samples were processed at the CAP and
CLIA accredited facilities of the Study Sponsor Datar Cancer Genetics, which also
adhere to quality standards ISO 9001:2015, ISO 27001:2013 and ISO 15189:2012.
11

**Enrichment of Circulating Tumor Cells from Peripheral Blood**
13  Aliquoted blood samples (5 mL) were processed for the enrichment of CTCs from
14  peripheral blood mononuclear cells (PBMC) as described previously (20).
15  Comprehensive details are provided in Supplementary Materials.
16

**Immunocytochemistry Profiling of CTCs**
The process of ICC profiling of CTCs was as described previously (19).
Comprehensive details are provided in Supplementary Materials. Figure 1 is a
Schema of the test showing the various steps in CTC enrichment and identification
by ICC profiling for various markers. The Decision Matrix for assigning samples as
Positive, Equivocal or Negative based on the findings of ICC profiling is provided in
Figure 2.
24

**Method Development and Optimization**
26  Comprehensive details of method development and optimization studies are
27  provided in the Supplementary Materials.
28

**Analytical Validation**
Analytical validation established the performance characteristics of the test with
standard analyte (SKBR3 cells), spiked into healthy donor blood to generate various
dilutions (cell densities). These dilutions were processed as per the described
procedures (proprietary differentially cytotoxic medium treatment and ICC profiling)

to determine the yield of spiked cells. Comprehensive details of analytical validation
studies are provided in the Supplementary Materials.

**Case Control Clinical Study**

The ability of the Test to discern / identify BrC from asymptomatic individuals was
initially ascertained and established in a case control study with 548 females who
were recently diagnosed, therapy naïve cases of BrC and 9,632 healthy females with
no prior diagnosis of any cancer, no current suspicion of any cancer and with
BIRADS-I on a mammogram, i.e., no evidence of breast cancer (Supplementary
Table S1). Initially, samples in the Asymptomatic cohort were randomized into
Training and Test Sets in a 70%:30% ratio. The BrC cases were first segregated by
Stage (0 – IV) and the samples per stage were then assigned to Training and Test
Sets in a 70%:30% ratio. The Training Set samples (384 BrC and 6742 cancer-free
females) was initially evaluated with the analysts unblinded to the status of the
samples to determine the concordance between the clinical status and the
interpretation of the marker status based on the Decision Matrix. Then the blinded
Test Set comprising of 164 BrC and 2890 cancer-free females' samples was
evaluated to determine the performance characteristics. Subsequently all Training
and Test samples (BrC and healthy) were shuffled and random 30% samples (with
stage-wise for cancer) were selected for analysis as Test Set Iteration 2. This
shuffling step was repeated to obtain 20 iterations of the Test Set. From these
iterative 20 sets, median and range of sensitivity, specificity and accuracy were
determined.

**Prospective Clinical Study**

The performance characteristics of the test was next ascertained and established in
a prospective blinded study of 141 individuals with clinical symptoms or radiological
findings, who were referred for a biopsy due to suspicion of breast cancer
(Supplementary Table S2). All participants provided blood sample prior to the biopsy.
The sponsor was blinded to the diagnosis, i.e., the findings of the histopathological
examination (HPE). Samples were prospectively accrued in this study until 24
samples each were obtained for Stage 0, I and II, 20 samples each were accrued for
Stage III and IV and 29 samples were accrued for individuals with benign findings.
Clinical status of samples (cancer / benign) was revealed to sponsor only after

sample analysis was complete and test findings shared with the clinical study

investigator. From these samples, performance characteristics including sensitivity,

specificity and accuracy were determined with equivocal findings considered as

positive and as negative, respectively.

**Molecular Concordance Study**

In a combined subset of 61 samples from the case control and prospective cohorts,

where matched tumor tissue and blood samples were available, a molecular

concordance study was performed. Tumor Tissue DNA (ttDNA) was isolated and

profiled by Next Generation Sequencing (NGS) using the Ion Proton Platform and

the Comprehensive Ampliseq Multi (409)-gene Cancer Panel. Simultaneously,

PBMCs were isolated from the matched blood samples and used for CTC

enrichment. On the 5$^{th}$ day, genomic DNA (gDNA) isolated from all surviving cells

was evaluated by a ddPCR assay specific to the driver mutation on a BioRad QX200

platform. Concordance between tumor tissue and CTCs was determined as the

proportion of the latter where the corresponding gene variant was detected by

ddPCR.

**RESULTS**

**Method Development and Optimization**

The method development and optimization studies showed the viability of

multiplexed fluorescence analysis of markers with minimal or no cross-interference

of markers as well as the ability to detect CTCs with much lower marker expression

than primary tumor cells or reference cell lines.  Additionally the studies also showed

the capability of the test in detecting CTCs irrespective of patient age, ethnicity,

cancer stage, tumor grade, subtype and hormone receptor status. Findings of

method development and optimization are provided in the Supplementary Materials.

**Analytical Validation**

The analytical validation studies established the analyte stability and also

demonstrated the high sensitivity and specificity of the test as well as significant

linear characteristics in addition to high precision. The sensitivity of the test was not

adversely affected in presence of potentially interfering substances or by controlled

variations to operating parameters. Findings of analytical validation which

established these performance characteristics of the test are provided in the Supplementary Materials. The summary of the analytical validation studies is provided in Table 1.

**Case Control Clinical Study**

We evaluated the performance characteristics of the test in two clinical studies. In the clinical study which was a case control cross-validation study, the median stage wise sensitivities were as follows: 70% for Stage 0, 89.36% for Stage I, 95.74% for Stage II, 100% for Stage III,100% for Stage IV and 92.07% overall. In absence of any positive of equivocal findings in the control (cancer free and asymptomatic) cohort, the specificity of the test (cancer v/s healthy) was 100%. Cancer samples (cases) with equivocal findings were considered as positive for determination of sensitivity and accuracy. Table 2 provides the specificity as well as median of stage-wise and cumulative sensitivity and accuracy across the 20 iterations. Details of this iteration analysis are provided in Supplementary Table S3. Sensitivity and accuracy were also determined with samples with equivocal findings being considered as negative. These findings are presented in Supplementary Table S4, which also indicates the stage-wise and cumulative range of sensitivity and accuracy.

**Prospective Clinical Study**

The second study was an independently conducted blinded prospective study. Of the total 141 individuals from whom samples were collected, there were 112 breast cancer cases (Stages 0 – IV) and 29 cases of various benign breast conditions. There were no samples with equivocal findings in the cancer cohort hence the overall sensitivity was 94.6% with stage wise sensitivities being 87.5% for Stage 0, 95.8% for Stage I, 95.8% for Stage II, 95.0% for Stage III and 100% for Stage IV. Two samples with equivocal findings were diagnosed with benign conditions of the breast. In absence of follow-up data indicating if these cases were indeed subsequently diagnosed with breast cancer, the samples were considered as false positives (worst case scenario), based on which the specificity of the test (cancer v/s benign) was determined to be 93.1%. When samples with equivocal findings were considered as negative, the specificity of the test (cancer v/s benign) was 100% (best case scenario). The sample-wise details of the prospective validation cohort findings are provided in Supplementary Table S5. The stage wise and cumulative

sensitivity and accuracy for both these scenarios are provided in Supplementary

Table S6.

**Molecular Concordance Study**

We identified a subset of 61 samples where driver mutations (allele frequency >

0.14) were detected by NGS in tumor tissue; for variants detected in 53 samples a

specific TaqMan ddPCR assay was available. CTC-enriched fraction from these

samples was used for gDNA isolation which in turn was evaluated by a ddPCR

assay specific to the driver mutation on a BioRad QX200 platform. Variants in ttDNA

detected by NGS were also detected by ddPCR in 81.1% of CTCs indicating

significant concordance (Supplementary Table S7).

**DISCUSSION**

We describe a blood test for BrC detection in asymptomatic women based on

multiplexed fluorescence ICC profiling of CTCs in peripheral blood. The test can

accurately determine the presence of CTCs in BrC irrespective of stage, grade,

subtype, age, ethnicity or hormone receptor status (Supplementary materials).

Analytical validation established high sensitivity, specificity, precision and robustness

in addition to non-interference from endogenous and exogenous factors

(Supplementary materials). Two separate clinical studies established 100%

specificity (cancer v/s asymptomatic) with 92% - 94% overall sensitivity and 70% -

87% Stage 0 sensitivity (Table 2). The test can differentiate samples from cancer

patients and healthy individuals with high (100%) specificity and can also identify

individuals with benign conditions with ≥93% specificity. Our test has (a) high

sensitivity especially for early stages including DCIS, so that the majority of cancers

are detected at localized stages which are amenable to curative resection, and (b)

high specificity so that vast majority of cancer-free individuals do not undergo

additional unnecessary procedures. Our test offers compelling advantages over

screening mammography and is hence a strong candidate for non-invasive BrC

screening in asymptomatic women.

Presently, any benefits of standard mammography screening are largely in

populations aged 50 years and above who have a higher age associated cancer risk

(21–27). Standard 2D digital mammography has been reported to have 73%-87.3%

sensitivity and 86%-96% specificity (28–30). The low accuracy of screening

mammography has been noted in younger women, particularly those aged below 40
years (31). In addition, challenges associated with screening mammography are the
high rates of false positives (7-12% at first mammogram (32) and 50-60% after ten
yearly mammograms (33)). Besides, mammography also has a lower sensitivity for
invasive cancers (76%-85%) than DCIS (83.0%-94.3%) (29,30). Prior studies have
also suggested a modest association between radiation exposure in mammograms
and elevated risk of cancer in BRCA1 mutation carriers (34).

A longitudinal study of screening mammograms in 69,025 women reported 705
cases of screen detected BrC (SBC) and 206 cases of Interval BrC (IBC, not
detected by mammography) (35); the latter were more likely to be of high grade as
well as have higher mortality than SBC. Niraula et al encourage a re-evaluation of
the concept of population-based screening mammography and recommend
exploring strategies beyond conventional screening mammography. In support, the
2018 data available at the US National Centre for Health Statistics, Centres for
Disease Control and Prevention indicates that approximately 27.1% of women in the
age group 50 to 74 years defaulted on SoC mammography (36). Improved BrC
screening and risk-mitigation strategies are hence vital to improve compliance and
BrC detection.

Recent efforts at developing non-invasive cancer screening technologies have
focussed on a multi- or a pan-cancer approach. Notably, GRAIL's Galleri has
introduced the pan-cancer screening test based on methylation profiling in ctDNA
(37). However, the Galleri test has very low sensitivity (<10% - 16%) for Stage I BrC
(38,39), with no data on its ability to detect DCIS. Similarly, the CancerSEEK test
based on simultaneous evaluation of serum proteins and gene variants has ~40%
cumulative sensitivity for early stage and overall ~30% sensitivity for BrC (40).
Purposeful screening for early cancer detection necessitates sensitivity for early
stages (0 – II), which has not been demonstrated by these tests. To the best of our
knowledge, there are no other platforms with high sensitivity and specificity for early
stages (0 – II) of BrC.

The present test is based on detection of CTCs, which are ubiquitously found in the
blood of patients with an underlying solid organ cancer and are undetectable in
healthy individuals (18,19). We show that functionally enriched CTCs can
differentiate malignant conditions from benign conditions as well as identify
asymptomatic individuals with an underlying malignant condition with high specificity.

Obtaining numerically sufficient CTCs is akin to a non-invasive biopsy of the breast

tumor without stromal or other non-tumor content. Since the present test is based on

detection of CTCs which represent the hematogenous phase of carcinomas

associated with a higher risk of progression / metastasis, it is likely that the test has a

higher sensitivity for detecting those sub-populations of DCIS where the risk of

progression is higher.

Traditionally, epitope capture with Anti-EpCAM has been the preferred method for

CTC enrichment. However, several studies have demonstrated the poor CTC

capture / detection rate of this platform (13,14,41). It would be pertinent to mention

that although technologies like CellSearch find frequent mention in research

pertaining to cancer detection, these have not been approved for detection of

cancers based on CTCs. Hence the limitations of these technologies for cancer

detection must be critically understood and proactively addressed to improve CTC

and cancer detection; our test has been developed on such a working hypothesis.

The label- and size-agnostic functional CTC enrichment technique in our test is

immune to the limitations of epitope capture platforms and hence may offer a more

realistic CTC detection rate. In our test, marker expression is determined by a

sensitive high content screening (HCS) system with standardized thresholds to

minimize false negatives. The detection thresholds of the test accommodate CTCs

with significantly lower marker expression (as compared to tumor cells or reference

cell lines), such as those undergoing epithelial to mesenchymal transition (42,43).

The potential benefits of the test include early BrC detection, especially in

asymptomatic women who decline guideline recommended screening

mammography as well as in asymptomatic women for whom the guidelines may not

recommend routine screening mammography (e.g., those below 50 years of age).

The high (>90%) cumulative sensitivity at Stage 0-II indicates a <10% risk of missing

these localized cases where the disease has not spread to other organs and where

the 5-year survival rate is ~99%. For the <10% cases that are not detected at local

stages, subsequent detection at Stage III (regional spread) with >95% reported

sensitivity is still associated with ~86% 5-year survival. The test also has a

significantly higher sensitivity for invasive carcinomas (all stages) than has been

reported for screening mammography (29,30), and can potentially mitigate risks of

IBC (this is yet to be prospectively established). The 100% specificity of the test

indicates a negligible risk of false positives. There are limited or no risks associated

with use of the test since it is non-invasive and is performed on a venous blood draw

of 5 mL of peripheral blood.

The strength of our study stems from the use of an adequately powered sample size

and avoidance of overfitting, since the findings of the iterative validation study

agreed well with that of the training set.

The test has certain limitations in the context of a universal BrC screening. The

sensitivity of the Test was lowest for Stage 0 disease. However, this does not

present any increased risk of false negatives as compared to screening

mammography. Since individuals with potentially false negative findings would not

be deprived of standard mammography screening, it would not add to the pre-

existing risk of the individual. While there is virtually no risk of false positives,

detection of BrAD-CTCs may be construed as false positive in individuals where the

malignancy may not be immediately evident on a standard screening mammogram

or in a biopsy (as was observed in the prospective validation cohort). This risk may

be mitigated by use of a more diagnostically relevant imaging modality or follow-up

among individuals with positive test findings. Minor non-(Adeno) carcinoma subtypes

of breast cancers are not detected by this test. The test has not been evaluated in a

prospective large cohort study in the intent to test asymptomatic population. Finally,

as inherent to any cancer screening test, our test could result in over-diagnosis and

over-treatment.

**CONCLUSION:**

We describe a blood-based, non-invasive test which detects breast AD associated

CTCs with high specificity and sensitivity. The test presents a superior alternative to

mammography screening of asymptomatic women for BrC detection. Approximately

38 million mammograms are performed every year in the US (44) of whom ~280,000

(~0.75%) of cases are diagnosed with BrC (1,45,46). Similarly, of the ~16 million

mammograms performed annually in Europe (47), ~500,000 (1) (~3.1%) are

diagnosed with BrC. Our test has the potential to minimize the need for

mammography screening in individuals with positive findings who could be referred

for diagnostic imaging and work up. The test may also minimize the need for

screening mammography in individuals with negative findings.

**ADDITIONAL INFORMATION**

## ACKNOWLEDGEMENTS

The authors are grateful to the staff of the Study Sponsor (DCG) for their contributions in managing various clinical, operational and laboratory aspects of the study.

## AUTHOR CONTRIBUTIONS

Timothy Crook: Conceptualization, Investigation, Methodology, Resources, Supervision, Writing - review & editing; Darshana Patil: Conceptualization, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing - review & editing; Dadasaheb Akolkar: Conceptualization, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing - review & editing; Vineet Datta: Conceptualization, Investigation, Methodology, Project administration, Resources, Supervision, Writing - review & editing; Pradeep Fulmali: Conceptualization, Formal analysis, Methodology, Project administration, Resources, Supervision, Validation, Visualization, Writing - review & editing; Sachin Apurwa: Data curation, Formal analysis, Methodology, Resources, Software, Validation, Visualization, Writing - review & editing; Stefan Schuster: Conceptualization, Methodology, Writing - review & editing; Ajay Srinivasan: Conceptualization, Data curation, Formal analysis, Methodology, Software, Supervision, Validation, Visualization, Writing - original draft, Writing - review & editing; Rajan Datar: Conceptualization, Funding acquisition, Investigation, Methodology, Resources, Software, Supervision, Validation, Visualization, Writing - original draft, Writing - review & editing

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

All biological samples were obtained from participants in two studies, TRUEBLOOD (WHO ICTRP https://trialsearch.who.int/?TrialID=CTRI/2019/03/017918, CTRI http://ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=31879) and RESOLUTE (WHO ICTRP https://trialsearch.who.int/?TrialID=CTRI/2019/01/017219, CTRI

http://ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=30733). Both studies were approved by Datar Cancer Genetics Limited Institutional Ethics Committee. All participants provided written informed consent. Both studies were performed in accordance with the Declaration of Helsinki.

## CONSENT FOR PUBLICATION

All study participants consented for publication of de-identified biological data. The present manuscript does not contain any personal or identifiable information or data of any participant.

## DATA AVAILABILITY

All relevant data are included in the manuscript and its Supplementary Information file.

## COMPETING INTERESTS

Timothy Crook has no competing interest. Darshana Patil, Dadasaheb Akolkar, Vineet Datta, Pradip Fulmali, Sachin Apurwa, Stefan Schuster and Ajay Srinivasan are employees of the Study Sponsor. Rajan Datar is the founder of the Study Sponsor.

## FUNDING

No external funding was obtained for this study. The entire study was funded by the Study Sponsor (DCG).

## REFERENCES

1.    Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin. 2021 May;71(3):209–49.

2.  Guidi AJ, Fischer L, Harris JR, Schnitt SJ. Microvessel density and distribution in ductal carcinoma in situ of the breast. J Natl Cancer Inst. 1994 Apr;86(8):614–9.

3.  Teo NB, Shoker BS, Jarvis C, Martin L, Sloane JP, Holcombe C. Vascular density and phenotype around ductal carcinoma in situ (DCIS) of the breast. Br J Cancer. 2002 Mar;86(6):905–11.

4.  Heffelfinger SC, Yassin R, Miller MA, Lower E. Vascularity of proliferative breast disease and carcinoma in situ correlates with histological features. Clin cancer Res an Off J Am Assoc Cancer Res. 1996 Nov;2(11):1873–8.

5.  Sänger N, Effenberger KE, Riethdorf S, Van Haasteren V, Gauwerky J, Wiegratz I, et al. Disseminated tumor cells in the bone marrow of patients with ductal carcinoma in situ. Int J cancer. 2011 Nov;129(10):2522–6.

6.  Gruber I V, Hartkopf AD, Hahn M, Taran F-A, Staebler A, Wallwiener D, et al. Relationship Between Hematogenous Tumor Cell Dissemination and Cellular Immunity in DCIS Patients. Anticancer Res. 2016 May;36(5):2345–51.

7.  Hosseini H, Obradović MMS, Hoffmann M, Harper KL, Sosa MS, Werner-Klein M, et al. Early dissemination seeds metastasis in breast cancer. Nature. 2016 Dec;540(7634):552–8.

8.  Krol I, Schwab FD, Carbone R, Ritter M, Picocci S, De Marni ML, et al. Detection of clustered circulating tumour cells in early breast cancer. Br J Cancer. 2021 Jul;125(1):23–7.

9.  Reduzzi C, Di Cosimo S, Gerratana L, Motta R, Martinetti A, Vingiani A, et al. Circulating Tumor Cell Clusters Are Frequently Detected in Women with Early-Stage Breast Cancer. Cancers (Basel). 2021 May;13(10).

10. Jin L, Zhao W, Zhang J, Chen W, Xie T, Wang L, et al. Evaluation of the diagnostic value of circulating tumor cells with CytoSorter(®) CTC capture system in patients with breast cancer. Cancer Med. 2020 Mar;9(5):1638–47.

11. Spizzo G, Fong D, Wurm M, Ensinger C, Obrist P, Hofer C, et al. EpCAM expression in primary tumour tissues and metastases: an immunohistochemical analysis. J Clin Pathol. 2011 May;64(5):415–20.

12. Soysal SD, Muenst S, Barbie T, Fleming T, Gao F, Spizzo G, et al. EpCAM expression varies significantly and is differentially associated with prognosis in the luminal B HER2(+), basal-like, and HER2 intrinsic subtypes of breast cancer. Br J Cancer. 2013 Apr;108(7):1480–7.

13. Rao CG, Chianese D, Doyle G V, Miller MC, Russell T, Sanders RAJ, et al. Expression of epithelial cell adhesion molecule in carcinoma cells present in blood and primary and metastatic tumors. Int J Oncol. 2005 Jul;27(1):49–57.

14. de Wit S, Manicone M, Rossi E, Lampignano R, Yang L, Zill B, et al. EpCAM(high) and EpCAM(low) circulating tumor cells in metastatic prostate and breast cancer patients. Oncotarget. 2018 Nov;9(86):35705–16.

15. Adams DL, Stefansson S, Haudenschild C, Martin SS, Charpentier M, Chumsri S, et al. Cytometric characterization of circulating tumor cells captured by microfiltration and their correlation to the CellSearch(®) CTC test. Cytometry A. 2015 Feb;87(2):137–44.

16. Deng G, Herrler M, Burgess D, Manna E, Krag D, Burke JF. Enrichment with anti-cytokeratin alone or combined with anti-EpCAM antibodies significantly increases the sensitivity for circulating tumor cell detection in metastatic breast cancer patients. Breast Cancer Res. 2008;10(4):R69.

17. Sheng Y, Wang T, Li H, Zhang Z, Chen J, He C, et al. Comparison of analytic performances of Cellsearch and iFISH approach in detecting circulating tumor cells. Oncotarget. 2017 Jan;8(5):8801–6.

18. Akolkar D, Patil D, Crook T, Limaye S, Page R, Datta V, et al. Circulating ensembles of tumor-associated cells: A redoubtable new systemic hallmark of cancer. Int J cancer. 2020 Jun;146(12):3485–94.

19. Gaya A, Crook T, Plowman N, Ranade A, Limaye S, Bhatt A, et al. Evaluation of circulating tumor cell clusters for pan-cancer noninvasive diagnostic triaging. Cancer Cytopathol. 2021 Mar;129(3):226–38.

20. Crook T, Gaya A, Page R, Limaye S, Ranade A, Bhatt A, et al. Clinical utility of circulating tumor-associated cells to predict and monitor chemo-response in solid tumors. Cancer Chemother Pharmacol. 2021 Feb;87(2):197–205.

21. WHO Position Paper on Mammography Screening [Internet]. Geneva; 2014. Available from: https://apps.who.int/iris/handle/10665/137339

22. Siu AL. Screening for Breast Cancer: U.S. Preventive Services Task Force Recommendation Statement. Ann Intern Med. 2016 Feb;164(4):279–96.

23. NCCN Guidelines: Breast Cancer Screening and Diagnosis version 1.2021 [Internet]. nccn.org. [cited 2021 Jul 25]. Available from: https://www.nccn.org/professionals/physician_gls/pdf/breast-screening.pdf

24. Cardoso F, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rubio IT, et

al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis,

treatment and  follow-up†. Ann Oncol  Off J Eur Soc Med Oncol. 2019

Aug;30(8):1194–220.

25.  National Guideline Alliance (UK). Early and locally advanced breast cancer:

diagnosis and management [Internet]. London; 2018. Available from:

https://www.nice.org.uk/guidance/ng101

26.  Wöckel A, Festl J, Stüber T, Brust K, Stangl S, Heuschmann PU, et al.

Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast

Cancer.  Guideline of the DGGG and the DKG (S3-Level, AWMF Registry

Number 032/045OL, December 2017) - Part 1 with Recommendations for the

Screening, Diagnosis and Therapy of Breast C. Geburtshilfe Frauenheilkd.

2018 Oct;78(10):927–48.

27.  Schünemann HJ, Lerda D, Quinn C, Follmann M, Alonso-Coello P, Rossi PG,

et al. Breast Cancer Screening and Diagnosis: A Synopsis of the European

Breast Guidelines. Ann Intern Med. 2020 Jan;172(1):46–56.

28.  Abdullah P, Alabousi M, Ramadan S, Zawawi I, Zawawi M, Bhogadi Y, et al.

Synthetic 2D Mammography Versus Standard 2D Digital Mammography: A

Diagnostic Test  Accuracy Systematic Review and Meta-Analysis. AJR Am J

Roentgenol. 2021 Aug;217(2):314–25.

29.  Salim M, Dembrower K, Eklund M, Lindholm P, Strand F. Range of Radiologist

Performance in a Population-based Screening Cohort of 1 Million  Digital

Mammography Examinations. Radiology. 2020 Oct;297(1):33–9.

30.  Lehman CD, Wellman RD, Buist DSM, Kerlikowske K, Tosteson ANA,

Miglioretti DL. Diagnostic Accuracy of Digital Screening Mammography With

and Without Computer-Aided  Detection. JAMA Intern Med. 2015

Nov;175(11):1828–37.

31.  Screening Mammograms in Younger Women Have Low Accuracy and Detect

Few Cancers. JNCI J Natl Cancer Inst [Internet]. 2010;102(10):NP-NP.

Available from: https://doi.org/10.1093/jnci/djq176

32.  Nelson HD, Fu R, Cantor A, Pappas M, Daeges M, Humphrey L. Effectiveness

of Breast Cancer Screening: Systematic Review and Meta-analysis to  Update

the 2009 U.S. Preventive Services Task Force Recommendation. Ann Intern

Med. 2016 Feb;164(4):244–55.

33.  Hubbard RA, Kerlikowske K, Flowers CI, Yankaskas BC, Zhu W, Miglioretti

DL. Cumulative probability of false-positive recall or biopsy recommendation after 10 years of screening mammography: a cohort study. Ann Intern Med. 2011 Oct;155(8):481–92.

34. Goldfrank D, Chuai S, Bernstein JL, Ramon Y Cajal T, Lee JB, Alonso MC, et al. Effect of mammography on breast cancer risk in women with mutations in BRCA1 or BRCA2. Cancer Epidemiol biomarkers Prev a Publ Am Assoc Cancer Res cosponsored by Am Soc Prev Oncol. 2006 Nov;15(11):2311–3.

35. Niraula S, Biswanger N, Hu P, Lambert P, Decker K. Incidence, Characteristics, and Outcomes of Interval Breast Cancers Compared With Screening-Detected Breast Cancers. JAMA Netw open. 2020 Sep;3(9):e2018179.

36. Health, United States, 2019 – Data Finder [Internet]. cdc.gov. [cited 2021 Jul 26]. Available from: https://www.cdc.gov/nchs/data/hus/2019/033-508.pdf

37. Locke WJ, Guanzon D, Ma C, Liew YJ, Duesing KR, Fung KYC, et al. DNA Methylation Cancer Biomarkers: Translation to the Clinic. Front Genet. 2019;10:1150.

38. Liu MC, Oxnard GR, Klein EA, Swanton C, Seiden M V. Sensitive and specific multi-cancer detection and localization using methylation signatures in cell-free DNA. Ann Oncol Off J Eur Soc Med Oncol. 2020 Jun;31(6):745–59.

39. Klein EA, Richards D, Cohn A, Tummala M, Lapham R, Cosgrove D, et al. Clinical validation of a targeted methylation-based multi-cancer early detection test using an independent validation set. Ann Oncol Off J Eur Soc Med Oncol. 2021 Sep;32(9):1167–77.

40. Cohen JD, Li L, Wang Y, Thoburn C, Afsari B, Danilova L, et al. Detection and localization of surgically resectable cancers with a multi-analyte blood test. Science. 2018 Feb;359(6378):926–30.

41. Sieuwerts AM, Kraan J, Bolt J, van der Spoel P, Elstrodt F, Schutte M, et al. Anti-epithelial cell adhesion molecule antibodies and the detection of circulating normal-like breast tumor cells. J Natl Cancer Inst. 2009 Jan;101(1):61–6.

42. Aktas B, Tewes M, Fehm T, Hauch S, Kimmig R, Kasimir-Bauer S. Stem cell and epithelial-mesenchymal transition markers are frequently overexpressed in circulating tumor cells of metastatic breast cancer patients. Breast Cancer Res. 2009;11(4):R46.

43. Krawczyk N, Meier-Stiegen F, Banys M, Neubauer H, Ruckhaeberle E, Fehm T. Expression of stem cell and epithelial-mesenchymal transition markers in circulating tumor cells of breast cancer patients. Biomed Res Int. 2014;2014:415721.

44. MQSA National Statistics [Internet]. fda.gov. [cited 2021 Sep 17]. Available from: https://www.fda.gov/radiation-emitting-products/mqsa-insights/mqsa-national-statistics

45. John M. Eisenberg Center for Clinical Decisions and Communications Science. Core-Needle Biopsy for Breast Abnormalities [Internet]. In: Comparative Effectiveness Review Summary Guides for Clinicians [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2007-. 2016. Available from: https://www.ncbi.nlm.nih.gov/books/NBK368367/

46. Cancer Stat Facts: Female Breast Cancer [Internet]. seer.cancer.gov. [cited 2021 Sep 17]. Available from: https://seer.cancer.gov/statfacts/html/breast.html

47. Peintinger F. National Breast Screening Programs across Europe. Breast Care (Basel). 2019 Dec;14(6):354–8.

**LIST OF FIGURES**

**Figure 1. Schema of Test.**

Functional enrichment of circulating tumor cells (CTCs) is achieved using a cell culture medium that is cytotoxic towards all non-malignant cells and permits survival of tumor derived malignant cells. Peripheral blood mononuclear cells (PBMC) isolated from whole blood are treated with the custom proprietary medium for 120 h after which the surviving cells and cell clusters are harvested and evaluated by multiplexed immunocytochemistry (ICC) profiling to determine presence of breast adenocarcinoma associated CTCs (BrAD-CTCs) which are identified as CD45-negative cells which express GATA3, GCDFP15 and EpCAM in combination with PanCK.

**Figure 2. Decision Matrix for Classifying Samples.**

The detection threshold for breast adenocarcinoma associated CTCs (BrAD-CTCs)

is ≥ 15 PanCK cells / 5 mL, which is constituted by the detection of ≥ 5 GATA3+,

PanCK+, CD45- cells along with ≥ 5 GCDFP15+, PanCK+, CD45- cells as well as ≥

5 EpCAM+, PanCK+, CD45- cells in the respective aliquots. Depending on the

number of each type of marker positive cells, samples are marked as Positive,

Equivocal or Negative. The decision matrix bestows priority to GATA3 and

GCDFP15 over EpCAM while classifying samples to ensure specificity for BrAD over

other epithelial malignancies where EpCAM+ cells may be detected but breast-

specific markers would be absent. Thus, while the test can detect EpCAM+,

PanCK+, CD45- cells which may be present in various epithelial malignancies, it

specifically reports only BrAD-CTCs.

**LIST OF TABLES**

**Table 1. Summary of Analytical Validation Studies.**

The summary of findings of the Analytical Validation Studies indicate that the Test

provides consistent, accurate and reproducible results with little or no interference

from routine endogenous or exogenous factors when samples are obtained, stored

and processed under the recommended conditions.

**Table 2. Summary of Clinical Validation Studies.**

The table provides the summary of both clinical validation studies. The stringent

cross validation design of the case control (cancer v/s healthy) study yielded a range

of sensitivities and accuracies, the median of which are reported along with the 95%

confidence interval (CI) for the median. Cancer samples (cases) with equivocal

findings were considered as positive for determination of sensitivity and accuracy.

The prospective clinical study evaluated the performance of the test among a cohort

of symptomatic cases who were eventually diagnosed with breast cancer or benign

conditions of the breast. In this study, benign samples with equivocal findings were

considered as false positives for determination of specificity and accuracy. Additional

analyses are provided in Supplementary Tables S3 – S6.

**Whole Blood**

**PBMC Pellet**

**C-ETACs + CTCs**

RBC Lysis

Functional Enrichment

**Fluorescent Immunocytochemistry (ICC) Analysis**

Brightfield

DAPI

PanCK

EpCAM / GATA3 / GCDFP15

CD45

Overlay

Red Blood Cells (RBC)

Peripheral Blood Mononuclear Cells (PBMCs)

Circulating Ensembles of Tumor Associated Cells (C-ETACs)

Circulating Tumor Cell (CTC)

Fluorophore Conjugated Antibodies

| | | | | | |
|---|---|---|---|---|---|
| **GATA3+ Cells** | ≥ 5 | ≥ 5 | *or* | < 5 | < 5 |
| **GCDFP15+ Cells** | ≥ 5 | < 5 | | ≥ 5 | < 5 |

| | | | | |
|---|---|---|---|---|
| **EpCAM+ Cells** | ≥ 5 | < 5 | Any | Any |

| ≥ 15 | ≥ 12 | < 12 | Any |
|---|---|---|---|

**Positive**      **Equivocal**

**Table 1. Summary of Analytical Validation Studies.**

| | EpCAM, PanCK, CD45 | GATA3, PanCK, CD45 | GCDFP15, PanCK, CD45 | Overall |
|---|---|---|---|---|
| **Analyte Stability** | 48 h | | | |
| **Recovery[1]** | 94.6% | 86.4% | 88.6% | 89.9% |
| **Limit of detection** | 1 cell / mL | | | |
| **Linear Range** | 1 - 64 cells / mL | | | |
| **Linearity** | $R^2 \geq 0.98$ | $R^2 \geq 0.98$ | $R^2 \geq 0.98$ | $R^2 \geq 0.98$ |
| **Sensitivity** | 96.0% (86.3% - 99.5%) | 98.0% (89.4% - 99.9%) | 94.0% (83.5% - 98.8%) | 94.0% (83.5% - 98.8%) |
| **Specificity** | 100.0% (88.4% - 100.0%) | 100.0% (88.4% - 100.0%) | 100.0% (88.4% - 100.0%) | 100.0% (88.4% - 100.0%) |
| **Accuracy** | 97.5% (91.3% to 99.7%) | 98.8% (93.2% to 99.9%) | 96.3% (89.4% - 99.2%) | 96.3% (89.4% - 99.2%) |
| **Precision** | CV = 4.6% | CV = 3.9% | CV = 3.8% | CV = 4.1% |
| **Robustness** | >95% | | | |

[1] *Above 10 cells / 5 mL as determined from the Linearity experiment. Values within parentheses represent 95% CI.*

**Table 2. Summary of Clinical Validation Studies.**

| | Case Control Study, Cancer v/s Asymptomatic | | Prospective Study, Cancer v/s Benign | |
| :---: | :---: | :---: | :---: | :---: |
| | Specificity: 100.00% (95% CI: 99.87% - 100.00%) | | Specificity: 93.10% (95% CI: 77.23% - 99.15%) | |
| | **Sensitivity** | **Accuracy** | **Sensitivity** | **Accuracy** |
| **Cumulative** | 92.07% <br> 95%CI: 91.12% - 93.03% | 99.57% <br> 95%CI: 99.34% - 99.81% | 94.64% <br> 95%CI: 88.70% - 98.01% | 94.33% <br> 95%CI: 89.13% - 97.52% |
| *Stage 0* | 70.00% <br> 95%CI: 34.75% - 93.33% | 99.90% <br> 95%CI: 99.70% - 99.98% | 87.50% <br> 95%CI: 67.64% - 97.34% | 90.57% <br> 95%CI: 79.34% - 96.87% |
| *Stage I* | 89.36% <br> 95%CI: 76.90% - 96.45% | 99.81% <br> 95%CI: 99.60% - 99.94% | 95.83% <br> 95%CI: 78.88% - 99.89% | 94.34% <br> 95%CI: 84.34% - 98.82% |
| *Stage II* | 95.74% <br> 95%CI: 85.46% - 99.48% | 99.91% <br> 95%CI: 99.75% - 99.99% | 95.83% <br> 95%CI: 78.88% - 99.89% | 94.34% <br> 95%CI: 84.34% - 98.82% |
| *Stage III* | 100.0% <br> 95%CI: 88.43% - 100.00% | 100.0% <br> 95%CI: 99.87% - 100.00% | 95.00% <br> 95%CI: 75.13% - 99.87% | 93.88% <br> 95%CI: 83.13% - 98.72% |
| *Stage IV* | 100.0% <br> 95%CI: 88.43% - 100.00% | 100.0% <br> 95%CI: 99.87% - 100.00% | 100.00% <br> 95%CI: 83.16% - 100.00% | 95.92% <br> 95%CI: 86.02% - 99.50% |