TITLE PAGE:

Title: Assessment of the humoral response to the homologous Gam-COVID-Vac (Sputnik V) or heterologous Sputnik V/mRNA-1273 (Moderna) vaccination against SARS-CoV-2 in dialysis patients.

Running title: Humoral response to the Sputnik V and Sputnik V/Moderna in dialysis.

Authors: Gustavo LAHAM[a], Alfredo P. MARTÍNEZ[b], Wanda ROJAS GIMENEZ[a], Lucas AMAYA[b], Anabel ABIB[a], Natalia ECHEGOYEN[b], Carlos DÍAZ[a], Alicia LUCERO[b], Antonella MARTELLI[b], Cristina VIDELA[b], Karin NEUKAM[c,d*], Federico A. DI LELLO[e,f*]

Affiliations:

[a]Neprhology Section, Department of Medicine, Centro de Educación Médica e Investigaciones Clínicas Norberto Quirno "CEMIC". Buenos Aires, Argentina.

[b]Virology Section, Centro de Educación Médica e Investigaciones Clínicas Norberto Quirno "CEMIC". Buenos Aires, Argentina.

[c]Servicio de Enfermedades Infecciosas, UCEIMP. Hospital Universitario Virgen del Rocío. Seville, Spain.

[d]Instituto de Biomedicina de Sevilla/CSIC/Universidad de Sevilla. Seville, Spain.

[e]Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Instituto de Investigaciones en Bacteriología y Virología Molecular (IBaViM). Buenos Aires, Argentina.

[f]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina.

*Corresponding Author: Dr. Federico Alejandro Di Lello, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Junín 956, 4º piso, (1113), Ciudad de Buenos Aires, Argentina.

27   E-mail: fadilello@ffyb.uba.ar

29   Dr. Karin Neukam, Servicio de Enfermedades Infecciosas, UCEIMP. Hospital Universitario Virgen

30   del Rocío. Avenida Manuel Siurot s/n. 41013, Seville, Spain and Instituto de Biomedicina de

31   Sevilla/CSIC/Universidad de Sevilla. Seville, Spain.

32   E-mail: karin.neukam@gmail.com

ABSTRACT

Introduction: The humoral response to vaccines is the most used tool to evaluate the protection against SARS-CoV-2 infection. Dialysis patients are a high-risk population and have a reduced immune response to vaccination.

Objective: To assess the humoral response to homologous Gam-COVID-Vac (Sputnik V) and heterologous Sputnik V/mRNA-1273 (Moderna) vaccination in dialysis patients.

Methods: SARS-CoV-2 anti-spike IgG (RBD) concentration was estimated 3-16 weeks after complete vaccination. Reactogenicity was evaluated until day 7 by patients´ self-reported side events.

Results: 107 participants were enrolled [n=84 homologous (SpV/SpV), n=23 heterologous (SpV/Mod)]. Median (IQR) age was 64 (50-75) years old and 79 (73.8%) were male. Additionally, 19 (22.6%) of the SpV/SpV and 4 (17.4%) of the SpV/Mod group had a prior confirmed SARS-CoV-2 infection (p=0.589). In the overall population, 103 patients reached seroconversion (96.3%). Anti-S-RBD IgG median titers (IQR) were higher in the heterologous [1222 (288-5680) BAU/mL] than in the homologous scheme [447 (100-1551) BAU/mL], p=0.022. In a linear model adjusted for age and gender, previous SARS-COV-2 infection (B: 1944.3; CI95: 1136.2-2753.4; p<0.001), and SpV/Mod vaccination scheme (B: 1241.5; CI95: 420.39-2062.6; p=0.003) were independently associated with anti-S-RBD levels. Finally, a higher frequency of adverse effects was associated with the heterologous scheme, although they were well tolerated by all individuals.

Conclusion: The present study provides evidence that the homologous SpV/SpV and heterologous SpV/Mod schemes showed good efficacy and safety under dialysis conditions. These results could be useful for future vaccination strategies, especially aimed at this risk group.

Keywords: Dialysis; SARS-CoV-2; Vaccine; Heterologous scheme; Gam-COVID-Vac; mRNA-1273

INTRODUCTION

Patients under dialysis are at increased risk for severe coronavirus infectious disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) than the general population, with reported mortality rates of up to 28.3% [1-3-]. This situation mainly obeys to a considerably higher mean patients' age (approximately 65 years old) and the frequent presence of comorbid conditions such as obesity, diabetes, and high blood pressure that are linked with a more severe COVID-19 course. Fortunately, effective vaccination schemes against SARS-CoV-2 inducing reduction of both infection and the risk of severe COVID-19 have been rapidly developed [4-7-]. Moreover, when facing first dose serious adverse effects or dose supply shortcomings, introduced heterologous vaccination schemes have shown promising results [8-12-. Nevertheless, clinical trials do not provide information about vaccine efficacy in dialysis populations and possible differences in immunogenicity among SARS-CoV-2 vaccination schemes are poorly understood due to their novelty. Particularly, the homologous Gam-COVID-VAC (Sputnik V) and the heterologous vaccination scheme including Sputnik V as a prime dose and mRNA-1273 (Moderna) as a booster dose have been barely studied since Sputnik V is not approved in all countries and its implementation suffers geographic limitations. In Argentina, the National Administration of Medicines, Food, and Medical Technology (ANMAT) has approved Sputnik V use; however, due to the second dose shortage, mRNA-1273 (Moderna) has been frequently administered as the booster dose.

In this scenario, the present study aimed to assess the humoral response to homologous Sputnik V and heterologous Sputnik V/Moderna vaccination in dialysis patients.

MATERIAL AND METHODS

Study design and population

From March to October 2021, subjects who attended for hemodialysis at the Centro de Educación Médica e Investigaciones Clínicas "Norberto Quirno" (CEMIC), Buenos Aires, Argentina, were included in this prospective cohort study. Inclusion criteria were i) patients who had received Sputnik V prime immunization, ii) they had received a boost dose of either Sputnik V (SpV/SpV) or Moderna (SpV/Mod) vaccines within 18 weeks post-prime dose, and iii) they presented to monitor their humoral immune response three weeks after the boost dose. The vaccination scheme depended on dose availability and the prioritization of risk populations as established by the Argentine Ministry of Health.

Immunogenicity

Binding IgG antibodies against the spike (S) receptor-binding domain (RBD) of SARS-CoV-2 (anti-S-RBD) concentration was assessed at 3-16 weeks after boost. Anti-S-RBD antibodies were quantified using the Abbott Diagnostics SARS-CoV-2 IgG II Quant chemiluminescent microparticle immunoassay (CMIA) on an Architect i2000 SR and an Alinity I analyzer (Abbott Diagnostics, Abbott Park, Illinois, USA). To standardize the results to WHO binding antibody units (BAU), a correction factor for Abbott arbitrary units (AU) was applied where 1 BAU/mL equals 0.142 AU, as previously established by Abbott with the WHO international standard NIBSC 20–136 [13-. Following the manufacturer's recommendations, samples were considered reactive for anti-S-RBD when titers were above 50 AU/mL (7.2 BAU/mL). An 80% protective effect (PROT-80) against symptomatic SARS-CoV-2 infection was assumed when anti-S-RBD titers were 506 BAU/ml or higher [14-.

Reactogenicity

All patients were invited to complete an online questionnaire to report all possible post-boost

vaccination adverse events and required medical assistance. The intensity of adverse effects was

graded as mild, moderate, and severe depending on the interference with daily activities.

Statistical analysis

Descriptive statistics and univariate analyses were performed. The outcome variable was the anti-

S-RBD titer at least three weeks after the boost dose. Differences in anti-S-RBD levels and PROT-

80 according to demographic and clinical parameters were evaluated. Categorical variables were

expressed as numbers (percentage) and analyzed using the Chi-square test or the Fisher's test.

The student's t-test and the Mann-Whitney *U* test were used to compare independent continuous

variables, expressed as median (interquartile range, IQR). For related continuous variables, the

Wilcoxon signed-rank test was applied. Those factors associated with anti-S-RBD levels with a

$p<0.2$ in the univariate analysis, were evaluated in a generalized linear model adjusted for age

and sex. Likewise, multivariate logistic regression models were developed to identify factors

associated with PROT-80. Adjusted odds ratios (AOR) with their corresponding 95% confidence

intervals (CI95) were calculated. Statistical analyses were carried out using the SPSS statistical

software package release 23.0 (IBM SPSS Inc., Chicago, IL, USA).

Ethical Approval and Informed Consent

The study was designed and performed according to the Helsinki declaration and all participants

gave their written informed consent (Study protocol EX-2021-06438339-UBA DME#SSA_FFYB,

Ethics committee of the Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires).

RESULTS

Study population

A total of 107 subjects were included in the study, 84 (78.5%) received SpV/SpV, and the remaining 23 (21.5%) the SpV/Mod scheme. Seventy-nine (73.8%) participants were male, and the median (IQR) age was 64 (50-75) years old. Overall, median time intervals were 91 (77-116) days from prime to boost dose (ΔP-B) and 32 (24-47) days from the boost dose to the anti-S-RBD IgG serological determination (ΔB-antiSRBD). Eighty-four (78.5%) individuals were *naïve* to SARS-CoV-2 infection at the time of prime vaccination. Then, 19 (22.6%) of those who received the SpV/SpV scheme and 4 (17.4%) who received the SpV/Mod one had a prior confirmed SARS-CoV-2 infection (p=0.589). Table 1 shows detailed characteristics of the study population.

Immunogenicity

In the overall population, anti-S-RBD IgG was reactive in 103 (96.3%) persons, 80 (95.2%) immunized with the SpV/SpV vaccine, and 23 (100%) with the SpV/Mod one (p=0.286). Median (IQR) anti-S-RBD titers were 42.5 (4-1297) BAU/mL after the first dose and 502 (110-1993) BAU/mL after the boost dose (p<0.001). In participants with a confirmed SARS-CoV-2 infection before vaccination receiving the SpV/SpV scheme, the humoral response as measured by anti-S-RBD levels was 6.5-fold higher than that observed in *naïve* ones. Similarly, people with a confirmed SARS-CoV-2 infection before the SpV/Mod scheme administration presented 11-fold higher anti-S-RBD levels when compared to participants without prior infection (Figure 1). Anti-S-RBD levels according to epidemiological and clinical parameters are shown in Table 2. Previous COVID-19 (B: 1944.3; CI95: 1136.2-2753.4; p<0.001) and SpV/Mod vaccination scheme (B:1241.5; CI95: 420.39-2062.6; p=0.003) were independently associated with anti-S-RBD levels in a linear model adjusted for age (B:-1.297; CI05: -23.20-20.61; p=0.907), sex (B:-371.92; 95CI -1131.5-387.66; p=0.334), ΔP-B (B:54.911; 95CI: -36.641-146.46; p=0.237), and ΔB-antiSRB (B:-7.828; CI95: -107.04-91.383; p=0.876).

A total of 53 (49.5%) individuals achieved PROT-80. Among participants without prior COVID-19

who received SpV/SpV or SpV/Mod and those with confirmed COVID-19 who received the

homologous and the heterologous schemes, PROT-80 rates were 36.9%, 57.9%, 73.7%, and

100% ($p_{linear\ association}$<0.001), respectively. Corresponding values according to epidemiological and

clinical parameters are displayed in Table 2. In the multivariate analysis, an independent

association with PROT-80 was observed for prior COVID-19 (AOR: 8.663; CI95: 2.541-29.540;

p=0.001), a heterologous vaccination scheme (AOR: 3.767; CI95: 1.229-10.921; p=0.015) and

age (AOR: 0.962; CI95: 0.933-0.992; p=0.012), in a model adjusted for these parameters, as well

as sex (AOR: 0.883; CI95: 0.331-2.361; p=0.805).

Reactogenicity

The homologous and heterologous immunization schemes were well tolerated, and no medical

assistance or potentially fatal events were reported. Adverse events, including local and systemic

symptoms, were higher for SpV/Mod (47.6%) than for SpV/SpV (23.7%) schemes, p=0.031. In

general, the most frequent systemic adverse events were fatigue (9.9%), myalgia (5.9%), and

fever (2.0%). No patients reported headaches, chills, nausea/vomiting, arthralgia, or diarrhea. The

heterologous vaccine scheme tends to induce more systemic adverse effects than the

homologous one (28.6% vs 15.0%, p=0.148). Regarding local adverse events, pain at the

injection site was reported by 11 patients (10.9%) and tended to be more frequent for the

heterologous scheme than for the homologous one (19.0% vs 8.8%, p=0.178). Figure 2 shows

the reactogenicity by adverse effects (local and systemic) according to the vaccination scheme.

DISCUSSION

The present work shows that the implementation of SpV/SpV or SpV/Mod vaccination schemes against SARS-CoV-2 in patients under dialysis was effective leading to a seroconversion rate of 96.3%. Overall, the heterologous scheme showed an anti-S-RBD level almost 3-fold higher than the homologous one. Moreover, both vaccination schemes were well tolerated, and no medical assistance was required.

To our knowledge, this is the first study on the heterologous scheme including Sputnik V and Moderna vaccines in *naïve* and previously SARS-CoV-2 infected patients under dialysis conditions. Additionally, the study allowed results comparison with the homologous SpV/SpV scheme. The seroconversion rate of our population under dialysis was similar to that reported by previous studies showing values between 80-98% [15-20-. Rosa-Diez (2021) detected a 98% seroconversion rate in a study including 102 dialysis patients vaccinated with the Sputnik V scheme [16-. On the other hand, studies performed on patients under dialysis conditions immunized with Moderna vaccines have shown seroconversion rates ranging between 95% and 97.9% [15-,18-,21-. Recent evidence suggested that the mRNA vaccine has a better humoral response when compared to adenovirus-based vaccines [8-,9-,22-24-. In fact, the higher anti-S-RBD titer achieved in this study with the SpV/Mod group agrees with previous works performed on the general population [12-26-]. In this context, our results support the use of Sputnik V or its combination with the Moderna vaccine as an alternative for dialysis patients. However, additional studies are necessary to assess the cellular response since the combination of vaccines appears to enhance the characteristic immune response generated by each vaccination scheme [22-,27-. Results from our study showed the development of a strong humoral response in the dialysis setting, even when some works have shown that patients under dialysis present a lower seroconversion rate and anti-S-RBD titers than healthy controls [28-.

Regarding PROT-80, almost 50% of the studied population achieved this threshold while

heterologous scheme and prior SARS-CoV-2 infection were associated with higher PROT-80

proportions. Furthermore, the multivariate analysis showed that the 80% vaccine efficacy was not

associated with age, gender, ΔP-B, and ΔB-antiSRBD. Obtained results were in accordance with

our previous work performed on general population immunized with the SpV/SpV and SpV/MOD

schemes [12-].

In the present study, there were no anti-S-RBD IgG significant differences between genders.

These findings are consistent with several studies showing that gender seems not to influence

anti-S-RBD IgG titers achieved with both analyzed vaccine schemes [28-32-].

No anti-S-RBD IgG difference according to the age of dialyzed patients was observed in our

sample. Available data on the relationship between age and response to vaccines are scarce and

controversial [30-,32-34-]; importantly, older ages were associated with lower rates of PROT-80.

More studies on this issue are warranted as most reports analyzing the age-associated responses

are conducted in healthy populations [30-,32-34-].

Regarding previous infection with SARS-CoV-2, a significant correlation was found between

higher anti-S-RBD IgG titers and a confirmed past infection for both the SpV/SpV and SpV/MOD

schemes, with a notable difference (11 times) for the heterologous scheme. This finding is in

accordance with our previously reported values for the same vaccination schemes [12-, for the

sputnik V homologous scheme [29-30-] and for RNAm vaccines [35-.

Additionally, the current study found that the body mass index, time on dialysis, weekly total Kt/V,

as well as other evaluated laboratory parameters showed no association with Anti-S-RBD or

PROT-80 levels. There are previous contradictory results regarding this issue. While some

authors, as we did, could not find an association between these variables and the anti-S-RBD IgG

titers [36-, other studies suggest that less comorbidity presence leads to a higher anti-S-RBD IgG

titer [37-38-].

There is little data on SpV/SpV or SpV/Mod reactogenicity under dialysis conditions and this work

offers a contribution to the understanding of SARS-CoV-2 vaccination in this setting. In our

sample, adverse events including local and systemic symptoms were higher in dialyzed patients

who received the SpV/Mod scheme as we have previously described for the general population

[12-]. In agreement with other studies, fatigue and myalgia were the most frequent systemic

reactions [12-,39-. The higher trend in the rate of adverse events in the SpV/Mod group was

expected since it is well proven that stronger side effects were associated with mRNA vaccines

in the general population [11-,12-27-40-,41-. It is important to note that the frequency of adverse

events was lower than that observed in the general population and no patients required medical

support [12-]. However, the median age of the sample analyzed in the present work was higher

and the effect of age on reactogenicity has already been described in previous studies [4-,5-,42-

. Moreover, in a Polewska's (2021) study, adverse events were less frequently observed in

dialyzed patients than in the age and sex-matched control group [39-.

This study has a limitation. It should be considered that serious adverse effects have been

reported in a very low frequency and the small size of the analyzed sample might have influenced

this aspect.

In conclusion, both analyzed vaccine schemes were immunogenic and showed a high

seroconversion rate. In addition, a significant correlation was found between higher anti-S-RBD

IgG titers and a confirmed prior infection with SARS-CoV-2 for both schemes. Moreover, the

heterologous scheme was also associated with a better humoral response. Finally, local and

systemic adverse effects were scarce and mostly mild, demonstrating that both schemes are safe

and well-tolerated. These findings should promote patients on dialysis to receive these

immunization schemes.

Contributors and authorship

FAD, GL and KN: Study concept and design, analysis, and interpretation of data, drafting of the

manuscript and statistical analysis and study supervision.

WRG, AA, NE, CD, AL, AM and **CV:** Acquisition of data, critical revision of the manuscript for

important intellectual content and final approval of the version to be submitted. **APM:** Acquisition

of data, analysis and interpretation of data, critical revision of the manuscript for important

intellectual content and final approval of the version to be submitted.

Declaration of Competing Interest: The authors declare that they have no known competing

financial interests or personal relationships that could have appeared to influence the work

reported in this paper. All authors have approved the final version of the article.

ACKNOWLEDGMENT

FAD is a member of the National Research Council (CONICET) Research Career Program. K.N.

is the recipient of a Miguel Servet contract by the Instituto de Salud Carlos III (grant number

CPII18/00033). We would like to thank Mrs. Silvina Heisecke, from CEMIC-CONICET, for the

copyediting of the manuscript

Funding: This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Conflict of Interest: The authors have no conflicts of interest to declare.

REFERENCES

1-   Suleyman G, Fadel RA, Malette KM, et al. Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. JAMA Netw Open 2020; 3 (6): e2012270. doi: 10.1001/jamanetworkopen.2020.12270.PMID: 32543702.

2-   Ozturk S, Turgutalp K, Arici M, et al. Mortality analysis of COVID-19 infection in chronic kidney disease, haemodialysis and renal transplant patients compared with patients without kidney disease: a nationwide analysis from Turkey. Nephrol Dial Transplant 2020; 35 (12): 2083-2095. doi: 10.1093/ndt/gfaa271.PMID: 33275763.

3-   Hsu CM, Weiner DE, Aweh G, et al. COVID-19 Among US Dialysis Patients: Risk Factors and Outcomes From a National Dialysis Provider. Am J Kidney Dis 2021; 77 (5): 748-756.e1. doi: 10.1053/j.ajkd.2021.01.003. PMID: 33465417

4-   Ramasamy MN, Minassian AM, Ewer KJ, et al. Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine administered in a prime-boost regimen in young and old adults (COV002): a single-blind, randomised, controlled, phase 2/3 trial. Lancet 2021; 396 (10267): 1979-1993. doi: 10.1016/S0140-6736(20) 32466. PMID: 33220855.

5-   Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med 2021; 384 (5): 403-416. doi: 10.1056/NEJMoa2035389. PMID: 33378609.

6-   Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. N Engl J Med 2020; 383 (27): 2603–2615.

7-   Logunov DY, Dolzhikova IV, Shcheblyakov DV, et al. Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. Lancet 2021; 397 (10275), 671–681. https://doi.org/10.1016/S0140-6736(21)00234-8.

8-   Borobia AM, Carcas AJ, Pérez-Olmeda M, et al. Immunogenicity and reactogenicity of BNT162b2 booster in ChAdOx1-S-primed participants (CombiVacS): a multicentre, open-label,

randomised, controlled, phase 2 trial. Lancet 2021; *398* (10295), 121–130. https://doi.org/10.1016/S0140-6736(21)01420-3.

9- Fabricius D, Ludwig C, Scholz J, et al. mRNA Vaccines Enhance Neutralizing Immunity against SARS-CoV-2 Variants in Convalescent and ChAdOx1-Primed Subjects. Vaccines 2021; 9 (8), 918. https://doi.org/10.3390/vaccines9080918

10- Hillus D, Schwarz T, Tober-Lau P, et al. Safety, reactogenicity, and immunogenicity of homologous and heterologous prime-boost immunisation with ChAdOx1 nCoV-19 and BNT162b2: a prospective cohort study. The Lancet Respiratory medicine 2021; 9 (11), 1255–1265. Advance online publication. https://doi.org/10.1016/S2213-2600(21)00357-X

11- Normark J, Vikström L, Gwon YD, et al. Heterologous ChAdOx1 nCoV-19 and mRNA-1273 Vaccination. N Engl J Med 2021; 385 (11), 1049–1051. https://doi.org/10.1056/NEJMc2110716

12- Pereson JM, Amaya L, Neukam K, et al. Heterologous Gam-COVID-Vac (Sputnik V) / mRNA-1273 (Moderna) vaccination induces a stronger humoral response than homologous Sputnik V in a real-world data analysis. Clin Microbiol Inf 2022; in press.27-

13- Galli C, Daghfal D, Averhoff F. Antibody testing for SARS-CoV-2 infection, quantitative determination, response to vaccines and viral variability. https://cdn.pepperapps.io/diagnostics-cms/public/60dcbed551c1ff090981ed95?signature=eyJhbGciOiJkaXIiLCJlbmMiOiJBMTI4Q0JD LUhTMjU2In0..9nFGX43vdCD-Qd2XE-NzdA.e5mgnWdULSy2PGkSwfQ10kEG1UQzLUxIzkdUvU7F1xv06WNo-c47joEI46OgfiQdEoako-TvRl4CwkLYtVIVYRR7v2jcqnkBx9SFQIzw-nqvFqHkx_WIydBAcI4ZA_wEKCPydLqBtvFu7APi9pVFVDt-WE7028r1nMWpvAe5CiYb2tzNgCGlvM09-oxpfdsY.vWBKl-boWE1UxgZgSc1K6Q (3 April 2022, date last accessed).

14-   Feng S, Phillips DJ, White T, et al. Correlates of protection against symptomatic and

asymptomatic SARS-CoV-2 infection. Nature medicine 2021; 10.1038/s41591-021-01540-1.

Advance online publication. https://doi.org/10.1038/s41591-021-01540-1.

15-   Beilhack G, Monteforte R, Frommlet F, et al. Antibody Response and Safety After mRNA-

1273 SARS-CoV-2 Vaccination in Peritoneal Dialysis Patients - the Vienna Cohort. Front Immunol

2021; 12: 780594. doi: 10.3389/fimmu.2021.780594. PMID: 34925359.

16-   Rosa-Diez G, Papaginovic Leiva MM, Lombi F, et al. Safety and Effectiveness of COVID-

19   SPUTNIK   V   Vaccine   in   Dialysis   Patients.   MedRxiv.   **doi:**

https://doi.org/10.1101/2021.10.21.21265349

17-   Grupper A, Sharon N, Finn T, et al. Humoral Response to the Pfizer BNT162b2 Vaccine

in Patients Undergoing Maintenance Hemodialysis. Clin J Am Soc Nephrol 2021; 16 (7): 1037–

42. doi: 10.2215/CJN.03500321.

18-   Yau K, Abe KT, Naimark D, et al. Evaluation of the SARS-CoV-2 Antibody Response to

the BNT162b2 Vaccine in Patients Undergoing Hemodialysis. JAMA Netw Open 2021; 4:

e2123622. doi: 10.1001/jamanetworkopen.2021.23622.

19-   Anand S, Montez-Rath M, Han J, et al. Antibody Response to COVID-19 Vaccination in

Patients   Receiving   Dialysis.   J   Am   Soc   Nephrol   2021;   32(10):2435–8.   doi:

10.1101/2021.05.06.21256768.

20-   Chan L, Fuca N, Zeldis E, et al. Antibody Response to mRNA-1273 SARS-CoV-2 Vaccine

in Hemodialysis Patients With and Without Prior COVID-19. Clin J Am Soc Nephrol 2021;

16:1258–60. doi: 10.2215/CJN.04080321.

21-   Ionita C, Marcelli D, Nita C, et al. Comparison of antibody response to two different mRNA

Covid-19 vaccines in patients on hemodialysis. J Nephrol 2022; 35 (1): 143-151. doi:

10.1007/s40620-021-01195-8. PMID: 34978050.

22- Pozzetto B, Legros V, Djebali S et al. Immunogenicity and efficacy of heterologous

ChadOx1/BNT162b2 vaccination. Nature 2021, 10.1038/s41586-021-04120-y. Advance online

publication. https://doi.org/10.1038/s41586-021-04120-y

23- Schmidt T, Klemis V, Schub D, et al. Immunogenicity and reactogenicity of heterologous

ChAdOx1 nCoV-19/mRNA vaccination. Nat Med 2021a*; 27* (9), 1530–1535.

https://doi.org/10.1038/s41591-021-01464-w

24- Fiolet T, Kherabi Y, MacDonald CJ, et al. Comparing COVID-19 vaccines for their

characteristics, efficacy and effectiveness against SARS-CoV-2 and variants of concern: A

narrative review. Clin Microbiol Infect 2021, S1198-743X (21)00604-2.

https://doi.org/10.1016/j.cmi.2021.10.005

25- Mahallawi WH, Ibrahim NA, Mumena WA. Effectiveness of COVID-19 Vaccines in Patients

Under Maintenance Hemodialysis. Risk Manag Healthc Policy 2021; 14: 5081-5088. doi:

10.2147/RMHP.S345686.PMID: 35002344.

26- Macchia A, Ferrante D, Bouzas MB, et al. Immunogenicity induced by the use of

alternative vaccine platforms to deal with vaccine shortages in a low- to middle-income country:

Results of two randomized clinical trials. Lancet Reg Health Am. 2022; 9: 100196. doi:

10.1016/j.lana.2022.100196.PMID:35128512

27- Schmidt T, Klemis V, Schub D, et al. Cellular immunity predominates over humoral

immunity after homologous and heterologous mRNA and vector-based COVID-19 vaccine

regimens in solid organ transplant recipients. Am J transplant 2021b, 10.1111/ajt.16818.

https://doi.org/10.1111/ajt.16818.

28- Yen CC, Lin SY, Chen SC, et al. COVID 19 Vaccines in Patients with Maintenance

Hemodialysis. J Pers Med 2021;11(8):789. doi: 10.3390/jpm11080789.PMID: 34442432.

29- Claro F, Silva D, Rodriguez M, et al. Immunoglobulin G antibody response to the Sputnik

V vaccine: previous SARS-CoV-2 seropositive individuals may need just one vaccine dose. Int j

Inf Dis 2021. 111, 261–266. https://doi.org/10.1016/j.ijid.2021.07.070

30- Rovere P, Laurelli A, Diaz A, et al. Seroprevalence of anti S1 SARS-CoV-2 antibodies in

workers vaccinated with Sputnik V at a public hospital in Buenos Aires. Medicina (B Aires) 2021,

81 (6):895-901.

31- Babamahmoodi F, Saeedi M, Alizadeh-Navaei R, et al. Side effects and Immunogenicity

following administration of the Sputnik V COVID-19 vaccine in health care workers in Iran.

Scientific reports 2021; 11(1), 21464. https://doi.org/10.1038/s41598-021-00963-7

32- Wheeler SE, Shurin GV, Yost M, et al. Differential Antibody Response to mRNA COVID-

19 Vaccines in Healthy Subjects. Microbiology spectrum 2021; 9 (1), e0034121.

https://doi.org/10.1128/Spectrum.00341-21

33- Steensels D, Pierlet N, Penders J, et al. Comparison of SARS-CoV-2 Antibody Response

Following Vaccination With BNT162b2 and mRNA-1273. JAMA 2021; 326(15), 1533–1535.

https://doi.org/10.1001/jama.2021.15125

34- Yang HS, Costa V, Racine-Brzostek SE, et al. Association of Age With SARS-CoV-2

Antibody Response. JAMA network open 2021; 4 (3), e214302.

https://doi.org/10.1001/jamanetworkopen.2021.4302

35- Ebinger JE, Fert-Bober J, Printsev I, et al. Antibody responses to the BNT162b2 mRNA

vaccine in individuals previously infected with SARS-CoV-2. Nat Med 2021; 27(6), 981–984.

https://doi.org/10.1038/s41591-021-01325-6

36- Rodriguez-Espinosa D, Broseta JJ, Maduell F, et al. Humoral Response of the mRNA-

1273 SARS-CoV-2 Vaccine in Peritoneal Dialysis Patients. Kidney Int 2021; 100:476–7. doi:

10.1016/j.kint. 2021.05.018.

37- Tylicki L, Piotrowska M, Biedunkiewicz B, et al. Humoral Response to COVID-19

Vaccination in Patients Treated With Peritoneal Dialysis: The COViNEPH Project. Pol Arch Intern

Med 2021; 131(10). doi: 10.20452/pamw.16091

38-   Agur T, Ben-Dor N, Goldman S, et al. Antibody Response to mRNA SARS-CoV-2 Vaccine

Among Dialysis Patients - A Prospectivecohort Study. Nephrol Dial Transplant 2021; gfab155.

doi: 10.1093/ndt/gfab155.

39-   Polewska K, Tylicki P, Biedunkiewicz B, et al. Safety and Tolerability of the BNT162b2

mRNA COVID-19 Vaccine in Dialyzed Patients. COViNEPH Project. Medicina (Kaunas) 2021;

57(7): 732. doi: 10.3390/medicina57070732.PMID: 34357013.

40-   Powell AA, Power L, Westrop S, et al. Real-world data shows increased reactogenicity in

adults after heterologous compared to homologous prime-boost COVID-19 vaccination, March-

June 2021, England. Euro surveillance: bulletin Europeen sur les maladies transmissibles =

European communicable disease bulletin, 2021 26 (28), 2100634. https://doi.org/10.2807/1560-

7917.ES.2021.26.28.2100634

41-   Haase M, Lesny P, Anderson M, et al. Humoral immunogenicity and tolerability of

heterologous ChAd/BNT compared with homologous BNT/BNT and ChAd/ChAd SARS-CoV-2

vaccination in hemodialysis patients: A multicenter prospective observational study. J Nephrol.

2022: 1-12. doi: 10.1007/s40620-022-01247-7. PMID: 35084719.

42-   Nachtigall I, Bonsignore M, Hohenstein S, et al. Effect of gender, age and vaccine on

reactogenicity and incapacity to work after COVID-19 vaccination: a survey among health care

workers. BMC Infect Dis 2022; 22 (1): 291. doi: 10.1186/s12879-022-07284-8.PMID: 35346089.

TABLES

Table 1. Population epidemiological characteristics (n=107).

Characteristic	Total, n=107
Age* (years)	64 (50-75)
Male gender, n (%)	79 (73.8)
Primary kidney diseases	
Diabetes, n (%)	15 (14)
High blood pressure, n (%)	13 (12.1)
Polycystic kidney disease, n (%)	8 (7.5)
Glomerular diseases, n (%)	10 (9.3)
Other, n (%)	23 (21.5)
Unknown, n (%)	38 (35.6)
Time on dialysis* (years)	4 (2-7)
Weekly total Kt/V median*	1.5 (1.3-1.9)
Laboratory	
Hemoglobin* (g/dL)	10.8 (9.7-11.6)
Albumin* (g/L)	4.1 (3.7-4.3)
C-reactive protein* (mg/dL)	4.6 (1.4-10.0)
Transferrin saturation*	28 (23-36)
Body mass index (BMI)* (kg/m2)	25.3 (22.4-27.9)
Obesity (BMI >30 kg/m2)	13 (12.1)

429 *Median (interquartile range).

Table 2. Anti-S-RBD IgG and PROT-80 according to epidemiological and vaccination-specific parameters (N=107)

Parameter	Anti-S-RBD			PROT-80	
	N	(BAU/mL)*	*p*	N (%)	*p*
Age					
30-49 years	25	605 (285-1396)		17 (68.0%)	
50-60 years	19	501 (88-914)		7 (36.8%)	
>60 years	63	342 (100-2873)	0.770	29 (46%)	0.084
Gender					
Female	28	431 (105-2419)		14 (50.0%)	
Male	79	502 (110-1740)	0.859	39 (49.4%)	0.954
Time on dialysis					
≤3 years	42	345 (100-1551)		18 (42.8)	
≥4 years	65	555 (112-2634)	0.346	35 (53.8)	0.267
Vaccination scheme					
SpV/SpV	84	447 (100-1442)		38 (45.2%)	
SpV/Mod	23	1222 (288-5680)	0.022	15 (65.2%)	0.090
Prior SARS-CoV-2 infection					
No	84	339 (99-911)		35 (41.7%)	
Yes	23	2872 (508-5680)	<0.001	18 (78.3%)	0.002
ΔP-B					
3-10 weeks	19	288 (94-612)		7 (36.8%)	
11-18 weeks	88	518 (117-2761)	0.141	46 (52.3%)	0.312
ΔB-antiSRBD					
3-4 weeks	51	443 (114-1303)		23 (45.1%)	

5-8 weeks	41	584 (105-3852)		24 (58.5%)	
9-12 weeks	9	502 (126-2436)		4 (44.4%)	
13-16 weeks	6	120 (53-608)	0.187	2 (33.3%)	0.484

433   SpV = Sputnik V, Mod = Moderna

434   *Median (interquartile range); time intervals from prime to boost (ΔP-B) and from boost to anti-S-

435   RBD IgG determination (ΔB-antiSRBD).

436   Weekly total Kt/V median, Hemoglobin, Albumin, C-reactive protein, Transferrin saturation, and

437   Body mass index showed no association with Anti-S-RBD or PROT-80.

  FIGURE

440 **Figure 1.** Anti-S-RBD IgG levels for homologous (SpV/SpV) and heterologous (SpV/Mod)

441 vaccination schemes in subjects who had (A) or had not (B) a confirmed SARS-CoV-2 infection

442 prior to immunization.

457   **Figure 2.** Reactogenicity: frequency of local and systemic adverse effects as reported by the

458   participants for the homologous (SpV/SpV) and heterologous (SpV/Mod) schemes, classified by

459   severity.