Resolving Discordant Variant Classifications using Clinical Guideline Evidence

Vineel Bhat[1], Ivan A. Adzhubei[2], James D. Fife[1], Matthew Lebo[3,4], Christopher A. Cassa[1,4*]

[1] Division of Genetics, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts

[2] Department of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts

[3] Laboratory for Molecular Medicine, Mass General Brigham Personalized Medicine, Boston, Massachusetts

[4] Harvard Medical School, Boston, Massachusetts

*Please address correspondence to: Christopher Cassa cassa@alum.mit.edu

Disclosures: CC has served as a consultant or received honoraria from gWell Health, athenahealth, and Data Sentry Solutions. The remaining authors have no disclosures.

Acknowledgements: We gratefully acknowledge support from the National Human Genome Research Institute (R01HG010372; VB, IA, JF, ML, CC) and for manuscript comments and helpful suggestions from Melissa Landrum, Team Lead of ClinVar at the National Center for Biotechnology Information (NCBI).

Author Contributions: Data curation: V.B.; Formal analysis: V.B.; Software: V.B., I.A.A., J.D.F.; Writing-original draft: V.B., C.A.C.; Writing-review & editing: V.B., C.A.C., M.L., I.A.A., J.D.F.

Abstract

Purpose: To resolve variants of uncertain significance using evidence of pathogenicity from prior variant classifications using the ACMG/AMP clinical guidelines.

Methods: We identify distinct SNVs which are either similar in location or in functional consequence to pathogenic variants, and analyze evidence in support of pathogenicity using three diagnostic criteria.

Results: Thousands of variants, including many in clinically actionable disease genes (ACMG SFv3.0), have evidence of pathogenicity from existing variant classifications, accounting for 4.66% of non-synonymous SNVs within ClinVar. Notably, there are many variants with uncertain or conflicting classifications which cause the same amino acid substitution as other pathogenic variants (PS1, N=323), variants which lead to different amino acid substitutions in the same codon as pathogenic variants (PM5, N=17,563), and LOF variants which are present in genes where many LOF variants are classified as pathogenic (PVS1, N=3,635). The majority of these variants have similar computational predictions of pathogenicity and splicing impact as their associated pathogenic variants.

Conclusion: Broadly, over 1.5 million SNVs exome-wide could make use of information from previously classified pathogenic variants. We have developed a pipeline to identify variants meeting these criteria. This may be used to inform variant interpretation efforts and improve visibility of related cases.

Introduction

Understanding how coding variants affect clinical risk for disease is a major goal in precision medicine.[1] While diagnostic sequencing has advanced dramatically, the interpretation of monogenic variants remains challenging, even in established disease genes.[2] Despite extensive diagnostic testing, the majority of variants have only been observed in a few cases or controls, if at all.[3] An analysis of over 60,000 exomes revealed that over 50% of coding variants were found only in a single individual.[4] On the basis of their low frequencies, lack of observations in clinical cases, and/or availability of functional data, most of these variants have been classified as Variants of Uncertain Significance (VUS). These VUSs pose a substantial challenge in clinical management, and are a major barrier to realizing precision medicine.[1]

Observations in multiple cases and familial segregation data (informative meiosis events) are often considered to be gold-standard evidence of the causality of individual variants.[5] Given that recurrent cases of the same sequence variant are generally infrequent, and large pedigrees are often required to detect significant co-segregation between a variant and phenotype, it can be challenging to confirm pathogenicity from available epidemiological data. In addition, inference from disease segregation analysis can be complicated by the existence of multiple rare variants on the same haplotype. This dearth of human genetics evidence makes it difficult to classify the pathogenicity of an individual variant.[3]

Because of the inherent complexities in variant interpretation and the value of many sources of evidence, the American College of Medical Genetics and Genomics (ACMG) and the Association for Molecular Pathology (AMP) published clinical guidelines about sequence variant interpretation in 2015.[2] These guidelines have provided opportunities to standardize and increase the consistency of variant assessment across clinical laboratories, increasing the utility

of reported findings in publicly accessible databases.[6,7] These guidelines describe sources of evidence and criteria which would help classify a variant as pathogenic or benign. When there are no or few reports describing an identified variant, labs must rely primarily on these sources of evidence, including computational predictive techniques, functional assays, and published gene-level evidence to predict functional effect.[6]

The clinical knowledge base of disease variants is growing rapidly: there are over 460,000 non-synonymous single nucleotide variants (nsSNVs) which have been reviewed by diagnostic labs and deposited to ClinVar.[8] Among these variants, a majority are classified as VUS (66.4%), have conflicting interpretations (5.7%), or have no assertion provided (1.3%), accounting for 73.4% of variants which do not have clear guidance to communicate to patients.[9] A recent study found that fewer than 1% of VUS were reclassified in ClinVar over a 3.5 year period, which demonstrates that this problem is being resolved very slowly.[10] These conflicting or uncertain assessments limit opportunities to improve clinical management and have been associated with overtreatment (*e.g.*, unnecessary surveillance or prophylactic surgery).[11]

Here, we evaluate the extent to which information from prior variant classifications might be used to inform assessments of existing or novel variants. We make use of variant classifications which have been previously submitted to ClinVar with pathogenic or likely pathogenic (P/LP) assertions, which can provide evidence of pathogenicity for other variants which are similar in location or functional consequence. Specifically, we evaluate how frequently multiple variants affect the same codon (whether causing the same amino acid substitution or not), and also evaluate how frequently loss-of-function (LOF) variants are present in genes where LOF variants are known to be pathogenic. We note that all submissions are being correctly processed by ClinVar and diagnostic labs may have intentionally classified a variant as VUS even if there is evidence of pathogenicity from the clinical guidelines.

Materials and Methods

Variant inclusion criteria and annotation

We analyze all nsSNVs in ClinVar[8] and in dbNSFP[12], which are filtered and annotated as described in the supplementary methods. When analyzing clinical assessments from ClinVar, our P/LP category includes 'Pathogenic', 'Likely pathogenic', and 'Pathogenic/Likely pathogenic' classifications, and B/LB includes 'Benign', 'Likely benign', and 'Benign/Likely benign' classifications. Importantly, we use the term VUS inclusively to capture all variants with an inconclusive classification, including 'Uncertain significance', 'Conflicting interpretations of pathogenicity', or 'not provided'.

Evaluation of ACMG/AMP diagnostic criteria

We evaluate predictive evidence for three ACMG/AMP diagnostic criteria:

- **Pathogenic Strong 1 - PS1:** The diagnostic guidelines specify that a variant which leads to the same amino acid substitution as a pathogenic variant has strong evidence of pathogenicity, as it would be expected to lead to the same change in protein structure or function.

- **Pathogenic Moderate 5 - PM5:** The diagnostic guidelines specify that a variant which leads to a different amino acid substitution in the same codon as a pathogenic variant has moderate evidence of pathogenicity, as changes to the same protein codon may be similarly impactful in protein structure or function.

- **Pathogenic Very Strong 1 - PVS1:** The diagnostic guidelines specify that a variant which is expected to lead to loss-of-function in a gene where loss-of-function variants are known to be pathogenic has very strong evidence of pathogenicity.

For these three criteria, we analyze how often a variant which is annotated as uncertain, missing, or conflicting can benefit from knowledge of variants with existing clinical classifications (**Figure 1**). For the PS1 diagnostic criterion, we identify how often two variants which lead to the same amino acid substitution carry inconsistent classifications within ClinVar, and also analyze the frequency at which two VUS encode the same change. For the PM5 criterion, we identify how often two variants which lead to distinct amino acid substitutions in the same codon carry inconsistent classifications within ClinVar. Finally, for PVS1 evidence, we make use of LOFTEE to identify variants which might not be expected to impact function (*e.g.*, present in the last 50 bp, low expected splicing impact, or within NAGNAG sites)[13]. LOFTEE labels variants which pass all filtering criteria as 'high confidence'. We restrict our analysis to LOF which are described in ClinVar as VUS and are present in genes where greater than 50% of LOFTEE 'high confidence' variants are classified as pathogenic.

Additionally, we evaluate these criteria for all non-synonymous SNVs not present in ClinVar. We also consider whether previously classified benign variants may provide evidence in support of benign classification, though the ACMG/AMP does not describe analogous criteria for this case.

Measuring the similarity of variant pairs
We use the term 'variant pair' to describe two distinct SNVs which are related to one another because they alter the same protein codon. We measure the extent to which variant pairs have similar computationally predicted impacts, experimental functional assay values, and allele frequencies. For each variant pair, we compare CADD[14] scores of predicted functional impact, SpliceAI[15] scores of splicing impact, and functional data in *BRCA1*[16] for PS1 and PM5 criteria. For the PVS1 criterion, we analyze computational predictions for loss-of-function SNVs (nonsense, canonical splice acceptor, and canonical splice donor variants) using LOFTEE classifications and flags[13].

Results

Distinct sequence variants which lead to the same amino acid substitution

Within ClinVar, we find many instances where the same amino acid substitution is encoded by more than one distinct sequence variant – in total 7,639 SNVs encoding 3,770 unique amino acid substitutions. Variants with matching amino acid substitutions would generally be expected to have the same functional impact, however we find that some of the sequence variants which lead to the same amino acid substitution have discordant clinical assertions in ClinVar. Among the codons, 1,728 have at least one variant which has been previously classified as P/LP of which 81.3% (N=1,405) are classified exclusively as P/LP, while 18.4% (N=318) contain variants classified as both P/LP and VUS, and 0.3% (N=5) contain variants classified as both P/LP and B/LB (**Table 1**). There are no codons in which P/LP, B/LB, and VUS variants are present.

Overall, this translates to hundreds of unique ClinVar VUS which have strong evidence of pathogenicity as described by the PS1 criterion from the ACMG/AMP classification guidelines while not being in the presence of B/LB variants in the same codon. Among these VUS variants with strong evidence of pathogenicity, we find 212 unique ClinVar submissions across 116 variants in the 73 genes specified for review of secondary findings (ACMG SF v3.0)[17]. This could result in hundreds of patients receiving clinically actionable information related to updated variant classifications.

We further analyze these 'P/LP-VUS variant pairs' (P/LP and VUS variants which encode the same amino acid substitution, N=333) to determine whether each VUS variant is expected to have a similar functional effect as its related P/LP variant. We find that the VUS variants largely appear to have similar computational predictions of functional effects when compared to their

pathogenic partners. When comparing CADD scores for the PS1 criterion, we find very strong correlation (R^2=0.942, MSE=0.223, y=0.997x, **Figure 2a**) with a mean difference of 0.24 (σ = 0.41). Additionally, we find no significant difference between the CADD scores of P/LP and VUS variants (Mann-Whitney U P=0.980).

Two distinct sequence variants could cause the same amino acid substitution yet have different splicing effects (e.g., the P/LP variant has a splicing effect while the VUS does not). To determine whether this may be a primary driver of discordant variant classifications observed in the PS1 variant pairs, we analyze the splicing impact for each variant using SpliceAI[15]. We find that this is unlikely to be the explanation in 98.8% of discordant P/LP-VUS variant pairs (**Figure 3a**). Among 329 distinct P/LP-VUS pairs with SpliceAI annotations, only 2 (0.6%) of the P/LP variants were predicted to have a high impact on splicing (score >0.8) when their VUS partners were predicted to have a weak or low impact on splicing (score <= 0.5), and only 2 (0.6%) of the P/LP variants were predicted to have an intermediate impact on splicing (score (0.5, 0.8]) when their VUS partners had a weak or low impact.

Applicability to benign classifications: While there is no ACMG/AMP guideline which is analogous to the PS1 criterion for benign variation, we measured the extent to which the same principles could potentially provide diagnostic evidence. Similar to evidence in favor of a pathogenic classification, VUS-B/LB variant pairs (two distinct sequence variants leading to the same amino acid substitution with one B/LB and one VUS classification [in the absence of P/LP variants in the same codon], N=126) show strong correlation in CADD scores (R^2=0.860, MSE=0.366, y=0.994x, **Figure 2c**), with a mean difference of 0.21 (σ = 0.57). There is also no significant difference between the CADD scores of B/LB and VUS variants (Mann-Whitney U P=0.994. Additionally, among 125 variant pairs with SpliceAI annotations, 100.0% (N=125) did

not have discordant SpliceAI scores (**Figure 3c**). This suggests the potential utility of such evidence in the variant assessment process.

Potential for aggregating VUS cases causing the same substitution: Variants may be classified as VUS based on insufficient numbers of independent cases. We sought to evaluate the frequency that two distinct sequence variants which lead to the same amino acid substitution were each classified as VUS. Clinical labs could potentially consider such additional case observations when assessing both of these variants. We note that the ACMG/AMP guidelines do not detail such associations to be evidence of pathogenicity, but we analyze the potential for aggregating cases which cause the same amino acid substitution. Overall, we find 3,790 VUS which are in the presence of another VUS which encodes the exact same amino acid substitution while not being in the presence of any P/LP or B/LB variants in the same codon, representing 1,985 unique VUS-VUS variant pairs affecting 1,865 distinct codons. These VUS-VUS variant pairs are often submitted by more than 2 labs in total (mean of 3.3 and a standard deviation of 2.1, **Supplementary Figure 1**), indicating that aggregating this evidence may provide substantial new evidence in favor of pathogenicity.

Variants which lead to distinct amino acid substitutions in the same codon
When considering distinct amino acid substitutions in the same codon (PM5), we find 151,572 single nucleotide variants spanning 64,436 unique codons. Among the 18,073 codons which have at least one variant which has been previously classified as P/LP, only 25.0% are classified exclusively as P/LP (N=4,513). The proportion of codons which contain variants classified as VUS and P/LP is larger at 71.3% (N=12,890) and the proportion of codons which contain variants classified as P/LP and B/LB is 2.9% (N=519). The proportion of codons with variants of all three classifications is 0.8% (N=151, **Table 1**).

In total, this translates to 17,563 unique ClinVar VUS which have moderate evidence of pathogenicity according to the PM5 criterion from the ACMG/AMP guidelines while not being in the presence of B/LB variants in the same codon. Among those which are in variant pairs containing only missense variants, 32.9% (N=2,475, 5,439 submissions) are in clinically actionable disease genes (ACMG SF v3.0). We restrict all further analysis to variant pairs containing only missense variants (Original N=20,950, Restricted N=9,998).

For these PM5 P/LP-VUS variant pairs (P/LP and VUS missense variants which encode different amino acid substitutions in the same codon), the CADD scores of P/LP and VUS components are correlated (R^2=0.184, MSE=0.623, y=0.943x, **Figure 2b**), and the distance between their CADD scores is relatively low with a mean of 0.52 (σ = 0.64).

Additionally, splicing continues to explain little discordance in variant classifications. Among 9,968 P/LP-VUS variant pairs with SpliceAI annotations, 97.2% (N=9,687) did not fall in categories where the P/LP variant was predicted to have a high impact on splicing (score >0.8) when the VUS partner was predicted to have a weak or low impact on splicing (score <0.5), or where the P/LP variant was predicted to have an intermediate impact on splicing (score (0.5, 0.8]) when the VUS partner had a weak or low impact (**Figure 3b**).

Applicability to benign classifications: When considering evidence in favor of a benign classification (which we again note is not included in the ACMG/AMP guidelines), we evaluate the case when two variants, one classified as B/LB and the other classified as VUS, encode different amino acid substitutions at the same codon, while not being in the presence of P/LP variants in the same codon. Among 6,158 missense variant pairs, we find correlation in CADD scores (R^2=0.376, MSE=1.212, y=0.953x, **Figure 2d**), which also have a relatively low distance between one another with a mean of 0.81 (σ = 0.75). Additionally, among 6,083 variant pairs

with SpliceAI annotations, 99.3% (N=6,042) did not have discordant SpliceAI scores (**Figure 3d**). As with PS1, this suggests that evidence analogous to PM5 for the benign classification may have utility for variant interpretation.

LOF variants in genes where LOF is known to be pathogenic

We identify genes where LOF variants (canonical splice-affecting and nonsense) are known to be pathogenic based on the proportion of variants classified as P/LP. Among 4,014 genes in ClinVar which contain LOF SNVs, 2,768 have greater than 50% of the predicted high impact LOFs classified as P/LP (**Figure 4a**). When using this threshold to describe genes where LOF variants are known to be pathogenic (one possible PVS1 approach), we find 3,635 LOF variants which are classified as VUS in ClinVar, and 100 classified as B/LB, accounting for 7.5% and 0.2% of all such variants, respectively (**Table 1**). To determine whether these are variants which are likely to result in loss of function, we used LOFTEE (**Methods**) to identify any variants which might not be functionally impactful (e.g., nonsense mediated decay in last 50 bp). We find that 92.7% of LOF variants (N=3,371) are classified as 'high confidence' by LOFTEE, suggesting that many of these variants are likely to be impactful (**Figure 4b**). Many LOF VUS with strong evidence of pathogenicity (PVS1) are also in clinically actionable disease genes (27.5%, N=1,000, 1,625 submissions, ACMG SF v3.0).

Submission dates of VUS which have evidence of pathogenicity

As the ACMG/AMP guidelines were published in 2015, it is possible that some discordant VUS were last evaluated on a date prior to 2015. However, we find that this is not a common explanation for the majority of discordant VUS. Only 6.4% (N=15) of VUS variants with PS1 evidence, 3.1% (N=206) of VUS variants with PM5 evidence, and 2.3% (N=68) of VUS variants with PVS1 evidence were last evaluated before 2015, among variants which had an available

date of last evaluation. These proportions are similar to the 3.9% of all ClinVar nsSNVs which were last evaluated before 2015, among those with an available date of last evaluation.

It is also notable that for the PS1 and PM5 criteria, a large proportion of VUS were evaluated prior to their pathogenic partners. Among variant pairs with available dates of last evaluation, 44.7% (N=97) and 42.7% (N=2,700) of VUS were last evaluated before their pathogenic partners for the PS1 and PM5 criteria, respectively. These large percentages highlight that many discordant VUS classifications were made when evidence of pathogenicity was not known, and are likely strong candidates for reassessment.

Prioritizing variants for reassessment with the strongest supporting evidence
In addition to submission dates, we provide metrics which may be useful for prioritizing variants for reassessment. These characteristics include allele frequencies (PS1, PM5, PVS1), SpliceAI scores (PS1, PM5), LOFTEE classifications (PVS1), CADD scores (PM5), and ClinVar review status [or 'star' rating] (PS1, PM5, PVS1).

For the PS1 criterion, we find that 43.2% (142 of 329 variant pairs with SpliceAI scores) pass all filters (**Supplementary Methods, Supplementary Table 1**). The vast majority of variants are filtered due to ClinVar review status. For PM5 evidence, this proportion drops to 13.0%, primarily due to additional filtering based on CADD scores. Finally, 25.0% of variants with PVS1 evidence of pathogenicity met all criteria.

Evaluating all non-synonymous SNVs not present in ClinVar
Extending our analysis, we evaluate all nsSNVs in genes within ClinVar which could be informed by evidence of pathogenicity based on PS1, PM5, and PVS1 criteria. In total, we find over 1.5 million variants with such evidence, and provide information about their frequency and similarity

to related pathogenic variants (**Supplementary Table 2, Supplementary Methods and Materials**).

Analyzing the similarity of functional effects of PS1 and PM5 variant pairs in BRCA1

Finally, using functional assay data from nearly 4,000 SNVs in *BRCA1*[16], we analyze the extent to which experimental measurements of functional impact differ between missense variants in the same codon, regardless of their status in ClinVar.

First, we analyze variant pairs which encode the same amino acid substitution (PS1). We find that 98.0% of PS1 missense variant pairs have functional assay measurements that are within one functional classification (e.g. 'functional', 'intermediate', 'loss-of-function'), 92.0% had functional scores within one standard deviation of each other, and 85.9% shared the same functional classification. Overall, we observe no significant difference between functional assay values for PS1 variant pairs, when compared with a null model of differences (generated using all differences in functional scores among variants in separate codons, **Supplementary Figure 2**; Kolmogorov-Smirnov P=0.866). Further, we find a significant difference between PS1 variant pairs and a null model of approximately 2 million missense variant pairs which are not in the same codon (**Supplementary Figure 2**; Kolmogorov-Smirnov $P=1.496 \times 10^{-26}$), indicating that these score differences are significantly different from neutral expectation.

When reviewing variant pairs which encode different amino acid substitutions in the same codon (PM5), 85.7% of all PM5 missense variants pairs had functional scores within one functional classification, 78.6% had functional scores within one standard deviation of each other, and 73.1% shared the same functional class. Compared to a null distribution of functional scores, we also find the distribution of PM5 variant functional scores to be significantly different (**Supplementary Figure 2**; Kolmogorov-Smirnov $P=7.441 \times 10^{-55}$).

This suggests that distinct variants in the same codon often have similar measurements of functional effect, supporting the use of such guidelines based on location.

Discussion

The 2015 ACMG/AMP sequence variant interpretation guidelines describe how to make use of evidence from previously classified variants during variant assessment. For the three diagnostic criteria we analyzed, it appears that rules which make use of prior variant classifications may be underutilized in variant assessment. This may be because systems to review, annotate, and interpret variants often refer to nucleotide substitutions rather than amino acid substitutions. It may also relate to frequency of variant reassessment, as a substantial proportion of VUS variant assertions were not re-evaluated after new information became available about related variants which could have influenced variant interpretation. Other possible reasons include that there have been historically different nomenclature to describe sequence variants, or that variants within the same codon or of a related type are not identified by existing variant analysis tools.

Variant curators should be aware of how valuable these rules are and expand efforts to identify related variant data. A substantial number of SNVs are classified as uncertain, conflicting, or have missing interpretations, but have some form of supporting evidence from an existing classified variant in ClinVar. This information could be broadly useful for novel variant assessment by diagnostic labs, and could inform future guideline development. To make such information more accessible, we provide pre-computed lists of variants within ClinVar and exome-wide which meet these criteria.

The information could also be made more visible for individuals who search or submit to variant databases such as ClinVar. One opportunity to make this information easier to discover would be to have a section which describes 'linked variants', whether affecting the same codon (e.g., PS1, PM5) or having equivalent functional consequence in the appropriate context (e.g., PVS1). Separately, it may be possible to reduce variants which are submitted that have not made use of such data. While there is a process to notify a clinical submitter about a discordant classification during ClinVar submission, we note that there is no such process to notify submitters about discordant classifications or related variant data within the same codon or gene. Lines of conflicting or supporting evidence (such as the PS1, PM5, and PVS1 criteria) would be useful considerations for diagnostic labs to be made aware of during the submission process. Finally, as there are a substantial number of VUSs which are not re-evaluated once new evidence becomes available, there may be an opportunity to trigger re-evaluation of such variants as new structured evidence arises.

Standardizing how and when to apply the PVS1 criterion could be helpful to improve adherence to this form of evidence. We have selected one possible threshold (greater than 50% of all high confidence LOFs have been previously described as pathogenic). Future work may seek to identify the most appropriate threshold for this classification, and the extent to which such data may be supported by hypothesis-free functional assays or clinical data.

Additionally, while the ACMG/AMP guidelines do not have equivalent tiers of evidence in favor of benign classification for PS1 and PM5 criteria, it appears that computational predictions of variant effects, both VUS and unclassified, are similar to those of their B/LB partners to a degree that it is worth evaluating such rules in the future. If sufficiently strong, this evidence could help reduce the number of individuals who would learn about carrying a VUS in a clinically actionable disease gene, avoiding potential overtreatment and unnecessary surveillance.

Although we have identified many occurrences of variants with uncertain, missing, or conflicting classifications which have evidence of pathogenicity, we emphasize that these variants are not necessarily incorrectly classified, and may be influenced by other evidence. Overall, we believe that our tool Tocayo (a pipeline to identify related and discordant variants) and pre-computed variant lists can improve variant interpretation accuracy and efficiency.

Data Availability

A pipeline which can be used to analyze variants from ClinVar and dbNSFP, as well as relevant input files, are provided online (https://github.com/vbhat4/tocayo). Pre-computed lists of variants within ClinVar and exome-wide which meet PS1, PM5, and PVS1 criteria can be found here (https://figshare.com/s/a8ed691e7985afdbb0a1).

Tables

Table 1

Evidence of Impact	Description	Classification	Unique Sites	Unique P/LP Sites	Unique VUS Sites	Unique B/LB Sites	Unique VUS and Confl. Sites
PS1	Same amino acid change as a known pathogenic variant	P/LP	1,728	**1,405**	318	5	0
PM5	Different amino acid change in the same amino acid position as a known pathogenic variant	P/LP	18,073	**4,513**	12,890	519	151
PVS1	LOF SNV in a gene where they are known to be pathogenic	P/LP	48,298	44,563	3,635	100	N/A
PS1*	Same amino acid change as a known benign variant	B/LB	182	5	123	**54**	0
PM5*	Different amino acid change in the same amino acid position as a known benign variant	B/LB	6,256	519	4,928	**658**	151

Table 1: Number of unique sites in ClinVar by evidence and classification. For PS1 and PM5, sites refer to unique codons, while for PVS1, they refer to unique variants. In the case of

PS1 and PM5 specifically, P/LP, VUS, and B/LB sites represent codons where only variants with

that classification exist in the presence of pathogenic or benign variants within the codon (*e.g.*

PS1 Unique B/LB Sites are codons in which at least one P/LP variant and one B/LB variant

encode the same amino acid substitution, but no VUS variants exist in the same codon).

*Evidence not explicitly mentioned in ACMG guidelines.

Figures

Figure 1:

Figure 1: Examples of evidence in favor of pathogenicity in *BRCA1*. 'PS1' evidence applies

to a variant which encodes the same amino acid substitution as a variant previously classified

as pathogenic, 'PM5' evidence applies to a variant which encodes a distinct amino acid

substitution in same codon as a variant previously classified as pathogenic, and 'PVS1'

evidence applies to a LOF variant which is in a gene where LOF variants are known to be

pathogenic. In the first example, variant '17-43115745-A-T' within codon 35 in *BRCA1* is classified as VUS, has strong evidence of pathogenicity (PS1) based on the P/LP classification of variant '17-43115744-C-G' which encodes the same amino acid substitution, as well as moderate evidence of pathogenicity (PM5) from several other variants within the same codon. In the second example, variant '17-43092070-A-C' within codon 1154 in *BRCA1* does not have a classification in ClinVar, and has strong evidence of pathogenicity (PS1) based on the P/LP classification of variant '17-43092070-A-T' which encodes the same amino acid substitution. The variant also has very strong evidence of pathogenicity (PVS1) as it is a LOF variant and LOF variants are frequently assessed as pathogenic in *BRCA1*. In both examples, the VUS or unclassified variant with evidence of pathogenicity has a CADD score similar to its P/LP partner which encodes the same amino acid substitution.

Figure 2

Figure 2: CADD scores of variant pairs are correlated. Individual points represent variant pairs where one variant has a known annotation (P/LP or B/LB) and there is a related VUS affecting the same codon. Each dot compares the CADD score of the variant with an annotation to that of the related VUS variant. **[a]** CADD scores for P/LP-VUS PS1 variant pairs. The variant pairs have highly correlated scores (R^2=0.942) and are not significantly different (Mann-Whitney U p=0.980). **[b]** CADD scores for P/LP-VUS PM5 missense variant pairs. The variant pairs are correlated (R^2=0.184). **[c]** CADD scores for B/LB-VUS PS1 variant pairs. The variant pairs have highly correlated scores (R^2=0.860) and are not significantly different (Mann-Whitney U

p=0.994). **[d]** CADD scores for B/LB-VUS PM5 missense variant pairs. The variant pairs are correlated (R^2=0.376).

Figure 3

Figure 3: SpliceAI scores of related variants. To evaluate how frequently variants with discordant classifications may be attributable to different splicing impacts, we use SpliceAI. Score ranges of >0.8, >0.5 and <=0.8, >0.2 and <=0.5, and <=0.2 represent high, intermediate, weak, and low predicted impacts on splicing, respectively. **[a]** SpliceAI scores for P/LP-VUS PS1 variant pairs. In 86.3% of pairs (N=284), both variants have low predicted splicing impact. **[b]** SpliceAI scores for P/LP-VUS PM5 variant pairs. In 89.4% of pairs (N=18,670), both variants have scores <=0.2. **[c]** SpliceAI scores for B/LB-VUS PS1 variant pairs. In 96.0% of pairs (N=120), both variants have scores <=0.2. **[d]** SpliceAI scores for B/LB-VUS PM5 variant pairs. In 96.4% of pairs (N=5,970), both variants have scores <=0.2.

Figure 4

[a]

Proportion of HC LoF SNVs Classified as P/LP in ClinVar Genes

[b]

LoF Confidence: PVS1 Very Strong Evidence

Consequence

	Splice Affecting	Stop Gained
HC 44051 (99%)	35.1%	64.9%
LC 512 (1%)	10.4%	89.6%
HC 3371 (93%)	42.6%	57.4%
LC 264 (7%)	39.0%	61.0%
HC 84 (84%)	46.4%	53.6%
LC 16 (16%)	43.8%	56.2%

Total 48298

P/LP 44563 (92%)

VUS 3635 (8%)

B/LB 100 (0%)

Figure 4: Evaluating whether LOF SNVs are predicted to be impactful in ClinVar genes.

Using Loss-Of-Function Transcript Effect Estimator (LOFTEE), we classify functional impact for LOF variants (canonical splice-affecting and nonsense) as Low Confidence (LC) or High Confidence (HC). **[a]** For each gene in ClinVar with HC LOF variants, we measure the proportion of such variants which are pathogenic (P/LP) and find 40.1% of genes have a proportion of exactly 1.0, while 21.6% of genes have a proportion of exactly 0.0. **[b]** LOF SNV classifications in genes passing our >50% P/LP proportion threshold. The vast majority of all LOF SNVs are estimated to be HC and impactful.

References

1. Green, E. D. *et al.* Strategic vision for improving human health at The Forefront of Genomics. *Nature* **586**, 683–692 (2020).

2. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine* **17**, 405–423 (2015).

3. Rehm, H. L. *et al.* ClinGen — The Clinical Genome Resource. *New England Journal of Medicine* **372**, 2235–2242 (2015).

4. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291 (2016).

5. MacArthur, D. G. *et al.* Guidelines for investigating causality of sequence variants in human disease. *Nature* **508**, 469–476 (2014).

6. Walsh, R. *et al.* Reassessment of Mendelian gene pathogenicity using 7,855 cardiomyopathy cases and 60,706 reference samples. *bioRxiv* (2016).

7. Amendola, L. M. *et al.* Performance of ACMG-AMP Variant-Interpretation Guidelines among Nine Laboratories in the Clinical Sequencing Exploratory Research Consortium. *American*

journal of human genetics **99**, 247 (2016).

8. Landrum, M. J. *et al.* ClinVar: Public archive of relationships among sequence variation and human phenotype. *Nucleic Acids Research* **42**, (2014).

9. Fife, J. D. *et al. A framework for integrated clinical risk assessment using population sequencing data*. http://medrxiv.org/lookup/doi/10.1101/2021.08.12.21261563 (2021) doi:10.1101/2021.08.12.21261563.

10. Harrison, S. M. & Rehm, H. L. Is 'likely pathogenic' really 90% likely? Reclassification data in ClinVar. *Genome Medicine* **11**, 72 (2019).

11. Bonadies, D. C. *et al.* Adverse events in cancer genetic testing: the third case series. *Cancer J* **20**, 246–253 (2014).

12. Liu, X., Li, C., Mou, C., Dong, Y. & Tu, Y. dbNSFP v4: a comprehensive database of transcript-specific functional predictions and annotations for human nonsynonymous and splice-site SNVs. *Genome Med* **12**, 103 (2020).

13. Karczewski, K. *LOFTEE (Loss-Of-Function Transcript Effect Estimator)*. (2015).

14. Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Res* **47**, D886–D894 (2019).

15. Jaganathan, K. *et al.* Predicting Splicing from Primary Sequence with Deep Learning. *Cell* **176**, 535-548.e24 (2019).

16. Findlay, G. M. *et al.* Accurate classification of BRCA1 variants with saturation genome editing. *Nature* **562**, 217–222 (2018).

17. Miller, D. T. *et al.* ACMG SF v3.0 list for reporting of secondary findings in clinical exome and genome sequencing: a policy statement of the American College of Medical Genetics and Genomics (ACMG). *Genet Med* **23**, 1381–1390 (2021).

Ethics Declaration

This study does not involve human subjects, live vertebrates, higher invertebrates, or any individual-level data. All data used were publicly available and aggregated at the variant level.

Supplementary Methods and Materials

Data processing and labeling

Using the ClinVar variant summary file (dated 12/19/2021), we restricted to single nucleotide variants (SNVs) in assembly GRCh38 and calculated functional consequences using VEP v105. All amino acid substitutions of moderate or high impact were considered for PS1 and PM5 criteria, and LOF variants (functional consequences of 'stop_gained', 'splice_donor_variant', 'splice_acceptor_variant') for the PVS1 criterion. Variants were matched by transcript for functional effects when a variant fell within more than one transcript or gene.

Variants with clinical significance values of 'Pathogenic', 'Likely pathogenic', or 'Pathogenic/Likely pathogenic' were labeled P/LP; 'Benign', 'Likely benign', or 'Benign/Likely benign' were labeled B/LB, and 'Uncertain significance', 'Conflicting interpretations of pathogenicity', and 'not provided' were labeled as VUS.

Computational Score, Allele Frequency, and Other Data

All RefSeq annotations, computational predictions (CADD, PolyPhen2, SIFT), gnomAD allele frequencies (N=125,748 exomes), and SpliceAI scores (masked) were obtained from VEP v105.

Identifying all possible related nsSNVs within ClinVar

To identify all possible variants within ClinVar which might have ACMG/AMP evidence, we evaluate every nsSNV. For PS1 and PM5 criteria, variants were mapped using matching RefSeq transcripts, and if no matching transcript ID was found due to annotation inconsistencies, any transcript with a moderate or high impact was used with a priority for canonical transcripts. ClinVar variants with no moderate or high impact transcripts in VEP, as well as those in the presence of conflicting evidence in the same codon, were filtered. For PVS1 criteria, the annotation process was the same but, instead of impact, required transcripts with a consequence of either 'splice_donor_variant', 'splice_acceptor_variant', or 'stop_gained'.

Identifying all possible related nsSNVs in the coding genome

To identify all possible variants which might have ACMG/AMP forms of evidence, regardless of whether they are already present in ClinVar, we enumerated all possible nsSNVs in dbNSFP v4.2a with potential PS1, PM5, or PVS1 evidence. For PS1 and PM5, this specifically included any variant which affects the same codon as an existing ClinVar variant, and for PVS1, included any essential splice site or nonsense variant. We filtered any variants which have inconsistent transcripts between dbNSFP (Ensembl) and ClinVar (RefSeq). For PS1 or PM5 evidence, all dbNSFP variants were matched to the RefSeq transcript of the ClinVar variant. Otherwise, or when no matching transcript was found, transcripts with moderate or high impact were used with a priority for canonical transcripts. Similar to ClinVar, dbNSFP variants without a transcript meeting these conditions or which were in the presence of conflicting evidence in the same codon were removed from datasets. For the PVS1 criterion, transcripts were matched based on having a matching consequence (e.g., 'splice_donor_variant', 'splice_acceptor_variant', or 'stop_gained'), with a priority for canonical transcripts.

Amino acid substitution and matching of LOF variants

For the PS1 criterion, variants are considered to have the same amino acid substitution and form a 'variant pair' if they meet either of the following criteria:

- The variants have matching amino acid substitutions in ClinVar/dbNSFP and VEP, matching RefSeq transcripts (from VEP), and are in the same gene, regardless of assembly positions.
- The variants have the same amino acid substitution in ClinVar/dbNSFP and VEP, and have matching assembly positions in the same frame, and are in the same gene, regardless of transcript.

For PM5 evidence, this criterion is modified, checking for variants which have the same amino acid position in ClinVar/dbNSFP and VEP but do not have the same amino acid substitution.

For PVS1 evidence, any stop gain (Ter/X) or canonical splice site variant (-1, -2, +1, or +2 base pairs from splice junction) is included if it is within a gene where >50% of such variants annotated as High Confidence by LOFTEE (Loss-Of-Function Transcript Effect Estimator) are classified as P/LP in ClinVar.

Assembly positions are determined to be in the same frame based on codon position and strand direction.

ACMG SF v3.0 analysis

We analyze VUS which are in genes specified for review of secondary findings (ACMG SF v3.0) to identify a subset in these clinically actionable disease genes. For this analysis, we filter to this set of 73 unique genes, and include any VUS with evidence of pathogenicity.

Evaluation of ClinVar submission dates

Dates of last evaluation were analyzed for variants with evidence of pathogenicity as well as their pathogenic partners. When checking for submissions before 2015, only the year of submission was considered, regardless of month or day, while the exact date was considered when comparing the submission dates of related P/LP and VUS variants. For variants with PM5 evidence, all analysis was restricted to missense variant pairs. A substantial number of variants, especially those with PS1 evidence, had missing annotations for the date of last evaluation. Any variant pairs with missing annotations were removed for PS1 and PM5 evidence, and any variants with missing annotations were removed for PVS1.

Annotations for prioritization of variant reassessment

When prioritizing VUS with the strongest evidence, we use various metrics which provide supporting context. These characteristics include the following:

- Missense components; i.e., the P/LP variant pair component is a missense variant and the VUS variant pair components is a missense variant (PM5)
- Allele frequencies less than 10^{-3}; i.e., the P/LP variant pair component has an allele frequency less than 10^{-3} and the VUS variant pair component has an allele frequency less than 10^{-3} (PS1, PM5, PVS1).
- SpliceAI scores which are not discordant; i.e., the SpliceAI score of the P/LP variant pair component is less than or equal to 0.5 when the SpliceAI score of the VUS variant pair component is less than or equal to 0.5 (PS1, PM5).
- LOFTEE classification of High Confidence [HC] with no flags (PVS1).
- CADD scores within one standard deviation of each other; i.e., the absolute value of the difference in CADD scores between the P/LP variant pair component and the VUS variant pair component is less than or equal to the standard deviation of the distribution of score differences when restricted to missense components which meet the filtering criteria based on allele frequencies and SpliceAI scores (PM5).

- Star rating of the VUS variant pair component is less than two (PS1, PM5, PVS1) and the star rating of the P/LP variant pair component is not zero (PS1, PM5).
- Star rating of the VUS variant pair component is not zero (PS1, PM5, PVS1) and the star rating of the P/LP variant pair component is greater than or equal to two (PS1, PM5).

For variants with evidence analogous to PS1 and PM5 for the benign classification, analogous criteria are checked with the exception of the allele frequency criteria, which is not checked, and the SpliceAI criterion, which is modified to check for variant pairs in which the B/LB component has a score greater than 0.5 when the VUS component has a score greater than 0.5.

All filters are cumulative, meaning each subsequent filtering step is applied on the set of variants which passed all previous criteria.

Frequency and similarity of related non-synonymous variants not present in ClinVar
We find 11,981 unique variants not present in ClinVar which have the same amino acid substitution as a P/LP variant in ClinVar while not being in the presence of B/LB variants in the same codon (PS1), translating to 11,186 unique sites. We also find 206,654 unique variants not present in ClinVar which have a different amino acid substitution in the same codon as a P/LP variant in ClinVar [in the absence of B/LB variants in the codon] (PM5), translating to 47,605 unique sites. Finally, there are 1,295,274 unique nsSNV LOF variants (including nonsense and canonical splice site variants) not present in ClinVar which are in genes where more than 50% of all predicted high confidence LOF variants are classified as P/LP (**Supplementary Table 2**).

These variants also have similar predicted impacts: CADD scores of paired P/LP and VUS variants are highly correlated (R^2=0.973, MSE=0.156, y=0.997x) and are not significantly different (Mann-Whitney U P=0.361) for PS1 evidence. All possible substitutions with missense

components meeting the PM5 criterion (N=110,029) show a weaker correlation, and the groups are significantly different from one another (R^2=0.299, MSE=0.581, y=0.940x, Mann-Whitney U P=0.0). With respect to splicing impacts, 98.2% of PS1 variant pairs (N=11,756) and 97.9% of PM5 variant pairs (N=106,889) did not have discordant SpliceAI scores, among variants pairs with SpliceAI annotations. B/LB-VUS variant pairs showed similar or higher numbers for correlation (R^2=0.923, MSE=0.150, y=0.974x), similarity (Mann-Whitney U P=0.529), and splicing (99.5%) for PS1, as well as for PM5 [similarly restricted to missense components] (R^2=0.402, MSE=1.094, y=0.971x, Mann-Whitney U P=0.0, 99.4%). LOF variants with PVS1 evidence were overwhelmingly classified as High Confidence by LOFTEE, at a rate of 98.0%.

Identifying associations in BRCA1 functional data

To analyze the extent to which variants within the same codon differ from one another, we used a dataset of 3,893 variants in *BRCA1* and their associated functional scores. From this dataset, variant pairs were formed to represent PS1, PM5, synonymous, and null models.

A variant pair was included in the PS1 model if two missense variants encoded the same amino acid substitution, and a variant pair was included in the PM5 model if two missense variants encoded distinct amino acid substitutions in the same codon. Variant pairs were included in the synonymous model if two distinct variants encoded the same amino acid substitution as the reference. For the null model, all unique combinations of two missense variants were included if they were not already part of another model (PS1, PM5, or synonymous). Transcripts and assembly positions were ignored in all matching processes.

Differences in functional scores were calculated by taking the absolute value of the difference between the mean function scores of variant pair components.

Statistical tests and data processing

Linear regressions and related statistics (R^2, MSE) were produced using Scikit-learn v0.23.2.

Statistical tests including the Mann-Whitney U and Kolmogorov-Smirnov tests were calculated

using SciPy v1.7.3. Figures were built using Matplotlib v3.3.2, Seaborn v0.11.0, and PowerPoint

2019. Other data processing used Pandas v1.1.4, NumPy v1.21.1, Sqlite v3.30.0, Natsort

v8.0.2, and Python v3.8.5.

Supplementary Figures

Supplementary Figure 1

Supplementary Figure 1: Total number of ClinVar submitters for each variant, among

exclusively VUS substitutions. The sum of the number of submitters was taken for all amino

acid substitutions encoded by multiple VUS variants while not being in the presence of any

other P/LP or B/LB variants in the same codon. The distribution of these sums is strongly right

skewed. 49.7% of substitutions (N=926) have 2 submitters collectively, while 18.3% (N=342)

have 5 or more submitters collectively.

Supplementary Figure 2

[a]

[b]

Tier	PS1	Synonymous	PM5	Null
PS1	1.0	0.866	8.329×10^{-6}	1.496×10^{-26}
Synonymous	0.866	1.0	1.589×10^{-5}	2.608×10^{-28}
PM5	8.329×10^{-6}	1.589×10^{-5}	1.0	7.441×10^{-55}
Null	1.496×10^{-26}	2.608×10^{-28}	7.441×10^{-55}	1.0

Supplementary Figure 2: Distributions of *BRCA1* functional score differences. Of the 4 evidence tiers, PS1 (N=268) represents two missense variants encoding the same amino acid change, Synonymous (N=284) represents two synonymous variants encoding the same amino acid change, PM5 (N=1,703) represents two missense variants encoding different amino acid changes at the same position, and Null (N=2,172,703) represents all possible missense variant combinations of two excluding ones included in previous tiers. All data is based on a functional study of *BRCA1* variants (N=3,893). **[a]** Density curves of the distributions of functional score differences in different evidence tiers. All distributions are right-skewed, but to varying degrees.

[b] Kolmogorov-Smirnov p-value matrix table comparing the functional score difference distributions of different evidence tiers. The distributions of PS1 and Synonymous are most similar, while the distributions of PM5 and Null are most different.

Supplementary Tables

Supplementary Table 1

[a]

Criteria (P/LP)	PS1	PM5	PVS1
None	329	20,877	3,635
Missense components	-	9,968	-
& Allele frequencies <10^-3	329	9,926	3,614
& SpliceAI score of P/LP variant not >0.5 when VUS variant <=0.5	325	9,645	-
& LOFTEE classification of HC as well as no flags	-	-	3,270
& CADD scores within 1 std. deviation of each other	-	6,975	-
& Star rating of VUS variant <2 and P/LP variant star rating not 0	261	5,075	3,124
& Star rating of P/LP variant >=2 or star rating of VUS variant is 0	142	2,710	908

[b]

Criteria (B/LB)	PS1	PM5
None	125	6,193
Missense Components	-	6,083
& SpliceAI score of B/LB variant not <=0.5 when VUS variant >0.5	125	6,042
& CADD scores within 1 std. deviation of each other	-	3,529
& Star rating of VUS variant <2 and B/LB variant star rating not 0	110	2,828
& Star rating of B/LB variant >=2 or star rating of VUS variant is 0	52	1,524

Supplementary Table 1: Number of variants by evidence with cumulative filters. All numbers are among variant pairs with SpliceAI annotations. **[a]** P/LP-VUS variant pairs with PS1 and PM5 evidence and variants with PVS1 evidence cumulatively filtered by relevant criteria. Filters regarding the star rating of the P/LP variant do not apply to the PVS1 column. **[b]** B/LB-VUS variant pairs with PS1 and PM5 evidence cumulatively filtered by relevant criteria.

Supplementary Table 2

	PS1	PM5	PVS1
P/LP	11,186	47,605	1,295,274
B/LB*	5,247	33,105	-

Supplementary Table 2: Number of unique sites outside of ClinVar by evidence and classification. For PS1 and PM5, sites refer to unique codons, while for PVS1, they refer to unique variants. PS1 and PM5 sites exclude unclassified variants which are in the presence of both P/LP and B/LB variants in the same codon. *B/LB evidence not explicitly described in ACMG guidelines.