

1 **The health effects of 14 weeks of physical activity in a real-life setting for**
2 **adults with intellectual disabilities**

3

4 Laurits Munk Højberg^{1*}, Eva Wulff Helge¹, Jessica Pingel², Jacob Wienecke¹

5

6 ¹Department of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen,
7 Copenhagen, Denmark

8 ²Department of Neuroscience, Faculty of Health and Medical Sciences, University
9 of Copenhagen, Copenhagen, Denmark

10

11 *Corresponding author: Laurits Munk Højberg, Department of Nutrition Exercise and Sports,
12 University of Copenhagen, Copenhagen, Nørre Allé 51, 2100 København N, Denmark

13 E-mail: lmh@nexs.ku.dk

14 Phone: +45 91532178

15

16 Word count: 7913

17 Tables: 3

18 Figures: 3

19 Appendices: 1

20

21 **Ethics approval statement**

22 This study was carried out with approval of The Danish Committee on Health and Research
23 Ethics (H-19017828). All participants gave informed written consent and the study followed
24 the Helsinki declaration. All participants were thoroughly informed about the study (one-to-
25 one) by a member of the research team, where they were made aware of their rights as
26 participants, and were able to ask questions before giving their written consent to participate in
27 the present study. Family members or guardians were present when needed to support the
28 conversation.

29

30

31 Abstract

32 *Background:* Individuals with intellectual disabilities (ID) have a reduced physical health
33 compared to the general population, and one of the main contributors is inactivity.

34 *Aim:* To investigate how 14 weeks of physical activity (PA) in a real-life setting affects
35 cardiovascular fitness, body composition and bone health of adults with ID.

36 *Methods:* Adults with ID were recruited into a PA-group or a control group (CON). The PA-
37 group participated in 14 weeks of PA. Body composition, cardiovascular fitness and bone
38 health were assessed before and after the intervention.

39 *Outcomes & results:* Cardiovascular fitness and body composition improved from pre to post
40 within the PA-group: Heart rates (HR) during the last 30 seconds of two increments of a
41 treadmill test, were reduced (3.2 km/h: -4.4 bpm, $p < 0.05$; 4.8 km/h: -7.5 bpm, $p < 0.001$) and
42 fat mass was reduced (-1.02 kg, $p < 0.05$). Between-group differences in favour of the PA-group,
43 were observed in whole body bone mineral density (BMD) (0.024 g/cm^2 , $p < 0.05$) and in BMD
44 of the left femur neck (0.043 g/cm^2 , $p < 0.05$).

45 *Conclusions & Implications:* Fourteen weeks of PA increase cardiovascular fitness, reduced
46 fat mass and improved BMD in the weight-bearing skeleton in the PA-group. Increased and
47 regular PA seems to be a promising tool to promote physical health in adults with ID.

48

49 What this paper adds

50 This paper underlines the importance of including physical activity in the everyday lives of
51 individuals with intellectual disabilities. The health impact of physical activity performed
52 outside controlled research laboratories needs further investigation, and therefore this paper
53 sheds light on physical activity performed in a real-life setting. Our results indicate a positive
54 impact of physical activity on the cardiovascular system and body composition, as well as bone
55 health. The question of how of bone health of adults with intellectual disabilities responds to
56 physical activity is largely understudied in intervention designs, and this paper includes state-
57 of-the art investigation of development in bone mineral density after participation in varied
58 physical activities. These results give promise and indicate a positive impact of the intervention
59 on the bone health of the participants.

60

61 **Keywords:** Intellectual disabilities, Physical activity, Cardiovascular fitness, Bone health,
62 Body composition

63

64 **1. Background**

65 Individuals with intellectual disabilities (ID) are subjected to increased mortality, in Denmark,
66 individuals with ID die 14.5 years earlier than the general population (Flachs, Uldall, & Juel,
67 2014). The increased mortality is in part attributed to decreased physical health, high levels of
68 inactivity and obesity which can lead to lifestyle diseases such as type II diabetes,
69 cardiovascular disease and poor bone health (Anderson et al., 2013; Bartlo & Klein, 2011;
70 Melville, Hamilton, Hankey, Miller, & Boyle, 2007; Robertson et al., 2000; Srikanth, Cassidy,
71 Joiner, & Teeluckdharry, 2011; Zylstra, Porter, Shapiro, & Prater, 2008). Physical activity (PA)
72 has been suggested as an effective way to improve the health in individuals with ID (Bartlo &
73 Klein, 2011; Bouzas, Ayán, & Martínez-Lemos, 2018; Robertson et al., 2000). Previous
74 investigations of individuals with ID have shown that the group respond to physical activity
75 similarly to the general population. Evidence from controlled trials and systematic reviews
76 show improvements in cardiovascular fitness following 80-135 minutes of aerobic training per
77 week for 8-16 weeks (Boer et al., 2014; Cluphf, O'Connor, & Vanin, 2001; Dodd & Shields,
78 2005). The evidence regarding resistance training indicates that 8-24 weeks of resistance
79 training increases muscle strength of persons with ID (Cowley et al., 2011; Obrusnikova,
80 Firkin, Cavalier, & Suminski, 2021; Shields et al., 2013). Interventions with both aerobic and
81 resistance training has shown that the participants experienced the benefits from both these
82 modalities, within 10-20 weeks (Calders et al., 2011; Elmahgoub et al., 2011; Oviedo, Guerra-
83 Balic, Baynard, & Javierre, 2014). PA for individuals with ID seem to induce reductions in
84 body weight and fat mass (Elmahgoub et al., 2011; Son, Jeon, & Kim, 2016), and improved
85 bone health (González-Agüero et al., 2012; Lizondo, Caplliure-Llopis, Escrivá, De La Rubia,
86 & Barrios, 2019), but the results are equivocal within these areas, and more research is
87 warranted. Improvements in the above-mentioned domains have a positive influence on general
88 health. Improved cardiovascular fitness is associated with reduced risk of early death from
89 cardiovascular and respiratory diseases (Steell et al., 2019), and increased muscle strength in
90 the lower body can prevent falls and fractures (Benichou & Lord, 2016). The mentioned studies
91 have generally used supervised exercise interventions, with researchers controlling the PA
92 performed by the participants. The evidence from these studies are fundamental, but the
93 translation into real-life, health-promoting exercise programs for the population with ID is
94 difficult and demand further research (Bartlo & Klein, 2011; Bouzas et al., 2018). Hence, the
95 aim of the present study was to investigate how 14 weeks of varied and adapted PA performed
96 in a real-life setting, 5 days per week affects the health of adults with ID. To fulfil this aim, we
97 collaborated with a local daily activity centre, where adults with ID is enrolled for a year, and

98 participating in PA at the centre 5 days a week. Thus, an already feasible intervention could be
99 investigated on a large number of participants. We hypothesised that a PA-group would
100 improve their health from pre- to post-test compared with a control group. In addition, we
101 hypothesized that a group of participants, who had already attended the activity centre at least
102 one school year prior to the study (2nd-year participants), would have a better health profile at
103 PRE-tests than the participants, who started their first year at the centre when entering the study
104 (1st-year participants), due to impact of the PA that the 2nd-year participants had already
105 performed. For the same reason, we expected to find the largest improvements in the selected
106 health parameters in the 1st-year group over the course of the study.

107

2. Methods

108

2.1 Intervention and setting

109

110 The intervention consisted of varied forms of PA, performed in four periods of 3-4 weeks
111 in duration, for a total of 14 weeks. Period 1: Dance, cycling, basic ballgames, outdoor fitness
112 and athletics; Period 2: Adventure, motor training, football (soccer) and maritime outdoor
113 activities. Period 3: Body theatre, swimming, floorball, I and E-sport. Period 4: Body theatre,
114 racket sports, volleyball and fitness. The activities were performed at the activity centre; the
115 Sports School for Adults with Developmental Disabilities (SSADD) in Copenhagen, Denmark.
116 Adults with ID attending the SSADD, referred to as “participants”, were recruited into the PA-
117 group. Each participant chose one of the above activities, in each period. Five days per week,
118 they had 150 minutes of scheduled PA, in two blocks of 75 minutes with 45 minutes of break
119 time in between. The SSADD has a capacity of ~50 adults with different grades of ID, per
120 school year. Each year approximately half of the participants continue at SSADD for an
121 additional year (2nd year participants), and ~25 new participants start at the school (1st year
122 participants). The personnel at SSADD aims at improving the physical, social and personal
123 competences of the individuals attending the institution, through the use of inclusive PA and
124 sports, and the activities are adjusted to meet the capabilities of the individuals attending the
125 SSADD.

126

2.2 Participants

127

128 Fifty-two adults with ID (26.7 ± 1.2 years) were recruited from the SSADD (PA-group).
129 The PA-group was further divided into a 1st-year group and a 2nd-year group. A control group

130 (CON-group) of 14 adults (29.1 ± 1.7 years) were recruited from other institutions and activity
131 centres in Copenhagen. The institutions which the individuals in the CON-group attended did
132 not focus on physical activity, but were recreationally physically active (Figure 1). This study
133 was carried out with approval of The Danish Committee on Health and Research Ethics (H-
134 19017828). All participants were thoroughly informed about the study (one-to-one) by a
135 member of the research team, where they were made aware of their rights as participants, and
136 were able to ask questions before giving their written consent to participate in the present study.
137 Family members or guardians were present when needed to support the conversation. Being
138 under legal guardianship was an exclusion criteria. The study followed with the Helsinki
139 declaration. Since the project ran for two years, some participants were tested PRE and POST
140 twice, with a year between the intervention periods. Between the two rounds of measurements,
141 the subjects were allowed to change groups; for example, many participants the 1st-year group
142 in the first round continued at SSADD, and were then included in the project in the 2nd-year
143 group (Figure 1). Some participants who had initially given their consent to participate in the
144 study chose to withdraw their consent, during the course of the study. These circumstances
145 resulted in different numbers of participants both between the different tests and between the
146 PRE and POST measurements. An overview of the number of measurements performed in the
147 different tests can be found in Appendix 1. The grade of ID of the participants varied from mild
148 to moderate and consisted of individuals with diagnoses such as Downs' syndrome, cerebral
149 palsy and other variants of ID, which the researchers did not have information on.

150

151 [Figure 1]

152

153 **2.3 Testing procedure**

154 The individuals in the PA-group were recruited during their first week at SSADD over
155 two intervention rounds, in 2019 and 2020. After recruitment, the participants in the PA-group
156 underwent the tests described below over the course of two weeks. Participants in the CON-
157 group were recruited during this period, and were tested after the PA-group. From PRE- to
158 POST-tests, 14 weeks passed on average in both groups. All tests were carried out at the
159 facilities of the Department of Nutrition Exercise and Sports, University of Copenhagen.

160

161 **2.4 Assessment of physical activity**

162 To measure the total amount of daily PA performed by the participants, accelerometry
163 using the Axivity AX3 system were performed. Accelerometers (Axivity, Newcastle UK) were

164 attached to the right thigh of the participants for periods of seven days, using medical tape and
165 waterproof artificial skin (Opsite Flexifix ®). The accelerometers were checked daily in the
166 weekdays to ensure correct positioning. If the accelerometer had detached it was reattached
167 and recalibrated. Based on the threshold values of standard deviation of acceleration and the
168 derived deviation (Skotte, Korshøj, Kristiansen, Hanisch, & Holtermann, 2014), the software
169 ACTi4 (ACTI Corp., Copenhagen, Denmark) was used to discriminate the accelerometer data
170 into different PA types (sitting, standing, walking, walking on stairs, running, cycling and total
171 number of steps). This enabled the measurement of steps taken pr. hour and minutes of activity
172 pr. week. The participants were asked to register their sleep schedules (hours slept) with the
173 help of family or personnel at their home. Forty-three participants were divided between six
174 different activities; football (n=9), motor training (n=5), fitness activities (n=9), outdoor
175 activities (n=13) and cycling (n=7). Not all participants were able to wear the accelerometer
176 for seven days. In these cases, the days they had completed, were included in the data.

177

178 **2.5 Dual X-ray Absorptiometry (DXA) scans**

179 The outcome measures of interest to the present study was areal bone mineral density
180 (BMD, g/cm²), and body composition measures (e.g. fat mass (kg) and fat free mass (kg)). The
181 DXA-protocol consisted of a whole body scan (Albanese, Diessel, & Genant, 2003) and scans
182 of the lumbar vertebrae (L1-L4) and bilateral proximal femur, in that order (Lunar iDXA, GE
183 Healthcare). DXA-scans of the lumbar and femur regions are routinely used as clinical tools
184 for diagnosing osteoporosis (Blake & Fogelman, 2007).. The participants' weight and height
185 were recorded prior to the scans (Soehnle Professional Scale, Soehnle, Germany). The
186 participants were asked to remove all metal objects (jewellery, buttons etc.) from their body
187 and clothing prior to the scans. The participants were asked to arrive to the tests in a fasting
188 state. Not all participants were able to meet this request, so their fasting state was recorded at
189 the PRE-tests (fasting/non-fasting). The participants were asked to arrive to the POST-test in
190 the same fasting-state as at the PRE-test. The participants were placed in the middle of the
191 scanning bed with their arms held to the sides of the body, their hands in neutral position and
192 their palms on the side of their thighs. With the knees bent in a 90 degrees angle, a foam block
193 was placed under the participant's calves for the lumbar scans to ensure that the lower back
194 was placed correctly on the scanning bed. During the femur scans, the participants were asked
195 to rotate their legs inwards from the hips, while their feet were strapped to a plastic trapezoid.
196 This was done to ensure the right position of the femur neck, and to promote reproducibility of
197 the measurements.

198

199 **2.6 Oxygen uptake protocol and heart rate**

200 The submaximal oxygen uptake (VO_2)-protocol of the present study was inspired by
201 study by Ebbeling and colleagues (Ebbeling, Ward, Puleo, Widrick, & Rippe, 1991), where a
202 submaximal treadmill test for estimation of VO_2max was developed. The test consisted of a 4
203 minute warm up with 0% incline at two different speeds, two minutes at 3.2 km/h followed by
204 two minutes at 4.8 km/h. The warm-up was followed by 1 minute at standing rest. After the
205 rest the participants performed the test, a bout of 8 minutes with two intensities. First, the
206 participants walked at 3.2 km/h and 5% incline for 4 minutes, immediately followed by 4
207 minutes at 4.8 km/h and 5% incline. Online breath-by-breath analysis was performed during
208 the exercise protocol using a metabolic cart (Masterscreen CPX, CareFusion). Participants
209 wore a mask with ~70 ml dead space, which was directly connected to the gas analyser and
210 flow sensor. The system was calibrated before each trial. The heart rates (HR) were sampled
211 using POLAR H3 heart rate sensors. An experimenter noted the heart rates of the participants
212 every 10 seconds during the exercise protocol. The area under the curve (AUC) of the VO_2 -
213 curves during intensity transitions is an estimate of oxygen uptake kinetics. The first 20 seconds
214 of the transitions were omitted from the analyses to exclude phase 1 kinetics and investigate
215 phase 2 kinetics. AUC were then calculated for the subsequent 100s. An increase in AUC of
216 the VO_2 -curve during similar intensity transitions and a fixed time frame, is a measure of
217 increased total oxygen consumption.

218

219 **2.7 Data analysis and statistics**

220 Data was analysed using R statistics (*R statistics 4.0.0*, 2020). Linear mixed models
221 (LLM's) were used to analyse changes within and between groups ("lme4" R-package (Bates,
222 Mächler, Bolker, & Walker, 2015)). In the first LLM, the group variable had two levels; PA
223 and CON. In the second LLM, the group variable had three levels; 1st-year group, 2nd-year
224 group and CON. The LLM's included the age, gender and diagnosis of the participants as well
225 as the test round (i.e. 2019 and 2020), as fixed effects, and the participant ID's as a random
226 effect. LLM's were chosen, for their ability to handle missing data. Post hoc analysis of the
227 models were used to investigate the changes from PRE- to POST-tests between and within the
228 groups ("multcomp" R-package (Hothorn, Bretz, & Westfall, 2008)). The p-values were
229 corrected for multiple comparisons with the single step method. The critical value for statistical
230 significance were assumed at an alpha level $p \leq 0.05$. Unless otherwise stated, data is presented

231 as mean \pm standard error (SE). Percent changes within groups are calculated as the change from
232 PRE- to POST-test, and percentwise differences between groups are calculated as the change
233 score between the PA groups and the CON group, relative to the CON-group PRE-test value.

234 **3. Results**

235 **3.1 Demographic data**

236 The demographic data collected at the PRE-measurements is presented in Table 1. Some
237 participants were tested in both 2019 and 2020. They have contributed to the demographic data
238 once, in 2019. An overview of the total number of data points included in the measurements,
239 can be found in Appendix 1.

240

241 **3.2 Daily physical activity**

242 During one week, the participants in the PA-group on average took more steps/hour
243 ($p < 0.001$) while at SSADD (1453 ± 60 steps/h) compared to leisure hours, both on weekdays
244 (738 ± 67 steps/h) and in the weekends (564 ± 58 steps/h) (Figure 2). The PA-group performed
245 on average 407 minutes of PA/week during the total of 1500 minutes they spent at the school.
246 This included activities other than standing, sitting or lying down. As mentioned, they had 150
247 minutes of scheduled PA/day (equivalent of 750 minutes/week).

248

249 [Figure 2]

250

251 **3.3 PRE-test measurements**

252 Raw PRE-test values are presented in Table 2. The comparison between the 1st-year
253 group and the 2nd-year group showed a higher total body weight (4.7%, $p < 0.05$) and a higher
254 fat mass (11%, $p < 0.001$) in the 1st-year group. In addition, the 1st-year group had a higher
255 visceral (16.2%, $p < 0.001$) and gynoid fat mass (8.9%, $p < 0.05$) than the 2nd-year group. The 1st-
256 year-group had a higher heart rate (HR) compared to the 2nd-year group during the last 30s at
257 3.2 km/h (5.0%, $p > 0.05$) (Figure 3). The PA-group had lower BMD compared to the CON-
258 group in L1-L4 (-6.8%, $p > 0.05$). The T-score distribution of the right femur neck of the PA-
259 group was: $T \leq -2.5$ (osteoporosis): 0%, $-2.5 > T \leq -1$ (osteopenia): 23.5%, $-1 > T < 0$: 29.4%
260 and $T \geq 0$: 43.1 %. The CON-group distribution was: $T \leq -2.5$ (osteoporosis): 0%, $-2.5 > T \leq -$
261 1 (osteopenia): 28.6%, $-1 > T < 0$: 35.7% and $T \geq 0$: 35.7%. As no differences were observed
262 between the right and left sides at PRE-tests, only data from the right side is presented.

263

264 [Table 1]

265 [Table 2]

266

267 **3.4 Pre-to posttest changes**

268 **4.4.1 Body composition.** Table 3 presents the model estimated Δ values (POST – PRE)
269 within the groups, and the differences in change scores between the groups (e.g. PA-group Δ
270 value – CON-group Δ value). Analyses of body composition outcomes from the whole body
271 DXA scans revealed that the total PA group lost -4.0% fat mass ($p < 0.05$), with the 1st-year
272 group lost -5.8% fat mass ($p < 0.05$). For the 1st-year group, parts of the fat loss were located at
273 the visceral and gynoid regions (-7.4% and -7.1%, $p < 0.05$, respectively). However, no between
274 group differences were observed ($p > 0.05$).

275

276 **3.4.2 Bone mineral density.** A within-group change was observed in the PA-group, thus
277 the BMD of the right femur neck increased ($p < 0.05$) (Table 3). Tendencies indicating reduced
278 BMD within the CON-group over the course of the study was evident as borderline significant
279 reductions of the L1 and left femur neck ($0.5 < p \leq 0.1$). Differences in change scores for whole
280 body BMD (2.0%, $p < 0.05$) and BMD of the left femur neck (4.3%, $p < 0.05$) between the PA-
281 and CON-group, in favor of the PA-group, were observed.

282

283 **3.4.3 Cardiovascular fitness.** Within-group changes was observed in the PA-group as a
284 reduction in HR during the last 30s at both 3.2 km/h (-3.8%, $p < 0.05$) and 4.8 km/h (-5.4%,
285 $p > 0.001$) (Figure 3). This decrease was more pronounced within the 1st-year group (3.2 km/h:
286 -7.0%, $p < 0.01$. 4.8 km/h: 7.2% $p > 0.001$). However, no between group differences were
287 observed in the HR measurements, and VO_2 -kinetics during the transitions of the treadmill test
288 did not change ($p > 0.05$).

289 [Table 3]

290 [Figure 3]

291 **3.5 Post-test differences**

292 Between-group differences at POST-tests were analysed to investigate if the differences
293 observed at PRE-tests were still present (data not shown). The differences in body composition
294 observed at the PRE-tests between the 1st- and 2nd-year group, and the difference in BMD of
295 L1-L4 between the PA-group and the CON-group (Table 2), were not observed at the POST-

296 tests. The only statistically significant finding from this analysis was a significant 8.9 % lower
297 HR at 3.2 km/h in the PA- compared to the CON-group (Figure 2).

298 **4. Discussion**

299 The results of the present study indicates that adults with ID by participation in PA in a real-
300 life setting for 14 weeks can improve their body composition, bone health and cardiovascular
301 fitness. The amount of PA performed by the PA-group is above the weekly 300 minutes of PA,
302 recommended for adults by the World Health Organization (Bull et al., 2020).

303

304 **4.1 Body composition**

305 The PA-group lost 1.02 kg of fat mass and the 1st-year group lost 1.7 kg of fat mass, with
306 no changes observed in body weight. The change in the total PA-group seems to be driven by
307 the 1st-year group, as no changes were found for the 2nd-year group. The results of previous
308 studies on the effect of physical activity on body composition for adults with ID are equivocal.
309 Some studies on individuals with ID have reported reductions in fat mass of similar magnitude
310 as the present study (Elmahgoub et al., 2011; Ordoñez, Rosety, & Rosety-Rodriguez, 2006;
311 Wu et al., 2017) while other studies did not (Calders et al., 2011; Oviedo et al., 2014). In a
312 meta-analysis of 6 RCT's assessing the effects of PA interventions on body composition in
313 young adults with ID, Harris and colleagues reported no overall effect of physical activity
314 interventions on body composition (Harris, Hankey, Murray, & Melville, 2015). The studies
315 included in the meta-analysis used interventions of 80-195 minutes of activity/week (Donnelly
316 et al., 2009; Harris et al., 2015). As Harris and colleagues suggested, this might be too little to
317 elicit changes in body composition. In the present study, the participants in the PA-group
318 performed ~400 minutes of PA/week, of variable intensity. Thus, the training load imposed on
319 our participants, should be enough to elicit the adaptations in body composition observed in
320 the present study. This is supported by the findings from a systematic review on the effects of
321 exercise on fat mass in the general population, where ~10 weeks of either moderate- or high-
322 intensity exercise, induced ~2 kg fat mass reduction (Wewege, van den Berg, Ward, & Keech,
323 2017). Food intake is another important variable when studying changes in body composition
324 (Clifton, Condo, & Keogh, 2014; Miller et al., 2013). Studies on individuals with ID have
325 successfully improved body composition through multi-component interventions with
326 exercise, diet and motivation (Martínez-Zaragoza, Campillo-Martínez, & Ato-García, 2016).
327 The diet of the participants were not monitored in the present study. Thus, the results indicate
328 that it is possible to reduce fat mass in this population, solely with increased PA without a diet

329 intervention. Significant parts of the fat mass reduction seen in the present study could be
330 located to the gynoid and visceral areas (Table 3). The visceral fat mass reduction for the 1st-
331 year group is of significance in a health perspective, as the amount of visceral fat is a risk factor
332 in relation to life style diseases such as type II diabetes mellitus and cardiovascular diseases
333 (Finelli, Sommella, Gioia, La, & Tarantino, 2013; Kuk et al., 2006; Lee et al., 2018). Thus, our
334 results show a positive effect of 14 weeks of PA performed at SSADD on fat mass.

335

336 **4.2 Bone health**

337 All together, the between group differences in BMD observed in whole body, L1-L4 and
338 the left femur neck indicate that the PA-group benefitted from 14 weeks of PA, compared to
339 the CON-group (Table 3). The differences between the groups seem to be driven by a decrease
340 in BMD in the CON-group, while the PA-group maintained their BMD in these regions.
341 Another indication of this, is the higher BMD of L1 -L4 of the CON-group compared to the
342 PA-group at PRE-tests (Table 2), which were not present at POST-tests (data not shown).

343 To our knowledge, the present study is one of the first to investigate the bone health in
344 individuals with ID after a PA intervention, and the first to do so in adults. A previous study
345 on 7-10 year old boys with ID found an increase in femoral neck BMD after 6 months of
346 exercise (165 min/week) (Hemayattalab, 2010). Another study observed an increase in bone
347 mineral content of the femur in a group of subjects with DS between 10 and 19 years, after 21
348 weeks of exercise (50-60 min/week) (González-Agüero et al., 2012). The results of the present
349 study is thus in line with the previous research, showing that PA has an osteogenic effect on
350 individuals with ID. In the present study, the osteogenic effect of PA is evident as an increase
351 in right femur neck BMD in the PA-group, and as a counteraction of decrease in lumbar BMD
352 in the PA-group relative to the CON-group. There are differences between the present study
353 and earlier research regarding bone adaptations to exercise. First, it is common in this line of
354 research to include activities aimed at inducing an osteogenic response; activities which
355 imposes strain and impact to the bones (Ehrlich & Lanyon, 2002; Hinton, Nigh, & Thyfault,
356 2015; Kato et al., 2006). The study by Gonzalez-Agüero and colleagues used plyometric
357 training and the study by Hemayatta's lab used running, hopping and skipping exercises. In the
358 present study, we did not monitor what type of activities were performed, and the
359 accelerometers used were not able to distinguish between activities at this level. Additionally,
360 the present intervention had a duration of 14 weeks, which is a short time to observe substantial
361 adaptations in bone structure as a result of training (Ahola, Korpelainen, Vainionpää,
362 Leppäluoto, & Jämsä, 2009; Harding & Beck, 2017). Finally, previous studies investigated

363 children and adolescents with ID, while this study concerned adults. With regards to the clinical
364 relevance of the results, the increase of 0.017 g/cm^2 of the right femur neck is within the range
365 seen in systematic reviews of exercise studies on adults from the general population (Bolam,
366 Van Uffelen, & Taaffe, 2013; James & Carroll, 2010), and it may be of significance to the
367 fracture risk of the subjects (Kemmler, Bebenek, Kohl, & von Stengel, 2015).
368 At PRE-tests, the bone health of the participants in the present study was in agreement with
369 previous research, regarding the BMD of the whole body (Geijer, Stanish, Draheim, & Dengel,
370 2014) and of the lumbar vertebrae (Angelopoulou, Souftas, Sakadamis, & Mandroukas, 1999).
371 Our study population had higher T-scores and healthier distribution of T-scores of the right
372 femur neck, than what was observed in earlier cross-sectional studies on similar populations
373 (Chen, Chen, Lin, & Chen, 2015; Zylstra et al., 2008). The lowest average T-score of the right
374 femur neck in the present study was seen in the 1st-year group (-0.40), which was higher than
375 the value reported by Zylstra and colleagues (-1.51) (Zylstra et al., 2008). Zylstra and
376 colleagues reported an osteoporosis prevalence ($T \leq -2.5$) in the femur region of 22% and an
377 osteopenia prevalence ($-2.5 > T \geq -1$) of 33% in 36 subjects with ID aged 19-29 years. The
378 prevalence distribution of all participants in the present study was combined for comparison
379 with the data reported by Zylstra and colleagues. We observed an osteoporosis prevalence of
380 0% and an osteopenia prevalence of 25.4%. Zylstra and colleagues did not report which model
381 of DXA-scanner they used in their study, which could influence the results. In summary, adults
382 with ID is a population with low T-scores of the femur and lumbar regions, but the population
383 in the present study are not in as high risk of developing osteoporosis, as earlier studies have
384 reported (Jasien, Daimon, Maudsley, Shapiro, & Martin, 2012; Zylstra et al., 2008).

385

386 **4.3 Cardiovascular fitness**

387 The PA-group reduced their heart rate during the last 30 seconds on both speeds during
388 the treadmill-test (Figure 3), but we did not find any changes in the amount of O_2 consumed
389 per kg body weight during the submaximal exercise transitions on the treadmill test. The
390 reductions in HR point towards an improved cardiovascular function; the subjects' hearts were
391 able to meet the energy demand of the working muscles during the walking test with less beats
392 per minute, possibly due to an increase in stroke volume (Hartley et al., 1969; Wilmore et al.,
393 2001). The increase in cardiovascular fitness observed in the present study is in line with what
394 was reported by earlier studies with similar target groups and intervention durations. Barwick
395 and colleagues reported a decrease in HR in a after a 10 week functional training program
396 (Barwick et al., 2012) and studies employing combined training programs observed increases

397 in peak O₂ uptake velocity (VO₂peak) post intervention (Calders et al., 2011; Oviedo et al.,
398 2014). The set-up of the present study complicates the comparison with the mentioned
399 traditional training studies, especially with regards to the physical load imposed on the subjects.
400 In the present study, we investigated the amount of PA performed by the PA-group while at
401 SSADD and in their leisure (Figure 2), while traditional training studies delivered and
402 monitored the PA conducted by the subjects. The knowledge obtained from these earlier studies
403 about the effect of PA on cardiovascular fitness, together with the results of the present study,
404 suggests a link between the PA performed at SSADD and the cardiovascular adaptations
405 observed. Measures of cardiac function such as reduced resting HR and improved HR recovery
406 reduces all-cause mortality from cardiovascular diseases in the general population (Adabag et
407 al., 2008; Reil et al., 2011). We did not obtain these measures in the present study, but a reduced
408 HR at the same absolute intensities could point towards an improved health-enhancing cardiac
409 function.

410

411 **4.4 Limitations**

412 The quasi-randomized design of the present study poses a selection bias. Individuals who
413 choose to attend SSADD could have an overall higher interest in PA, than the general
414 population with ID. We tried to meet this challenge by including individuals in our control
415 group, who were recreationally physically active in their leisure. The researchers observed the
416 PA occasionally, but did not supervise the PA performed by the participants in the PA-group.
417 Instead, this study focused on measuring the effects of an already existing PA option for the
418 target group. Other studies have employed a similar approach, with regards to investigating
419 already existing and well established offers and programs with PA for individuals with ID, and
420 have found health-enhancing effects as well (Barak & Gan, 2019; Messiah et al., 2019;
421 Rubenstein et al., 2020). This approach is valuable, when investigating the effects of existing
422 solutions aimed upon the societal problem of the poor health status of individuals with ID. It
423 would have been preferable to conduct accelerometry in the CON-group too, but this was not
424 possible due to logistical challenges.

425

426 **5. Conclusion**

427 This study on the effects of 14 weeks of PA and sports participation for adults with ID observed
428 positive health effects in the domains of body composition, bone health and cardiovascular
429 fitness. The PA performed at SSADD seems to be most efficient in improving the health the of

430 1st-year group. For participants who have spent one or more years at the school (2nd-year
431 group), the PA performed is likely to maintain their health in the parameters investigated in the
432 present study. A continuously progressive increase in the intensity of the PA performed at
433 SSADD could be a way to ensure continued health benefits for the participants in the 2nd year-
434 group. Improvements in the health parameters investigated in the present study could reduce
435 the groups' risk of developing lifestyle diseases.

436

437 **Acknowledgements**

438 The foundation of the project was the collaboration between the Department of Nutrition,
439 Exercise and Sports, University of Copenhagen and the Sports School for Adults with
440 Developmental Disabilities. Without the teamwork between the two institutions, this study
441 would not have been possible. The data collection in this study, relied on the great help of Jens
442 Jung Nielsen (academic staff), Rosita Olvhøj (scientific assistant), Kathrine Elsig Jensen
443 (student) and Jonas Rud Bjørndal (PhD-student). Thanks to Peter Møller Christensen (Team
444 Denmark), for guidance in the investigation of VO₂ kinetics.

445

446 **Authors' contributions**

447 JW and LMH conceived the study. EWH supervised all work related to the DXA-scans. All
448 authors contributed to the conception of the manuscript and approved the study before
449 publication.

450

451 **Conflicts of interest disclosure**

452 None of the authors have any financial or non-financial conflicts of interests, in relation to this
453 study.

454

References

- 455
456
457 Adabag, A. S., Grandits, G. A., Prineas, R. J., Crow, R. S., Bloomfield, H. E., & Neaton, J. D. (2008).
458 Relation of Heart Rate Parameters During Exercise Test to Sudden Death and All-Cause
459 Mortality in Asymptomatic Men. *American Journal of Cardiology*, *101*(10), 1437–1443.
460 <https://doi.org/10.1016/j.amjcard.2008.01.021>
- 461 Ahola, R., Korpelainen, R., Vainionpää, A., Leppäluoto, J., & Jämsä, T. (2009). Time-course of
462 exercise and its association with 12-month bone changes. *BMC Musculoskeletal Disorders*,
463 *10*(1), 8–15. <https://doi.org/10.1186/1471-2474-10-138>
- 464 Albanese, C. V., Diessel, E., & Genant, H. K. (2003). Clinical applications of body composition
465 measurements using DXA. *Journal of Clinical Densitometry*, *6*(2), 75–85.
466 <https://doi.org/10.1385/JCD:6:2:75>
- 467 Anderson, L. L., Humphries, K., McDermott, S., Marks, B., Sisarak, J., & Larson, S. (2013). The state
468 of the science of health and wellness for adults with intellectual and developmental disabilities.
469 *Intellectual and Developmental Disabilities*, *51*(5), 385–398. [https://doi.org/10.1352/1934-9556-](https://doi.org/10.1352/1934-9556-51.5.385)
470 [51.5.385](https://doi.org/10.1352/1934-9556-51.5.385)
- 471 Angelopoulou, N., Souftas, V., Sakadamis, A., & Mandroukas, K. (1999). Bone mineral density in
472 adults with Down's syndrome. *European Radiology*, *9*(4), 648–651.
473 <https://doi.org/10.1007/s003300050726>
- 474 Barak, S., & Gan, R. (2019). *The Game of Life soccer program : Effect on skills , physical fitness and*
475 *mobility in persons with intellectual disability and autism spectrum disorder*. (July 2018), 1401–
476 1411. <https://doi.org/10.1111/jar.12620>
- 477 Bartlo, P., & Klein, P. J. (2011). Physical activity benefits and needs in adults with intellectual
478 disabilities: Systematic review of the literature. *American Journal on Intellectual and*
479 *Developmental Disabilities*, *116*(3), 220–232. <https://doi.org/10.1352/1944-7558-116.3.220>
- 480 Barwick, R. B., Tillman, M. D., Stopka, C. B., Dipnarine, K., Delisle, A., & Sayedul Huq, M. (2012).
481 Physical capacity and functional abilities improve in young adults with intellectual disabilities
482 after functional training. *Journal of Strength and Conditioning Research*, *26*(6), 1638–1643.
483 <https://doi.org/10.1519/JSC.0b013e318234e860>
- 484 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
485 lme4. *Journal of Statistical Software*, *67*(1). <https://doi.org/10.18637/jss.v067.i01>
- 486 Benichou, O., & Lord, S. R. (2016). Rationale for Strengthening Muscle to Prevent Falls and
487 Fractures: A Review of the Evidence. *Calcified Tissue International*, *98*(6), 531–545.
488 <https://doi.org/10.1007/s00223-016-0107-9>
- 489 Blake, G. M., & Fogelman, I. (2007). The role of DXA bone density scans in the diagnosis and
490 treatment of osteoporosis. *Postgraduate Medical Journal*, *83*(982), 509–517.
491 <https://doi.org/10.1136/pgmj.2007.057505>

- 492 Boer, P. H., Meeus, M., Terblanche, E., Rombaut, L., De Wandele, I., Hermans, L., ... Calders, P.
493 (2014). The influence of sprint interval training on body composition, physical and metabolic
494 fitness in adolescents and young adults with intellectual disability: A randomized controlled
495 trial. *Clinical Rehabilitation*, 28(3), 221–231. <https://doi.org/10.1177/0269215513498609>
- 496 Bolam, K. A., Van Uffelen, J. G. Z., & Taaffe, D. R. (2013). The effect of physical exercise on bone
497 density in middle-aged and older men: A systematic review. *Osteoporosis International*, 24(11),
498 2749–2762. <https://doi.org/10.1007/s00198-013-2346-1>
- 499 Bouzas, S., Ayán, C., & Martínez-Lemos, R. I. (2018). Effects of exercise on the physical fitness level
500 of adults with intellectual disability: a systematic review. *Disability and Rehabilitation*, 8288, 1–
501 23. <https://doi.org/10.1080/09638288.2018.1491646>
- 502 Bull, F. C., Al-Ansari, S. S., Biddle, S., Borodulin, K., Buman, M. P., Cardon, G., ... Willumsen, J. F.
503 (2020). World Health Organization 2020 guidelines on physical activity and sedentary
504 behaviour. *British Journal of Sports Medicine*, 54(24), 1451–1462.
505 <https://doi.org/10.1136/bjsports-2020-102955>
- 506 Calders, P., Elmahgoub, S., De Mettelinge, T. R., Vandebroek, C., Dewandele, I., Rombaut, L., ...
507 Cambier, D. (2011). Effect of combined exercise training on physical and metabolic fitness in
508 adults with intellectual disability: A controlled trial. *Clinical Rehabilitation*, 25(12), 1097–1108.
509 <https://doi.org/10.1177/0269215511407221>
- 510 Chen, P. H., Chen, C. Y., Lin, Y. C., & Chen, M. Y. (2015). Low bone mineral density among adults
511 with disabilities in Taiwan: A cross-sectional descriptive study. *Disability and Health Journal*,
512 8(4), 635–641. <https://doi.org/10.1016/j.dhjo.2015.03.010>
- 513 Clifton, P. M., Condo, D., & Keogh, J. B. (2014). Long term weight maintenance after advice to
514 consume low carbohydrate, higher protein diets - A systematic review and meta analysis.
515 *Nutrition, Metabolism and Cardiovascular Diseases*, 24(3), 224–235.
516 <https://doi.org/10.1016/j.numecd.2013.11.006>
- 517 Cluphf, D., O'Connor, J., & Vanin, S. (2001). Effects of aerobic dance on the cardiovascular
518 endurance of adults with intellectual disabilities. *Adapted Physical Activity Quarterly*, 18(1), 60–
519 71. <https://doi.org/10.1123/apaq.18.1.60>
- 520 Cowley, P. M., Ploutz-Snyder, L. L., Baynard, T., Heffernan, K. S., Young Jae, S., Hsu, S., ...
521 Fernhall, B. (2011). The effect of progressive resistance training on leg strength, aerobic
522 capacity and functional tasks of daily living in persons with Down syndrome. *Disability and*
523 *Rehabilitation*, 33(23–24), 2229–2236. <https://doi.org/10.3109/09638288.2011.563820>
- 524 Dodd, K. J., & Shields, N. (2005). A systematic review of the outcomes of cardiovascular exercise
525 programs for people with Down syndrome. *Archives of Physical Medicine and Rehabilitation*,
526 86(10), 2051–2058. <https://doi.org/10.1016/j.apmr.2005.06.003>
- 527 Donnelly, J. E., BLAIR, S. N., JAKICIC, J. M., MANORE, M. M., RANKIN, J. W., & SMITH, B. K.
528 (2009). Appropriate Physical Activity Intervention Strategies for Weight Loss and Prevention of

- 529 Weight Regain for Adults. *Medicine & Science in Sports & Exercise*, 41(2). Retrieved from
530 [https://journals.lww.com/acsm-](https://journals.lww.com/acsm-msse/Fulltext/2009/02000/Appropriate_Physical_Activity_Intervention.26.aspx)
531 [msse/Fulltext/2009/02000/Appropriate_Physical_Activity_Intervention.26.aspx](https://journals.lww.com/acsm-msse/Fulltext/2009/02000/Appropriate_Physical_Activity_Intervention.26.aspx)
- 532 Ebbeling, C. B., Ward, A., Puleo, E. M., Widrick, J., & Rippe, J. M. (1991). Development of a single-
533 stage submaximal treadmill walking test. *Medicine and Science in Sports and Exercise*, 23(8),
534 966–973. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1956273>
- 535 Ehrlich, P. J., & Lanyon, L. E. (2002). Mechanical Strain and Bone Cell Function: A Review
536 Introduction: The Strain-Adaptive Response of Bone. *Osteoporosis International*, 13, 688–700.
537 Retrieved from [https://link.springer-](https://link.springer-com.proxy.mul.missouri.edu/content/pdf/10.1007%2Fs001980200095.pdf)
538 [com.proxy.mul.missouri.edu/content/pdf/10.1007%2Fs001980200095.pdf](https://link.springer-com.proxy.mul.missouri.edu/content/pdf/10.1007%2Fs001980200095.pdf)
- 539 Elmahgoub, S. S., Calders, P., Lambers, S., Stegen, S. M., Van Laethem, C., & Cambier, D. C.
540 (2011). The effect of combined exercise training in adolescents who are overweight or obese
541 with intellectual disability: the role of training frequency. *Journal of Strength and Conditioning*
542 *Research*, 25(8), 2274–2282. <https://doi.org/10.1519/JSC.0b013e3181f11c41>
- 543 Finelli, C., Sommella, L., Gioia, S., La, N., & Tarantino, G. (2013). Should visceral fat be reduced to
544 increase longevity? *Ageing Research Reviews*, 12(4), 996–1004.
545 <https://doi.org/10.1016/j.arr.2013.05.007>
- 546 Flachs, E. M., Uldall, P., & Juel, K. (2014). *Sundhedstilstanden blandt voksne med*
547 *udviklingshæmning*. København K.
- 548 Geijer, J. R., Stanish, H. I., Draheim, C. C., & Dengel, D. R. (2014). Bone mineral density in adults
549 with down syndrome, intellectual disability, and nondisabled adults. *American Journal on*
550 *Intellectual and Developmental Disabilities*, 119(2), 107–114. [https://doi.org/10.1352/1944-](https://doi.org/10.1352/1944-7558-119.2.107)
551 [7558-119.2.107](https://doi.org/10.1352/1944-7558-119.2.107)
- 552 González-Agüero, A., Vicente-Rodríguez, G., Gómez-Cabello, A., Ara, I., Moreno, L. A., & Casajús,
553 J. A. (2012). A 21-week bone deposition promoting exercise programme increases bone mass in
554 young people with Down syndrome. *Developmental Medicine and Child Neurology*, 54(6), 552–
555 556. <https://doi.org/10.1111/j.1469-8749.2012.04262.x>
- 556 Harding, A., & Beck, B. (2017). Exercise, Osteoporosis, and Bone Geometry. *Sports*, 5(2), 29.
557 <https://doi.org/10.3390/sports5020029>
- 558 Harris, L., Hankey, C., Murray, H., & Melville, C. (2015). The effects of physical activity
559 interventions on preventing weight gain and the effects on body composition in young adults
560 with intellectual disabilities: systematic review and meta-analysis of randomized controlled
561 trials. *Clinical Obesity*, 5(4), 198–210. <https://doi.org/10.1111/cob.12103>
- 562 Hartley, L. H., Grimby, G., Kilbom, Å., Nilsson, N. J., Åstrand, I., Bjure, J., ... Saltin, B. (1969).
563 Physical training in sedentary middle-aged and older men III. Cardiac output and gas exchange
564 at submaximal and maximal exercise. *Scandinavian Journal of Clinical and Laboratory*
565 *Investigation*, 24(4), 335–344. <https://doi.org/10.3109/00365516909080170>

- 566 Hemayattalab, R. (2010). Effects of physical training and calcium intake on bone mineral density of
567 participants with mental retardation. *Research in Developmental Disabilities, 31*(3), 784–789.
568 <https://doi.org/10.1016/j.ridd.2010.02.002>
- 569 Hinton, P. S., Nigh, P., & Thyfault, J. (2015). Effectiveness of resistance training or jumping-exercise
570 to increase bone mineral density in men with low bone mass: A 12-month randomized, clinical
571 trial. *Bone, 79*, 203–212. <https://doi.org/10.1016/j.bone.2015.06.008>
- 572 Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models.
573 *Biometrical Journal. Biometrische Zeitschrift, 50*(3), 346–363.
574 <https://doi.org/10.1002/bimj.200810425>
- 575 James, M. M. S., & Carroll, S. (2010). Effects of different impact exercise modalities on bone mineral
576 density in premenopausal women: A meta-analysis. *Journal of Bone and Mineral Metabolism,*
577 *28*(3), 251–267. <https://doi.org/10.1007/s00774-009-0139-6>
- 578 Jasien, J., Daimon, C. M., Maudsley, S., Shapiro, B. K., & Martin, B. (2012). Aging and bone health
579 in individuals with developmental disabilities. *International Journal of Endocrinology, 2012*.
580 <https://doi.org/10.1155/2012/469235>
- 581 Kato, T., Terashima, T., Yamashita, T., Hatanaka, Y., Honda, A., & Umemura, Y. (2006). Effect of
582 low-repetition jump training on bone mineral density in young women. *Journal of Applied*
583 *Physiology, 100*(3), 839–843. <https://doi.org/10.1152/jappphysiol.00666.2005>
- 584 Kemmler, W., Bebenek, M., Kohl, M., & von Stengel, S. (2015). Exercise and fractures in
585 postmenopausal women. Final results of the controlled Erlangen Fitness and Osteoporosis
586 Prevention Study (EFOPS). *Osteoporosis International, 26*(10), 2491–2499.
587 <https://doi.org/10.1007/s00198-015-3165-3>
- 588 Kuk, J. L., Katzmarzyk, P. T., Nichaman, M. Z., Church, T. S., Blair, S. N., Ross, R., ... Ross, R.
589 (2006). *Visceral Fat Is an Independent Predictor of All-cause Mortality in Men. 14*(2).
- 590 Lee, S. W., Young, J., Jeong, S., Kim, M., Hwang, S., Han, J. S., & Heo, N. J. (2018). *Body fat*
591 *distribution is more predictive of all-cause mortality than overall adiposity.* (April 2017), 141–
592 147. <https://doi.org/10.1111/dom.13050>
- 593 Lizondo, V., Caplliure-Llopis, J., Escrivá, D., De La Rubia, J. E., & Barrios, C. (2019). Bone quality
594 in young adults with intellectual disability involved in adapted competitive football. *European*
595 *Journal of Sport Science, 19*(6), 850–859. <https://doi.org/10.1080/17461391.2018.1563633>
- 596 Martínez-Zaragoza, F., Campillo-Martínez, J. M., & Ato-García, M. (2016). Effects on Physical
597 Health of a Multicomponent Programme for Overweight and Obesity for Adults with Intellectual
598 Disabilities. *Journal of Applied Research in Intellectual Disabilities, 29*(3), 250–265.
599 <https://doi.org/10.1111/jar.12177>
- 600 Melville, C. A., Hamilton, S., Hankey, C. R., Miller, S., & Boyle, S. (2007). The prevalence and
601 determinants of obesity in adults with intellectual disabilities. *Obesity Reviews, 8*(3), 223–230.
602 <https://doi.org/10.1111/j.1467-789X.2006.00296.x>

- 603 Messiah, S. E., D'Agostino, E. M., Patel, H. H., Hansen, E., Mathew, M. S., & Arheart, K. L. (2019).
604 Changes in cardiovascular health and physical fitness in ethnic youth with intellectual
605 disabilities participating in a park-based afterschool programme for two years. *Journal of*
606 *Applied Research in Intellectual Disabilities*, 32(6), 1478–1489.
607 <https://doi.org/10.1111/jar.12642>
- 608 Miller, C. T., Fraser, S. F., Straznicki, N. E., Dixon, J. B., Selig, S. E., & Levinger, I. (2013). Effect
609 of Diet versus Diet and Exercise on Weight Loss and Body Composition in Class II and III
610 Obesity: A Systematic Review. *Journal of Diabetes & Metabolism*, 04(06), 2–7.
611 <https://doi.org/10.4172/2155-6156.1000281>
- 612 Obrusnikova, I., Firkin, C. J., Cavalier, A. R., & Suminski, R. R. (2021). Effects of resistance training
613 interventions on muscular strength in adults with intellectual disability: a systematic review and
614 meta-analysis. *Disability and Rehabilitation*, 0(0), 1–14.
615 <https://doi.org/10.1080/09638288.2021.1910738>
- 616 Ordoñez, F. J., Rosety, M., & Rosety-Rodriguez, M. (2006). Influence of 12-week exercise training
617 on fat mass percentage in adolescents with Down syndrome. *Medical Science Monitor*, 12(10),
618 2004–2007.
- 619 Oviedo, G. R., Guerra-Balic, M., Baynard, T., & Javierre, C. (2014). Effects of aerobic, resistance and
620 balance training in adults with intellectual disabilities. *Research in Developmental Disabilities*,
621 35(11), 2624–2634. <https://doi.org/10.1016/j.ridd.2014.06.025>
- 622 *R statistics 4.0.0*. (2020). Retrieved from <https://www.r-project.org/>
- 623 Reil, J. C., Custodis, F., Swedberg, K., Komajda, M., Borer, J. S., Ford, I., ... Böhm, M. (2011). Heart
624 rate reduction in cardiovascular disease and therapy. *Clinical Research in Cardiology*, 100(1),
625 11–19. <https://doi.org/10.1007/s00392-010-0207-x>
- 626 Robertson, J., Emerson, E., Gregory, N., Hatton, C., Turner, S., Kessissoglou, S., & Hallam, A.
627 (2000). Lifestyle related risk factors for poor health in residential settings for people with
628 intellectual disabilities. *Research in Developmental Disabilities*, 21(6), 469–486.
629 [https://doi.org/10.1016/S0891-4222\(00\)00053-6](https://doi.org/10.1016/S0891-4222(00)00053-6)
- 630 Rubenstein, E., DuBois, L., Sadowsky, M., Washburn, K., Forquer, M., Stanish, H., & Shriver, T.
631 (2020). Evaluating the potential of Special Olympics fitness models as a health intervention for
632 adults with intellectual disabilities. *Disability and Health Journal*, 13(2), 100850.
633 <https://doi.org/10.1016/j.dhjo.2019.100850>
- 634 Shields, N., Taylor, N. F., Wee, E., Wollersheim, D., O'Shea, S. D., & Fernhall, B. (2013). A
635 community-based strength training programme increases muscle strength and physical activity
636 in young people with Down syndrome: A randomised controlled trial. *Research in*
637 *Developmental Disabilities*, 34(12), 4385–4394. <https://doi.org/10.1016/j.ridd.2013.09.022>
- 638 Skotte, J., Korshøj, M., Kristiansen, J., Hanisch, C., & Holtermann, A. (2014). Detection of physical
639 activity types using triaxial accelerometers. *Journal of Physical Activity and Health*, 11(1), 76–

- 640 84. <https://doi.org/10.1123/jpah.2011-0347>
- 641 Son, S., Jeon, B., & Kim, H. (2016). Effects of a walking exercise program for obese individuals with
642 intellectual disability staying in a residential care facility. *Journal of Physical Therapy Science*,
643 28(3), 788–793. <https://doi.org/10.1589/jpts.28.788>
- 644 Srikanth, R., Cassidy, G., Joiner, C., & Teeluckdharry, S. (2011). Osteoporosis in people with
645 intellectual disabilities: A review and a brief study of risk factors for osteoporosis in a
646 community sample of people with intellectual disabilities. *Journal of Intellectual Disability*
647 *Research*, 55(1), 53–62. <https://doi.org/10.1111/j.1365-2788.2010.01346.x>
- 648 Steell, L., Ho, F. K., Sillars, A., Petermann-Rocha, F., Li, H., Lyall, D. M., ... Celis-Morales, C. A.
649 (2019). Dose-response associations of cardiorespiratory fitness with all-cause mortality and
650 incidence and mortality of cancer and cardiovascular and respiratory diseases: The UK Biobank
651 cohort study. *British Journal of Sports Medicine*, 53(21), 1371–1378.
652 <https://doi.org/10.1136/bjsports-2018-099093>
- 653 Wewege, M., van den Berg, R., Ward, R. E., & Keech, A. (2017). The effects of high-intensity
654 interval training vs. moderate-intensity continuous training on body composition in overweight
655 and obese adults: a systematic review and meta-analysis. *Obesity Reviews*, 18(6), 635–646.
656 <https://doi.org/10.1111/obr.12532>
- 657 Wilmore, J. H., Stanforth, P. R., Gagnon, J., Rice, T., Mandel, S., Leon, A. S., ... Bouchard, C.
658 (2001). Cardiac output and stroke volume changes with endurance training: the HERITAGE
659 Family Study. *Medicine and Science in Sports and Exercise*, 33(1), 99–106.
660 <https://doi.org/10.1097/00005768-200101000-00016>
- 661 Wu, W. L., Yang, Y. F., Chu, I. H., Hsu, H. T., Tsai, F. H., & Liang, J. M. (2017). Effectiveness of a
662 cross-circuit exercise training program in improving the fitness of overweight or obese
663 adolescents with intellectual disability enrolled in special education schools. *Research in*
664 *Developmental Disabilities*, 60, 83–95. <https://doi.org/10.1016/j.ridd.2016.11.005>
- 665 Zylstra, R. G., Porter, L. L., Shapiro, J. L., & Prater, C. D. (2008). Prevalence of Osteoporosis in
666 Community-Dwelling Individuals with Intellectual and/or Developmental Disabilities. *Journal*
667 *of the American Medical Directors Association*, 9(2), 109–113.
668 <https://doi.org/10.1016/j.jamda.2007.09.008>
- 669
- 670

Table 1

Demographic data of participants from their first measurement

Variables	PA	1st	2nd	CON
Demographic data				
Number of participants	52	32	20	14
Gender (f/m)	28/24	17/15	11/9	5/9
Age (years)	26.7 ± 1.2	26.7 ± 1.5	26.7 ± 1.4	29.1 ± 1.7
Height (cm)	163.3 ± 1.8	162.3 ± 2.2	165.1 ± 3.0	163.3 ± 2.4
Developmental disorder (DS/CP/UD)	15/6/31	12/3/19	5/3/12	4/1/9

671 The number of participants in the study groups. The participants' age was recorded at PRE-
672 tests. Abbreviations: f: female, m: male, cm: centimetres, DS: Downs' Syndrome, CP: Cerebral
673 Palsy, UD: Undefin

Table 2
Body composition and bone health measures at PRE-tests

Variables	PA	PRE		
		1st	2nd	Control
Body composition				
Body weight (kg)	73.90 ± 2.11	75.98 ± 3.46^b	72.42 ± 2.65	75.12 ± 5.72
Fat mass (kg)	25.74 ± 1.55	29.25 ± 2.45^{bb}	25.96 ± 3.87	25.74 ± 4.3
Fat free mass (kg)	44.14 ± 1.06	44.24 ± 1.62	44.08 ± 1.42	47.11 ± 1.96
Visceral fat (kg)	2.46 ± 0.18	2.71 ± 0.31^{bb}	2.27 ± 0.21	2.27 ± 0.46
Gynoid fat (kg)	4.70 ± 0.26	4.96 ± 0.40^b	4.52 ± 0.51	4.48 ± 0.70
BMI (kg/m ²)	28.19 ± 0.88	29.05 ± 1.36^b	27.58 ± 1.16	28.05 ± 2.45
Whole body fat percent	36.72 ± 1.35	38.12 ± 2.14^{bb}	35.73 ± 1.74	32.74 ± 2.51
Bone health				
Whole body (g/cm ²)	1.155 ± 0.014	1.163 ± 0.026	1.149 ± 0.015	1.208 ± 0.021
Whole body T-score	0.26 ± 0.126	0.40 ± 0.239	0.16 ± 0.136	0.47 ± 0.245
L1-L4 (g/cm ²)	1.140 ± 0.016^a	1.129 ± 0.026	1.150 ± 0.020	1.217 ± 0.031
L1-L4 T-score	-0.53 ± 0.140	-0.56 ± 0.262	-0.51 ± 0.158	-0.43 ± 0.288
Left total femur (g/cm ²)	0.981 ± 0.015	0.980 ± 0.028	0.981 ± 0.017	1.023 ± 0.028
Left total femur T-score	-0.44 ± 0.119	-0.43 ± 0.225	-0.45 ± 0.133	-0.27 ± 0.259
Right total femur (g/cm ²)	0.981 ± 0.015	0.987 ± 0.025	0.976 ± 0.018	1.027 ± 0.028
Right total femur T-score	-0.46 ± 0.114	-0.40 ± 0.203	-0.50 ± 0.134	-0.24 ± 0.257
Left femur neck (g/cm ²)	0.972 ± 0.015	0.960 ± 0.028	0.981 ± 0.018	1.007 ± 0.027
Left femur neck T-score	-0.36 ± 0.116	-0.46 ± 0.210	-0.30 ± 0.134	-0.24 ± 0.257
Right femur neck (g/cm ²)	0.980 ± 0.015	0.971 ± 0.024	0.986 ± 0.019	0.995 ± 0.029
Right femur neck T-score	-0.32 ± 0.109	-0.40 ± 0.178	-0.26 ± 0.140	-0.33 ± 0.271

674 PRE-test values as raw means ± standard error of the mean (SEM). Abbreviations:

675 Significance codes: **a**: Difference compared to the Control group. **b**: Difference compared to

676 2nd year participants. p<0.001: "aaa / bbb" 0.001>p<0.01: "aa/bb ", 0.01>p<0.05: "" a/b ".
677


Table 3

Changes within and between groups over time in body composition and bone health.

Variables	Within group Δ -values from PRE to POST tests				Between group change scores from PRE to POST tests			
	Δ PA	Δ 1st	Δ 2nd	Δ Con	Δ PA - Δ Con	Δ 1st - Δ Con	Δ 2nd - Δ Con	Δ 1st - Δ 2nd
Body composition								
Body weight (kg)	-0.44 \pm 0.52	-1.10 \pm 0.72	0.01 \pm 0.69	-0.31 \pm 1.16	0.14 \pm 1.25	-1.05 \pm 1.31	0.06 \pm 1.27	-1.11 \pm 0.99
Fat mass (kg)	-1.02 \pm 0.43*	-1.71 \pm 0.59*	-0.53 \pm 0.56	0.23 \pm 0.94	-1.24 \pm 1.02	-2.18 \pm 1.07	-1.00 \pm 1.04	-1.18 \pm 0.81
Fat free mass (kg)	0.29 \pm 0.17	0.22 \pm 0.25	0.31 \pm 0.24	0.24 \pm 0.38	0.04 \pm 0.41	-0.06 \pm 0.45	0.04 \pm 0.44	-0.09 \pm 0.35
Visceral fat (kg)	-0.08 \pm 0.06	-0.20 \pm 0.08*	0.01 \pm 0.07	0.04 \pm 0.13	-0.12 \pm 0.14	-0.028 \pm 0.14	-0.07 \pm 0.14	-0.21 \pm 0.11
Gynoid fat (kg)	-0.27 \pm 0.08*	-0.35 \pm 0.11*	-0.23 \pm 0.11	-0.06 \pm 0.18	-0.22 \pm 0.20	-0.32 \pm 0.21	-0.21 \pm 0.20	-0.11 \pm 0.16
BMI (kg/m ²)	-0.18 \pm 0.19	-0.40 \pm 0.26	0.00 \pm 0.25	0.07 \pm 0.42	-0.11 \pm 0.45	-0.43 \pm 0.48	-0.01 \pm 0.47	-0.42 \pm 0.36
Whole body fat percent	-0.97 \pm 0.32**	-1.46 \pm 0.45**	-0.62 \pm 0.43	-0.16 \pm 0.72	-0.81 \pm 0.78	-1.47 \pm 0.82	-0.63 \pm 0.80	-0.84 \pm 0.63
Bone health (BMD)								
Whole body (g/cm ²)	0.008 \pm 0.004	0.004 \pm 0.006	0.010 \pm 0.005	-0.016 \pm 0.008	0.024 \pm 0.009*	0.019 \pm 0.010	0.025 \pm 0.010*	-0.006 \pm 0.008
L1-L4 (g/cm ²)	0.005 \pm 0.008	0.003 \pm 0.011	0.006 \pm 0.011	-0.038 \pm 0.017	0.043 \pm 0.019*	0.040 \pm 0.021	0.043 \pm 0.020	-0.003 \pm 0.016
Left total femur (g/cm ²)	0.002 \pm 0.008	0.001 \pm 0.005	0.003 \pm 0.005	-0.007 \pm 0.007	0.009 \pm 0.008	0.008 \pm 0.009	0.010 \pm 0.009	-0.002 \pm 0.007
Right total femur (g/cm ²)	0.002 \pm 0.003	0.003 \pm 0.006	0.011 \pm 0.006	-0.007 \pm 0.007	0.009 \pm 0.008	0.003 \pm 0.011	0.012 \pm 0.011	-0.009 \pm 0.008
Left femur neck (g/cm ²)	0.005 \pm 0.008	0.003 \pm 0.011	0.006 \pm 0.011	-0.038 \pm 0.017	0.043 \pm 0.102*	0.040 \pm 0.021	0.043 \pm 0.016	-0.003 \pm 0.016
Rigt femur neck (g/cm ²)	0.017 \pm 0.007*	0.017 \pm 0.010	0.017 \pm 0.009	0.007 \pm 0.015	0.010 \pm 0.016	0.012 \pm 0.0178	0.012 \pm 0.017	0.000 \pm 0.013

678 Changes within the groups are reported as delta values from PRE to POST tests (POST - PRE) \pm standard error of estimate. Differences between
 679 groups are reported as change scores (i.e. the differences between the delta-values of the groups) \pm standard error of estimate. Abbreviations: BMI:
 680 Body Mass Index, BMD: Bone Mineral Density, L1- L4: Lumbar vertebrae 1 to 4, kg: kilogram, g/cm²: .gram per square centimetre. Significance
 681 codes: p<0.001: "****" 0.001>p<0.01: "***", 0.01>p<0.05: "**", 0.05>p<0.1: "."

682




683

684 Figure 1: Flow chart of the movements between the groups of the participants between the two
685 rounds of experiments, from August 2019 to August 2020.

686

687




688


689 Figure 2: The amount of steps taken pr. hour while the participants were at SSADD, during
690 leisure on weekdays after the school day ends and during weekends. Measurements were made
691 over a 7 day period. Error bars: Standard error of the mean. Significance codes: $p < 0.001$: "****"
692 $0.001 > p < 0.01$: "***", $0.01 > p < 0.05$: "*", $0.05 > p < 0.1$: "."

693

A



B



694

695 Figure 3: The average heart rates during the last 30 s. at 3.2 km/h (A) and 4.8 km/h (B) for the

696 PA-, 1st-year, 2nd-year and CON-group. Significance codes: $p < 0.001$: "****" $0.001 > p < 0.01$:

697 "***", $0.01 > p < 0.05$: "**", $0.05 > p < 0.1$: ".

Appendix 1

Participants, characteristics and distribution on the different outcomes.

Tests	Numbers of measurements							
	PA		1st		2nd		Control	
	PRE	POST	PRE	POST	PRE	POST	PRE	POST
DXA scans	76	73	31	30	45	43	18	15
HR (3.2 km/t)	70	72	26	30	44	42	17	13
HR (4.8 km/t)	64	62	23	22	41	40	15	12
Accelerometry	43	-	21	-	22	-	-	-

	Age (years ± SD)							
	PA		1st		2nd		Control	
	PRE	POST	PRE	POST	PRE	POST	PRE	POST
DXA scans	26.7±8.3	26.8±8.0	26.8±9.0	27.0±9.2	26.7±8.0	26.4±7.1	28.9±5.8	29.7±6.2
HR (3.2 km/t)	26.7±7.8	26.5±8.0	26.7±7.6	27.1±9.1	26.7±8.0	26.0±7.1	28.9±6.0	28.7±4.7
HR (4.8 km/t)	27.1±8.1	26.5±7.5	27.3±7.9	27.3 ±8.1	26.9±8.3	26.1±7.2	29.7±5.9	28.1±4.5

	Gender (f/m)							
	PA		1st		2nd		Control	
	PRE	POST	PRE	POST	PRE	POST	PRE	POST
DXA scans	42/34	40/33	16/15	15/15	26/19	25/18	6/12	6/9
HR (3.2 km/h)	38/32	38/34	13/13	15/15	25/19	23/19	5/12	4/9
HR (4.8 km/h)	34/30	31/31	11/12	9/13	23/18	22/18	3/12	4/8

698 An overview of how the Abbreviations: DXA: Dual X-ray Absorptiometry HR: heart rate,

699 km/h: kilometres per hour SD: standard deviation, f: female, m: male.