Modelling the impact of combining HIV prevention interventions on HIV dynamics in fishing communities in Uganda

Cécile Kremer[1,*], Anatoli Kamali[2], Monica Kuteesa[3], Janet Seeley[3,4] , Niel Hens[1,5] , Rebecca N Nsubuga[4]

[1] Interuniversity Institute for Biostatistics and Bioinformatics (I-BioStat), Data Science Institute, Hasselt University, Hasselt, Belgium

[2] International AIDS Vaccine Initiative, Nairobi, Kenya

[3] London School of Hygiene and Tropical Medicine, London, United Kingdom

[4] Medical Research Council/Uganda Virus Research Unit & LSHTM Uganda Research Unit (MRC/UVRI & LSHTM), Entebbe, Uganda

[5] Centre for Health Economics Research and Modelling Infectious Diseases (CHERMID), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

[*] Corresponding author: cecile.kremer@uhasselt.be

Abstract

Background. In countries with mature generalized HIV epidemics such as Uganda, there are still groups of individuals that are disproportionately affected. These most at risk populations (MARPs) have high HIV prevalence compared to the general population. Among the MARPs in Uganda are fishing communities, which make up about 10% of the population.

Methods. We investigated the impact of combined HIV prevention interventions on HIV transmission dynamics in high-risk fishing communities in Uganda using a deterministic compartmental model. The model was calibrated to seroprevalence data from a census performed in 2014. To account for remaining uncertainty in the calibrated model parameters, an additional 50 000 simulated scenarios were modelled to investigate the impact of combined interventions.

Results. The projected HIV incidence decreased from 1.87 per 100 PY without intervention scale-up to 0.25 per 100 PY after 15 years (2014-2029) of intervention scale-up. A potential combination achieving this 87% reduction in incidence included condom use in about 60% of sexual acts, 23% of susceptible men circumcised, 87% of people living with HIV aware of their status, 75% of those on ART, and about 3% of susceptible individuals on oral PrEP. Uncertainty analysis revealed relative reductions in incidence ranging from 30.9% to 86.8%. Sensitivity analyses suggested that condom use and early ART were the most important intervention components.

Conclusions. Reducing HIV incidence, and consequently prevalence and AIDS-related mortality, in these high-risk fishing communities in Uganda is attainable in the long-run with a combination prevention package. Our projected intervention coverage levels are well within the national targets set by the Uganda government and close to reaching the UNAIDS 95-95-95 goal to end AIDS by 2030.

Keywords

HIV; combination prevention; fishing communities; mathematical modelling

Background

With 1.5 million new infections in 2021, HIV-1 remains a major global health problem. Of these new infections, 45% occurred in Eastern and Southern Africa (1). In countries with mature generalized epidemics such as Uganda, there are still groups of individuals that are disproportionately affected by HIV. These most at risk populations (MARPs) have high HIV prevalence compared to the general population, where prevalence has remained stable between 6 and 7% since 2001 (2). Among the MARPs in Uganda are fishing communities (FCs), which make up about 10% of the population and were until recently, due to their high mobility (3), believed to serve as reservoirs of infection for the general population (4,5). Several studies have indicated that HIV incidence in Ugandan FCs is up to 5 times, or more than 10 times depending on the community, higher than the national average, and therefore greatly contributes to the overall HIV burden in Uganda (6,7). This high incidence in FCs has been attributed to several factors including limited access to HIV prevention and treatment services as well as high-risk sexual behaviour (6), such as frequent partner change and low condom use (3,4).

Several HIV prevention interventions have been proven to effectively reduce HIV transmission. It has been shown that voluntary medical male circumcision (VMMC) decreases male susceptibility by 60%, while circumcised men living with HIV are as infectious as non-circumcised men (8). HIV counselling and testing (HCT) may lead to adoption of less risky behaviour, such as increased condom use and reduced sexual activity, as well as increased uptake of VMMC and oral pre-exposure prophylaxis (PrEP). On the other hand however, HIV-negative persons who repeatedly accept HCT are more likely to engage in high-risk sexual behaviour (9). Oral PrEP has been shown to reduce susceptibility by 67-75% (10), but high adherence to the prescribed doses is necessary for effective HIV prevention.

Another important intervention strategy is antiretroviral treatment (ART) for individuals

living with HIV. In addition to prolonged life expectancy, persons living with HIV that are on

ART have been shown to be less infectious (11). According to UNAIDS figures in 2017 (12),

68-77% of Ugandans living with HIV were receiving treatment. However, coverage in FCs is

expected to be much lower than in the general population (13), because of barriers to

treatment uptake such as stigma and inconvenience due to high mobility (14,15).

These HIV prevention strategies are however only partly effective and often not readily

available to MARPs. It has been shown that minimal overlap of partly effective interventions

increases the chances of synergistic effects (16). Cox et al. (17) found that using a

combination of complementary interventions may be more practical for achieving substantial

reductions in HIV incidence. Therefore, to achieve effective HIV control and have the

greatest sustained impact on reducing new infections, several intervention strategies need to

be combined to present a mix of effective behavioural, biomedical, and structural

interventions. The Uganda national guidelines for HIV care incorporated the UNAIDS 90-90-

90 (90% of people living with HIV aware of their status, 90% in care, and 90% virally

suppressed) targets in 2016 (18). However, HIV prevention and treatment services may not be

easily accessed by individuals living in FCs. Common barriers to accessing HIV prevention

services include a lack of social support, stigma, long distances to healthcare facilities, and

high population migration rates. Regarding HIV testing and ART, varied coverage was found

among FCs but in general was not fulfilling the UNAIDS 95-95-95 targets in order to

eliminate HIV by 2030. Hence to achieve global targets on HIV elimination, these FCs

require special attention by implementing a combination of services tailored to the specific

needs of this community (19).

Several modelling studies have shown the possible benefits of implementing combination

prevention packages for reducing HIV incidence in high-risk populations (17,20,21). In the

present study we investigated the impact of combining HIV prevention interventions on HIV

dynamics in four FCs in Uganda. The interventions considered were VMMC, condom use,

HCT, ART, and oral PrEP. The study findings provide additional knowledge on the use of

targeted intervention combinations in the fight against the HIV epidemic.

Methods

Model description

We developed a population-level, deterministic, compartmental model of heterosexual HIV

transmission incorporating different risk groups as is done in similar HIV transmission

models (20,21). In our model the population aged 16-60 years is stratified into subgroups

based on gender and sexual risk behaviour. The model consists of 14 compartments for

females and 16 for males, representing susceptibility (with or without oral PrEP, and

circumcision status for males) and treatment status (undiagnosed, diagnosed untreated, and

diagnosed treated) in four disease stages. The rate at which susceptible individuals acquire

HIV (i.e. force of infection) depends on their rate of partner change and PrEP/circumcision

status, as well as their partner's risk group, disease stage, and treatment status. It further

depends on the type of partnership (by the frequency of coital acts) and probability of condom

use per coital act (see Additional File 1). Once an individual is infected with HIV, they move

through four disease stages: (i) an initial high viremia stage (acute infection), (ii) a low

viremia stage (chronic infection), (iii) a pre-AIDS high viremia stage, and (iv) the final AIDS

stage in which they remain until death from AIDS-related or other causes. It is assumed that

individuals in the high viremia stages of infection are more infectious than those in the low

viremia stage (22). Furthermore we assume that individuals in the untreated AIDS stage cease

all sexual activity, while those in the treated AIDS stage greatly reduce their sexual activity.

Three sexual activity risk groups are defined: (i) low – individuals having only main partners,

(ii) medium – individuals having main and non-main regular partners, and (iii) high –

individuals having main, non-main regular, and casual partners. Hence an additional layer of

heterogeneity is added by including different types of partnerships, with different associated

risks, within each risk group. Although these partnerships are instantaneous in the model, the

risk of infection throughout a partnership is a function of the frequency of coital acts per

partnership, reflecting the duration and associated risk of a partnership (i.e. less coital acts

indicates shorter duration but with a higher risk of infection due to a higher background HIV

prevalence in these groups) (20). A detailed description of the model structure and parameters

is presented in Figure S1 and Additional File 1. Below we describe the intervention measures.

124 **Target population**

The target population of this study were four FCs around Lake Victoria, Uganda.

Sociodemographic and prevalence data used to inform and calibrate the model came from the

2014 census and baseline serosurvey of the HIVCOMB study, respectively (23). Briefly, the

HIVCOMB study was a pilot cluster-randomized trial assessing the feasibility of conducting

combination HIV prevention interventions. Two rural and two urban FCs were included and

in each urban-rural pair one cluster was assigned to receive the combination prevention

package while the other cluster served as a control, receiving a standard prevention package.

Before the community-wide intervention was introduced, a serosurvey was undertaken in a

simple random sample from each community to determine HIV status. HIV prevalence

estimates from this survey were used to calibrate our model to the actual situation in 2014.

135 **Model calibration**

Most of the model parameters were unknown and had to be calibrated. Uncertainty ranges for

all unknown parameters were constructed based on literature or, if unavailable, reasonably

wide ranges were assumed (Table S2, see Additional File 1). Parameters that were calibrated

included the transmission probability per coital act, parameters on baseline intervention

uptake, and all behavioural parameters. The initial population size was set to 2000 and the

proportion of individuals in each sexual risk group was fixed as those observed in the

HIVCOMB survey: 33% of males and 43% of females in the low risk group, 7% of males and

2% of females in the medium risk group, 10% of males and 5% of females in the high risk

group. The initial HIV prevalence was set to 1% in each risk group, with all of those in the

initial high viremia stage.

The model was calibrated to the 2014 HIV prevalence in each sexual risk group, obtained

from the 2014 HIVCOMB survey described above. Using Latin hypercube sampling (LHS)

(24), 10 000 parameter input sets θ were sampled from the uncertainty ranges of the unknown

parameters (Table S2, see Additional File 1). For each of these parameter sets, the model was

run for 45 years (i.e. 1969-2014) to simulate an epidemic. Since we calibrated the model to

only one time point, because no further data are available, overall goodness of fit (GOF) was

measured as the sum of the squared deviations from the target prevalence in each risk group

(25). An active learning approach, in which the parameter space is optimized in an iterative

manner (26), was used to reduce uncertainty ranges for the unknown parameters by

investigating the top 1% (based on GOF) of model outputs, $\Theta_{1\%}$ (see Additional File 1).

Interventions

HIV testing was assumed to have been available since 1987 (27). Persons living with HIV are

diagnosed according to a rate of testing uptake that changes over time such that the proportion

of infected individuals aware of their status increases linearly until reaching a specified level

(21). When tested, infected individuals move to the 'diagnosed' compartment (representing

awareness of their status) corresponding to their disease stage at that time. While diagnosed

but untreated, individuals are assumed to slightly reduce their risk behaviour due to HIV

awareness and/or counselling. Since results from studies about the effects of counselling on

risk behaviour reduction are inconsistent (28), this was implemented as a conservative

decrease of 10% in the number of sexual partners and a 10% increase in condom use.

ART was introduced in the pre-AIDS and AIDS stages from 2004 onward (29), and

additionally in the other stages in 2011 (2). Depending on disease stage, a proportion of

diagnosed individuals will initiate ART on average one year after diagnosis, and this

proportion was assumed to increase over time. ART was assumed to reduce infectiousness by

96% (11). Treatment failure was included in the ART dropout rate which was set to an annual

dropout of 20%. After ART dropout, individuals move back to the diagnosed untreated

compartment and can re-initiate treatment at the same rate as treatment-naïve individuals.

Individuals on ART were assumed to return to their baseline risk behaviour. ART and testing

uptake were assumed to be lower in males compared to females (14).

Susceptible men opting for VMMC were assumed to have accepted HIV testing. Hence,

VMMC uptake was modelled as a combination of the rate of testing uptake and the proportion

of men getting circumcised on average one year after testing, which was assumed to increase

over time. Circumcised males were assumed to have a relative susceptibility of 40% (8). Due

to possible risk compensation, no reduction in sexual risk behaviour was assumed for

circumcised males. We assumed that there was no VMMC before the introduction of HIV

testing in 1987, although a constant proportion of males enter the population already

circumcised (e.g. due to religion). The level of condom use was assumed to differ for each

type of partnership, and to have increased over time until reaching a specified level.

Oral PrEP has been available since 2015 and hence was not included in the model calibration

(2). Susceptible individuals initiating PrEP were assumed to have accepted HCT, and uptake

was modelled as the combination of an annual probability of accepting HCT and a probability

of initiating PrEP on average one year after testing, which was assumed to increase over time.

Individuals on oral PrEP were assumed to have a relative susceptibility of 33% (10), further

depending on the level of adherence which was assumed to be 60%. PrEP discontinuation was

included as a fixed annual dropout of 20%. As for VMMC, no reduction in risk behaviour was

assumed.

Assessing the impact of combined interventions

As a baseline model, intervention uptake rates were kept constant for 15 years starting in

2014, although we slightly increased the proportion of individuals aware of their HIV-positive

status to 65% to avoid zero rates of VMMC uptake. Since we had no information on the

current level of oral PrEP uptake in these FCs but which is assumed to be very low, we did

not include this in our baseline model.

The impact of intervention scale-up on HIV dynamics was investigated using the 'best fit'

calibrated model. LHS was used to sample 500 input parameter sets θ_I from the uptake ranges

of the intervention parameters (Table S3, see Additional File 1). To estimate long-term

intervention impact, the calibrated model was run for an additional 15 years (i.e. 2014 –

2029). Starting in 2014, intervention uptake was gradually increased for 6 years (i.e. until

2020), after which a more steep increase was implemented for 9 years to mimic real-life

combination prevention efforts. Different combinations of intervention uptake levels were

then compared in terms of their impact on the relative reduction in HIV incidence and

prevalence, AIDS-related mortality, and the total number of new infections, by comparing the

model with scale-up of interventions to our baseline model without increased intervention

uptake (i.e. 'best fit' intervention parameters as in Table S2, see Additional File 1).

Uncertainty analysis

210   To account for the remaining uncertainty in the parameters of the 'best fit' calibrated model,

211   which reflect only one possible epidemic trajectory, parameter ranges of simulated epidemics

212   belonging to the top 1% of model outputs were also used to investigate the impact of

213   combined interventions, resulting in 50 000 simulated scenarios (i.e. the 500 intervention

214   parameter sets θ_I applied to 100 sets of model parameters $\Theta_{1\%}$).

Sensitivity analyses

We performed various sensitivity analyses to assess the influence of certain assumptions on

the projected impact of intervention scale-up. In our best case scenario (i.e. calibrated model

resulting in the highest reduction in incidence), we lowered the annual rate of PrEP

discontinuation and ART dropout to 5%, increased the annual rate of ART dropout to 70%,

assumed no increase in condom use after 2014, no PrEP use, PrEP adherence of 90%, and

PrEP adherence of 30%. We also investigated the impact of equal HCT and ART uptake in

males and females.

Results

Model calibration

Tables 1 and 2 show the HIV prevalence estimates and intervention coverage levels,

respectively, from the 'best fit' model after completing calibration. The corresponding model

parameters are shown in Table S2 (see Additional File 1). HIV dynamics in terms of

prevalence and incidence in the different sexual risk groups for the baseline model (i.e.

without intervention scale-up) are shown in Figure S3 (see Additional File 1). It can be seen

that even without increasing the intervention uptake, HIV prevalence and incidence would

already decline after 2014 (overall, a decline of 19.89% in prevalence and 30.50% in

incidence when comparing 2029 to 2014), with steeper declines observed for females in the

medium and high risk groups.

Impact of combined interventions

Using the parameters from the 'best fit' calibrated model, the intervention uptake levels

shown in Table S3 (highest impact scenario; see Additional File 1) resulted in a relative

reduction in HIV incidence of 86.68% at the end of 15 years of intervention scale-up,

compared to a baseline scenario without scale-up. The corresponding coverage after 15 years

for each of the interventions is shown in Table 3 (see Figure S4 in Additional File 1 for

coverage levels over time), and the relative reduction in prevalence, incidence, AIDS-related

mortality, and number of new infections per risk group are shown in Table 4. Overall

incidence after 15 years was 1.87 per 100 PY in the baseline scenario compared to 0.25 per

100 PY after intervention scale-up. The relative reduction in incidence was highest for males

in the low risk group (88.15%) and lowest for females in the high risk group (83.15%). In

each risk group, relative reductions were slightly greater for males.

Figure 1 shows the evolution of HIV incidence and prevalence in the total population for the

baseline model and the model with intervention scale-up. After intervention scale-up, a much

steeper decline in prevalence and incidence is observed. The results also indicate that

implementation of combined interventions would yield a total of 408 infections averted over

15 years (989 new infections under the baseline scenario, vs. 581 new infections under the

highest impact scale-up scenario).

Table 1. Prevalence (%) observed in the HIVCOMB study and estimates of prevalence (%) and incidence (per 100 PY) from the 'best fit' model

(based on GOF) after completing model calibration.

	Observed prevalence (95% CI)		Prevalence (95% CI) best fit model		Incidence (per 100 PY) best fit model	
Total	18.13 (15.35 – 20.91)		17.65 (16.48 – 18.85)		2.69	
Risk group	**Male**	**Female**	**Male**	**Female**	**Male**	**Female**
Low risk	11.74 (8.30 – 16.35)	20.89 (16.77 – 25.70)	13.76 (12.05 – 15.60)	19.72 (17.86 – 21.65)	1.67	2.55
Medium risk	20.83 (11.73 – 34.26)	33.33 (15.18 – 58.29)	20.87 (15.91 – 26.44)	33.51 (21.52 – 48.27)	2.50	4.57
High risk	14.29 (8.17 – 23.80)	36.11 (22.48 – 52.43)	13.84 (10.67 – 17.74)	36.64 (27.97 – 46.62)	1.50	5.02

255 **Table 2.** Intervention coverage (%) in 2014 for the calibrated 'best fit' model.

	Aware of HIV+ status	On ART of all HIV infected	On ART of those aware of HIV+ status	Circumcised
Overall	60.92%	14.81%	24.31%	-
Female	65.36%	17.29%	26.45%	-
Male	55.19%	11.62%	21.05%	15.29%

258   **Table 3.** Intervention coverage that had the highest impact on incidence after 15 years of scale-up. Coverage over time is shown in Figure S4

259   (see Additional File 1).

	Aware of HIV+ status	On ART of all HIV infected	On ART of those aware of HIV+ status	Circumcised	Oral PrEP
Overall	87.13%	65.28%	74.93%	-	2.65%
Female	89.40%	68.92%	77.09%	-	3.16%
Male	83.87%	60.09%	71.65%	22.58%	2.24%

268 **Table 4.** Highest intervention impact estimates of HIV prevalence (%) without (baseline) and with scale-up of interventions; relative reduction in

269 prevalence, incidence, AIDS-related mortality, and number of new infections over 15 years (obtained by comparing the model with intervention

270 scale-up to the baseline model after 15 years). LR = low risk, MR = medium risk, HR = high risk.

	Baseline	Intervention scale-up	Relative reduction (compared to baseline)			
			Prevalence	Incidence	AIDS-related mortality	New infections over 15 years
Overall	**14.14% (13.17 – 15.13)**	**8.75% (7.98 – 9.55)**	**0.3810**	**0.8668**	**0.5376**	**0.4123**
Female LR	16.78% (15.18 – 18.42)	10.51% (9.26 – 11.89)	0.3739	0.8672	0.5478	0.4092
Female MR	27.90% (16.14 – 40.96)	18.34% (9.87 – 31.41)	0.3425	0.8500	0.5320	0.3848
Female HR	29.85% (21.82 – 39.42)	20.33% (13.26 – 28.28)	0.3192	0.8315	0.5170	0.3651
Male LR	10.88% (9.50 – 12.35)	6.37% (5.33 – 7.55)	0.4144	0.8815	0.5389	0.4299
Male MR	15.67% (11.81 – 20.52)	9.64% (6.59 – 13.63)	0.3848	0.8651	0.5222	0.4104
Male HR	9.53% (7.12 – 12.45)	6.07% (4.15 – 8.48)	0.3636	0.8443	0.5056	0.3983

Figure 1. Overall HIV prevalence (a) and incidence (b) obtained under the baseline model

(full black line) and the 'best fit' model with intervention scale-up (dashed red line) giving the

highest reduction in incidence, 1969-2029. Intervention scale-up started in 2014 (dotted grey

vertical line). The impact of decreasing the yearly rate of ART dropout, assuming no increase

in condom use, and lowering the probability of ART initiation in the chronic stage are also

shown. Parameter estimates for the baseline model are shown in Table S2, while parameter

estimates for the intervention uptake under the highest-impact scenario are shown in Table S3

(see Additional File 1).

Uncertainty analysis

When accounting for parameter uncertainty, the relative reductions in incidence ranged from

30.90% to 86.80% (median 65.68%, IQR 55.48-76.66%). The relative reductions in

prevalence ranged from 10.05% to 38.66% (median 26.24%, IQR 20.17-32.02%). Figure 2

shows that condom use in the low-risk group, ART initiation during the chronic stage, and the

proportion of diagnosed individuals confer the greatest impact on the predicted relative

reduction in HIV incidence, with higher uptake levels leading to higher reductions (Figure 3),

when implemented in combination with the other interventions.

Figure 2. Correlation between each intervention component and the relative reduction in HIV incidence based on the 50 000 combinations used in the uncertainty analysis.

Figure 3. Relative reduction in incidence, prevalence, and AIDS-related mortality for the

three most important intervention components (condom use in the low risk group, ART

initiation during the chronic stage, and the proportion of diagnosed individuals).

Sensitivity analysis

Results of the sensitivity analyses are shown in Table S4 (see Additional File 1) and Figure 1

(selected scenarios). Scenarios varying the assumptions regarding oral PrEP use did not affect

the projected impact of the interventions, probably due to the low coverage levels. Assuming

equal testing and treatment uptake in males and females slightly increased the projected

reduction in incidence from 86.68% to 88.11%. Reducing the annual rate of ART dropout

from 20% to 5% resulted in an increase of the projected reduction in incidence from 86.68%

to 91.16%. Under this scenario, a substantially greater reduction in AIDS-related mortality

was seen, as well as in the number of new infections over 15 years. In contrast, increasing the

annual rate of ART dropout to 70% resulted in a decrease of the projected reduction in

incidence from 86.68% to 77.78%, as well as a substantial decrease of the projected reduction

in AIDS-related mortality and number of new infections over 15 years. Assuming no increase

in condom use after 2014 had a large impact, with the relative reduction in incidence dropping

to 56.19%. To assess the impact of early ART initiation, we lowered the probability of

initiating ART in the chronic stage to 30%. This resulted in 81.54% reduction in incidence

and 41.54% reduction in AIDS-related mortality (Table S4, see Additional File 1). In this

scenario, only 55.94% of those aware of their HIV status were on ART, compared to 74.93%

on ART in the highest impact scenario that assumed a 72.51% probability of initiating ART in

the chronic stage.

Discussion

Results of our modelling study have indicated that we can achieve as high as 87% relative

reduction in HIV incidence through combination prevention. We have projected that this

reduction in incidence can be achieved by 2029 using the following combination of

intervention coverage levels: (i) condom use in about 60% of sexual acts, (ii) 23% of

susceptible men being circumcised, (iii) 87% of people living with HIV aware of their status,

(iv) 75% of those aware of their status on ART, and (v) about 3% of susceptible individuals

on PrEP. In our best case scenario, reaching the above coverage levels, we projected that HIV

incidence may reduce to 2.5 per 1000 PY, and overall prevalence to 8.75%. These numbers

are still higher than for the general population in Uganda, and incidence is still above the

threshold of 1 per 1000 PY that can be seen as HIV elimination (30). It should be noted that

other combinations of coverage levels of the intervention components may lead to similar

reductions in incidence. The decline in prevalence and incidence observed in our baseline

model is in line with the globally declining number of new HIV infections since 1996 (1).

Our projected coverage levels are modest in comparison to the Uganda national targets and

hence attainable. For instance the national target for ART coverage by 2020 was 80% (31).

Moreover, the Uganda AIDS Commission laid down strategies to improve HIV testing and

linkage to care in FCs by partnering with the Ministry of Agriculture and Fisheries (31). The

Uganda government in her HIV and AIDS Strategic Plan 2015/2016-2019/2020 emphasized

the need to scale up combination prevention strategies to critical levels by 2020 (2). Our

projected levels are well within these 2020 national targets. The prevention components we

considered in our study are key in objective two of this National Strategic Plan.

Our projections indicate that condom use and ART will have the largest impact on HIV

transmission in these FCs. Without an increase in condom use, the projected reduction in

incidence was only 56.19% compared to our baseline scenario. Sensitivity analysis also

showed that initiating ART early plays an important role in achieving substantial reductions

in incidence and AIDS-related mortality. Lowering the probability of ART initiation during

the chronic stage decreased the projected impact to an incidence reduction of 81.54%. A

previous modelling study in a hyperendemic setting with relatively low levels of condom use

found that increasing early ART coverage rather than providing PrEP to uninfected

individuals would be more cost-effective, but that early ART alone is not sufficient for

reducing HIV incidence to very low levels (20). In our projections, early ART did have a

substantial impact on reductions in HIV incidence, although combined with relatively high

levels of condom use. Uncertainty analyses using 50 000 possible scenarios also showed that

condom use in the low risk group was the most important component of the prevention

package. As it is known that achieving high levels of condom use in these FCs is difficult,

public health initiatives should focus on adequate ART delivery in this hard-to-reach

population and ensure sufficient adherence, combined with increased efforts regarding

condom promotion and distribution.

A systematic review in 2020 found that fisher folk have a preference for PrEP and VMMC

over other prevention services such as condom use, and that there is an urgent need for

innovative approaches to HIV care delivery in FCs (19). In our projections, oral PrEP use did

not seem to play a very important role in reducing HIV incidence, but coverage in the best

case scenario was very low (around 3%). However, increasing oral PrEP uptake is expected to

further reduce HIV incidence. Oral PrEP uptake and adherence to a daily regimen have not

been optimal in high-risk populations (32). This may be mitigated by the use of long-acting

injectables that do not require a daily regimen (33), although a previous study has found that

fisher folk would prefer oral PrEP (34). In addition, several barriers to PrEP initiation such as

travel distance to health facilities and community stigma should be overcome first (35).

Furthermore, efforts should be made to avoid risk-compensating behaviour in susceptible

individuals on PrEP, which could cancel out its effect.

An advantage of the compartmental model used in this study is that it accounts for a large part

of the heterogeneity in HIV risk behaviour through the inclusion of different sexual risk

groups that can form different types of partnerships, each with its own associated risk. An

alternative approach would be to use an individual-based model (IBM) which can incorporate

a higher degree of heterogeneity at the individual level, such as more detailed information on

sexual mixing behaviour (35). When model structure becomes too complex to capture in a

compartmental modelling framework, an IBM may be the solution. In addition, when using an

IBM, intervention policies can be tailored to the individual, but this was beyond the scope of

the present study and would require more detailed data.

This study has several limitations. First, the model was calibrated to observed HIV prevalence

at only one time point, since these were the only data available. Although the 'best fit' model

represents only one possible epidemic trajectory, we performed an uncertainty analysis using

a range of model parameters. Second, the model does not explicitly account for other STIs

that are known to be co-factors in HIV infection. These data were not available from the

HIVCOMB study, and it was not the focus in this work to investigate the contribution of other

STIs. However, since the model was calibrated to the observed prevalence data, we have

assumed that it does implicitly account for the role of other STIs in HIV transmission. Third,

the model does not account for behavioural change; we assumed that people stay in the same

sexual risk group over time. Fourth, the model does not incorporate stochasticity, which could

provide model flexibility to accommodate changes in the transmission rates that might occur

due to unobserved stochastic processes (36). We also did not account for the fact that some

interventions may induce risk compensating behaviour such as reduced condom use and more

sexual risk-taking. In addition, we have not addressed the complex relationship between

alcohol use, intimate partner violence (IPV), and HIV infection. IPV may increase a woman's

risk of HIV infection, for example due to forced unprotected sexual intercourse or because of

their lack of power to negotiate condom use (37). Lastly, our model was calibrated to

prevalence data due to the lack of incidence data. However, although prevalence is influenced

by new infections as well as deaths, we also assessed the impact of the combined

interventions on incidence.

Conclusions

Despite the highlighted limitations, we have projected that a substantial reduction in HIV

incidence can be achieved through combination prevention without necessarily reaching high

coverage of all the specific intervention package components. Although Ugandan FCs may

have extensive sexual networks bridging into the general population, Ratmann et al. (5) have

recently reported that only 1.3% of HIV transmissions occurred from lakeside to inland areas.

However, HIV transmission from inland to lakeside areas was slightly higher (3.7%), hence

future work could incorporate partnerships formed outside of the FCs, to investigate the

interplay between reducing HIV prevalence in either inland or lakeside areas (4). It should be

noted that the projections obtained in the present study are specific to the included Ugandan

FCs and may not necessarily be generalizable to other settings with a different underlying

HIV incidence or epidemic stage, as for instance the general population in Uganda.

List of abbreviations

AIDS   Acquired Immunodeficiency Syndrome

ART   Antiretroviral therapy

FC   Fishing community

GOF   Goodness of fit

HCT   HIV counselling and testing

HIV   Human Immunodeficiency Virus

IPV   Intimate partner violence

LHS   Latin hypercube sampling

MARP   Most at risk population

PrEP   Pre-exposure prophylaxis

STI   Sexually transmitted infection

VMMC   Voluntary medical male circumcision

Declarations

Ethics approval and consent to participate

The HIVCOMB study was approved by the Uganda Virus Research Institute, the Uganda
National Council of Science and Technology, and the London School of Hygiene and
Tropical Medicine ethics committees.

Consent for publication

429   Not applicable.

Availability of data and materials

431   All data generated or analysed during this study are included in this published article and its

432   supplementary information files. Computer code accompanying this manuscript is available at

433   https://github.com/cecilekremer/hivug.

Competing interests

435   The authors declare that they have no competing interests.

Funding

RNN was funded to visit Hasselt by Hasselt University Special Research Fund (BOF)

Programme for incoming mobility. The HIVCOMB Study was jointly funded by the UK

Medical Research Council (MRC) and the UK Department for International Development

(DFID) under the MRC/DFID Concordat agreement. Award Number MR/L004305/1. The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Authors' contributions

444   RNN, NH, and CK contributed to conceptualization of the study. RNN, AK, MK, and JS were

445   involved in the HIVCOMB study. CK and RNN contributed to the analysis code. CK

446   performed the analysis. NH, RNN, CK, AK, MK, and JS contributed to the interpretation of

447   results. CK and RNN drafted the manuscript. All co-authors critically reviewed and revised

448   the manuscript.

Acknowledgements

Project carried out with the support of the Fund AIDS managed by the King Baudouin Foundation. The computational resources and services used in this work were provided by the VSC (Flemish Supercomputer Center), funded by the Research Foundation Flanders (FWO) and the Flemish Government department EWI. The authors would like to thank Helen A. Weiss (LSHTM) for her insightful comments on an earlier draft of this manuscript.

References

(1) Global HIV & AIDS statistics – Fact sheet. 2022; Geneva: UNAIDS.

(2) National HIV and AIDS strategic plan 2015/2016-2019/2020. 2015; Kampala, Uganda: Uganda AIDS Commission.

(3) Seeley J, Nakiyingi-Miiro J, Kamali A, Mpendo J, Asiki G, Abaasa A, et al. High HIV incidence and socio-behavioral risk patterns in fishing communities on the shores of Lake Victoria, Uganda. Sex Transm Dis 2012;39(6):433-9. doi: 10.1097/OLQ.0b013e318251555d.

(4) Bbosa N, Ssemwanga D, Nsubuga RN, Salazar-Gonzalez JF, Salazar MG, Nanyonjo M, et al. Phylogeography of HIV-1 suggests that Ugandan fishing communities are a sink for, not a source of, virus from general populations. Sci Rep 2019;9(1):1051. doi: 10.1038/s41598-018-37458-x.

(5) Ratmann O, Kagaayi J, Hall M, Golubchick T, Kigozi G, Xi X, et al. Quantifying HIV transmission flow between high-prevalence hotspots and surrounding communities: a population-based study in Rakai, Uganda. Lancet HIV 2020;7:e173-83. doi: 10.1016/S2352-3018(19)30378-9.

(6) Asiki G, Mpendo J, Abaasa A, Agaba C, Nanvubya A, Nielsen L, et al. HIV and syphilis prevalence and associated risk factors among fishing communities of Lake Victoria, Uganda. Sex Transm Infect 2011;87:511-5. doi: 10.1136/sti.2010.046805.

(7) Kamali A, Nsubuga RN, Ruzagira E, Bahemuka U, Asiki G, Price MA, et al.

Heterogeneity of HIV incidence: a comparative analysis between fishing communities and in

a neighbouring rural general population, Uganda, and implications for HIV control. Sex

Transm Infect 2016;92:447–54. doi: 10.1136/sextrans-2015-052179.

(8) Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, et al. Male

circumcision for HIV prevention in men in Rakai, Uganda: a randomized trial. Lancet

2007;369:657-66. doi: 10.1016/S0140-6736(07)60313-4.

(9) Matovu JKB, Gray RH, Kiwanuka N, Kigozi G, Wabwire-Mangen F, Nalugoda F, et al.

Repeat voluntary HIV counseling and testing (VCT), sexual risk behavior and HIV incidence

in Rakai, Uganda. AIDS Behav 2007;11:71-8. doi: 10.1007/s10461-006-9170-y.

(10) Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, Wangisi J, et al. Antiretroviral

prophylaxis for HIV prevention in heterosexual men and women. N Engl J Med

2012;367:399-410. doi: 10.1056/NEJMoa1108524.

(11) Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al.

Antiretroviral therapy for the prevention of HIV-1 transmission. N Engl J Med 2016;375:830-

9. doi: 10.1056/NEJMoa1600693.

(12) UNAIDS. Fact sheet - latest global and regional statistics on the status of the AIDS

epidemic. Geneva: UNAIDS 2018.

(13) Chang LW, Grabowski MK, Ssekubugu R, Nalugoda F, Kigozi G, Nantume B, et al.

Heterogeneity of the HIV epidemic in agrarian, trading, and fishing communities in Rakai,

Uganda: an observational epidemiological study. Lancet HIV 2016;3(8):e388-96.

(14) Mafigiri R, Matovu JKB, Makumbi FE, Ndyanabo A, Nabukalu D, Sakor M, et al. HIV

prevalence and uptake of HIV/AIDS services among youths (15-24 years) in fishing and

neighboring communities of Kasensero, Rakai District, South Western Uganda. BMC Public

Health 2017;17(1):251. doi. 10.1186/s12889-017-4166-2

(15) Musumari PM, Techasrivichien T, Srithanaviboonchai K, Wanyenze RK, Matovu JK,

Poudyal H, et al. HIV epidemic in fishing communities in Uganda: a scoping review. PloS

One 2021;16(4):e0249465. doi: 10.1371/journal.pone.0249465.

(16) Dodd PJ, White PJ, Garnett GP. Notions of synergy for combinations of interventions

against infectious diseases in heterogeneously mixing populations. Math Biosci 2010;227:94-

104. doi: 10.1016/j.mbs.2010.06.004.

(17) Cox AP, Foss AM, Shafer LA, Nsubuga RN, Vickerman P, Hayes RJ, et al. Attaining

realistic and substantial reductions in HIV incidence: model projections of combining

microbicide and male circumcision interventions in rural Uganda. Sex Transm Infect

2011;87:635-9. doi: 10.1136/sti.2010.046227.

(18) McRobie E, Matovu F, Nanyiti A, Nonvignon J, Abankwah DNY, Case KK, et al.

National responses to global health targets: exploring policy transfer in the context of the

UNAIDS "90-90-90" treatment targets in Ghana and Uganda. Health Policy Plan 2018;33:17-

33.

(19) Toms K, Potter H, Balaba M, Parkes-Ratanshi R. Efficacy of HIV interventions in

African fishing communities: a systematic review and qualitative synthesis. Int J Infect Dis

2020;101:326-33. doi: 10.1016/j.ijid.2020.09.1476.

(20) Cremin I, Alsallaq R, Dybul M, Piot P, Garnett G, Hallett TB. The new role of

antiretrovirals in combination HIV prevention: a mathematical modelling analysis. AIDS

2013;27:447-58.

(21) Li J, Peng L, Gilmour S, Gu J, Ruan Y, Zou H, et al. A mathematical model of

biomedical interventions for HIV prevention among men who have sex with men in China.

BMC Inf Dis 2018;18:600. doi: 10.1186/s12879-018-3516-8.

(22) Hollingsworth TD, Anderson RM, Fraser C. HIV-1 transmission, by stage of infection.

JID 2008;198:687-93. doi: 10.1086/59050.

(23) Kuteesa MO, Weiss HA, Abaasa A, Nash S, Nsubuga RN, Newton R, et al. Feasibility of

conducting HIV combination prevention interventions in fishing communities in Uganda: a

pilot cluster randomized trial. PLoS One 2019;14:e0210719. doi:

10.1371/journal.pone.0210719.

(24) McKay MD, Beckman RJ, Conover WJ. A comparison of three methods for selecting

values of input variables in the analysis of output from a computer code. Technometrics

1979;21:239-245. doi: 10.2307/1268522.

(25) Green IRA, Stephenson D. Criteria for comparison of single event models. Hydrolog Sci

J 1986;31(3):395-411. doi: 10.1080/02626668609491056.

(26) Willem L, Stijven S, Vladislavleva E, Broeckhove J, Beutels P, Hens N. Active learning

to understand infectious disease models and improve policy making. PLoS Comput Biol

2014;10(4):e1003563. doi: 10.1371/journal.pcbi.1003563.

(27) Alexander TS. Human immunodeficiency virus diagnostic testing: 30 years of evolution.

Clin Vaccine Immunol 2016:23:249-53.

(28) Rosenberg NE, Hauser BM, Ryan J, Miller WC. The effect of HIV counseling and

testing on HIV acquisition in Sub-Saharan Africa: a systematic review. Sex Transm Infect

2016;92:579-86. doi: 10.1136/sextrans-2016-052651.

(29) Ministry of Health Uganda. Uganda national antiretroviral treatment and care guidelines

for adults, adolescents, and children. Released 2003.

(30) Granich RM, Gilks CF, Dye C, De Cock KM, Williams BG. Universal voluntary HIV

testing with immediate antiretroviral therapy as a strategy for elimination of HIV

transmission: a mathematical model. Lancet 2009;373:48-57. doi: 10.1016/S0140-

6736(08)61697-9.

(31) Acceleration of HIV prevention in Uganda, a road map towards zero new infections by

2030. 2018; Kampala, Uganda: Uganda AIDS Commission.

(32) Mugwanya KK, Pintye J, Kinuthia J, Abuna F, Lagat H, Begnel ER, et al. Integrating

preexposure prophylaxis delivery in routine family planning clinics: a feasibility

programmatic evaluation in Kenya. PLoS Med 2019; 1002885. doi:

10.1371/journal.pmed.1002885.

(33) Trezza C, Ford SL, Spreen W, Pan R, Piscitelli S. Formulation and pharmacology of

long-acting cabotegravir. Curr Opin HIV AIDS 2015;10:239-45. doi:

10.1097/COH.0000000000000168.

(34) Kuteesa MO, Quaife M, Biraro S, Katumba KR, Seeley J, Kamali A, et al. Acceptability

and predictors of uptake of anti-retroviral pre-exposure prophylaxis (PrEP) among fishing

communities in Uganda: a cross-sectional discrete choice experiment survey. AIDS Behav

2019;23(10):2674-86. doi: 10.1007/s10461-019-02418-7.

(35) Willem L, Verelst F, Bilcke J, Hens N, Beutels P. Lessons from a decade of individual-

based models for infectious disease transmission: a systematic review (2006-2015). BMC

Infect Dis 2018;17:612. doi: 10.1186/s12879-017-2699-8.

(36) Garnett G, Cousens S, Hallet TB, Steketee R, Walker N. Mathematical models in the

evaluation of health programmes. Lancet 2011;378:515-25. doi: 10.1016/S0140-

6736(10)61505-X.

(37) Miller AP, Ddaaki WG, Bloom BE, Wirtz AL, Nakyanjo N, Kigozi G, et al. Perspectives

of women living with HIV on addressing violence and use of alcohol during HIV services:

qualitative findings from fishing communities in Uganda. Violence against women

2021;28(6-7):1483-1504.

569 Additional material

Additional File 1. Detailed model description.

Figure S1. Model structure for males in risk group j (which is the same for females,

without circumcision compartments).

Figure S2. Parameters that are highly informed by the data. Dashed lines represent the

density f_U of the uniform distribution on the uncertainty range of the input parameter. Solid

lines represent the density $f_{\Theta 1\%}$ corresponding to the values within the subspace $\Theta_{1\%}$. A

peaked unimodal density $f_{\Theta 1\%}$ indicates that the parameter was highly informed by the data.

Figure S3. HIV transmission dynamics. HIV prevalence (a) and incidence (b) in the

different sexual risk groups obtained from the 'best fit' model. Filled symbols on the dotted

grey vertical line represent the observed prevalence in 2014 obtained from the HIVCOMB

study, to which the model was calibrated.

Figure S4. Intervention coverage over time. (a) HIV infected individuals on ART, (b) HIV

infected individuals aware of their status on ART, (c) circumcised susceptible males, and (d)

individuals aware of their HIV+ status. Dashed lines indicate the year 2011 when ART was

implemented in the chronic stage, dotted lines indicate the year 2020 when intervention scale-

up was started, dash-dotted lines indicate 1987 when HIV testing was introduced.

Table S1. Probability of transmission per coital act. Baseline corresponds to the chronic

low viral load stage.

Table S2. Uncertainty ranges and fixed values for the baseline model parameters and

value in the 'best fit' solution.

Table S3. Uptake ranges for the intervention parameters and their values in the highest

impact scenario.

Table S4. Sensitivity analyses for the impact of combined interventions.