Low testing rates limit the ability of genomic surveillance programs to

monitor SARS-CoV-2 variants: a mathematical modelling study

Alvin X. Han, Ph.D.[1,*], Amy Toporowski, M.D.[2], Jilian A. Sacks, Ph.D.[3], Mark Perkins,

M.D.[3], Sylvie Briand, M.D.[3], Maria van Kerkhove, Ph.D.[3], Emma Hannay, Ph.D.[2], Sergio

Carmona, M.D.[2], Bill Rodriguez, M.D.[2], Edyth Parker, Ph.D.[4], Brooke E. Nichols,

Ph.D.[1,2,5,†], Colin A. Russell, Ph.D.[1,5,†,*]

[1]Department of Medical Microbiology & Infection Prevention, Amsterdam University

Medical Center, University of Amsterdam, Amsterdam, The Netherlands

[2]Foundation for Innovative New Diagnostics (FIND), Geneva, Switzerland

[3]Department of Epidemic and Pandemic Preparedness and Prevention, Emergency

Preparedness Programme, World Health Organization, Geneva, Switzerland

[4]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA,

USA

[5]Department of Global Health, School of Public Health, Boston University, Boston, MA,

USA

†Contributed equally

*Correspondence to Alvin X. Han (x.han@amsterdamumc.nl) and Colin A. Russell

(c.a.russell@amsterdamumc.nl)

Summary (293/300 words)

Background

Genomic surveillance is essential for monitoring the emergence and spread of SARS-CoV-2

variants. SARS-CoV-2 diagnostic testing is the starting point for SARS-CoV-2 genomic

sequencing. However, testing rates in many low- and middle-income countries (LMICs) are

low (mean = 27 tests/100,000 people/day) and global testing rates are falling in the post-crisis

phase of the pandemic, leading to spatiotemporal biases in sample collection. Various public

health agencies and academic groups have produced recommendations on sample sizes and

sequencing strategies for effective genomic surveillance. However, these recommendations

assume very high volumes of diagnostic testing that are currently well beyond reach in most

LMICs.

Methods

To investigate how testing rates, sequencing strategies and the degree of spatiotemporal bias

in sample collection impact variant detection and monitoring outcomes, we used an

individual-based model to simulate COVID-19 epidemics in a prototypical LMIC. Within the

model, we simulated a range of testing rates, accounted for likely testing demand and applied

various genomic surveillance strategies, including sentinel surveillance.

Findings

Diagnostic testing rates play a substantially larger role in monitoring the prevalence and

emergence of new variants than the proportion of samples sequenced. To enable timely

detection and monitoring of emerging variants, programs should achieve average testing rates

of at least 100 tests/100,000 people/day and sequence 5-10% of test-positive specimens,

which may be accomplished through sentinel or other routine surveillance systems. Under

realistic assumptions, this averages to ~10 samples for sequencing/1,000,000 people/week.

Interpretation

For countries where testing capacities are low and sample collection is spatiotemporally

biased, surveillance programs should prioritize investments in wider access to diagnostic

testing to enable more representative sampling, ahead of simply increasing quantities of

sequenced samples.

Funding

European Research Council, the Rockefeller Foundation, and the Governments of Germany,

Canada, UK, Australia, Norway, Saudi Arabia, Kuwait, Netherlands and Portugal.

Research in context

Evidence before this study

Genomic sequencing has been an integral part of the COVID-19 pandemic response, critical

to monitoring the evolution of SARS-CoV-2 and identifying novel variants of interest and

variants of concern (VOCs). As of March 2022, more than 10 million unique sequences had

been submitted to GISAID. However, SARS-CoV-2 sequences have been disproportionately

submitted from high-income countries (HICs), with large surveillance gaps existing in most

LMICs. To strengthen genomic surveillance of SARS-CoV-2, previous studies focused on

estimating a minimal number of positive SARS-CoV-2 tests to reflex for sequencing for

effective variant detection and monitoring. We searched PubMed and Google Scholar using

combinations of search terms (i.e., "SARS-CoV-2", "COVID-19", "diagnostic", "genomic

surveillance", "sequencing", "LIC", "LMIC") and critically considered published articles and

preprints that studied or reviewed SARS-CoV-2 testing and genomic surveillance, especially

in the LMIC context. We also reviewed SARS-CoV-2 sequencing recommendations

published by the World Health Organization (WHO) and European Centre for Disease

Prevention and Control (ECDC). We reviewed all studies and the latest recommendations

published in English up to February 2022. We found that prevailing recommendations for

estimating sequencing sample size to identify or monitor the prevalence of new variants

assume that COVID-19 testing is performed at high rates per capita and in high absolute

numbers, such that the sequenced samples are largely representative of the circulating SARS-

CoV-2 viral diversity. This is, however, not the case in many countries, particularly in many

LMICs, and can vary dramatically depending on the epidemiological situation.

Added value of this study

To our knowledge, this is the first study that quantitatively estimates the joint impact of

COVID-19 testing rates and sequencing strategies on SARS-CoV-2 variant detection and

monitoring. We developed an individual-based COVID-19 transmission model that was

specifically designed to simulate VOC emergence in LMICs under a wide range of test

availability and sampling strategies for sequencing. We showed that given the current

average COVID-19 testing rate of 27 tests per 100,000 people per day across LMICs, the

sequencing sample size recommendations for early variant detection from WHO/ECDC and

other academic groups would likely result in delayed detection of a new VOC until it had

spread through a substantial portion of the population. We quantitatively demonstrated that

increasing COVID-19 testing rates to at least 100 tests per 100,000 people per day, including

through sentinel surveillance sites, and sampling as broadly as possible, yields far earlier

VOC detection and greater accuracy of variant prevalence estimates than simply increasing

the proportion of samples to be sequenced.

Implications of the available evidence

Spatiotemporal representativeness of SARS-CoV-2 positive samples being sequenced, which

can be accomplished by increasing diagnostic testing rates, and widening the geographic

coverage from where samples are collected, as well as shortening sequencing turnaround time

are the key features of an effective genomic surveillance program aimed at detection and

monitoring of novel SARS-CoV-2 variants. Only once these areas have been strengthened

does increasing the volume of sequenced samples have significant impact.

Main Text (~3,500/3,500 words)

Introduction

Since the start of the COVID-19 pandemic in 2019, unprecedented expansion of genomic surveillance efforts has led to the generation of more than 10 million SARS-CoV-2 sequences deposited in the publicly accessible GISAID database (https://www.gisaid.org/) as of May 2022. These efforts have been integral to understanding the COVID-19 pandemic,[1] including the identification of the Alpha variant in the United Kingdom during the fall 2020,[2] the Delta variant in India in late 2020,[3] and the Omicron variant in Southern Africa in November 2021.[4] Despite the value of these efforts for monitoring the evolution of SARS-CoV-2, the intensity of genomic surveillance is highly heterogenous across countries. High-income countries (HICs) on average produced 16 times more SARS-CoV-2 sequences per reported case than low- and middle-income countries (LMICs) as a result of longstanding socioeconomic inequalities and consequent underfunding of laboratory and surveillance infrastructures.[5] To strengthen global pandemic preparedness, initiatives such as the Access to COVID-19 Tools Accelerator Global Risk Monitoring Framework, the Pan American Health Organization COIVD-19 Genomic Surveillance Regional Network and Africa Pathogen Genomics Initiative, among others, have supported LMICs in developing pathogen genomic surveillance programs.

As resources are finite, it is critical that sequencing sample sizes, and the diagnostic testing needed to obtain samples for sequencing, are carefully set for genomic surveillance programs to detect and monitor variants as efficiently as possible. Current recommended sample sizes are based on sampling theory[5–8] and assume that the volume of diagnostic testing is large enough such that the diversity of sampled viruses is representative of the diversity of viruses circulating in the population. However, LMICs test at a mean rate of 27 tests per 100,000 persons per day (tests/100k/day) as opposed to >800 tests/100k/day across HICs between January 2020 and March 2022 (https://www.finddx.org/covid-19/test-tracker/), with even higher testing rates in some HICs (Figure 1). Low testing rates lead to spotty information and smaller virus specimen pools available for sequencing, resulting in strong sampling biases. These factors can render efforts to monitor the emergence of new variants or prevalence of existing variants highly unreliable.

Figure 1: **Global disparities in SARS-CoV-2 testing rates**. Each country is colored by the average
total number of SARS-CoV-2 tests performed per 100,000 persons per day (/100K/day) (**A**) between
1 December 2021 and 31 March 2022 when the Omicron variant-of-concern spread around the world;
(**B**) between 1 April 2022 and 6 May 2022 when most countries were past peak Omicron wave of
infections (https://www.finddx.org/covid-19/test-tracker/).

Here, we studied how different testing rates can impact genomic surveillance outcomes.

Specifically, we used an individual-based modelling framework to simulate concurrently-

circulating wild-type SARS-CoV-2/Alpha- as well as Delta-/Omicron-like epidemics in

Zambia as a representative LMIC archetype. We then applied different genomic surveillance

sampling strategies (i.e., sources of sample collection and varying proportion of specimens to

sequence) to elucidate how testing, sequencing volumes and the degree of sampling bias

arising from sources of specimens jointly impact the timeliness of variant detection and the

accuracy of variant monitoring.

Methods

Simulating SARS-CoV-2 epidemics with the Propelling Action for Testing And Treating (PATAT) model

We used PATAT, a stochastic individual-based model to simulate SARS-CoV-2 epidemics in a community with demographic profiles, contact mixing patterns, and level of public health resources mirroring those typically observed in LMICs. Here, the model was based on Zambia. Briefly, PATAT creates an age-structured population, linking individuals within contact networks of multi-generational households, schools, workplaces, and churches (i.e., regular mass gatherings) (Table S1). Healthcare facilities (i.e., community clinics and tertiary hospitals) where individuals with mild symptoms seek symptomatic testing and have their virus specimens collected are simulated to approximate localized community structures based on an empirical clinic-to-population ratio. Households are proximally distributed around these facilities based on the given empirical distance distribution that correlates with probabilities of symptomatic individuals seeking testing at clinics (Table S1).

We then simulated SARS-CoV-2 infection waves in a population of 1,000,000 individuals over a 90-day period that begins with an initial 1% prevalence of an extant SARS-CoV-2 variant and the introduction of a mutant variant at 0·01%. We assumed that clinic-based professional-use Antigen Rapid Diagnostic Tests (Ag-RDTs) form the basis of testing given persistent reports that polymerase chain reaction (PCR) tests are poorly accessible for detection of symptomatic cases in most LMICs.[9] As Ag-RDT sensitivity depends on within-host viral loads,[10] PATAT generates viral load trajectories, measured in cycle threshold (Ct) values, for infected individuals by randomly sampling from known viral load distributions of different SARS-CoV-2 variants.[11,12] We performed simulations for two variant replacement scenarios – Alpha variant introduction while the wild-type virus was circulating (wild-type/Alpha) and Omicron (BA.1) variant introduction while Delta was circulating (Delta/Omicron), applying known distributions of their peak viral load, incubation, and virus clearance periods[11,13] (Table S1). Before simulating the two-variant epidemic, we first calibrated the transmission probability parameter for the extant variant such that it would spread in a completely susceptible population at $R_0 = 2·5$-$3·0$. We then assumed Alpha and Omicron (BA.1) were more transmissible that the respective extant virus to achieve growth rates of ~0·15/day and ~0·35/day respectively.[2,14]

For both sets of simulations, we assumed that 10% of the population had infection-acquired

immunity against the extant strain initially with some level of protection against infection by

the mutant virus (wild-type SARS-CoV-2: 80% protection against Alpha;[15] Delta: 20%

protection against Omicron[14]). We also investigated the scenario where 40% of the

population had infection-acquired immunity as part of sensitivity analyses (see below). We

did not investigate scenarios involving vaccine-acquired immunity due to low vaccine uptake

in most LMICs.[16]

PATAT uses the SEIRD (Susceptible-Exposed-Infected-Recovered/Death) epidemic model

for disease progression and stratifies infected individuals based on their symptom

presentation (asymptomatic, mild, or severe). After an assumed random delay post-symptom

onset (mean = 1 day; s.d. = 0.5 day), symptomatic individuals who seek testing would do so

at their nearest healthcare facility, where test-positive samples may be reflexively collected

for sequencing. We assumed that symptomatic individuals sought testing based on a

probability distribution of health services-seeking behaviour that inversely correlates with the

distance between the individual's household and the nearest healthcare facility (Table S1)[17].

We varied levels of Ag-RDT stocks per day (i.e., 27, 100, and 200-1,000 (in increments of

200) tests/100k/day), running 10 bootstrap simulations for each testing rate. Given the start of

a week on Monday, we assumed that a week's worth of tests are delivered to healthcare

facilities every Monday and unused Ag-RDTs in the previous week are carried forward into

the next week. Due to overlapping symptoms between COVID-19 and other respiratory

diseases, a proportion of available Ag-RDTs would be used by individuals who are not

infected with SARS-CoV-2. Based on test positivity rates reported by various countries in the

second half of 2021,[18] we assumed 10% test positivity rate at the start and end of the

simulated epidemic, and 20% test positivity at its peak, linearly interpolating the rates

between these timepoints. We also assumed that false positive specimens could be sampled

based on reported Ag-RDT specificity of 98·9%.[10]

We assumed that any specimens collected for genomic surveillance after positive detection

through Ag-RDT would be reflexively confirmed with PCR. We also assumed that all

symptomatic individuals who have severe symptoms require hospitalization, and are tested

separately from mild symptomatic persons who sought testing. Given that likely only ~10-

20% of COVID-19 deaths in Zambia were tested for the disease in life,[19,20] we assumed that

only 20% of individuals with severe disease would be tested by Ag-RDT or PCR upon

presenting severe symptoms and have specimens collected for sequencing.

The list of parameters and full technical details on PATAT can be found in the

Supplementary Appendix. The model source code is available at

https://github.com/AMC-LAEB/PATAT-sim.

Genomic surveillance strategies

Twenty percent of healthcare facilities were assumed to be tertiary facilities based on

empirical data collected from Zambia.[21,22] We assumed that tertiary facilities provide testing

for mild symptomatic individuals as well as hospitalized patients with severe symptoms. We

also assumed that a proportion of tertiary facilities serve as sentinel surveillance sites that

reflexively collect SARS-CoV-2 positive samples for sequencing. We then simulated six

strategies with varying degrees of sampling coverage where positive specimens collected

from testing sites would be consolidated and sampled for sequencing: (1) all samples from

community clinics and tertiary hospitals are sent to a centralized facility and further sampled

for sequencing (i.e. *population-wide* strategy); (2) only *one* tertiary sentinel facility for the

population of 1,000,000 simulated people would sequence a portion of positive specimens it

has collected, both from mild individuals seeking symptomatic testing and severe patients

who sought tertiary care at the facility; or only (3) 10%, (4) 25%, (5) 50%, and (6) 100% of

all tertiary sentinel facilities would sample and sequence a proportion of the specimens they

have collected.

For all strategies, we assumed that a proportion (1%-100%; in 2% increments between 1%

and 5%, in 5% increments between 5% and 100%) of positive specimens are collected daily

for sequencing. We also assumed that positive specimens sampled within each week for

sequencing are consolidated into a batch before they are referred for sequencing. Turnaround

time refers to the time between collection of each weekly consolidated batch of positive

specimens to the acquisition of its corresponding sequencing data. Since the within-host viral

loads of infected individuals were simulated, we assumed that only high-quality samples

where Ct values < 30 could be sequenced and that sequencing success rate is 80% as assumed

in other studies.[6]

For each strategy and sequencing proportion, we performed 100 bootstrap simulations for

each epidemic simulation with a given test stock availability, thus totaling to 1,000 random

simulations for each set of variables (i.e., testing rate, sequencing proportion, and strategy).

Results

Performance of current guidance

We first applied current guidance from different stakeholder and academic groups on the

number of positive specimens to sequence to detect SARS-CoV-2 variants at low prevalence

(Table 1) for simulated wild-type/Alpha and Delta/Omicron epidemics in Zambia with a

mean testing rate of 27 tests/100k/day (Figure 2). Even when assuming negligible turnaround

time (i.e. time from specimen collection to acquisition of sequencing data), the current

recommended approaches were insufficient to detect the variant on their respective target

detection day when testing rates were low, regardless of the genomic surveillance sampling

strategy. The first strategy of sampling specimens collected from the whole population that

were sent to one sequencing facility (i.e. population-wide strategy) led to the best

performance (closest to target detection day) for all recommendations, as it involves random

uniform sampling of all available samples, a fundamental assumption made by all current

guidance. However, if the specimen pools available for sequencing are restricted to those

collected from a subset of sentinel tertiary facilities only, the non-uniformity in sampling

coverage results in spatiotemporal bias within the sequenced samples, and leads to delayed

detection of VOCs, which gets progressively worse as the proportion of tertiary facilities

performing sequencing decreases to one facility.

Table 1: **Current guidance by various stakeholder and academic groups on the number of**
specimens to sequence for detection of novel variants at low prevalence.

	Recommendation on number/proportion of positive cases to sequence		Critical considerations
World Health Organization (WHO)/European Centre for Disease Prevention and Control (ECDC)[7,8]	No. of positive cases	No. of sequences to detect at 1% with 95% confidence	• Agnostic to variant properties • Assumes specimen pool to be sampled for sequencing is representative of circulating diversity but acknowledges that unless testing coverage is evenly distributed this will be a biased sample • Notes that in countries with limited sequencing capacity monitoring relative prevalence of variants should be prioritized
	<1000 cases	141	
	1001 – 2,500	196	
	2,500 – 5,000	243	
	5,001 – 10,000	270	
	>10,000	285	
Brito et al.[5]	At least 0·5% of all cases with a turnaround time of 21 days to detect novel lineage before it reaches 100 cases at 20% probability		• Based on sequencing data from Denmark which is testing at >2,000 tests per 100,000 persons per day (https://www.finddx.org/covid-19/test-tracker/)
Wohl et al.[6]	1-29 sequences per day to detect an Alpha-like variant based on 0·03% initial introduction for a population of 10,000 (assuming growth rate of 0·1/day) at 1% with 95% confidence. We used the spreadsheet (https://github.com/HopkinsIDD/VOCsamplesize) provided and input appropriate parameters to obtain the recommendation relevant to the simulated epidemics.		• Assumes that the *observed* variant proportion in the positive specimens collected is representative of the *circulating* variant proportions among the infected population • Assumes asymptomatic patients are tested as well which may not be applicable for many LMICs where self-testing and asymptomatic community testing programs are currently rare. Widespread asymptomatic testing has also been substantially reduced in most HICs in the post-crisis phase of the pandemic. • Requires a large number of specimens that are randomly collected for assumption to hold true at low circulating variant proportions.

Figure 2: **Performance of current guidance on number of positive specimens to sequence for**
variant detection with testing rate at 27 tests per 100,000 persons per day. First day of detection
since variant introduction at 95% confidence and the corresponding circulating variant proportion
using guidance from the World Health Organization (WHO)/European Centre for Disease Prevention
and Control (ECDC)[7,8], Brito et al.[5], and Wohl et al.[6] (Table 1) under different genomic surveillance
strategies with varying sampling coverage (i.e. all collected specimens from all healthcare facilities
are sent to one facility to be sampled for sequencing (*population-wide* strategy); only *one*, 10%, 25%,
50%, or 100% of tertiary sentinel facilities would sample the specimens they collected for
sequencing). Turnaround time (i.e. time from specimen collection to acquisition of sequencing data)
was assumed to be negligible. 1,000 random bootstrap simulations were performed for each
guidance/surveillance strategy. We simulated epidemics for (**A**) Wild-type SARS-CoV-2/Alpha. (**B**)
Delta/Omicron. Grey regions denote that we could not reliably detect the variant virus with 95%
confidence using the guidance in question under the assumed genomic surveillance strategy.
293

Variant detection

To elucidate how SARS-CoV-2 testing rates and the proportion of positive specimens

sequenced impact the speed of variant detection, we simulated wild-type SARS-CoV-2/Alpha

and Delta/Omicron epidemics at different Ag-RDT availability ranging from 27

tests/100k/day to 1,000 tests/100k/day (Figure 3). We assumed that specimens to be

sequenced are sampled on their collection day, and varied the proportion of positive

specimens to sample for sequencing each day between 1% and 100%. We analyzed the

impact of testing rates and sequencing proportions on the expected day when the first

specimen sampled for sequencing containing the variant was collected as a measure of

variant detection speed. In Figure 3, we did not consider the time between sample collection

and sequencing nor the turnaround time to obtaining sequencing results as they would only

delay the actual day of variant detection by the assumed turnaround time.

Figure 3: **Impact of SARS-CoV-2 testing rates and proportion of positive specimens to sequence**
on variant detection. For each hypothetical daily test availability, the expected day when the first
variant specimen is sampled for sequencing since its introduction is plotted against the proportion of
positive specimens to be sampled for sequencing daily. Different genomic surveillance strategies with
varying sampling coverage (i.e. all specimens collected from all healthcare facilities sent to one
facility to be sampled for sequencing (*population-wide* strategy); only *one*, 10%, 25%, 50%, or 100%
of tertiary sentinel facilities would sample the specimens they collected for sequencing) were
simulated. (**A**) Wild-type SARS-CoV-2/Alpha. (**B**) Delta/Omicron.

For all testing rates, the relationship between the expected day when the first sample

containing the variant was collected and the proportion of positive specimens sequenced per

day can be described by a convex operating curve, reflecting rapidly diminishing returns in

the speed of variant detection as more specimens are sampled for sequencing. Across all

genomic surveillance sampling strategies, relatively larger marginal improvements to the

speed of variant detection are generally made when the sequencing proportion is increased

from 5 to 20% of all samples collected. Further sequencing only minimally shortens the

expected time to variant detection, as the operating curve asymptotically approaches the

earliest possible day of detection.

Importantly, increasing SARS-CoV-2 testing allows smaller sequencing proportions to attain

similar detection day targets, and higher testing rates lower the earliest possible detection

day. For both the Alpha and Omicron variants, increasing testing rates from 27 tests/100k/day

to 100 tests/100k/day brings forward the expected day of sampling the first variant sequence

by at least one week (Figure 3).

For the same level of testing and sequencing proportion, the population-wide strategy led to

the earliest initial detection of a variant sequence. If sequencing were restricted to samples

collected at a subset of tertiary sentinel facilities only, increasing the number of facilities

sending samples for sequencing reduced the spatiotemporal bias in the specimen pool,

thereby shaping the operating curves closer to the ones observed for the population-wide

strategy. Interestingly, results similar to the population-wide strategy could be attained if all

tertiary facilities acted as sentinel sites and sent the samples they collected for sequencing to

increase the representativeness of sampling.

Observed variant proportion

Test availability and sampling coverage also affect the accuracy of the observed variant

proportion (Figures 4 and S1). At a testing rate of 27 tests/100k/day, the observed variant

proportion maximally differs from the true circulating proportion by >30% for both the Alpha

and Omicron variants and for more than 15% of the time, the proportional difference between

the observed and true variation was greater than 20%. Both the maximum absolute difference

and percentage of timepoints where the difference is >20% can be lowered to <20% and <5%

respectively if testing rate is increased to 100 or more tests/100k/day.

Figure 4: **Impact of SARS-CoV-2 testing rates on the capacity to monitor changes in variant**
prevalence based on diagnostic test availability and proportion of test-positive samples
sequenced. Different genomic surveillance strategies (i.e. all specimens collected from all healthcare
facilities sent to one facility to be sampled for sequencing (*population-wide* strategy); only *one*, 10%,
25%, 50%, or 100% of tertiary sentinel facilities would sample the specimens they collected for
sequencing) were simulated. (**A**) Maximum absolute difference between observed and circulating
variant proportions. (**B**) Proportion of timepoints when sequencing was performed that the absolute
difference between observed and circulating variant proportions is greater than 20%.

Critically, when the representativeness of the specimen pool is spatiotemporally biased by

sequencing samples collected at tertiary sentinel facilities only, increasing the proportion of

specimens to be sequenced only marginally lowers the maximum absolute difference or

lessens the number of times where observed variant proportion deviates less than 20% from

true circulating proportions (Figure 4, near vertical isoclines at low daily rates of testing).

Increasing testing rates at sentinel surveillance sites provides more accurate detection in

changes to circulating prevalence than sequencing more samples in the context of low testing

rates.

Sensitivity analyses

We repeated our analyses using virus properties (i.e., incubation period, maximum viral load,

protection against infection by the mutant virus after extant virus infection) of the Omicron

variant but varied different relative transmissibility to the Delta variant (1·0 to 4·0) as well as

the initial proportion of individuals who had been infected by the Delta variant (10% and

40%). The variant growth rates simulated for these hypothetical Delta/Omicron epidemics

ranged from 0·17/day to 0·42/day.

Under these varied conditions, the expected day when the specimen of the first variant

sequence is collected still follows a convex-shaped operating curve against the daily

proportion of positive specimens to sequence. For all curves, the larger marginal

improvements in shortening variant detection are still in sequencing proportions of up to 5-

20% (Figure S2). In terms of the accuracy of observed variant to true circulating proportions,

the maximum absolute difference and percentage of timepoints where difference is >20% are

both substantially lowered if testing rate is increased to at least 100 tests/100k/day (Figures

S3-4).

We also varied the prevalence of extant Delta infections when the Omicron variant was

introduced (Figure 5). We found that lower test availability causes a delay in sampling the

first variant specimen if the variant is introduced when pre-existing extant variant circulation

is high. At 27 tests/100k/day, regardless of specimen proportions sequenced, detection could

be delayed by ~1 week if Omicron was introduced when Delta was circulating at 10%

prevalence as opposed to 1%. This is because a greater share of tests would be used to

diagnose the more prevalent extant virus infections which in turn decreases the likelihood of

detecting the newly introduced variant at low proportions.

Figure 5: **Impact of prevalence of extant variant of concern (   ) at the time of new variant**
introduction. For each Ag-RDT availability, the expected day when the first Omicron variant
specimen (in the background of Delta) is sampled for sequencing since its introduction is plotted
against the proportion of positive specimens to be sampled for sequencing daily. Each panel shows a
different prevalence of the Delta variant (   ) at the point of Omicron introduction. Sampling for
sequencing was drawn from the *population-wide* scenario.

Discussion

Our findings show that the emphasis on the size of the sample referred for genomic

surveillance is misplaced if testing capacity is insufficient and sample sources are highly

spatiotemporally biased. As such, at the current mean rate of testing in LMICs (27

tests/100k/day), current guidance[5–8] on sequencing sample size estimation could likely lead to

later-than-predicted detection of novel variants at best or, at worst, leave new variants

undetected until they have infected a majority of a population.

Based on our work, we identified three major areas of improvement that should be prioritized

to enhance the robustness of genomic surveillance programs (Figure 6). First, the most

substantial improvements are likely to come from increasing the mean testing rate in LMICs

from 27 tests/100k/day (Figure 6A) to at least 100 tests/100k/day (Figure 6B). Even if one

were to only doing sentinel surveillance at one tertiary facility, this increase in testing rate for

the catchment area of the facility would speed up variant detection by 1-2 weeks.

Figure 6: **Recommended approach to enhance genomic surveillance robustness**. In each plot, the
operating curves of the expected detection day of the Alpha variant with wild-type SARS-CoV-2 in
the background circulating at 1% are plotted for different proportion of specimens to sample for
sequencing per day and turnaround times. The vertical axes denote the number of days passed since
the introduction of the Alpha variant (left) and its corresponding circulating proportion (right). The
horizontal axes denote the proportion of positive specimens to sample for sequencing per day
(bottom) and the corresponding mean number of sequences to be generated per week per 1,000,000
people over a 90-day epidemic period. (**A**) Specimen pools for sequencing from *one* tertiary sentinel
facility with testing rate at 27 tests per 100,000 persons per day (tests/100k/day). (**B**) Specimen pools
for sequencing from *one* tertiary sentinel facility with testing rate at 100 tests/100k/day. (**C**) Specimen
pools for sequencing from 25% of all tertiary sentinel facilities with testing rate at 100 tests/100k/day.
(**D**) Zoomed-in plot of (C) to highlight sequencing proportions varying between 1-25%. Sequencing
5-10% of positive specimens (blue shaded region) would ensure that we would expectedly detect
Alpha in 30 days if turnaround time is kept within one week.

Second, the representativeness of a specimen pool for sequencing can be further improved by

expanding sampling coverage. In our model, variant detection was further sped up by 1-3

weeks by increasing the percentage of tertiary sentinel facilities sending the samples they had

collected for sequencing to 25% of facilities (Figure 6C). Additionally, in terms of prevalence

monitoring, if 25% of tertiary facilities sequenced 5% of all positive specimens they had

collected to detect and monitor an Alpha-like variant, the maximum absolute difference to

true circulating proportion is expected to decrease from >50% (assuming a single sentinel

facility) to no more than 20%.

Third, reducing turnaround time of sample referred for sequencing results in a 1:1 decrease in

time to new variant detection regardless of the proportion of sequenced samples, test

availability or sampling coverage (Figure 6). These gains require scale up in sample transport

networks, access to sequencing machinery, trained personnel, and/or increases in numbers of

sequenced samples to make the most efficient use of each sequencing run.

After reducing spatiotemporal bias in the specimen pool through increased testing and

sampling coverage, sequencing up to 5-10% of the positive specimens collected would return

the greatest information gains while minimizing resource wastage. For an Alpha-like variant,

at 100 tests/100k/day with sampling from 25% of tertiary sentinel facilities for sequencing,

this amounts to an estimated 5-10 sequences per week averaged over a 90-day period per

1,000,000 people. If turnaround time is kept within one week, the variant would likely be

detected within one month at ~4% circulating proportion (Figure 6D). Similarly, at the same

testing rate, sampling coverage and turnaround time (i.e. average 5-11 sequences per week

per 1,000,000 people), an Omicron-like variant would be detected before the first month

since its introduction but at ~23% circulating proportion owing to its faster transmission

(Figure S5).

Our findings here serve to inform expectations as genomic surveillance programs are being

developed and should be interpreted according to the public health objectives of each

program. If the objective is to serve as an early warning system for the emergence of new

variants of concern before they are likely to have spread widely, then all factors above are

essential and will likely require substantially more than 100 tests/100k/day. Critically,

determining that a new variant is a threat requires not only detection of the variant itself but

also the capacity to reliably monitor changes in its prevalence and potential clinical impact on

short timescales. The results presented here also inform the design of programs for the

sensitive and reliable detection of changes in variant prevalence.

The emergence and detection of each VOC to date represents interesting case studies for the

work described here (Supplementary Appendix). For example, at the time of first detection of

the Omicron variant, in South Africa in November 2021, the daily SARS-CoV-2 testing rate

was 51 tests/100,000 people/day (https://www.finddx.org/covid-19/test-tracker/), which was

among the highest testing rates in Africa. The Omicron variant was however detected 6-8

weeks after its likely emergence.[4] While this is commendable, Omicron had already infected

a substantial portion of the population in Gauteng, South Africa (i.e., the estimated

circulating variant proportion was >80% by mid-November).[4] Not only had the variant

already spread across the rest of South Africa and neighboring Botswana,[4] Omicron samples

were also collected in multiple other countries, including Hong Kong,[23] Denmark,[24] and the

Netherlands[25] before the initial reports of the existence of the Omicron variant. This situation

is consistent with the modelling findings, where novel variant detection is possible with <100

tests/100k/day but only after the new variant has spread widely across the population,[5]

abrogating any possibility of containment.

Expanding genomic sequencing capabilities, especially in LMICs, is a global priority[26] and

current investments in sequencing must continue.[27,28] Simultaneously, sustained investments

in public health systems are required to expand access to, and availability of, diagnostic

testing to underpin SARS-CoV-2 surveillance programs. Here, we primarily focused on

LMICs but our findings on the impact of testing rates and representativeness on genomic

surveillance programs are equally important for HICs as they consider dismantling parts of

their testing and surveillance infrastructure in the post-crisis phase of the pandemic. While

we find that routine representative sampling is vital for monitoring SARS-CoV-2 evolution,

additional surveillance systems, including targeted surveillance of particular populations and

settings (such as immunocompromised individuals or unusual events) could enable increased

sensitivity.[29] Ultimately, detecting the next SARS-CoV-2 variant or pathogen that causes the

next pandemic requires fundamental clinical diagnostic capacity to detect infections in the

first place.

Data sharing

All data relevant to the study are included in the Article, the Supplementary Appendix and

the github repository (https://github.com/AMC-LAEB/PATAT-sim). The PATAT model

source code is also available at https://github.com/AMC-LAEB/PATAT-sim.

Declaration of interests

A.T., E.H., S.C., B.R. and B.E.N. declare that they are employed by FIND, the global

alliance for diagnostics.

Acknowledgements

A.X.H. and C.A.R. were supported by ERC NaviFlu (No. 818353). C.A.R. was also

supported by NIH R01 (5R01AI132362-04) and an NWO Vici Award (09150182010027).

The authors are pleased to acknowledge that all computational work reported in this paper

was performed on the Shared Computing Cluster which is administered by Boston

University's Research Computing Services (www.bu.edu/tech/support/research/).

Authors' contributions

A.X.H. contributed to the conceptualization, data curation, formal analysis, investigation,
methodology, software, validation and visualization of the study. B.E.N. and C.A.R.
contributed to the conceptualization, data curation, funding acquisition, investigation,
methodology, project administration, resources, validation and supervision of the study.
J.A.S., M.P., S.B. and M.V.K. contributed to the conceptualization and interpretation of the
study. A.T., E.H., S.C., and B.R. contributed to the conceptualization, funding acquisition,
project administration, and resources of the study. E.P. contributed to the validation and
visualization of the study. A.X.H. and C.A.R. wrote the original draft of the manuscript. All
authors are involved in the review and editing of the manuscript. All authors had full access
to all data of the study and the final responsibility for the decision to submit for publication.

References

1   Robishaw JD, Alter SM, Solano JJ, *et al.* Genomic surveillance to combat COVID-19:
challenges and opportunities. *The Lancet Microbe* 2021; **2**: e481–4.
2   Davies NG, Abbott S, Barnard RC, *et al.* Estimated transmissibility and impact of
SARS-CoV-2 lineage B.1.1.7 in England. *Science (1979)* 2021; **372**: eabg3055.
3   Cherian S, Potdar V, Jadhav S, *et al.* SARS-CoV-2 Spike Mutations, L452R, T478K,
E484Q and P681R, in the Second Wave of COVID-19 in Maharashtra, India.
Microorganisms 2021, Vol 9, Page 1542 2021; **9**: 1542.
4   Viana R, Moyo S, Amoako DG, *et al.* Rapid epidemic expansion of the SARS-CoV-2
Omicron variant in southern Africa. *Nature 2022* 2022; : 1–10.
5   Brito AF, Semenova E, Dudas G, *et al.* Global disparities in SARS-CoV-2 genomic
surveillance. *medRxiv* 2021; : 2021.08.21.21262393.
6   Wohl S, Lee EC, DiPrete BL, Lessler J. Sample Size Calculations for Variant
Surveillance in the Presence of Biological and Systematic Biases. *medRxiv* 2022; :
2021.12.30.21268453.
7   Sequencing of SARS-CoV-2 - first update.
https://www.ecdc.europa.eu/en/publications-data/sequencing-sars-cov-2 (accessed
April 27, 2022).
8   Guidance for surveillance of SARS-CoV-2 variants: interim guidance, 9 August 2021.
https://apps.who.int/iris/handle/10665/343775 (accessed Feb 25, 2022).

9   Adepoju P. Closing Africa's wide COVID-19 testing and vaccination gaps. *The Lancet*
Microbe 2021; **2**: e573.

10 Brümmer LE, Katzenschlager S, Gaeddert M, *et al.* Accuracy of novel antigen rapid
diagnostics for SARS-CoV-2: A living systematic review and meta-analysis. *PLOS*
Medicine 2021; **18**: e1003735-.

11 Linton NM, Kobayashi T, Yang Y, *et al.* Incubation Period and Other Epidemiological
Characteristics of 2019 Novel Coronavirus Infections with Right Truncation: A
Statistical Analysis of Publicly Available Case Data. *Journal of Clinical Medicine*
2020, Vol 9, Page 538 2020; **9**: 538.

12 Kissler SM, Fauver JR, Mack C, *et al.* Viral dynamics of acute SARS-CoV-2 infection
and applications to diagnostic and public health strategies. *PLOS Biology* 2021; **19**:
e3001333-.

13 Hay JA, Kissler SM, Fauver JR, *et al.* Viral dynamics and duration of PCR positivity
of the SARS-CoV-2 Omicron variant. *medRxiv* 2022; : 2022.01.13.22269257.

14 Report 49 - Growth, population distribution and immune escape of Omicron in
England | Faculty of Medicine | Imperial College London.
https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-49-
Omicron/ (accessed Feb 25, 2022).

15 Pouwels KB, Pritchard E, Matthews PC, *et al.* Impact of Delta on viral burden and
vaccine effectiveness against new SARS-CoV-2 infections in the UK. *medRxiv* 2021; :
2021.08.18.21262237.

16 Mathieu E, Ritchie H, Ortiz-Ospina E, *et al.* A global database of COVID-19
vaccinations. *Nature Human Behaviour 2021 5:7* 2021; **5**: 947–53.

17 Dovel K, Balakasi K, Gupta S, *et al.* Frequency of visits to health facilities and HIV
services offered to men, Malawi. *Bull World Health Organ* 2021; **99**: 618–26.

18 Hasell J, Mathieu E, Beltekian D, *et al.* A cross-country database of COVID-19
testing. *Scientific Data 2020 7:1* 2020; **7**: 1–7.

19 Mwananyanda L, Gill CJ, Macleod W, *et al.* Covid-19 deaths in Africa: prospective
systematic postmortem surveillance study. *BMJ* 2021; **372**. DOI:10.1136/BMJ.N334.

20 Gill CJ, Mwananyanda L, MacLeod W, *et al.* Sustained high prevalence of COVID-19
deaths from a systematic post-mortem study in Lusaka, Zambia: one year later.
medRxiv 2022; : 2022.03.08.22272087.

21 Nichols BE, Girdwood SJ, Crompton T, *et al.* Monitoring viral load for the last mile:
what will it cost? *J Int AIDS Soc* 2019; **22**: e25337.

22   Girdwood SJ, Nichols BE, Moyo C, Crompton T, Chimhamhiwa D, Rosen S.

583   Optimizing viral load testing access for the last mile: Geospatial cost model for point

584   of care instrument placement. *PLOS ONE* 2019; **14**: e0221586.

585   23   Gu H, Krishnan P, Ng DYM, *et al.* Probable Transmission of SARS-CoV-2 Omicron

586   Variant in Quarantine Hotel, Hong Kong, China, November 2021. *Emerging Infectious*

587   *Diseases* 2022; **28**: 460.

588   24   Espenhain L, Funk T, Overvad M, *et al.* Epidemiological characterisation of the first

589   785 SARS-CoV-2 Omicron variant cases in Denmark, December 2021.

590   *Eurosurveillance* 2021; **26**: 2101146.

591   25   Omicron variant found in two previous test samples | RIVM.

592   https://www.rivm.nl/en/news/omicron-variant-found-in-two-previous-test-samples

593   (accessed March 17, 2022).

594   26   Adepoju P. Challenges of SARS-CoV-2 genomic surveillance in Africa. *The Lancet*

595   *Microbe* 2021; **2**: e139.

596   27   Tegally H, San JE, Cotten M, *et al.* The evolving SARS-CoV-2 epidemic in Africa:

597   Insights from rapidly expanding genomic surveillance. *medRxiv* 2022; published

598   online April 22. DOI:https://doi.org/10.1101/2022.04.17.22273906.

599   28   Leite JA, Vicari A, Perez E, *et al.* Implementation of a COVID-19 Genomic

600   Surveillance Regional Network for Latin America and Caribbean region. *PLOS ONE*

601   2022; **17**: e0252526.

602   29   Knyazev S, Chhugani K, Sarwal V, *et al.* Unlocking capacities of genomics for the

603   COVID-19 response and future pandemics. *Nature Methods 2022 19:4* 2022; **19**: 374–

604   80.

606