Plant to animal protein ratio in the diet: nutrient adequacy, long-term health and environmental pressure

Hélène Fouillet[1]*, Alison Dussiot[1], Elie Perraud[1], Juhui Wang[1], Jean-François Huneau[1], Emmanuelle Kesse-Guyot[2], François Mariotti[1]

[1]Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, 75005, Paris, France

[2]Sorbonne Paris Nord University, Inserm, Inrae, Cnam, Nutritional Epidemiology Research Team (EREN), Epidemiology and Statistics Research Center (CRESS) – University of Paris, 93017 Bobigny, France

*Address correspondence to Dr. Helene Fouillet, helene.fouillet@agroparistech.fr or Prof. François Mariotti, francois.mariotti@agroparistech.fr.

Sources of support: The authors report that no funding was received for this study.

Author disclosures: HF has received a research grant by INRAE from Roquette. FM has received research grants as PhD fellowships under his direction by AgroParisTech and INRAE from Terres Univia and Ecotone foundation, under the aegis of Fondation de France.

Availability Statement: Data described in the manuscript, code book, and analytic code will be made available upon request pending application and approval.

3 Figures, 2 Tables

16 Supplementary Data (2 Supplemental Methods, 8 Supplemental Tables, 6 Supplemental Figures)

Running title: Diet plant to animal protein ratio

Abbreviations used: ANSES, French Agency for Food, Environmental and Occupational Health and Safety; CIQUAL, French Information Centre on Food Quality; DALYs, Disability-adjusted life-years; *DD*, Diet departure criterion; GBD, Global Burden of Diseases; GHGe, greenhouse gas emissions; *HR*, Health risk criterion; INCA3, Third Individual and National Study on Food Consumption French Survey; %PP, Percentage of plant protein in the diet; TMREL, Theoretical minimum risk exposure level.

Abstract

Background: Animal and plant protein sources have contrasting relationships with nutrient adequacy

and long-term health, and their adequate ratio is highly debated.

Objective: We aimed to explore how the percentage of plant protein in the diet (%PP) relates to

nutrient adequacy and long-term health but also to environmental pressures, to determine the

adequate and potentially optimal %PP values.

Design: Observed diets were extracted from the dietary intakes of French adults (INCA3, n=1 125),

and optimized by non-linear multi-criteria optimization. Using reference values for nutrients and

disease burden risks for foods, and accounting for nutrient bioavailability and food cultural

acceptability, we identified modeled diets with graded %PP values that simultaneously ensure

nutrient adequacy, minimize long-term health risk and preserve at best dietary habits. We explored

the tensions between objectives and identified the most critical nutrients and influential constraints

by sensitivity analysis. Environmental pressures related to the modeled diets were estimated using

the Agribalyse database.

Results: Nutrient-adequate diets must fall within the ~15%-80% %PP range, a slightly wider range

being nevertheless identifiable by waiving the food acceptability constraints. Fully healthy diets, also

achieving the minimum-risk exposure levels for both unhealthy and healthy foods, must fall within

the 25%-70% %PP range. All of these healthy diets were very distant from current typical diet.

Those with higher %PP had lower environmental impacts, notably on climate change and land use,

while being as far from current diet.

Conclusions: There is no single optimal %PP value when considering only nutrition and health, but

high %PP diets are more sustainable. For %PP>80%, nutrient fortification/supplementation and/or

new foods are required.

Keywords: Plant-based diets; Healthy dietary patterns; Nutrient adequacy; Environmental

footprints; Diet optimization

Introduction

Historical and current nutritional transitions are coupled with changes in the relative contribution of

dietary animal and plant proteins. This has been studied from hunter-gatherers to post-agricultural

societies (1), from traditional diets that enabled thriving civilizations to diets in post-industrialized

countries (2, 3), and more recently, in western countries with the emerging trend toward more plant-

predominant diets.

Changes in plant and animal protein intake raise classic nutritional questions. One in particular

concerns the possible risk of deficiency with diets too low in animal protein, since animal protein

foods contribute significantly to the intake of indispensable nutrients (4). However, plant proteins are

also important for the intake of some indispensable nutrients and fiber, and are also lower in

saturated fats (5). Beyond the relationship to nutrient adequacy, animal/plant proteins and their

packages largely affect the metabolome and the microbiota and physiological functions that are

crucial for long-term health (5-8). Accordingly, there have been many contrasting associations

reported recently between plant and animal protein intake and mortality, especially regarding

cardiovascular diseases (5, 9, 10).

More globally, plant (such as legumes, nuts and whole grains) and animal (such as red and

processed meats) protein sources have heterogeneous relationships to nutrient adequacy (11) and

to long-term health regarding cardiovascular diseases (7, 12-14) and cancers (15, 16). There is

indeed a challenge for food-based dietary guidelines to point out what proportions of plant and

animal protein foods should be recommended (17, 18). However, the plant to animal protein ratio

remains a poor, summarizing descriptor of dietary patterns, since two diets with the same plant to

animal protein ratio can actually be very different (7). There is thus a need to analyze the overall

proportion of plant protein in the diet in view of the related dietary profiles and their nutritional

adequacy and healthiness.

Furthermore, current interest in the proportions of plant and animal proteins in the diet also stems

from their differential association with environmental pressures, in particular greenhouse gas

emissions (GHGe) and land use (19-22). Altogether, the plant to animal protein ratio in the diet

appears central to the sustainability of the food systems (19, 23). This has implications for dietary

guidelines that aim to encompass both human and planetary health (24-26).

Here, using advanced diet modeling and optimization, we studied whether an optimal proportion of

plant protein in the diet (%PP, the percentage of plant protein in total protein intake) can be

identified when taking into account the reference values for nutrients and the disease burden risks

for food categories. We characterized modeled diets that departed as little as possible from

prevailing diets at all levels of adequate %PP values for nutrient adequacy and long-term health to

identify nutritional issues (i.e., limiting nutrients) and dietary levers (i.e., effective foods). We

furthermore estimated the environmental pressures associated with modeled diets along the whole

range of adequate %PP values.

Methods

<u>Input of dietary data</u>

The data used for this study were extracted from the French Individual and National Study on Food

Consumption Survey 3 (INCA3) conducted in 2014-2015. The INCA3 survey is a representative

cross-sectional survey of the French population; its method and design have been fully described

elsewhere (27). Males aged 18-64 y (n=564) and pre-menopausal females aged 18-54 y (n=561),

not identified as under-reporters, were included in the present study; the final sample contained

1125 adults.

Dietary data were collected by professional investigators assisted by a dietary software from three

unplanned, non-consecutive, 24h dietary recalls spread over a three-week period. Portion sizes

were estimated using validated photographs (27), and the nutrient contents of different food items

came from the 2016 food composition database operated by the French Information Centre on Food

Quality (CIQUAL) (28). Mixed foods were broken down into ingredients and then gathered into 45

food groups (Supplemental Table 1). For each sex, the nutrient content of each food group was

calculated as the mean nutrient content of food items constituting the food group weighted by their

mean intake by the sex considered, as previously described (29).

<u>Multi-criteria diet optimization under constraints</u>

Using multi-criteria optimization, we identified modeled diets (i.e., modeled consumptions of the 45

food groups) with a minimal long-term health risk and a minimal departure from the observed diet

(taking into account cultural acceptability and inertia), under constraints that would ensure adequate

nutrient intakes and remain within current consumption limits. In this context, we investigated the

role of %PP to identify its adequate range of variations and to characterize the dietary, nutritional

and environmental consequences of these variations.

This non-linear optimization problem was performed using the NLP solver of the OPTMODEL

procedure of SAS software version 9.4 (SAS Institute Inc., Cary, NC, USA.). Optimization was

implemented at the population level but in males and females, separately. The optimized diets of

males and females were then averaged to derive optimized diets for the adult population.

Objectives

The main optimization objective was to minimize the long-term health risk of the modeled diet, as

assessed by the Heath Risk (*HR*) criterion. The *HR* criterion was set to target the dietary

recommendations from the Global Burden of Diseases (GBD) based on epidemiological studies

about the associations between consumption of different food groups and risk of chronic diseases

(30). The *HR* criterion thus aimed to limit the consumption of three unhealthy food groups or

categories (red meat, processed meat and sweetened beverages), while promoting that of six

healthy food groups or categories (whole grain products, fruits, vegetables, legumes, nuts and

seeds, and milk) until their minimum risk exposure levels (TMREL) values were reached. According

to the most recent (2019) estimates from the GBD, TMREL values were 0 g/d for red meat,

processed meat and sweetened beverages, and 150, 325, 300, 95, 14.5 and 430 g/d, respectively,

for whole grain products, fruits, vegetables, legumes, nuts and seeds, and milk (30). In our study,

the *HR* criterion was thus expressed and minimized as:

$$\text{Min } HR = \sum_{i=1}^{3} \left(\frac{\text{Opt}(i)}{\text{Max}(i)} \times \frac{\text{DALYs}(i)}{\text{DALYs}(\text{all})} \right) + \sum_{j=1}^{6} \left(\max \left[\frac{\text{TMREL}(j) - \text{Opt}(j)}{\text{TMREL}(j)} ; 0 \right] \times \frac{\text{DALYs}(j)}{\text{DALYs}(\text{all})} \right)$$

Where i denotes the food groups to be decreased (red meat, processed meat and sweetened

beverages), j denotes the food groups to be increased (whole grain products, fruits, vegetables,

legumes, nuts and seeds, and milk), Opt(i) and Opt(j) are the optimized consumptions of food

groups i and j, respectively (in g/d), Max(i) is the upper limit of consumption of food group i (in g/d),

TMREL(j) is the TMREL value of food group j (in g/d), DALYs(i) and DALYs(j) are the disability-

adjusted life-years associated with excessive or insufficient consumptions of food groups i and j,

respectively (in y), and DALYs(all) is the sum of all DALYs(i) and DALYs(j). The Max values used

were the maximal recommended consumption of unhealthy foods in line with the French dietary

guidelines (31): 71g/d for red meat, 25g/d for processed meat and 263 g/d (corresponding to the

average portion size) for sweetened beverages intake. The TMREL and DALYs values used were

issued from the most recent (2019) estimates from the GBD (30) adapted to our study context (by

using sex-specific and French DALYs values, Supplemental Table 2).

We also evaluated how the modeled diets deviated from current diets, in order to consider inertia to

changes in food consumption, which is one way to account for social/cultural acceptability. The *Diet*

Departure (*DD)* criterion was defined as the sum of the squares of the differences between

observed and optimized food group consumption, standardized by their observed standard

deviations, as previously explained (29). *DD* was thus expressed and minimized as:

$$\text{Min}\,DD = \sum_{k=1}^{n} \left[\frac{\text{Obs(k) - Opt(k)}}{\text{SD(k)}} \right]^2$$

Where k is the number of food groups (n=45), Obs(k) and Opt(k) are respectively the observed and

optimized consumption of food group k (in g/d) and SD(k) is the current standard deviation of the

consumption of food group k.

<u>Constraints</u>

During diet optimization, the total energy intake was constrained to stay within ±5% of its observed

value. Thirty-five nutritional constraints were applied to ensure adequate nutrient intake in the male

and female populations (Supplemental Table 3), based on the most recent reference values from

the French Agency for Food, Environmental and Occupational Health & Safety (ANSES) (32). We

did not consider any constraints for vitamin D, because its reference value is known to be much too

high to be reached by a non-fortified diet alone (29, 31). As the absorption of iron and zinc is

dependent on dietary factors, the requirements were based on bioavailable iron and zinc calculated

from the dietary intake using equations that predict their absorption (33-35), as detailed in a

previous study by our group (29). This previous study had demonstrated that current

recommendations regarding bioavailable iron and zinc are very constraining when trying to model

healthier diets, these recommendations being much higher than current intakes (e.g., there is a

current iron-deficiency anemia prevalence of 4.1% in French women) (29). Therefore, like in this

previous study, we used threshold values lower than current reference values. They correspond to a

deficiency prevalence of 5%, because such flexibility enables the identification of diets that are

apparently healthier overall, with a better balance in disability-adjusted life-years (DALYs) due to

less cardiometabolic disease, despite a higher prevalence of iron-deficiency anemia (29). In

addition, to take into account the slightly lower digestibility of plant *vs* animal proteins regarding the

nutritional constraint on protein requirement, a 5% penalty was applied to protein intake from plant

protein food items, as previously described (36). As the intake of individual amino acids is generally

adequate when the protein intake is sufficient in a varied diet (37), only protein requirements were

considered in the model constraints, but we have *a posteriori* verified that modeled diets also

contained adequate intakes of indispensable amino acids by using a database of the amino acid

composition of food groups (Supplemental Method 1).

Moreover, some acceptability constraints were applied to the food group consumption

(Supplemental Table 4). Acceptability constraints aimed to keep the food group intakes within the

range of observed intakes, by bounding each food group intake between its 5[th] and 95[th] percentile of

observed consumption in males and females separately. We did not do this for the unhealthy food

groups or categories (red meat, processed meat and sweetened beverages), for which a dietary

constraint with an upper limit was already defined according to the French dietary guidelines.

Another exception was made for some healthy food groups (legumes and milk) that had 95[th]

percentile values slightly lower than TMREL values, and for which the upper limit has thus been

raised to their TMREL values.

Optimization strategy

We firstly aimed to determine the range of adequate %PP values in the diet that would ensure

nutrient adequacy with a minimal long-term health risk. This first problem of identifying the adequate

%PP range was addressed by optimizing the *HR* criterion under all the nutritional and acceptability

constraints, with an additional constraint on %PP that was iteratively parameterized according to a

grid search. This grid search constraint forced the %PP value to be equal to x%, with x% varying

from 0% to 100% by steps of 5% (or even 1% at the edges of the adequate %PP range). As this

problem was often non-uniquely identifiable, leading to different solutions with slightly distinct

dietary patterns but similar *HR* values (especially for the intermediate %PP values that allowed for a

variety of food group combinations with a similarly null *HR* value), we choose to systematically

select the dietary solution that was the most acceptable *a priori*, based on the lowest departure from

the current diet. According to the hierarchical method in multi-criteria optimization (38), this second

problem of diet selection was addressed in a second stage. This time it was done by optimizing the

DD criterion under the constraint that *HR* was equal to its previously identified minimal value, always

under all the nutritional and acceptability constraints, and the grid search constraint on %PP

covering its previously identified adequate range.

<u>Limiting nutrients and contribution of food groups to their intake</u>

We conducted a dual value analysis to better characterize the tensions between %PP, nutrient

adequacy and long-term health. We reported the dual values associated with the %PP equality

constraint and the nutritional constraints during *HR* optimization (obtained during the first problem

solving, as explained above), which represent the potential *HR* gain if the limiting bound (lower or

upper) of the considered constraint was relaxed by one unit. In order to compare the relative

influence of nutrients, their dual values were standardized to represent the potential *HR* gain if the

limiting bound was relaxed by 10%, to classify nutrients from the most limiting (higher absolute

standardized dual value) to the least limiting (lowest absolute standardized dual value).

For the most limiting nutrients in the different modeled diets (i.e., nutrients with the most active

constraints), we studied contributions of different food groups to intake of that particular nutrient in

each modeled diet identified for each adequate %PP value (i.e., in the modeled diets resulting from

the second problem solving, as explained above).

<u>Sensitivity analysis</u>

We also conducted a sensitivity analysis to assess the influence of some constraints of particular

interest. We thus compared the results obtained when requiring the deficiency prevalence to be

≤1% rather than ≤5% (main model) in the nutritional constraints for bioavailable iron and zinc (their

alternative threshold values are given in Supplemental Table 3), and when removing or not (main

model) all the dietary and acceptability constraints on food group intakes.

<u>Diet environmental impacts</u>

Finally, to assess environmental pressures related to the observed and modeled diets, we used the

French agricultural life cycle inventory database AGRIBALYSE®; its methodological approach

(summarized in Supplemental Method 2) has been described elsewhere (39-41). In particular, we

evaluated the food-related GHGe (in kg CO_2eq, with the non-CO_2 GHGe included and weighted

according to their relative impact on warming), land use (referring to the use and transformation of

land, dimensionless), water use (relating to the local scarcity of water, in m^3 water deprivation) and

fossil resource use (use of non-renewable fossil resources such as coal, oil, and gas, in MJ),

together with a single environmental footprint score (dimensionless) that aggregated 16 indicators

(42).

Results

<u>Range of adequate %PP values and identified tensions between %PP, nutrient adequacy and long-</u>

<u>term health</u>

The adequate %PP range compatible with nutrient adequacy was 16%-82% in males and 16%-77%

in females, and only the 25%-70% %PP range was additionally compatible with a minimal health

risk (*HR* criterion) for both sexes (**Table 1**). In this narrower range, a null *HR* value was attained by

the removal of unhealthy foods (red meat, processed meat and sweetened beverages) and an

increase in healthy foods (whole grain products, fruits, vegetables, legumes, nuts and seeds, and

milk) up to or above their TMREL values (30).

Among the %PP equality constraint and the nutritional constraints, none were found limiting for *HR*

minimization over the 25%-70% %PP range (**Table 2**). The %PP equality constraint was limiting

only for %PP values lower than 25% (strongly) and higher than 70% (more moderately, due to the

lower *HR* impact of the milk decrease for the highest %PP values than of the red meat increase and

whole grain product decrease for the lowest %PP values). Nutrients identified as increasingly

limiting as %PP decreased below 25% were fiber, sugar (excluding lactose), saturated fatty acids

and atherogenic fatty acids (lauric, myristic and palmitic acids). As %PP decreased below 25%, it

was hence increasingly challenging to maintain sufficient intake of fiber and non-excessive intakes

of sugar and fatty acids (as shown by the opposite sign of their dual values), which resulted in

dietary solutions of increasingly degraded *HR* values. Nutrients that were identified as increasingly

limiting when %PP increased above 70% were iodine, sodium, vitamin B2, calcium, EPA+DHA,

vitamin A and ⍺-linolenic acid in both sexes together with vitamin B12 in males and bioavailable iron

in females. As %PP increased above 70%, its was increasingly challenging to maintain sufficient

intakes of these nutrients and a non-excessive sodium intake. The other nutrients (n=20, those not

shown in Table 2) were never limiting over the adequate %PP range, including, of note, protein.

Sensitivity analysis showed that being more demanding for bioavailable iron and zinc (i.e.,

constraining their deficiency prevalence at ≤1% rather than ≤5% as in the main model) resulted in

slightly restricting the adequate %PP range on the right (16%-79% in males and 16%-70% in

females), and the %PP range ensuring a null *HR* value on both sides (30%-65% in males and 35%-

45% in females) (data not shown). Conversely, when suppressing all the food group consumption

limits (i.e., all the dietary and acceptability constraints) from the model (Supplemental Table 5), the

range of adequate %PP values was expanded on both sides (8%-94% in males and 8%-92% in

females), as was the %PP range ensuring a null *HR* value (16%-86% in males and 16%-84% in

females), but consistently with the same limiting nutrients as in the main model (in particular,

insufficient fiber intake for excessively low %PP values, or insufficient intakes of vitamin B12, iodine

and EPA+DHA for excessively high %PP values).

<u>Modeled diets</u>

All the modeled diets identified (**Figure 1** and Supplemental Figure 1) were very distant from the

current typical French diets, with departure values (*DD* criterion*)* equal to or greater than twice the

standard deviation observed in the population. Furthermore, most of the modeled diets with a null

HR value (i.e., in the 25%-70% %PP range) were all about equally distant from the observed diets,

with close *DD* values (differing by less than 20%) in the 35%-65% %PP range and similar *DD*

values (differing by less than 5%) in the 45%-60% %PP range.

Although the energy intake remained relatively stable between modeled and observed diets (by

construction), the total intakes of both animal-based and plant-based foods were increased in the

25%-70% %PP range, notably owing to the important increases in milk, fruits and vegetables up to

or above their TMREL values (Supplemental Table 6 and Supplemental Figure 2). Regarding plant

products, all the modeled diets exhibited dramatic increases in fruits and vegetables, whole grain

products and legumes and nuts. Regarding animal products, red and processed meats were readily

removed as %PP increased. These meats were replaced by poultry and eggs, which transiently

increased, the modeled diets then being meat-free from PP%=60%. Dairy and seafood were the

only remaining animal products at the right end of the adequate %PP range (Supplemental Figure

2).

Over the entire adequate %PP range, including meat-free diets, the intakes of protein and of each

indispensable amino acid were always much higher than their 98% safe intake thresholds (**Figure 2**

and Supplemental Table 7 and Supplemental Figure 3).

<u>Contributions of food groups to limiting nutrient intakes</u>

Regardless of their %PP value, all the modeled diets were nutrient-adequate, in contrast with

observed diets (Supplemental Table 8).

As %PP increased, it was increasingly difficult to maintain sufficient intakes of bioavailable iron,

vitamins B12, B2 and A, and iodine and calcium, owing to the decreases in the animal products that

were their main contributors (red meat, dairy products and eggs) (Supplemental Figure 4). The

EPA+DHA and ⍺-linolenic acid intakes, which are largely insufficient in the observed diets, were

made sufficient in all the modeled diets by increases in their main contributors, respectively,

seafood and added fats, with difficulties to maintain them sufficient for the highest %PP values

(Supplemental Figure 4). The sodium intake, which is dramatically excessive in the observed diets,

was reduced to its upper limit in all the modeled diets as a result of removing processed meat and

reducing refined grain products, with difficulties to maintain sodium not excessive for the highest

%PP values, due to increases in some starch and miscellaneous foods (Supplemental Figure 4).

Conversely, as %PP decreased, it was increasingly difficult to maintain a sufficient intake of fiber

and non-excessive intakes of sugar and saturated fatty acids, due to the meat and dairy increases

(Supplemental Figure 5).

<u>Environmental impacts of modeled diets</u>

Across modeled diets, GHGe gradually decreased as %PP increased, with a stagnation from

%PP=70% onwards, from which the GHGe were 43% lower than with the observed diet (**Figure 3**).

Similar trends were observed for land use and, to a lesser extent, fossil resource use (Supplemental

Figure 6), with 43% and 12% decreases, respectively, from the observed to the modeled diet with

%PP=70%. In contrast, water use was similarly ~40% higher for all the modeled diets, regardless of

their %PP value, than for the observed diet, due to their very high levels of fruits and vegetables

that were by far the most water-demanding food groups (Supplemental Figure 6). Overall, at the

level of the single environmental footprint score that aggregated 16 indicators, the same trend was

observed as for GHGe, with a 34% decrease in this aggregated score from the observed to the

modeled diet with %PP=70% (Supplemental Figure 6).

Discussion

Gathering all nutritional information over a large spectrum that covered nutrient reference values

and long-term health risks, our study formally establishes ranges of plant protein proportion (%PP)

for nutrient-adequate and healthy diets. One major finding is that there is no optimal %PP value, as

we found a spectrum of similarly healthy diets over the 25%-70% range. However, diets in the upper

end were associated with substantially lower GHGe and overall environmental impact.

A wide dietary %PP range, from ~15% to 80%, was found compatible with providing all nutrients in

adequate amounts. As shown by sensitivity analysis, this %PP range was slightly restrained by the

considered constraints for food acceptability, whereas the nutritional issues identified remained

broadly the same with or without these constraints. From a nutritional viewpoint, no diet with %PP <

~15% was able to provide enough fiber and non-excessive amounts of saturated fatty acids, while

also satisfying all constraints for nutrient intakes and food acceptability. More interestingly, given the

ongoing dietary transition, we could not find diets with %PP > ~80% that would provide sufficient

amounts of a large set of nutrients, particularly iodine, vitamin B12 (in males), bioavailable iron (in

females), calcium and EPA+DHA. These nutrients are considered to be at issue in vegetarian diets

(except calcium and iodine in lacto-ovo-vegetarian), notably calcium and B12 in predominantly

plant-based diets (43, 44). From a dietary viewpoint, as shown when approaching the critical value

of %PP=80%, dairy appeared to be key to preventing iodine and calcium shortages. Seafood,

meanwhile, appeared critical to providing EPA+DHA (with oily fishes as the main source) as well as

iodine and B12. Milk and seafood were the last remaining animal products at the highest %PP

values, confirming their importance as healthy, nutrient-dense protein sources (45). Healthy plant

protein sources such as legumes and nuts apparently could not replace milk or seafood. This is

because they actually reached their upper allowed intake very early (as soon as %PP=25% for

legumes), which indicates that they constitute an effective dietary lever. However, even when

removing all food intake limits (in sensitivity analysis), it remained impossible to obtain 100% plant-

based diets because of the same nutritional issues (insufficient intakes of vitamin B12, iodine and

EPA+DHA). Our findings do not indicate that vegetarian (without seafood) or vegan diets (without

seafood and dairy) cannot be nutritionally adequate. It means that solutions for diets that are

entirely or almost entirely plant-based should rely on additional food products than those presently

consumed by the general population, including fortified foods (46-48). This warrants further studies

about the potential of new foods to extend the limit of the %PP range identified as adequate here.

Within the wide range of nutrient-adequate %PP values, we did not find a single optimal diet, but a

large range of diets with %PP from 25% to 70%. These diets were all optimal when considering

their health value, because their food consumptions complied with minimum-risk exposure levels.

These consisted of no red meat and high levels of fruits and vegetables, whole grains, legumes,

nuts and milk, in line with dietary guidelines (49). Modeled healthy diets were variations of this

pattern, which explains why they were also similarly distant from current diets. Within this healthy

pattern spectrum, the increase in %PP was predominantly related to the decrease in total and

animal proteins. This occurred mostly in poultry and eggs and, to a lesser extent, dairy. This finding

aligns well with the current spectrum of observed diets, with plant-based diets being higher in plant

protein but especially low in total and animal protein (50, 51). This could simply be ascribed to the

higher protein density in animal protein sources compared to plant protein sources. Also, the

nutrients identified as limiting at the borders of the healthy %PP range appear to be related to the

nutrient density of animal vs plant protein sources when expressed relative to protein density.

However, the dietary protein amount was never limiting, even at the highest %PP levels. There is a

growing consensus that the protein package and not the protein per se are important to the question

of plant to animal protein ratio in the diet (5, 7). Likewise, indispensable amino acid amounts were

well above reference values based on requirements. In real diets, composed of a mix of different

types of proteins that complement each other, sufficient amounts of protein appear to guarantee

sufficient amounts of amino acids (20, 51, 52).

Distance from the prevailing diets is often used in diet modeling to take into account so-called

cultural acceptability (53-55), also referred to as dietary inertia (56). In this study, healthy diets in the

35-65% %PP range departed rather similarly from the prevailing diets, which are still at ~35% %PP.

This confirms that the plant to animal protein ratio is, by itself, a poor descriptor of diet

characteristics, and so blanket statements about the right %PP are not warranted. Given that

modeled healthy diets ranging from 35% %PP (the level of the current diets in western countries) to

65% %PP were all very distant from current diets, our study also shows that overcoming dietary

inertia is required for healthy diets, irrespective of the plant to animal protein target ratio (57).

The GHGe and overall composite score for environmental pressures were lower for healthy

modeled diets than observed diets, and all the more as %PP increased. A large body of literature

has reported that diets which are more plant-based are associated with lower environmental

pressure, and vice versa, whether diets were modeled (44, 55, 58, 59), observed (60-62) or

composite (23). However, until our study, this relationship had not yet been shown according to

%PP in healthy diets. In our setting, %PP was strongly associated with the environmental impact of

healthy diets. As compared to the prevailing diets, lower GHGe and composite score are firstly

explained by the removal of total red meat in all healthy diets, red meat accounting for ~1/3 of the

pressure in prevailing diets. This is in line with the literature that points to red meat and associated

sustainability concerns (59, 63). Finally, we found that other environmental pressures (land use and

fossil resource use), except water use, had similar patterns of change, in line with the literature (19,

44). The general relationship between %PP and environmental pressure can mostly be ascribed to

the fact that animal sources are rich in protein, and that livestock breeding is associated with higher

resource use, higher land use, and higher GHGe (21, 64-66). Nevertheless, further investigation of

the relationship between %PP and environmental impacts would require prioritizing the minimization

of environmental impacts over that of diet departure. Therefore, we cannot rule out the possibility

that moderate %PP diets, if well-designed, may have as low environmental impacts as high %PP

diets, at the cost of a larger diet departure.

This study has some limitations. We modeled diets according to changes in intakes of food groups,

based on the present food repertoire and current intake levels in the population. Food grouping is

critical in diet modeling (31), and food diversity and composition can change rapidly in western

countries, as seen by recent changes (67). A similar limitation applies to the assessment of a diet's

environmental impacts, for which also we did not consider variations related to food production

systems (68, 69). Nevertheless, we used a classical food grouping, which helps represent dietary

patterns at an appropriately high level of detail. We also believe that using standard/traditional foods

in modeling provides a good starting point to evaluate the situation before considering changes in

the food offer or food composition. Our study uses sources of information as background

parameters, including references/targets for nutrients and food categories. Clearly, there are many

uncertainties in this regard (31). Nonetheless, we believe that a strength of our study is our use of a

conceptual framework that aggregates most of the state of the art knowledge in nutrition.

To conclude, we identified that the range of equally optimal %PP values for nutrition and health is

wide (25%-70%), and that all of these healthy diets deviate greatly from prevailing diets. From a

public health perspective, there is no unique, optimal %PP value when considering nutrition and

health alone. However, significant changes in current eating habits are nonetheless required to

achieve healthier diets. Moreover, in the higher end of the plant protein percentage range, modeled

healthy diets have a lower environmental impact and are thus more sustainable than other healthy

diets. At %PP > ~80%, changes in food repertoire diversity, food composition, nutrient enrichment

or nutrient supplementation are required for fully nutrient-adequate diets.

Acknowledgments and statement of authors' contributions to the manuscript

The authors thank Vincent Colomb and Mélissa Cornélius (Ademe) for their important support in the

use of the Agribalyse database.

HF and FM designed the research; HF conducted the research and analyzed the data; AD, EP, FM,

JW, J-FH and EK-G provided methodological support and help with interpretation of the results; HF

and FM wrote the first draft of the manuscript; and all authors provided critical comments on the

manuscript. HF and FM had primary responsibility for the final content, and all authors have read

and approved the final manuscript.

References

1. Styring AK, Fraser RA, Arbogast R-M, Halstead P, Isaakidou V, Pearson JA, Schäfer M, Triantaphyllou S, Valamoti SM, Wallace M, et al. Refining human palaeodietary reconstruction using amino acid δ15N values of plants, animals and humans. J Archaeol Sci 2015;53:504-15. doi: 10.1016/j.jas.2014.11.009.

2. World Health Organization (WHO). Diet, nutrition and the prevention of chronic diseases. 3. Global and regional food consumption patterns and trends. Technical Report Series 916. 2003:1-149. Internet: http://apps.who.int/iris/bitstream/handle/10665/42665/WHO_TRS_916.pdf?sequence=1.

3. Winnie Gerbens-Leenes P. Dietary Transition: Longterm Trends, Animal Versus Plant Energy Intake, and Sustainability Issues. Edtion ed. In: Mariotti F, ed. Vegetarian and Plant-Based Diets in Health and Disease Prevention: Academic Press, 2017:117-34.

4. Phillips SM, Fulgoni VL, 3rd, Heaney RP, Nicklas TA, Slavin JL, Weaver CM. Commonly consumed protein foods contribute to nutrient intake, diet quality, and nutrient adequacy. Am J Clin Nutr 2015. doi: 10.3945/ajcn.114.084079.

5. Mariotti F. Animal and Plant Protein Sources and Cardiometabolic Health. Adv Nutr 2019;10(Suppl_4):S351-S66. doi: 10.1093/advances/nmy110.

6. Lepine G, Fouillet H, Remond D, Huneau JF, Mariotti F, Polakof S. A Scoping Review: Metabolomics Signatures Associated with Animal and Plant Protein Intake and Their Potential Relation with Cardiometabolic Risk. Adv Nutr 2021;12(6):2112-31. doi: 10.1093/advances/nmab073.

7. Richter CK, Skulas-Ray AC, Champagne CM, Kris-Etherton PM. Plant Protein and Animal Proteins: Do They Differentially Affect Cardiovascular Disease Risk? Adv Nutr 2015;6(6):712-28. doi: 10.3945/an.115.009654.

8. Dietrich S, Trefflich I, Ueland PM, Menzel J, Penczynski KJ, Abraham K, Weikert C. Amino acid intake and plasma concentrations and their interplay with gut microbiota in vegans and omnivores in Germany. Eur J Nutr 2022. doi: 10.1007/s00394-021-02790-y.

9. Naghshi S, Sadeghi O, Willett WC, Esmaillzadeh A. Dietary intake of total, animal, and plant proteins and risk of all cause, cardiovascular, and cancer mortality: systematic review and dose-response meta-analysis of prospective cohort studies. BMJ 2020;370:m2412. doi: 10.1136/bmj.m2412.

10. Zhong VW, Allen NB, Greenland P, Carnethon MR, Ning H, Wilkins JT, Lloyd-Jones DM, Van Horn L. Protein foods from animal sources, incident cardiovascular disease and all-cause mortality: a substitution analysis. Int J Epidemiol 2021. doi: 10.1093/ije/dyaa205.

11. Salomé M, de Gavelle E, Dufour A, Dubuisson C, Volatier JL, Fouillet H, Huneau JF, Mariotti F. Plant-Protein Diversity Is Critical to Ensuring the Nutritional Adequacy of Diets When Replacing Animal With Plant Protein: Observed and Modeled Diets of French Adults (INCA3). J Nutr 2020;150(3):536-45. doi: 10.1093/jn/nxz252.

12. Malik VS, Li Y, Tobias DK, Pan A, Hu FB. Dietary Protein Intake and Risk of Type 2 Diabetes in US Men and Women. Am J Epidemiol 2016;183(8):715-28. doi: 10.1093/aje/kwv268.

13. Tharrey M, Mariotti F, Mashchak A, Barbillon P, Delattre M, Fraser GE. Patterns of plant and animal protein intake are strongly associated with cardiovascular mortality: the Adventist Health Study-2 cohort. Int J Epidemiol 2018. doi: 10.1093/ije/dyy030.

14. Guasch-Ferre M, Satija A, Blondin SA, Janiszewski M, Emlen E, O'Connor LE, Campbell WW, Hu FB, Willett WC, Stampfer MJ. Meta-Analysis of Randomized Controlled Trials of Red Meat Consumption in Comparison With Various Comparison Diets on Cardiovascular Risk Factors. Circulation 2019;139(15):1828-45. doi: 10.1161/CIRCULATIONAHA.118.035225.

15. Farvid MS, Sidahmed E, Spence ND, Mante Angua K, Rosner BA, Barnett JB. Consumption of red meat and processed meat and cancer incidence: a systematic review and meta-analysis of prospective studies. Eur J Epidemiol 2021;36(9):937-51. doi: 10.1007/s10654-021-00741-9.

16. Huang Y, Cao D, Chen Z, Chen B, Li J, Guo J, Dong Q, Liu L, Wei Q. Red and processed meat consumption and cancer outcomes: Umbrella review. Food Chem 2021;356:129697. doi: 10.1016/j.foodchem.2021.129697.

17. Havemeier S, Erickson J, Slavin J. Dietary guidance for pulses: the challenge and opportunity to be part of both the vegetable and protein food groups. Ann N Y Acad Sci 2017;1392(1):58-66. doi: 10.1111/nyas.13308.

18. Comerford KB, Miller GD, Reinhardt Kapsak W, Brown KA. The Complementary Roles for Plant-Source and Animal-Source Foods in Sustainable Healthy Diets. Nutrients 2021;13(10):3469. doi: 10.3390/nu13103469.

19. Clark MA, Springmann M, Hill J, Tilman D. Multiple health and environmental impacts of foods. Proc Natl Acad Sci U S A 2019. doi: 10.1073/pnas.1906908116.

20. Katz DL, Doughty KN, Geagan K, Jenkins DA, Gardner CD. Perspective: The Public Health Case for Modernizing the Definition of Protein Quality. Adv Nutr 2019;10(5):755-64. doi: 10.1093/advances/nmz023.

21. Xu XM, Sharma P, Shu SJ, Lin TS, Ciais P, Tubiello FN, Smith P, Campbell N, Jain AK. Global greenhouse gas emissions from animal-based foods are twice those of plant-based foods. Nat Food 2021;2(9):724-+. doi: 10.1038/s43016-021-00358-x.

22. Jarmul S, Dangour AD, Green R, Liew Z, Haines A, Scheelbeek PFD. Climate change mitigation through dietary change: a systematic review of empirical and modelling studies on the environmental footprints and health effects of 'sustainable diets'. Envir Res Lett 2020;15(12):123014. doi: 10.1088/1748-9326/abc2f7.

23. Reinhardt SL, Boehm R, Blackstone NT, El-Abbadi NH, McNally Brandow JS, Taylor SF, DeLonge MS. Systematic Review of Dietary Patterns and Sustainability in the United States. Adv Nutr 2020. doi: 10.1093/advances/nmaa026.

24. Kesse-Guyot E, Rebouillat P, Brunin J, Langevin B, Alles B, Touvier M, Hercberg S, Fouillet H, Huneau JF, Mariotti F, et al. Environmental and nutritional analysis of the EAT-Lancet diet at the individual level: insights from the NutriNet-Sante study. Journal of Cleaner Production 2021;296:10. doi: 10.1016/j.jclepro.2021.126555.

25. Willett W, Rockstrom J, Loken B, Springmann M, Lang T, Vermeulen S, Garnett T, Tilman D, DeClerck F, Wood A, et al. Food in the Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems. Lancet 2019;393(10170):447-92. doi: 10.1016/S0140-6736(18)31788-4.

26. Tuomisto HL. Importance of considering environmental sustainability in dietary guidelines. Lancet Planet Health 2018;2(8):e331-e2. doi: 10.1016/S2542-5196(18)30174-8.

27. Dubuisson C, Dufour A, Carrillo S, Drouillet-Pinard P, Havard S, Volatier JL. The Third French Individual and National Food Consumption (INCA3) Survey 2014-2015: method, design and participation rate in the framework of a European harmonization process. Public Health Nutr 2019;22(4):584-600. doi: 10.1017/S1368980018002896.

28. The French Agency for Food Environmental and Occupational Health & Safety (ANSES). ANSES-CIQUAL French food composition table. Version 2016. Internet: https://ciqual.anses.fr/.

29. Dussiot A, Fouillet H, Wang J, Salome M, Huneau JF, Kesse-Guyot E, Mariotti F. Modeled healthy eating patterns are largely constrained by currently estimated requirements for bioavailable iron and zinc - a diet optimization study in French Adults. Am J Clin Nutr 2021. doi: 10.1093/ajcn/nqab373.

30. GBD Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet 2020;396(10258):1223-49. doi: 10.1016/S0140-6736(20)30752-2.

31. Mariotti F, Havard S, Morise A, Nadaud P, Sirot V, Wetzler S, Margaritis I. Perspective: Modeling Healthy Eating Patterns for Food-Based Dietary Guidelines-Scientific Concepts, Methodological Processes, Limitations, and Lessons. Adv Nutr 2021;12(3):590-9. doi: 10.1093/advances/nmaa176.

32. The French Agency for Food Environmental and Occupational Health & Safety (ANSES). Avis de l'ANSES relatif à «⬛Actualisation des références nutritionnelles françaises en vitamines et minéraux. Saisine n°2018-SA-0238, saisine liée n°2012-SA-0103. 2021. Internet: https://www.anses.fr/fr/system/files/NUT2018SA0238Ra.pdf.

33. Armah SM, Carriquiry A, Sullivan D, Cook JD, Reddy MB. A complete diet-based algorithm for predicting nonheme iron absorption in adults. J Nutr 2013;143(7):1136-40. doi: 10.3945/jn.112.169904.

34. Hallberg L, Hulthen L. Prediction of dietary iron absorption: an algorithm for calculating absorption and bioavailability of dietary iron. Am J Clin Nutr 2000;71(5):1147-60. doi: 10.1093/ajcn/71.5.1147.

35. Miller LV, Krebs NF, Hambidge KM. Mathematical model of zinc absorption: effects of dietary calcium, protein and iron on zinc absorption. Br J Nutr 2013;109(4):695-700. doi: 10.1017/S000711451200195X.

36. de Gavelle E, Huneau JF, Mariotti F. Patterns of Protein Food Intake Are Associated with Nutrient Adequacy in the General French Adult Population. Nutrients 2018;10(2). doi: 10.3390/nu10020226.

37. de Gavelle E, Huneau JF, Bianchi CM, Verger EO, Mariotti F. Protein Adequacy Is Primarily a Matter of Protein Quantity, Not Quality: Modeling an Increase in Plant:Animal Protein Ratio in French Adults. Nutrients 2017;9(12). doi: 10.3390/nu9121333.

38. Mausser H, Grodzevich O, Romanko O. Normalization and other topics in multi-objective optimization. Proceedings of the Fields–MITACS Industrial Problems Workshop, 2006:89-101.

39. ADEME. Agribalyse data v3.0. 2020. Internet: https://doc.agribalyse.fr/documentation-en/.

40. Colomb V, Amar SA, Mens CB, Gac A, Gaillard G, Koch P, Mousset J, Salou T, Tailleur A, van der Werf HMG. AGRIBALYSE (R), the French LCI Database for agricultural products: high quality data for producers and environmental labelling. Ocl-Oilseeds and Fats Crops and Lipids 2015;22(1). doi: 10.1051/ocl/20140047.

41. Koch P, Salou T. AGRIBALYSE®: Rapport Méthodologique - Volet Agriculture - Version 3.0. ADEME. 2020:p. 384. Internet: www.ademe.fr/agribalyse-en.

42. European Commission. Guidance for the development of Product Environmental Footprint Category Rules (PEFCRs). version 6.3, 2018. Internet: https://ec.europa.eu/environment/eussd/smgp/pdf/PEFCR_guidance_v6.3.pdf

43. Bakaloudi DR, Halloran A, Rippin HL, Oikonomidou AC, Dardavesis TI, Williams J, Wickramasinghe K, Breda J, Chourdakis M. Intake and adequacy of the vegan diet. A systematic review of the evidence. Clin Nutr 2021;40(5):3503-21. doi: 10.1016/j.clnu.2020.11.035.

44. Springmann M, Wiebe K, Mason-D'Croz D, Sulser TB, Rayner M, Scarborough P. Health and nutritional aspects of sustainable diet strategies and their association with environmental impacts: a global modelling analysis with country-level detail. Lancet Planet Health 2018;2(10):e451-e61. doi: 10.1016/S2542-5196(18)30206-7.

45. Cifelli CJ, Houchins JA, Demmer E, Fulgoni VL. Increasing Plant Based Foods or Dairy Foods Differentially Affects Nutrient Intakes: Dietary Scenarios Using NHANES 2007-2010. Nutrients 2016;8(7). doi: 10.3390/nu8070422.

46. Mangano KM, Tucker KL. Bone Health and Vegan Diets. Edtion ed. In: Mariotti F, ed. Vegetarian and Plant-Based Diets in Health and Disease Prevention: Academic Press, 2017:315-31.

47. Thorpe DL, Beeson WL, Knutsen R, Fraser GE, Knutsen SF. Dietary patterns and hip fracture in the Adventist Health Study 2: combined vitamin D and calcium supplementation mitigate increased hip fracture risk among vegans. Am J Clin Nutr 2021. doi: 10.1093/ajcn/nqab095.

48. Craig WJ, Mangels AR, Fresan U, Marsh K, Miles FL, Saunders AV, Haddad EH, Heskey CE, Johnston P, Larson-Meyer E, et al. The Safe and Effective Use of Plant-Based Diets with Guidelines for Health Professionals. Nutrients 2021;13(11). doi: 10.3390/nu13114144.

49. Herforth A, Arimond M, Alvarez-Sanchez C, Coates J, Christianson K, Muehlhoff E. A Global Review of Food-Based Dietary Guidelines. Adv Nutr 2019. doi: 10.1093/advances/nmy130.

50. Halkjaer J, Olsen A, Bjerregaard LJ, Deharveng G, Tjonneland A, Welch AA, Crowe FL, Wirfalt E, Hellstrom V, Niravong M, et al. Intake of total, animal and plant proteins, and their food sources in 10 countries in the European Prospective Investigation into Cancer and Nutrition. Eur J Clin Nutr 2009;63 Suppl 4:S16-36. doi: 10.1038/ejcn.2009.73.

51. Mariotti F, Gardner CD. Dietary Protein and Amino Acids in Vegetarian Diets-A Review. Nutrients 2019;11(11). doi: 10.3390/nu11112661.

52. Young VR, Pellett PL. Plant proteins in relation to human protein and amino acid nutrition. Am J Clin Nutr 1994;59(5 Suppl):1203S-12S. doi: 10.1093/ajcn/59.5.1203S.

53. Gazan R, Brouzes CMC, Vieux F, Maillot M, Lluch A, Darmon N. Mathematical Optimization to Explore Tomorrow's Sustainable Diets: A Narrative Review. Adv Nutr 2018;9(5):602-16. doi: 10.1093/advances/nmy049.

54. Macdiarmid JI, Kyle J, Horgan GW, Loe J, Fyfe C, Johnstone A, McNeill G. Sustainable diets for the future: Can we contribute to reducing greenhouse gas emissions by eating a healthy diet? Am J Clin Nutr 2012;96(3):632-9. doi: 10.3945/ajcn.112.038729.

55. Seconda L, Fouillet H, Huneau JF, Pointereau P, Baudry J, Langevin B, Lairon D, Alles B, Touvier M, Hercberg S, et al. Conservative to disruptive diets for optimizing nutrition, environmental impacts and cost in French adults from the NutriNet-Sante cohort. Nat Food 2021;2(3):174-82. doi: 10.1038/s43016-021-00227-7.

56. Webb D, Byrd-Bredbenner C. Overcoming Consumer Inertia to Dietary Guidance. Adv Nutr 2015;6(4):391-6. doi: 10.3945/an.115.008441.

57. Petersen KS, Flock MR, Richter CK, Mukherjea R, Slavin JL, Kris-Etherton PM. Healthy Dietary Patterns for Preventing Cardiometabolic Disease: The Role of Plant-Based Foods and Animal Products. Curr Dev Nutr 2017;1(12):cdn.117.001289-cdn.117. doi: 10.3945/cdn.117.001289.

58. Springmann M, Godfray HC, Rayner M, Scarborough P. Analysis and valuation of the health and climate change cobenefits of dietary change. Proc Natl Acad Sci U S A 2016;113(15):4146-51. doi: 10.1073/pnas.1523119113.

59. Wilson N, Cleghorn CL, Cobiac LJ, Mizdrak A, Nghiem N. Achieving Healthy and Sustainable Diets: A Review of the Results of Recent Mathematical Optimization Studies. Adv Nutr 2019;10(Suppl_4):S389-S403. doi: 10.1093/advances/nmz037.

60. Kesse-Guyot E, Chaltiel D, Wang J, Pointereau P, Langevin B, Allès B, Rebouillat P, Lairon D, Vidal R, Mariotti F, et al. Sustainability analysis of French dietary guidelines using multiple criteria. Nat Sustain 2020;3(5):377-85. doi: 10.1038/s41893-020-0495-8.

61. Segovia-Siapco G, Sabate J. Health and sustainability outcomes of vegetarian dietary patterns: a revisit of the EPIC-Oxford and the Adventist Health Study-2 cohorts. Eur J Clin Nutr 2018. doi: 10.1038/s41430-018-0310-z.

62. Baudry J, Pointereau P, Seconda L, Vidal R, Taupier-Letage B, Langevin B, Allès B, Galan P, Hercberg S, Amiot MJ, et al. Improvement of diet sustainability with increased level of organic food in the diet: findings from the BioNutriNet cohort. Am J Clin Nutr 2019;109(4):1173-88. doi: 10.1093/ajcn/nqy361.

63. Nelson ME, Hamm MW, Hu FB, Abrams SA, Griffin TS. Alignment of Healthy Dietary Patterns and Environmental Sustainability: A Systematic Review. Adv Nutr 2016;7(6):1005-25. doi: 10.3945/an.116.012567.

64. Crippa M, Solazzo E, Guizzardi D, Monforti-Ferrario F, Tubiello FN, Leip A. Food systems are responsible for a third of global anthropogenic GHG emissions. Nat Food 2021;2(3):198-209. doi: 10.1038/s43016-021-00225-9.

65. Poore J, Nemecek T. Reducing food's environmental impacts through producers and consumers. Science 2018;360(6392):987-92. doi: 10.1126/science.aaq0216.

66. Godfray HCJ, Aveyard P, Garnett T, Hall JW, Key TJ, Lorimer J, Pierrehumbert RT, Scarborough P, Springmann M, Jebb SA. Meat consumption, health, and the environment. Science 2018;361(6399). doi: 10.1126/science.aam5324.

67. Wickramasinghe K, Breda J, Berdzuli N, Rippin H, Farrand C, Halloran A. The shift to plant-based diets: are we missing the point? Global Food Security 2021;29:100530. doi: 10.1016/j.gfs.2021.100530.

68. Reganold JP, Wachter JM. Organic agriculture in the twenty-first century. Nat Plants 2016;2(2). doi: 10.1038/NPLANTS.2015.221.

69. Nijdam D, Rood T, Westhoek H. The price of protein: Review of land use and carbon footprints from life cycle assessments of animal food products and their substitutes. Food Policy 2012;37(6):760-70. doi: 10.1016/j.foodpol.2012.08.002.

Table 1. Range of adequate values of the percentage of plant protein in the diet (%PP) and corresponding minimal values of long-term health risk (*HR*) in French males and females.

	Males							Females					
	Observed diet	Modeled diets						Observed diet	Modeled diets				
%PP	33%	16%	20%	25%-70%	75%	80%	82%	34%	16%	20%	25%-70%	75%	77%
<u>*HR* value</u>	0.983	0.602	0.180	0.000	0.004	0.024	0.049	0.736	0.516	0.065	0.000	0.039	0.052
<u>*HR* components:</u>													
Risk of excessive intake of unhealthy foods:													
Red meat	0.236	0.211	0.065	0.000	0.000	0.000	0.023	0.207	0.347	0.000	0.000	0.000	0.000
Processed meat	0.213	0.021	0.000	0.000	0.000	0.000	0.000	0.126	0.000	0.000	0.000	0.000	0.000
Sweetened beverages	0.026	0.000	0.000	0.000	0.000	0.000	0.000	0.024	0.000	0.000	0.000	0.002	0.013
Risk of insufficient intake of healthy foods:													
Whole grain products	0.213	0.195	0.000	0.000	0.000	0.000	0.000	0.157	0.090	0.000	0.000	0.000	0.000
Legumes	0.111	0.127	0.067	0.000	0.000	0.000	0.000	0.053	0.056	0.042	0.000	0.000	0.000
Fruits	0.091	0.000	0.000	0.000	0.000	0.000	0.000	0.093	0.000	0.000	0.000	0.000	0.000
Vegetables	0.033	0.000	0.000	0.000	0.000	0.000	0.000	0.021	0.000	0.000	0.000	0.000	0.000
Nuts & seeds	0.039	0.048	0.048	0.000	0.000	0.000	0.000	0.019	0.023	0.023	0.000	0.000	0.000
Milk	0.022	0.000	0.000	0.000	0.004	0.024	0.026	0.037	0.000	0.000	0.000	0.037	0.039

Values of the *HR* criterion and its individual components when optimizing *HR* under all the nutritional, dietary and acceptability constraints and an additional constraint imposing the %PP value according to an iterative grid search. Results are reported for the %PP values allowing model convergence (i.e., nutrient-adequate diets) for each sex (16%-82% and 16%-77% %PP ranges in males and females, respectively). Within these %PP values, those outside the 25%-70% range have non-null *HR* values because of excessive intake of unhealthy foods and/or insufficient intake of healthy foods.

Table 2. Dual values of the active constraints identified during minimization of the long-term health risk (*HR*) in French males and females[1].

	Males						Females				
	Modeled diets						Modeled diets				
%PP	16%	20%	25%-70%	75%	80%	82%	16%	20%	25%-70%	75%	77%
%PP constraint[2]	-0.246	-0.081	NS	0.003	0.004	0.037	-0.166	-0.042	NS	0.004	0.017
Nutrients that are more limiting as %PP increase[3]											
Iodine	NS	NS	NS	0,002	0,005	0,058	NS	NS	NS	0.006	0.043
Sodium	NS	-0.023	NS	-0,002	-0,005	-0,041	NS	NS	NS	-0.006	-0.027
Bioavailable iron	NS	NS	NS	NS	NS	NS	NS	0.096	NS	0.014	0.027
Vitamin B12	NS	NS	NS	0.002	0.004	0.035	NS	NS	NS	NS	NS
Vitamin B2	NS	NS	NS	NS	0.001	0.006	NS	NS	NS	NS	0.021
Calcium	NS	NS	NS	NS	0.001	0.009	NS	NS	NS	0.001	0.013
EPA+DHA	NS	NS	NS	0.001	0.001	0.007	NS	NS	NS	0.002	0.009
Vitamin A	NS	NS	NS	0.001	0.002	0.009	NS	NS	NS	0.004	0.001
⍺-linolenic acid	NS	NS	NS	0.001	0.001	0.001	NS	NS	NS	0.001	0.004
Nutrients that are more limiting as %PP decrease[3]											
Fiber	0.435	0.136	NS	NS	NS	NS	0.332	0.077	NS	NS	NS
Sugar excluding lactose	-0.125	-0.044	NS	NS	NS	-0.012	-0.106	-0.030	NS	NS	-0.002
Saturated fatty acids	-0.095	-0.074	NS	NS	NS	NS	-0.085	-0.036	NS	NS	NS
Atherogenic fatty acids	-0.083	-0.001	NS	NS	NS	NS	-0.094	-0.006	NS	NS	NS

[1]Dual values when optimizing *HR* under all the nutritional, dietary and acceptability constraints and an equality constraint imposing the percentage of plant protein in the diet (%PP) value according to an iterative grid search. Results are reported for the %PP values allowing nutrient-adequate diets (16%-82% and 16%-77% %PP ranges in males and females, respectively), which includes those also allowing a null *HR* value (25%-70% %PP range in both sexes). NS, not significant (<0.0001).

[2]For the %PP equality constraint, dual values represent the potential effect on *HR* of the relaxation by one unit of the limiting bound, with positive (negative) values if the lower (upper) bound is limiting (e.g., for %PP=82% in males, the dual value indicates that there would be a potential *HR* gain of 0.037 if %PP was decreased from 82% to 81%).

[3]For nutritional constraints, dual values have been standardized to represent the potential effect on *HR* of the relaxation by 10% of the limiting bound, to classify the nutrients from the most limiting (higher absolute value) to the least limiting (lowest absolute value). Limiting nutrients have a positive (negative) dual value if their lower (upper) bound is limiting (e.g., for %PP=82% in males, the dual value for iodine indicates that there would be a potential *HR* gain of 0.058 if the lower bound for iodine intake was decreased by 10%, from 150 to 135 μg/d). Only nutrients with an active constraint (i.e., with a non-null dual value) for at least one %PP value are presented here. For nutrients not presented in this table, dual values were always equal to zero, meaning that compliance with these constraints was not limiting.

Atherogenic fatty acids, lauric and myristic and palmitic acids; DHA, docosahexaenoic acid; EPA, eicosapentaenoic acid.

Figure legends

Figure 1. Daily food category consumption in the observed diets (obs) and modeled diets obtained by long-term health risk (*HR*) and diet departure (*DD*) minimization under imposed percentage of plant protein in the diet (%PP) in French adults. Results are reported for all the adequate %PP values ensuring nutrient adequacy (16%-77%), which includes those also ensuring a null *HR* value (25%-70%). The bar charts represent the cumulative consumptions of food categories (black axis on the left) and the curves represent the *HR* and *DD* values (blue and pink axes on the right, respectively). For clarity, the 45 modeled food groups are not represented here but grouped into broader categories that are included in *HR* (such as red and processed meats) or represent other protein sources (such as poultry and seafood). Consumption of water, hot beverages, alcohol and miscellaneous foods are not shown for clarity. Details about food grouping and consumptions of food categories not shown here are given in Supplemental Tables 1 and 6, respectively.

Figure 2. Contribution of food categories to protein intake in the observed diets (obs) and modeled diets obtained by long-term health risk (*HR*) and diet departure minimization under imposed percentage of plant protein in the diet (%PP) in French adults. Results are reported for all the adequate %PP values ensuring nutrient adequacy (16%-77%), which includes those ensuring also a null *HR* value (25%-70%). Sections inside the bars represent the contributions of food categories to protein intake (in g of protein/kg of BW/d). See Supplemental Table 1 for the detailed composition of food categories.

Figure 3. Greenhouse gas emissions (GHGe) associated with the observed (obs) and modeled diets obtained by long-term health risk (*HR*) and diet departure minimization under imposed percentage of plant protein in the diet (%PP) in French adults. Results are reported for all the adequate %PP values ensuring nutrient adequacy (16%-77%), which includes those ensuring also a null *HR* value (25%-70%). Sections inside the bars represent the contributions of food categories to GHGe (in kg CO_2 eq), and values above the bars represent the relative deviation in GHGe from its observed value (in %). See Supplemental Table 1 for the detailed composition of food categories.

Figure 1

Figure 2

Figure 3