medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

TITLE PAGE

The trends in the uninsured rates among non-elderly adult patients hospitalized with acute pancreatitis in the US during 2004-2019.

Kumar, Vivek MD 1; Barkoudah, Ebrahim MD MPH 1; Jin, David X MD MPH 2; Banks, Peter 2 MD; McNabb-Baltar, Julia MD MPH 2*.

1Department of General Internal Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA

2Division of Gastroenterology, Hepatology, and Endoscopy, Center for Pancreatic Disease Brigham and Women's Hospital, Boston, Massachusetts, USA

***Corresponding Author**

Correspondence

Dr. Julia McNabb-Baltar MD MPH

Division of Gastroenterology, Hepatology, and Endoscopy, Center for Pancreatic Disease, Brigham and Women's Hospital, 75 Francis Street, Boston, Massachusetts, USA 02115

Assistant Professor Harvard Medical School, Boston, Massachusetts, USA.

Email: Jmcnabb-baltar@bwh.harvard.edu

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Disclosures:

Guarantor of the article: Kumar, Vivek & McNabb-Baltar, Julia

Specific author contributions:

Kumar, Vivek: Study concept and design; acquisition of data; analysis and interpretation of data; drafting of the manuscript

Barkoudah, Ebrahim: Interpretation of data, Critical revision of the manuscript for important intellectual content

Jin, David X: Critical revision of the manuscript for important intellectual content

Banks, Peter: Critical revision of the manuscript for important intellectual content

McNabb-Baltar, Julia: Study concept and design; drafting of the manuscript; critical revision of the manuscript for important intellectual content; study supervision

Financial support: None

Potential competing interests

Kumar, Vivek: None

Barkoudah, Ebrahim: Dr. Barkoudah reports research support payments from National Institutes of Health/National Heart, Lung, and Blood Institute, Bristol Myers Squibb and Janssen, payments made to Brigham and Women's Hospital for performing clinical endpoints sponsored by various entities and Advisory Board fees from Bristol Myers Squibb, Janssen, Novartis, Pfizer and Portola, and travel expenses from Alexion.

Jin, David X: None

Banks, Peter: None

McNabb-Baltar, Julia: None

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Abstract

Background and Aim:
The impact of health reforms such as affordable care act (ACA) and Medicaid expansion program on health insurance coverage among acute pancreatitis (AP) patients in the United States (US) is unknown. We report the trends and forecasts for the uninsured rates among acute pancreatitis (AP) patients in the US.

Methods: We included non-elderly adult patients (aged 18 -64 years), hospitalized with AP in the nationwide inpatient sample database years 2004- 2019. We calculated the percentage of uninsured and Medicaid patients for each year and applied joinpoint (JP) regression model to study the trends.

Results The uninsured rates among patients hospitalized with AP were almost twice compared to the national average rates for all hospitalized patients. Uninsured rates were higher among Hispanic and African American races, rural location, lower income quartiles and in the southern US regions. A statistically significant decline was observed during 2013-2016 with an APC of -12.23 (-20- -4); ***p<0.01***). The decline was apparent among all racial groups, locations and income groups but not in the southern region where gap in the uninsured rates worsened compared to other geographical regions. The trends reversed more recently, and the uninsured rates surged in 2018 for the first time after 2010. In 2019, 12.5% AP patients were uninsured compared to 11.6% in 2017. The forecasts after taking unemployment rates into account showed that uninsured rates would peak in 2020 followed by a gentle decline in the following years but overall uninsured rates would remain higher compared to 2017.

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Conclusions: The ACA and Medicaid expansion programs resulted in overall decrease in the uninsured rates, particularly among racial minorities, rural location and in the lowest income group. However, there was a surge in uninsured rate in the most recent years (2017-2019) which will continue during the next four years

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Acute pancreatitis (AP) is one of the leading gastrointestinal causes of hospitalizations in the United States (US) with its incidence increasing[1]. In the year 2015, AP led to more than one million hospitalization days in the US, resulting in more than $2.5 billion in hospitalization charges[2]. A previous study has reported a higher incidence of uninsured patients in the year 2009-2012 compared to 2002-2005[1]. So far, no study has reported the trends in the rates of uninsured patients among AP patients. The lack of insurance has been associated with poor access to healthcare and subsequent adverse outcomes[3]. In the US, the number of uninsured patients decreased after health reforms such as the Affordable Care Act (ACA) and Medicaid expansion programs[4]. However, the impact of ACA among patients with AP is still unknown. The non-elderly (aged 18-64 years old) AP patients in the US are unique due to higher rates of alcohol and substance use, lower income groups, and a higher fraction of minority races; all these factors are also associated with a lack of health insurance in the US[5]. In this study, we aim to analyze the trends in the uninsured rates among patients hospitalized with a principal diagnosis of AP in the last 15 years, emphasizing the most recent trend changes. We examined the disparities in the health coverage among AP patients by geographical regions, race, rural/urban status, and income levels. We also project the uninsured rates among these patients for the next 4 years after adjusting for unemployment rates.

For this study, we included non-elderly adult patients, hospitalized with AP in the nationwide inpatient sample (NIS) database for the years 2004- 2019[12]. We collected data on uninsured and Medicaid patients, race, rural/urban residence, income, and geographical regions for each year. We calculated the percentage of uninsured and Medicaid patients for an individual year and applied Joinpoint (JP) regression model to study the trends[6]. This program enables us to test whether an apparent change in the trend is statistically significant by using Monte Carlo Permutation method

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

as the test of significance. The location of JP was not known prior and was estimated by the program. Annual percent change (APC) and average annual percent change (AAPC) were calculated with *'p'* values to test if apparent changes in the trends were statistically significant. For forecasting, we applied vector autoregression (VAR) model, a multivariate time series model, and adjusted uninsured rates with the unemployment rates in the US. The unemployment rates were obtained from the US Bureau of Labor Statistics of the US department of labor [7]. Statistical analysis was conducted using National Cancer Institute's Joinpoint regression software and 'R'.

The uninsured rates among patients hospitalized with AP were almost twice the national average rates for all hospitalized patients (Fig 1a). Uninsured rates were higher among Hispanic and African American races, rural locations, lower-income quartiles, and southern US regions (Table 1). The trends in the uninsured rates among AP patients mirrored the changes in the national rates, but the gap persisted (Fig 1a). Overall, the change in the uninsured rate from 2004 to 2019 was not statistically significant with an AAPC of -1 (-5.8-4.1); $p=0.7$). But a statistically significant decline was observed during 2013-2016 with an APC of -12.23 (-20- -4); $\boldsymbol{p<0.01}$). A simultaneous increase in Medicaid coverage was also noticed (fig 1b). The decline was apparent among all racial groups, locations, income groups and geographical regions, however uninsured rates in the southern region remained considerably higher (Table 1). Unfortunately, trends reversed more recently, and the uninsured rates surged in 2017 for the first time after 2010. In 2019, 12.5% AP patients were uninsured compared to 11.7% in 2017. After adjusting for unemployment rates, the uninsured rates are anticipated to increase drastically in the year 2020 to 16.04% (12.8-19.3), followed by some decline in the following years 13.9(10.7-17.13) in 2021, 13.97 (10.7-17.2) in 2022 and 13.5(10.3-16.8) in 2023.

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

The higher uninsured rates among AP patients are due to these patients' unique characteristics, as mentioned previously. On Joinpoint regression, we noted several periods with changes in the uninsured rates. The most prominent was a decline in the years 2013-2016. In the US, a major factor affecting the lack of insurance among the non-elderly population is unemployment[8]. While the pattern in the uninsured rates during the initial years of the study was parallel to the unemployment rates, the decline during the years 2013-2016 was out of proportion to what could be explained by declining unemployment rates alone. In fact, this corresponded to the rollout of ACA and Medicaid expansion in the US. By 2016, 33 out of 50 states had implemented Medicaid expansion. The Medicaid expansion was lower in the southern region, where only 7 out of 16 states had implemented Medicaid expansion by the year 2016[9]. This observation also explains the absence of a sharp decline among uninsured patients in the southern region, unlike other regions in the US. The higher rates of uninsured among Hispanics and African American is due to higher rates of unemployment and poverty, which is consistent with the overall rates of uninsured in the USA[10].

Unfortunately, we noted a rising trend in the uninsured in the years 2017-2019. This trend was despite the declining rates of unemployment in the US during this time and is most likely the result of several changes made in the year 2017 to roll back ACA. These changes included waiving or delaying the features of ACA that imposed financial or regulatory burden, reduction in the funding for ACA outreach and education program. Besides, there was also discontinuation of cost-sharing reduction to insurers which led to the loss of incentives to purchase the coverage and increased the uncertainty among the eligible individuals resulting in loss of health insurance coverage [11].

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

To forecast the uninsured rates for the next three years, we adjusted for the unemployment rates. Due to the pandemic, the unemployment rates increased drastically in the US in 2020 to 6.7, almost double of 3.6 in 2019. As unemployment is a major driving factor for insurance among the non-elderly population in the US, we anticipated a drastic rise in the uninsured rates, which was found to be true. The projected rates are similar to the rates prior to implementing ACA reforms which implies that we will probably lose most of the gains in terms of health coverage that were achieved due to ACA if there are no significant changes in the policy.

The limitations of this study include the use of an administrative database using ICD-9 and ICD-10 codes which is subject to misclassification bias. Besides unemployment other factors such as patient preference may also affect the uninsured rates, which could not be analyzed due to the limitations of the database. Nevertheless, we conclude that the ACA and Medicaid expansion programs positively impacted the health insurance coverage among patients hospitalized with AP, specifically on racial minorities and low-income groups. The impact was less apparent in the southern US, where many states did not implement these reforms. Despite the strong US economy and declining unemployment rates during 2017-2019, there was a surge in uninsured patients, likely due to the changes made in the Medicaid policies. This surge is projected to continue for the next few years. Therefore, new policies and reforms are warranted for the communities at the highest risk of losing coverage-ethnic minorities and lower-income groups.

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

1. Krishna SG, Kamboj AK, Hart PA, Hinton A, Conwell DL. The changing epidemiology of acute pancreatitis hospitalizations: a decade of trends and the impact of chronic pancreatitis. *Pancreas*. 2017;46(4):482.

2. Peery AF, Crockett SD, Murphy CC, et al. Burden and cost of gastrointestinal, liver, and pancreatic diseases in the United States: update 2018. *Gastroenterology*. 2019;156(1):254-272. e11.

3. Michael McWilliams J. Health consequences of uninsurance among adults in the United States: recent evidence and implications. *The Milbank Quarterly*. 2009;87(2):443-494.

4. Guth M, Garfield R, Rudowitz R. The effects of Medicaid expansion under the ACA: Studies from January 2014 to January 2020. *Kaiser Family Foundation*. 2020;

5. Substance Abuse and Mental Health Services Administration CfBHSaQF, 2014). The TEDS Report: Health Insurance Status of Adult Substance Abuse Treatment Admissions Aged 26 or Older: 2011. Rockville, MD. https://www.samhsa.gov/data/sites/default/files/sr134-health-insurance-2014/sr134-health-insurance-2014/sr134-health-insurance-2014.htm

6. Joinpoint Regression Program, Version 4.9.0.1 - February 2022; Statistical Methodology and Applications Branch, Surveillance Research Program, National Cancer Institute. https://surveillance.cancer.gov/joinpoint/

7. US BUREAU OF LABOR STATISTICS https://data.bls.gov/timeseries/LNS14000000

8. Broaddus M. *Research note: Medicaid enrollment decline among adults and children too large to be explained by falling unemployment*. Center on Budget and Policy Priorities.; 2019.

9. Courtemanche C, Fazlul I, Marton J, Ukert B, Yelowitz A, Zapata D. The Affordable Care Act's Coverage Impacts in the Trump Era. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*. 2021;58:00469580211042973.

10. Gonzales S, Sommers BD. Intra-ethnic coverage disparities among Latinos and the effects of health reform. *Health services research*. 2018;53(3):1373-1386.

11. Keith K. Uninsured rate rose in 2018, says Census Bureau report. *Health Aff*. 2019;

12. HCUP National Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2004-2019. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup us.ahrq.gov/nisoverview.jsp

medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Fig 1. Trends in the rates of uninsured patients hospitalized with acute pancreatitis

1a. Overall, among all hospitalized patients vs AP patients. 1b. Uninsured and Medicaid insurance among patients with acute pancreatitis

Fig. 1a



Fig. 1b



medRxiv preprint doi: https://doi.org/10.1101/2022.05.20.22275404; this version posted June 2, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

Fig 2. Trends in the rates of uninsured patients hospitalized with acute pancreatitis 2a. geographical regions and income groups 2b. stratified by race, and rural-urban location.

Fig 2a.



Fig 2b.