Small RNA sequencing reveals snoRNAs and piRNA-019825 as novel players in diabetic kidney disease

LM 't Hart[1,2,3,4] (ORCID 0000-0003-4401-2938), JA de Klerk[1,5,6], GA Bouland[1], JHD Peerlings[1], M.T. Blom[3,7], SJ Cramer[1], R Bijkerk[5,6], JWJ Beulens[2,3], RC Slieker[1]

[1] Department of Cell and Chemical Biology, Leiden University Medical Center, Leiden, The Netherlands

[2] Department of Epidemiology and Data Sciences, Amsterdam University Medical Center, Location Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[3] Health Behaviors & Chronic Diseases Research Program & Personalised Medicine Research Program, Amsterdam Public Health, Amsterdam, The Netherlands

[4] Department of Biomedical Data Sciences, Section Molecular Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

[5] Department of Internal Medicine, Division of Nephrology, Leiden University Medical Center, Leiden, The Netherlands

[6] Einthoven Laboratory for Vascular and Regenerative Medicine, Leiden University Medical Center, Leiden, The Netherlands

[7] Department of General Practice, Amsterdam University Medical Center, Location Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

Author for correspondence:

LM 't Hart, PhD

Leiden University Medical Center

Department of Cell and Chemical Biology

Albinusdreef 2

2333ZA Leiden, The Netherlands

+31 715269796

lmthart@lumc.nl

Abstract

Introduction Micro- and macrovascular complications are common among persons with type 2 diabetes. Recently there has been growing interest to investigate the potential of circulating small non-coding RNAs (sncRNAs) as contributors to the development of diabetic complications. In this study we investigate to what extent circulating sncRNAs levels associate with prevalent diabetic kidney disease (DKD) in persons with type 2 diabetes.

Methods Plasma sncRNAs levels were determined using small RNA-seq, allowing detection of miRNAs, snoRNAs, piRNAs, tRNA-fragments and various other sncRNA classes. We tested for differentially expressed sncRNAs in persons with type 2 diabetes, with DKD (n=69) or without DKD (n=405). In secondary analyses, we also tested the association with eGFR, albuminuria (UACR) and the plasma proteome.

Results In total seven sncRNAs were negatively associated with prevalent DKD (all $P_{FDR} \leq 0.05$). Including one microRNA (miR-143-5p), five snoRNAs (U8, SNORD118, SNORD24, SNORD107, SNORD87) and a piRNA (piR-019825|DQ597218). Proteomic analyses showed that the seven sncRNAs, and especially the piRNA piR-019825, were associated with plasma levels of 24 proteins of which several have known associations with kidney function including TNF sR-I (TNFRFS1A), DAN (NBL1) and cystatin C (CST3).

Conclusion We have identified novel small non-coding RNAs, primarily from classes other than microRNAs, that are associated with diabetic kidney disease. Our results shows that the involvement of small non-coding RNAs in DKD goes beyond the already known microRNAs and also involves other classes of sncRNA, in particular snoRNAs and the piRNA piR-019825, that have never been studied before in relation to kidney function.

Keywords: Type 2 diabetes, small RNA, microRNA, snoRNA, piRNA, Chronic kidney disease, Nephropathy

Introduction

Diabetic kidney disease (DKD) is one of the most common complications in type 2 diabetes.
Nonetheless, while some develop complications early in the disease, others will never. The
mechanisms underlying DKD are not fully understood though [1].

Circulating small non-coding RNAs (sncRNA) are mostly contained in extracellular vesicles
or are bound to RNA-binding proteins and lipoproteins and originate from various tissues including
tissues important in kidney function [2]. Whereas previously it was thought that sncRNAs merely
reflect degradation by-products, it has now been shown that they exert important functions in cell-to-
cell communication, the regulation of gene expression, splicing and ribosomal RNA maturation [2].

Although a substantial number of studies have examined circulating sncRNAs associated with
kidney function in persons with diabetes, most studies are relatively small (n<100) and focused on
microRNAs. From these studies a large number of miRNAs have emerged that are putatively
associated with kidney function. However, probably due to small sample sizes, heterogeneity in
endpoints and methodology used there is limited consistency between studies as shown in various
meta-analyses [3-5].

To the best of our knowledge there are currently no studies that have systematically examined
the other classes (biotypes) of small ncRNA. We thus go beyond the current state-of-the-art by not
only investigating microRNAs but also other sncRNAs classes, including besides miRNAs, piRNAs,
tRNA fragments (tRFs), lncRNA fragments, Y-RNAs and snoRNAs. For this we applied small RNA
sequencing in a large well-defined cohort (n=474) of persons with and without DKD to capture the
whole plasma sncRNA transcriptome allowing for a hypothesis-free, comprehensive analysis of the
role of sncRNAs in diabetic kidney disease [6].

Materials and Methods

A schematic overview of the study design can be found in Supplemental Figure 1.

Study sample

For this project we used data from the Hoorn Diabetes Care System cohort (DCS), a large prospective
study comprised of 14,000+ individuals with type 2 diabetes who receive structed diabetes care [7].
Participants visit the Hoorn diabetes research center annually for diabetes care including routine
biochemistry, medication review and clinical care. Patients were also invited to participate in DCS
biobanks in which we combine the annual phenotypic data with biobanking of liquid biopsies which
includes serum, plasma and urine. Blood samples were collected during two phases of biobanking in
2008/2009 and 2012-2014. For the current study we used a subsample from the whole DCS biobank
(n=475) using the following inclusion criteria: a plasma sample collected within four years after
diagnosis of diabetes and no-insulin use at the time of blood sampling.

All laboratory measurements were on samples taken in a fasted state on the same day as the

sample for RNA-seq. Routine anthropometric, biochemistry methods and definitions have been

described previously in detail in van der Heijden et al. [7]. Kidney function was assessed based on

eGFR (CKD-EPI) and urinary albumin to creatinine ratio (UACR) and included as continuous

variables in our analyses. Stages of chronic kidney disease (CKD) were defined according the KDIGO

criteria as previously described [8]. In short, this definition uses a combination of eGFR and UACR to

categorize persons based on kidney function. Persons in stage 0 do not show signs of kidney

dysfunction whereas the stages 1 to 3 have increasing kidney dysfunction as shown in Supplemental

Figure 2. Cases were defined as having CKD stage $\geq$ 1 (i.e. eGFR <60 ml/min/1.73 m^2 and/or UACR

>3 mg/mmol) at the time of blood sampling for sncRNA analysis which resulted in 69 cases.

Circulating small RNA sequencing

Circulating sncRNAs were isolated from 900ul of citrate plasma using a commercial cell-free RNA

isolation kit (Quick-cfRNA Serum & Plasma Kit, Zymo Research, Irvine, CA, USA). After quality

control 475 samples were submitted for RNA sequencing. Library preps were constructed using the

NEBNext® Multiplex Small RNA Library Prep Set for Illumina (New England Biolabs, Ipswich, MA,

USA). A Pippin Prep instrument (Sage Science, Beverly, MA,USA) was used for size selection to

further enrich for sncRNAs with a size of ~20 to 100 nucleotides. Sequencing was performed using the

Illumina NextSeq500 sequencer with v2.5 sequencing reagent kits and 35 cycle paired-end reads

(PE35). On average 11.3±2.8 million reads per sample were generated. The exceRpt pipeline [9],

developed by the NIH Extracellular RNA Communication Consortium (ERCC), was used to map the

obtained reads, for quality control, to process the data and generate RNA abundance estimates for

miRNA, piRNA, lncRNA, tRNA fragments, Y_RNA (fragments), snRNA, snoRNA, scaRNA and

various other sncRNA species. Using respectively miRBase v22 [10], piRNABank v1 [11], Gencode

(version 38) [12], circBase [13] and GtRNAdb [14]. One sample did not pass the QC thresholds.

Taqman validation of the RNA-seq results

We used Taqman qPCR assays (ThermoFisher Scientific, NL) to verify the validity and

reproducibility of the small RNA-seq results. First cDNA was generated from 5ng of RNA isolated

from plasma using the TaqMan™ Advanced miRNA cDNA Synthesis Kit (ThermoFisher Scientific,

NL) according to the instructions of the manufacturer. We chose four miRNA assays with a range

from low to relative high expression in our study sample (miR-542-3p, miR-323b-3p, miR-186-5p and

miR-423-5p) which were measured in 12 randomly selected plasma RNA samples for which sncRNA-

seq data were available. Clinical characteristics are shown in Supplemental Table 1. The Taqman

qPCR assays were run according to the manufacturer's instructions using TaqMan Fast Advanced

Master mix and TaqMan Advanced miRNA Assays (ThermoFisher Scientific, NL). miR-191-5p

expression was used as the reference (housekeeping gene) in both the Taqman and small RNA-seq

datasets as suggested by the manufacturer [15].

Expression patterns of small RNAs in tissues

Publicly available raw sncRNA-seq data for various metabolic tissues were downloaded from the

Gene Expression Omnibus (GEO) [16]. The tissues for which sncRNA-seq data was used included

kidney, thyroid, brain, liver, muscle, heart, colon, pancreas, subcutaneous white adipose tissue and

whole blood and urine. Accession numbers are given in Supplemental Table 2. Data were processed

using the exceRpt pipeline as described above. The tissue data were used to estimate the abundance of

the sncRNAs in these tissues. Data was z-scaled in figures.

Plasma proteomics.

Plasma proteomic data was available for 589 participants of the Hoorn DCS study and generated using

the SomaLogic SOMAscan platform (Boulder, CO) as previously described [17]. After quality control

1195 protein levels were available for analysis. For 199 subjects we also had the corresponding

sncRNA-seq data from the same sampling date. Clinical characteristics of the participants in this study

are shown in Supplemental Table 1.

Statistical analysis

The R-package edgeR (v3.42.4) was used to identify differentially expressed sncRNAs [18]. A

negative binomial generalized log-linear model was applied using edgeR's function

estimateGLMRobustDisp. For the determination of differential expression and p-values we used the

quasi-likelihood (QL) F-test using edgeR's *glmQLFTest* function according to the recommendations

of the developers. The base model was unadjusted. A fully adjusted model was adjusted for BMI,

diabetes duration, HbA1c, HDL, SBP and smoking unless otherwise stated. Age and sex were

excluded from the model as these are part of the definition of eGFR. UACR was log10-transformed

before analysis. In sensitivity analyses, we additionally included use of antihypertensive medication

and or eGFR or UACR. The correlation between the sncRNA transcriptome and plasma proteome was

tested using a linear model with the log transformed proteomic measurements as outcome and the

sncRNA transcriptome as predictors with adjustment for age, BMI, sex and technical covariates.

Associations between the proteome and eGFR or UACR were tested using the same model. The

Benjamini-Hochberg procedure was used to adjust for multiple hypothesis testing and an FDR below

0.05 was considered significant.

Results

Clinical characteristics

The clinical characteristics of our study group are shown in Table 1. The characteristics resemble

those of a typical T2D population with slightly more males than females, a mean age-at-diagnosis of

59.3 ± 9.0 years, BMI of 30.7 ± 6.0 kg/m2 and an average duration of diabetes at time of blood

sampling of 2.2 ± 1.1 years. Glycemic control was good with an average HbA1c of 46 (42, 51)

mmol/mol. Prevalence of DKD at the time of blood sampling was 15%.

sncRNA biotype abundance in our samples

After quality control, 938 sncRNAs were expressed above the threshold of a mean ≥5 copies

in our data, which included miRNAs, piRNA, Y_RNA fragments, lncRNA fragments,

tRNA_fragments, snoRNAs, snRNAs and various other sncRNA species (Table 2). miRNAs, Y-

RNAs and tRFs were the most abundant sncRNAs biotypes comprising respectively 57.6, 34.9 and 4.8

percent of all reads. Other sncRNA biotypes were much less abundant (≤1.4%). There were no

significant differences in the total number of reads per biotype between cases and controls (all P>0.4).

Taqman validation shows good reproducibility

We next sought to verify validity and reproducibility of the sncRNA-seq. Taqman qPCR

assays for sncRNAs with high, medium and low expression were used and these showed a good

correlation with the RNA-seq results (r^2= 0.86, Supplemental Figure 3).

sncRNAs associate with DKD, eGFR and UACR

Seven sncRNAs were significantly associated with prevalent DKD in the unadjusted model

(P_{FDR}≤0.05, Table 3, Figure 1A). All identified sncRNAs were downregulated in DKD cases. The top

hit was snoRNA U8 (log2 fold change=-0,859, P_{FDR}=1.90x10^{-3}, Figure 1B). Five of these sncRNAs

also showed a significant trend for lower levels in those with more advanced stages of CKD

(P_{FDR}≤0.05, Supplemental Figure 4).

Respectively, 107 and 220 sncRNAs were significantly associated with eGFR and UACR,

which are used for the DKD classification (Table 3, Supplemental Table 3, Supplemental Figures 5

and 6). The overlap between the sncRNAs associated with DKD and its components eGFR and UACR

is shown in figure 1C and Table 3. snoRNA U8 overlapped between all three endpoints used and

piR_019825 overlapped between eGFR and DKD. SNORD118, SNORD24, SNORD107, SNORD87

and miR-143-5p all overlapped between UACR and DKD.

In the full model respectively 7, 153 and 205 sncRNAs remained significantly associated with

DKD, eGFR and UACR (Supplemental Table 4, Supplemental Figure 7; 57%; 95% and 84% overlap

with the unadjusted model). U8, SNORD118, piR_019825 and miR-143-5p were significant in both

the base and full models. In a sensitivity analysis, additional adjustment for use of antihypertensive

medication and baseline eGFR or UACR (only for eGFR and UACR) did not have a major impact on

the effect sizes but reduced the number of significant associations (r^2>=0.92, Supplemental Figure 8).

Certain types of sncRNAs are enriched in DKD

Comparing the frequency of different types of sncRNAs identified showed that while snoRNAs represent 10% of the total small RNA transcriptome in our RNA-seq study they are the most abundant class associated with prevalent DKD (71%, OR=21.9, P=2.4x10^{-4}, Table 2). Enrichment for snoRNAs is also seen for UACR (OR=3.8, p=8.4x10^{-13}). Associations with miRNAs on the other hand are less frequent whilst piRNAs also seem enriched (Table 2). A similar pattern is again seen for UACR and to a lesser extend for eGFR. In the full model a largely similar picture is observed (Supplemental Table 5).

The pairwise correlation in expression between the seven sncRNAs associated with diabetic DKD was very modest (Figure 1D). The only exceptions were U8 and SNORD118, which is likely due to the fact that they are encoded in the same host gene (TMEM107) on chromosome 17. Results of these analyses for eGFR and UACR are shown in Supplemental Figure 9.

sncRNAs are tissue-enriched and correlate with plasma protein levels

In a next step, we examined the tissue expression patterns of the identified sncRNAs. We used publicly available expression data in 11 tissues or body fluids (Fig. 1E). This illustrated that some sncRNAs show tissue-enriched expression patterns. For the seven sncRNAs associated with DKD none had its highest expression in kidney tissue. piRNA-019825 was the only sncRNA that showed high expression in urine. Similar observations were made for eGFR and UACR associated sncRNAs (Supplemental Fig. 10).

Finally, since the function of most of the sncRNAs we found in our study is unknown we compared the expression levels of the sncRNAs with expression of 1195 proteins measured in plasma from the same date to identify potential targets or (co-)regulated proteins or pathways. As shown in Supplemental Table 6 we found for the seven sncRNAs 31 FDR significant pairwise correlations involving 24 proteins. The strongest association was observed between piR-019825 and TNF sR-1 (TNFRSF1A, Beta=-0.080 (-0.112 - -0.040), P$_{FDR}$=3.04x10^{-2}). piRNA-019825 also showed the largest number of correlations (n=11, Figure 2A). Interestingly, among the proteins associated with piR-019825 was Cystatin C, a known marker of kidney function. Next we assessed whether the 24 proteins were also significantly associated with eGFR or UACR (FDR≤0.05). Six were associated with eGFR and none with UACR. Four out of the six proteins that were negatively associated with piR-019825 expression were also negatively associated with eGFR (all P$_{FDR}$≤3.78x10^{-11}) (Figure 2A, Supplemental Table 7). Two proteins were associated with miR-143-5p expression and another eleven proteins were associated with snoRNA expression. Five proteins were correlated with two or more snoRNAs (NAMPT, UFM1, PDGFB, CCL27 and CCL28) and two were also significantly associated with eGFR (CCL27, RARRES2, Figure 2B). Pairwise correlations between the piR-019825 or snoRNA associated proteins is shown in figures 2C, D

Discussion

In this study we show that circulating sncRNAs are associated with prevalent DKD in people with type 2 diabetes. Importantly these sncRNAs do not only represent miRNAs but also other classes of small non-coding RNAs. In addition, these sncRNAs correlate with proteins that function in pathways known to be involved in DKD/kidney function.

Associations between kidney function and sncRNAs

miRNAs

MicroRNAs are involved in the post-transcriptional regulation of gene expression and are known to be involved in crosstalk between (distant) cells and tissues [2]. As such, there is an ever growing interest to study their role in disease development and as biomarkers of disease and disease progression.

In our study one miRNA, miR-143-5p, was associated with both DKD and UACR. Levels of this miRNA are lower in both DKD and persons with high UACR. miR-143-5p remained significant in the full models, suggesting it might have biomarker potential. Previous in vitro studies showed that this miRNA is associated with vascular pathology [19], an impaired glomerular filtration barrier [20] and with mesangial cell proliferation and fibrosis [21, 22]. miR-143-5p plasma levels associate with two proteins but neither of those have a clear relationship with kidney function nor were they predicted targets for miR-143-5p (miRDB, miRTarBase, MiRBase). In the full model three additional microRNAs were significant, miR-455-3p, miR-122-3p and miR-193b-3p. Previously it was shown that miR-455-3p was down regulated in human mesangial cells (HMC) and human proximal tubule epithelial cells (HK-2) under DKD stimulating conditions [23]. Interestingly, it was also shown that overexpression of this miR could alleviate the pathological changes such as renal fibrosis in a rat DN model [23]. In a mouse mesangial cell line miR-455-3p improved cell proliferation, inflammation and ECM accumulation induced by high glucose treatment [24]. Involvement of this miR in fibrosis in other tissues like liver and lung has also been reported. A previous report indicated that miR-122, a liver specific miR associated with liver injury, was reduced in patients with end stage renal disease but not in CKD when compared to healthy controls [25]. miR-193b-3p was identified as a potential biomarker for pre-diabetes and also for CKD after nephrectomy for renal cell carcinoma [26, 27].

In addition to the association with DKD we also found many miRNAs associated with eGFR and or UACR. A large number of which remained significant in the full models. Interestingly, of the 154 miRNAs found in the base models for eGFR and UACR only two of them were associated with both phenotypes suggesting that the associated miRNAs are specific to either defect in kidney function but not both.

piRNAs

PIWI-interacting RNAs are involved in silencing of transposons but they can also be involved
in the regulation of gene expression in a miRNA-like manner [28]. One piRNA, piRNA-019825
(DQ597218), was significantly associated with DKD. Previously, it was shown that this piRNA is one
of the most abundant piRNA species in urine and plasma [29]. In our proteomics analysis to identify
potential targets or pathways we found eleven proteins whose expression was correlated with the
levels of piR-019825 including cystatin C, a known marker for kidney function. Four of the
negatively associated proteins are also strongly associated with eGFR. This is in line with a putative
miRNA like function for piR-019825 in which lower piRNA expression would result in higher protein
levels, lower eGFR and increased risk of DKD. This is further corroborated by the fact that increased
levels of two of these proteins were previously found to increase risk of CKD or end stage renal
disease (ESRD) in prospective studies in persons with diabetes (TNF sR-I (TNFRSF1A), NBL1
(DAN) [30-33]. Whereas the other negatively associated proteins we found associated with this
piRNA (CST3, sCXCL16, CFD, PLA2G2A) have almost all also been implicated previously in kidney
dysfunction [34-36]. Interestingly, we also find various other piRNAs associated with either eGFR and
or UACR (10-14%). Most of which remained significant in the full models. Suggesting that they may
be useful early markers of DKD. Of note, eGFR-associated piRNAs often show their highest
expression in urine, whereas UACR associated piRNAs show low expression in urine, but often high
levels in tissues like pancreas, liver and colon.
294
snoRNAs
Small Nucleolar RNAs (snoRNA) are predominantly localized to the nucleolus where they
have a function in rRNA biogenesis, structure and activity [37]. There are two major types of
snoRNAs containing either box C/D or box H/ACA motifs. In our study we only detected box C/D
snoRNAs associated with DKD. Box C/D snoRNAs are involved in 2'-O-methylation of specific
sequences in target RNA sequences, mainly rRNA, thereby affecting RNA structure, stability and
interactions. In addition, other functions have been ascribed to the snoRNAs such as mRNA splicing
and editing. SnoRNA derived fragments have also been described and it has been speculated that they
exert piRNA or miRNA like functions [38].
Although snoRNAs represent only ten percent of the total circulating sncRNA transcriptome in our
study they represent seventy-one percent of the sncRNAs found associated with DKD suggesting that
snoRNAs may function as early biomarkers of DKD. SnoRNA U8 was associated with all three
phenotypes but has not been linked to kidney function previously. SnoRNAs SNORD118, SNORD24,
SNORD107 and SNORD87 where associated with DKD and UACR but not eGFR. SnoRNAs U8 and
SNORD118 both remained significant in the full models. SNORD87 and SNORD107 associate with
the same set of proteins, including UFM1, NAMPT, PDGFB, and the two chemokines CCL27 and
CCL28, which bind to the same receptor (CCR10). A recent report by Chen et al. showed that post-
transcriptional stabilization of the CCL28 mRNA by N6-methyladenosine (m6A) modification

increases protein levels of this chemokine and reduces kidney damage after ischemia reperfusion
injury [39]. Given the known role of snoRNAs in post-transcriptional RNA modification and
stabilization via 2'-O-methylation this might provide a putative mechanism and explanation for the
observed snoRNA-protein associations. The other proteins associated with these snoRNAs (UFM1,
NAMPT, PDGFB) have also been implicated in kidney function previously [40-43]. Further studies
are needed to investigate the role of these snoRNAs in DKD. In addition to the snoRNAs described
above various other snoRNAs are associated with eGFR and UACR (13-31% of total number of
significant sncRNA). Almost all remained significant in the full models. At present there are no other
studies that have investigated the role of snoRNAs in diabetic DKD.

Members of other sncRNA classes
Although other classes of sncRNAs, such as Y_RNA derived fragments, tRNA fragments (tRF),
lncRNA fragments and snRNAs represent roughly twenty percent of all circulating sncRNA detected
we did not find any significant associations with DKD. We did find a small number of associations
with eGFR and UACR though.

Strengths and limitations
Our study has several strengths. It is one of the largest circulating sncRNA profiling studies studying
the role of circulating sncRNAs in DKD. This improves power and reduces the chance of false
positives. In addition, it uses untargeted RNA sequencing allowing not only the study of miRNAs but
also several other classes of sncRNA showing that especially these other types of sncRNAs associated
with DKD and its components eGFR and UACR. The use of early, mainly mild cases of DKD
facilitates the identification of early markers of disease and may help to introduce early interventions
aiming to preserve kidney function in those at risk.
A limitation of our study is the use of a total RNA isolation procedure and as such we were not able to
investigate if the sncRNAs are carried by extracellular vesicles or are associated with carriers such as
HDL and Ago-2 [2]. The use of plasma for RNA isolation is another limitation. Although from a
biomarker perspective this is not a limitation, from a functional perspective the role of the sncRNAs
we identified needs to be established in the relevant cells and tissues. Another limitation of our study
is its cross-sectional, observational design. As such we cannot exclude that (part of) the observed
changes and associations are a reflection of the reduced kidney function in the participants.

In conclusion, in this study we have shown that several circulating sncRNAs are associated with DKD.
Interestingly, only a relatively small fraction belong to the well-studied microRNAs. This highlights
the potential involvement of other classes of sncRNAs, such as snoRNAs and piRNAs, that have so far
not been investigated in relation to kidney function before. Future prospective and functional studies

are needed to establish the role of these novel circulating sncRNAs in subjects with diabetes and

chronic kidney disease.

Statements and Declarations

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank all staff and participants of the Hoorn Diabetes Care System for their support

and participation.

Funding

This work was supported by a diabetes breakthrough grant from the Dutch Diabetes Research

Foundation and ZonMw (grant number 459001015) and a grant from the EFSD/Boehringer Ingelheim

European Research Programme on "Multi-System Challenges in Diabetes" 2021. The proteomic

analyses were funded by a grant from the Innovative Medicines Initiative 2 Joint Undertaking, under

grant agreement no. 115881 (RHAPSODY). This Joint Undertaking receives support from the

European Union's Horizon 2020 research and innovation program and EFPIA and is supported by the

Swiss State Secretariat for Education, Research and Innovation (SERI), under contract no. 16.0097.

The study sponsors/funders were not involved in the design of the study; the collection, analysis, and

interpretation of data; writing the report; and did not impose any restrictions regarding the publication

of the report.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

LMtH, JWJB and RCS designed the study. JAdK and SJC isolated RNA and prepared the samples for

RNAseq. MTB and JWJB were responsible for phenotypic data acquisition and preparation. LMtH,

GAB and RCS prepared, analyzed and interpreted the data. LMtH and RCS wrote the first draft of the

manuscript. RB contributed to the critical interpretation of the results. LMtH and RCS are the

guarantors of this work and, as such, had full access to all the data in the study and take responsibility

for the integrity of the data and the accuracy of the data analysis. All authors critically read and revised

the manuscript and approved the final version of the manuscript.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. The study has

been approved by the ethics committee of the Amsterdam University Medical Center, location VUmc

(approval number 2007/57; NL14692.029.07).

Consent to participate

All participants provided written informed consent prior to participating in this study.

Data availability

The datasets generated during and/or analyzed during the current study are not publicly available due restrictions in the informed consent but are available from the corresponding author on reasonable request.

References

[1]   Persson F, Rossing P (2018) Diagnosis of diabetic kidney disease: state of the art and future perspective. Kidney Int Suppl (2011) 8(1): 2-7. 10.1016/j.kisu.2017.10.003

[2]   van Zonneveld AJ, Zhao Q, Rotmans JI, Bijkerk R (2023) Circulating non-coding RNAs in chronic kidney disease and its complications. Nat Rev Nephrol 19(9): 573-586. 10.1038/s41581-023-00725-w

[3]   Loganathan TS, Sulaiman SA, Abdul Murad NA, et al. (2020) Interactions Among Non-Coding RNAs in Diabetic Nephropathy. Front Pharmacol 11: 191. 10.3389/fphar.2020.00191

[4]   Gholaminejad A, Abdul Tehrani H, Gholami Fesharaki M (2018) Identification of candidate microRNA biomarkers in diabetic nephropathy: a meta-analysis of profiling studies. J Nephrol. 10.1007/s40620-018-0511-5

[5]   Assmann TS, Recamonde-Mendoza M, de Souza BM, Bauer AC, Crispim D (2018) MicroRNAs and diabetic kidney disease: Systematic review and bioinformatic analysis. Mol Cell Endocrinol 477: 90-102. 10.1016/j.mce.2018.06.005

[6]   Freedman JE, Gerstein M, Mick E, et al. (2016) Diverse human extracellular RNAs are widely detected in human plasma. Nat Commun 7: 11106. 10.1038/ncomms11106

[7]   van der Heijden AA, Rauh SP, Dekker JM, et al. (2017) The Hoorn Diabetes Care System (DCS) cohort. A prospective cohort of persons with type 2 diabetes treated in primary care in the Netherlands. BMJ Open 7(5): e015599. 10.1136/bmjopen-2016-015599

[8]   KDIGO (2013) Kidney Disease: Improving Global Outcomes (KDIGO). CKD work group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. . Kidney Int 3: 1-150

[9]   Rozowsky J, Kitchen RR, Park JJ, et al. (2019) exceRpt: A Comprehensive Analytic Platform for Extracellular RNA Profiling. Cell Syst 8(4): 352-357.e353. 10.1016/j.cels.2019.03.004

[10]   Griffiths-Jones S (2004) The microRNA Registry. Nucleic Acids Res 32(Database issue): D109-111. 10.1093/nar/gkh023

[11]   Sai Lakshmi S, Agrawal S (2008) piRNABank: a web resource on classified and clustered Piwi-interacting RNAs. Nucleic Acids Res 36(Database issue): D173-177. 10.1093/nar/gkm696

[12]   Harrow J, Frankish A, Gonzalez JM, et al. (2012) GENCODE: the reference human genome annotation for The ENCODE Project. Genome Res 22(9): 1760-1774. 10.1101/gr.135350.111

[13]   Glažar P, Papavasileiou P, Rajewsky N (2014) circBase: a database for circular RNAs. Rna 20(11): 1666-1670. 10.1261/rna.043687.113

[14]   Chan PP, Lowe TM (2009) GtRNAdb: a database of transfer RNA genes detected in genomic sequence. Nucleic Acids Res 37(Database issue): D93-97. 10.1093/nar/gkn787

[15]   ThermoFisherScientific (2016) A technical guide to identifying miRNA normalizers using TaqMan Advanced miRNA Assays. Available from https://assets.thermofisher.com/TFS-Assets/GSD/Reference-Materials/identifying-mirna-normalizers-white-paper.pdf

[16]   Barrett T, Wilhite SE, Ledoux P, et al. (2012) NCBI GEO: archive for functional genomics data sets—update. Nucleic Acids Research 41(D1): D991-D995. 10.1093/nar/gks1193

[17]   Slieker RC, Donnelly LA, Fitipaldi H, et al. (2021) Distinct Molecular Signatures of Clinical Clusters in People with Type 2 Diabetes: an IMIRHAPSODY Study. Diabetes 70(11): 2683-2693. 10.2337/db20-1281

[18]   Robinson MD, McCarthy DJ, Smyth GK (2009) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26(1): 139-140.
10.1093/bioinformatics/btp616
[19]   Taïbi F, Metzinger-Le Meuth V, M'Baya-Moutoula E, et al. (2014) Possible involvement of
microRNAs in vascular damage in experimental chronic kidney disease. Biochim Biophys Acta
1842(1): 88-98. 10.1016/j.bbadis.2013.10.005
[20]   Müller-Deile J, Gellrich F, Schenk H, et al. (2016) Overexpression of TGF-β Inducible
microRNA-143 in Zebrafish Leads to Impairment of the Glomerular Filtration Barrier by Targeting
Proteoglycans. Cell Physiol Biochem 40(5): 819-830. 10.1159/000453142
[21]   Bai S, Xiong X, Tang B, et al. (2020) Exosomal circ_DLGAP4 promotes diabetic kidney
disease progression by sponging miR-143 and targeting ERBB3/NF-κB/MMP-2 axis. Cell Death Dis
11(11): 1008. 10.1038/s41419-020-03169-3
[22]   Ge X, Xi L, Wang Q, et al. (2020) Circular RNA Circ_0000064 promotes the proliferation and
fibrosis of mesangial cells via miR-143 in diabetic nephropathy. Gene 758: 144952.
10.1016/j.gene.2020.144952
[23]   Wu J, Liu J, Ding Y, et al. (2018) MiR-455-3p suppresses renal fibrosis through repression of
ROCK2 expression in diabetic nephropathy. Biochem Biophys Res Commun 503(2): 977-983.
10.1016/j.bbrc.2018.06.105
[24]   Zhu XJ, Gong Z, Li SJ, Jia HP, Li DL (2019) Long non-coding RNA Hottip modulates high-
glucose-induced inflammation and ECM accumulation through miR-455-3p/WNT2B in mouse
mesangial cells. Int J Clin Exp Pathol 12(7): 2435-2445
[25]   Rivoli L, Vliegenthart AD, de Potter CM, et al. (2017) The effect of renal dysfunction and
haemodialysis on circulating liver specific miR-122. Br J Clin Pharmacol 83(3): 584-592.
10.1111/bcp.13136
[26]   Mollet IG, Macedo MP (2023) Pre-Diabetes-Linked miRNA miR-193b-3p Targets
PPARGC1A, Disrupts Metabolic Gene Expression Profile and Increases Lipid Accumulation in
Hepatocytes: Relevance for MAFLD. Int J Mol Sci 24(4). 10.3390/ijms24043875
[27]   Trevisani F, Ghidini M, Larcher A, et al. (2016) MicroRNA 193b-3p as a predictive biomarker
of chronic kidney disease in patients undergoing radical nephrectomy for renal cell carcinoma. Br J
Cancer 115(11): 1343-1350. 10.1038/bjc.2016.329
[28]   Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD (2019) PIWI-interacting RNAs:
small RNAs with big functions. Nat Rev Genet 20(2): 89-108. 10.1038/s41576-018-0073-3
[29]   El-Mogy M, Lam B, Haj-Ahmad TA, et al. (2018) Diversity and signature of small RNA in
different bodily fluids using next generation sequencing. BMC Genomics 19(1): 408. 10.1186/s12864-
018-4785-8
[30]   Gohda T, Niewczas MA, Ficociello LH, et al. (2012) Circulating TNF receptors 1 and 2
predict stage 3 CKD in type 1 diabetes. J Am Soc Nephrol 23(3): 516-524. 10.1681/asn.2011060628
[31]   Niewczas MA, Gohda T, Skupien J, et al. (2012) Circulating TNF receptors 1 and 2 predict
ESRD in type 2 diabetes. J Am Soc Nephrol 23(3): 507-515. 10.1681/asn.2011060627
[32]   Kobayashi H, Looker HC, Satake E, et al. (2022) Neuroblastoma suppressor of tumorigenicity
1 is a circulating protein associated with progression to end-stage kidney disease in diabetes. Sci
Transl Med 14(657): eabj2109. 10.1126/scitranslmed.abj2109
[33]   Niewczas MA, Pavkov ME, Skupien J, et al. (2019) A signature of circulating inflammatory
proteins and development of end-stage renal disease in diabetes. Nat Med 25(5): 805-813.
10.1038/s41591-019-0415-5
[34]   Christensson A, Ash JA, DeLisle RK, et al. (2018) The Impact of the Glomerular Filtration
Rate on the Human Plasma Proteome. Proteomics Clin Appl 12(3): e1700067.
10.1002/prca.201700067
[35]   Jalal D, Sanford B, Renner B, et al. (2021) Detection of pro angiogenic and inflammatory
biomarkers in patients with CKD. Sci Rep 11(1): 8786. 10.1038/s41598-021-87710-0
[36]   Ngo D, Wen D, Gao Y, et al. (2020) Circulating testican-2 is a podocyte-derived marker of
kidney health. Proc Natl Acad Sci U S A 117(40): 25026-25035. 10.1073/pnas.2009606117
[37]   Deogharia M, Majumder M (2018) Guide snoRNAs: Drivers or Passengers in Human
Disease? Biology (Basel) 8(1). 10.3390/biology8010001

[38]   Bergeron D, Fafard-Couture É, Scott MS (2020) Small nucleolar RNAs: continuing
identification of novel members and increasing diversity of their molecular mechanisms of action.
Biochem Soc Trans 48(2): 645-656. 10.1042/bst20191046
[39]   Chen J, Xu C, Yang K, et al. (2023) Inhibition of ALKBH5 attenuates I/R-induced renal
injury in male mice by promoting Ccl28 m6A modification and increasing Treg recruitment. Nat
Commun 14(1): 1161. 10.1038/s41467-023-36747-y
[40]   Ostendorf T, Boor P, van Roeyen CR, Floege J (2014) Platelet-derived growth factors
(PDGFs) in glomerular and tubulointerstitial fibrosis. Kidney Int Suppl (2011) 4(1): 65-69.
10.1038/kisup.2014.12
[41]   Li H, Li M, Liu C, et al. (2023) Causal effects of systemic inflammatory regulators on chronic
kidney diseases and renal function: a bidirectional Mendelian randomization study. Front Immunol 14:
1229636. 10.3389/fimmu.2023.1229636
[42]   Hu X, Zhang H, Zhuang L, et al. (2020) Ubiquitin-Fold Modifier-1 Participates in the Diabetic
Inflammatory Response by Regulating NF-κB p65 Nuclear Translocation and the Ubiquitination and
Degradation of IκBα. Drug Des Devel Ther 14: 795-810. 10.2147/dddt.S238695
[43]   Muraoka H, Hasegawa K, Sakamaki Y, et al. (2019) Role of Nampt-Sirt6 Axis in Renal
Proximal Tubules in Extracellular Matrix Deposition in Diabetic Nephropathy. Cell Rep 27(1): 199-
212.e195. 10.1016/j.celrep.2019.03.024
512
513
514

515 **Table 1. Clinical characteristics of the study population.**

	all	No DKD	DKD
n	475	406	69
Gender (%Female)	45	46	42
Age (years)	61.4 ± 9.1	61.0 ± 8.8	63.8 ± 10.1*
BMI (Kg/m^2)	30.7 ± 6.0	30.3 ± 5.6	32.7 ± 7.3*
Age-at-Diagnosis (years)	59.3 ± 9.0	58.8 ± 8.7	61.8 ± 9.8*
Diabetes duration (years)	2.2 ± 1.1	2.2 ± 1.1	2.0 ± 1.1
Fasting glucose (mmol/L)	8.0 ± 1.6	8.0 ± 1.7	8.1 ± 1.4
HbA1c (mmol/mol)	46 (42, 51)	46 (42, 51)	46 (43, 54)
HDL (mmol/L)	1.19 ± 0.32	1.20 ± 0.33	1.11 ± 0.25
eGFR (ml/min/1,73 m^2)	84 ± 16	87 ± 13	70 ± 22*
eGFR ≤ 60 (%)	34 (7)	0	34 (50)
Urine Albumin-to-Creatinine ratio (mg/mmol)	0.48 (0.20, 1.00)	0.41 (0.18, 0.79)	3.99 (0.79, 10.5)*
Albuminuria (%)	43 (9)	0	43 (62)
Chronic Kidney Disease (%))	69 (15)	0	69 (100)
Diastolic blood pressure (mm Hg)	78 ± 9	78 ± 9	79 ± 10
Systolic blood pressure (mm Hg)	139 ± 18	139 ± 18	142 ± 20
Smoking (current/ever/never)	92/130/250	76/107/220	16/23/30
Medication use			
Metformin (%)	340 (72)	287 (71)	53 (77)
SU (%)	130 (27)	105 (26)	25 (36)
Other glucose-lowering (%)	13 (3)	12 (3)	1 (1)
Antihypertensive medication (%)	319 (67)	260 (64)	59 (86)*
Lipid lowering medication (%)	320 (67)	271 (67)	49 (71)

516 Data are mean ± SD, n (%) or median (IQR). All measures taken at the time of blood sampling.

517 * P≤0.05 versus those without DKD.

519 **Table 2. Frequency of different types of sncRNAs associated with the three endpoints.**

	Total # of sncRNA detected		DKD			eGFR			UACR		
	n	%	n	%	p*	n	%	p*	n	%	p*
Total # sncRNAs	**938**	**100**	**7**			**107**			**220**		
miRNA	602	64	1	14	0.011	63	59	0.29	93	42	4.5×10^{-9}
snoRNA	96	10	5	71	2.2×10^{-4}	14	13	0.40	67	31	8.4×10^{-13}
lncRNA	93	10				13	12	0.50	20	9	0.80
piRNA	45	5	1	14	0.30	11	10	0.037	23	11	4.0×10^{-3}
Y_RNA fragments	33	4							1	0.5	0.013
others	26	3							6	3	1.00
tRNA fragments	24	3				4	4	0.52	1	0.5	0.067
snRNA	12	1				1	1	1.00	6	3	0.13
scaRNA	7	1				1	1	0.58	3	1	0.41

520 * P-value for comparison against the distribution of all sncRNAs detected.

523 **Table 3. Overview of seven sncRNAs associated with prevalent DKD**

sncRNA	Prevalent DKD			eGFR			UACR	
	LogFC	P_{FDR}		logFC	P_{FDR}		logFC	P_{FDR}
U8	-0,859	$1,90 \times 10^{-03}$		0,012	$2,79 \times 10^{-02}$		-0,724	$6,74 \times 10^{-03}$
SNORD118	-0,861	$1,90 \times 10^{-03}$		0,009	$8,94 \times 10^{-02}$		-0,858	$1,73 \times 10^{-03}$
miR-143-5p	-0,429	$4,01 \times 10^{-02}$		0,000	$9,86 \times 10^{-01}$		-0,376	$3,30 \times 10^{-02}$
SNORD24	-0,785	$4,01 \times 10^{-02}$		0,011	$6,42 \times 10^{-02}$		-0,919	$1,73 \times 10^{-03}$
piR_019825|DQ597218	-0,485	$4,91 \times 10^{-02}$		0,012	$3,04 \times 10^{-03}$		-0,129	$5,71 \times 10^{-01}$
SNORD107	-1,194	$4,91 \times 10^{-02}$		0,013	$1,43 \times 10^{-01}$		-2,214	$1,58 \times 10^{-05}$
SNORD87	-1,129	$4,91 \times 10^{-02}$		0,012	$2,17 \times 10^{-01}$		-2,253	$1,18 \times 10^{-05}$

524 LogFC is the log2 of the fold change. P_{FDR} represents the P value after correction for multiple hypothesis testing using the Benjamini-Hochberg procedure.

 Figure 1. sncRNAs significantly associated with DKD.

Panel A. Volcano plot showing the association between sncRNAs and DKD in the base model. sncRNAs in blue are
significantly lower is cases with DKD after multiple hypothesis testing (FDR≤0.05). logFC is the Log2 of the fold change
between cases and controls. Panel B. U8 levels in those with or without DKD. CPM = counts per million reads. Panel C.
Overlap between sncRNAs significantly associated with DKD, eGFR or UACR in the base model. Panel D. Pairwise
correlations for the seven sncRNAs associated with DKD. Colors represent Spearman's rho. Panel E. Expression pattern for
sncRNAs associated with DKD in publicly available datasets for 11 tissues of interest. Colors represent the Z-scaled
expression of a sncRNA where blue indicates relative high expression and red relative low expression.

Figure 2. sncRNAs associated proteins.

Overview of the proteins that associate with sncRNA expression levels in plasma from persons with T2D. sncRNAs and

proteins in green are also significantly associated with eGFR and those in blue are associated with UACR ($P_{FDR} \leq 0.05$). Blue

line and text indicate a negative association and a red line and text indicates a positive association. Panel A. piR-019825

associated proteins ($P_{FDR} \leq 0.05$). Panel B. snoRNA associated proteins ($P_{FDR} \leq 0.05$). Panel C. Pairwise correlations between

the 11 piR-019825 associated proteins. Panel D. Pairwise correlations between the 11 snoRNA associated proteins.

545