Volitional brain regulation is related to improvements in affective characteristics of adolescents with autism spectrum disorder

Manfred Klöbl[1], Karin Prillinger[2], Robert Diehm[3], Kamer Doganay[2], Rupert Lanzenberger[1], Luise Poustka[4], Paul Plener[2,5], Lilian Konicar[2,*]

[1] Department of Psychiatry and Psychotherapy, Comprehensive Center for Clinical Neurosciences and Mental Health (C3NMH), Medical University of Vienna, Vienna, Austria
[2] Department of Child and Adolescence Psychiatry, Medical University of Vienna, Vienna, Austria
[3] Department of Pediatrics and Adolescent Medicine, Medical University of Vienna, Austria
[4] Department of Child and Adolescence Psychiatry, Medical University of Göttingen, Göttingen, Germany
[5] Department of Child and Adolescent Psychiatry and Psychotherapy, University of Ulm, Ulm, Germany

* Correspondence to:
Lilian Konicar
Department of Child and Adolescent Psychiatry
Medical University of Vienna
lilian.konicar@meduniwien.ac.at

Abstract

Intro: Even though emotions often play a role in neurofeedback (NF) regulation strategies, investigations of the relationship between the induced neuronal changes and improvements in affective domains are scare in electroencephalography-based studies. We thus extend the findings of the first study on slow cortical potential (SCP) NF in autism spectrum disorder (ASD) (Konicar, et al., 2021; NeuroImage: Clinical) by relating affective changes to whole-brain activity at rest and during regulation.

Methods: Forty-one male adolescents with ASD were scanned at rest using functional magnetic resonance imaging before and after half of them underwent NF training and half received treatment as usual. Furthermore, parents reported on affective characteristics at both times. The NF group had to alternatingly produce negative and positive SCP shifts (increase and decrease cortical excitability) during training and was additionally scanned while applying their developed regulation strategies.

Results: We found increases of resting activity in the anterior cingulate cortex and right inferior temporal gyrus as well as improvements in affective characteristics over both groups. Activation corresponding to SCP differentiation (difference of the activations caused by applying the strategies to produce negative and positive SCP shifts) in these regions correlated with the affective improvements. A further correlation was found for Rolandic operculum activation corresponding to positive SCP shifts. However, there were no significant correlations with the respective achieved SCP regulation during NF training.

Conclusion: SCP NF in ASD did not lead to superior improvements in neuronal or affective functioning compared to treatment as usual. However, the affective changes might be related to the individual regulation strategies as indicated by significant correlations to brain activation during the transfer run but not the achieved SCP regulation. Our study further corroborates the importance of the ACC and ITG as well as of affective functioning for the treatment of ASD.

Keywords

slow cortical potentials, neurofeedback, autism spectrum disorder, emotion regulation, functional magnetic resonance imaging

Introduction

Autism spectrum disorder (ASD) constitutes a pervasive neurodevelopmental disorder which has its onset during childhood and comprises deficits in communication and interaction abilities as well as atypical, repetitive behavior (McPartland & Volkmar, 2012). A variety of therapeutic approaches exists, each focusing on different symptoms (Correll et al., 2021; Famitafreshi & Karimian, 2018).

A particular family of treatment approaches based on a system- and network-level understanding of ASD comprises the application of non-invasive neuromodulation techniques (Khaleghi, Zarafshan, Vand, & Mohammadi, 2020). Especially transcranial direct current (Amatachaya et al., 2014; Amatachaya et al., 2015; Prillinger et al., 2021) and magnetic stimulation (Anninos, Chatzimichael, Adamopoulos, Kotini, & Tsagas, 2016; Doruk Camsari, Kirkovski, & Croarkin, 2018; Panerai et al., 2014) have been explored in the past. In contrast, neurofeedback (NF) is a neuromodulatory intervention without external stimulation.

When undergoing NF, patients train to gain volitional control over certain characteristics of their brain activity, aiming to improve aberrant patterns specific to their condition. While reviews on the application of electroencephalography-based (EEG) NF for the treatment of attention-deficit / hyperactivity disorder (ADHD) have yielded mixed results (Arns, Heinrich, & Strehl, 2014; Sonuga-Barke et al., 2013), there is a particular recommendation for slow cortical potentials (SCPs) NF (Mayer, Wyckoff, & Strehl, 2013).

SCP NF has been successfully applied for the treatment of ADHD in children (Drechsler et al., 2007; Strehl et al., 2017; Ute Strehl et al., 2006), adolescents (Albrecht et al., 2017) and adults (Mayer, Blume, Wyckoff, Brokmeier, & Strehl, 2016). These very slow EEG fluctuations (typically below 1 Hz) are related to the excitability threshold of the upper cortical layers, where SCP negativity corresponds to increased and SCP positivity to decreased cortical excitability (Birbaumer, Elbert, Canavan, & Rockstroh, 1990). SCPs further constitute suspected contributors to the blood-oxygenation-level-dependent (BOLD) signal measured in functional magnetic resonance imaging (fMRI) (He & Raichle, 2009; Khader, Schicke, Röder, & Rösler, 2008), which is now also explored as application modality for NF in ASD (Direito et al., 2019; Direito et al., 2021). However, the utility of NF for the treatment of ASD in general has also been questioned (Holtmann et al., 2011).

This criticism is based on the high comorbidity of ASD with ADHD with estimates ranging from 37 to 85% (Leitner, 2014) rendering it difficult to assign symptom improvements to either of the disorders in isolation. However, previous research using EEG NF in ADHD and ASD often focused on attention deficits (Arns et al., 2014; Hurt, Arnold, & Lofthouse, 2014) while studies on the emotional and empathic components are largely missing. The recently conducted first study on SCP NF in ASD reported positive effects on ASD-specific symptomatology (Konicar et al., 2021) but also showed the complex influence of the ADHD-related problems regarding attention, hyperactivity and impulsivity (Prillinger, Radev, Doganay, Poustka, & Konicar, 2022). Overall, transdiagnostic similarities between ASD and ADHD encompass deficits in emotion regulation (ER), emotion recognition, attention,

cognitive flexibility, inhibition, reward processing, working memory, organization and planning (Taurines et al., 2012; Vaidya et al., 2020).

Waddington et al. (2018) identified four classes of emotion recognition impairments in children and adolescents with and without diagnoses of ASD and/or ADHD. Subsequent polygenic risk score analyses revealed that higher scores for ASD were surprisingly related to faster visual emotion recognition (Waddington et al., 2021). Furthermore, a class defined by impulsive and imprecise visual emotion recognition was associated with lower genetic risk of ASD. On the contrary, slower emotion recognition was reported in adults with ASD or ADHD, related to mind wandering in ADHD but slower processing in ASD (Helfer et al., 2021). Another transdiagnostic analysis in preschool children revealed an increased probability for moderate ER and executive functioning deficits given an ASD or ADHD diagnosis, but for high ER deficits in ADHD and high executive functioning deficit in ASD (Ros & Graziano, 2020).

Regarding empathic capabilities, individuals with ASD showed deficits in cognitive empathy (CE) but not affective empathy (Mazza et al., 2014; Ziermans, de Bruijn, Dijkhuis, Staal, & Swaab, 2019). Reports of empathic abilities may further differ between objective and subjective assessments (McKenzie, Russell, Golm, & Fairchild, 2021). Interpretation of these results in synopsis is undoubtedly complex, which is indicative of the fact that ER research in ASD is still in its infancy (Cai, Richdale, Uljarević, Dissanayake, & Samson, 2018). However, the importance and efficacy of therapeutic approaches incorporating ER training for ASD are already evident (Reyes, Pickard, & Reaven, 2019).

Regarding neural signatures of ASD, resting-state (RS) functional connectivity was long believed to be characteristically altered, presenting with local over- and long-range under-connectivity but this view has been challenged by more recent studies (see Mohammad-Rezazadeh, Frohlich, Loo, and Jeste (2016) for a critical review). Moreover, it was shown that even the same regionally specific measures of functional connectivity were not reproducible across research sites (King et al., 2019). However, the average absolute global connectivity, which can be interpreted as a measure of the linear activation dependency across the brain, was found to be lower in children and adolescents with ASD compared to typically developing controls (Yerys et al., 2017). This might indicate global, rather than local alterations.

Besides functional connectivity, differences between individuals with and without ASD were found in measures of brain activity at rest. A meta-analysis combining results from studies using regional homogeneity (ReHo; a measure sensitive to brain activity and local connectivity), amplitude of low-frequency fluctuations (ALFF; activity within a certain frequency band) and cerebral blood flow reported robust over-activation in language-related and motor areas as well as under-activation in the default mode network (W. Wang et al., 2018). Moreover, a large-scale, multi-center study reported two clusters in the brain showing abnormalities in RS connectivity and activity across several measures (Di Martino et al., 2014): The first cluster reaches from the left posterior insula to the operculum with decreases in voxel-matched homotopic connectivity (i.e., symmetric connectivity), ReHo and degree centrality (a measure of global connectedness). The second cluster is located at the right dorsal superior

frontal cortex and shows increases in fractional ALFF (fALFF; an amplitude-normalized version of ALFF), ReHo and degree centrality.

Even though SCPs are a common NF target, only few studies investigated the neural processes accompanying SCP self-regulation using fMRI. Hinterberger et al. (2003) showed that during transfer runs (training runs without feedback for decoupling regulation success from the feedback procedure) SCP negativity is related to wide-spread activation in fMRI. In turn, SCP positivity was associated with widespread deactivation. It was further shown in patients with epilepsy that deactivation during transfer runs only occurred for individuals who successfully learned to self-induce SCP positivity (U. Strehl et al., 2006).

In order to gain further knowledge on ER in ASD, we investigated potential treatment-induced changes in several affective characteristics based on parent questionnaires. Since previous studies reported ASD-related changes in brain connectivity and activity, we further tested for treatment effects on these measures using RS fMRI and whether these are related to the questionnaire data. Finally, based on suggested links between SCPs and BOLD signal, we conducted an adapted transfer run in the MRI scanner and searched for relationships between whole-brain activation corresponding to SCP regulation and affective changes.

Methods

The overall clinical trial and all parts presented here were conducted in accordance with the Declaration of Helsinki and the good scientific practice guidelines of the Medical University of Vienna. The project was approved by the institutional review board (EK-Nr.: 1850/2015) and registered at drks.de (DRKS00012339).

Experimental Design

Participants were randomly allocated to either 24 sessions of SCP NF (n_1 = 21) or treatment as usual (TAU; n_2 = 20). SCP feedback was calculated from the fronto-central electrode located according to the extended 10-20 EEG system and presented on a screen as graphical object of the participants' choice. The participants' goal was to gain volitional control over their brain activity, moving the object up or down via changes in SCP positivity or negativity (see Konicar et al. (2021) for details on the NF training). In order to assess neuronal and subjective changes, fMRI and psychometric data were acquired before the first and after the last NF session in both groups.

Participants

Potential male adolescent (12-17 years) participants with a diagnosis of ASD (according to the German version of the "Autism Diagnostic Interview – Revised" (ADI-R) (Bölte, Poustka, Rühl, & Schmötzer, 2006) and the "Autism Diagnostic Observation Schedule" (ADOS-2) (Poustka et al., 2015) were recruited and invited to a pre-screening. The further inclusion criteria were: right-handedness and an IQ above 70 (Wechsler & Petermann, 2012; Wechsler,

Petermann, & Petermann, 2010). Informed assent was given by the participants and informed consent was provided by the parents / legal guardian. Participants were excluded in case of relevant psychiatric, neurological or internal conditions (head injuries, major axis I diagnosis of psychosis, obsessive-compulsive disorder, severe motor or vocal tics, Tourette syndrome, severe depression with suicidality) or MRI contraindications. Previous NF experience and current participation in pharmacological studies were not allowed. Concomitant psychosocial and pharmacological treatments were permitted if kept constant throughout the study participation.

Psychometric assessment

We used the Emotion Regulation Checklist (ERC; Shields and Cicchetti (1997)) and the Griffith Empathy Measure (GEM; Dadds et al. (2008)) to gather parent reports of the participants' development regarding their affective and empathic abilities. The emotion regulation (ER) and lability / negativity (LN) subscales of the ERC were analyzed separately, with the former quantifying expression and self-awareness of emotions as well as empathy, and the latter mood lability and anger dysregulation. For the GEM, we concentrated on the cognitive empathy (CE) subscale, since this ability was shown to be diminished in adolescents with ASD in contrast to affective empathy (Mazza et al., 2014).

Since the frequent co-occurrence of ADHD constitutes the major point of discussion to the application of NF to patients with ASD, the Diagnostic System for Psychiatric Disease in Children and Adolescent, parent-rated version 2 for ADHD (DISYPS-II, Döpfner, Görtz-Dorten, Lehmkuhl, Breuer, and Goletz (2008)) was used to quantify the respective characteristics (hereafter referred to as "ADHD score"). The ADHD score was calculated, age-corrected and transformed to the "standard nine" (stanine) score.

FMRI paradigms and acquisition

RS data was acquired with eyes open fixated on a fixation cross for 8 min.

In the SCP group only, after completing the training, a short transfer run was recorded in 2 min 42 s. The visual cues for up- and down-regulation were the same as for the SCP transfer run: A triangle pointing upwards ("Up" condition) indicated application of the strategy developed to induce negative SCP shifts and a triangle pointing downwards ("Down" condition) indicated application of the strategy developed to produce positive SCP shifts. The conditions were presented five times each for 8 s in a pseudo-randomized order interleaved with baselines of the same duration. Since the baseline in the SCP transfer run was only 2 s long and had no visual indicator, we added a crosshair of the same size and color scheme as the triangles for the fMRI run. As usual in transfer runs after NF training, there was no indication of the currently achieved regulation.

To check for structural anomalies, a T1-weighted image was acquired at the end of each MRI session while the participants were shown a relaxing video with natural scenes.

Measurements were performed on a Siemens Magnetom Prisma 3 T machine (Siemens, Erlangen, Germany) with the same sequence as in Moessnang et al. (2016) due to the previous

successful application in a comparable population: echo / repetition time = 30 / 2000 ms, 3 mm isotropic resolution (+25% gap), 33 slices with 64 x 64 voxels (field of view = 192 x 192 x 123 mm), bandwidth = 2365 Hz/Px. Prospective acquisition correction (PACE) was used for online motion correction.

FMRI preprocessing

As children and adolescents show considerably more in-scanner head movement than adults, additional motion-correction measures were taken. Unless mentioned otherwise, preprocessing was conducted using Statistical Parametric Mapping, version 12 (SPM12; www.fil.ion.ucl.ac.uk/spm/software/spm12/). In a combined first step, physiological artifacts were reduced using PESTICA (Beall & Lowe, 2007) and slice-wise motion correction was performed with SLOMOCO (Beall & Lowe, 2014). Slice-timing was corrected to the temporally middle slice. All acquisitions per paradigm were realigned together for each participant. A population-specific normalization template was created using the CerebroMatic toolbox (Wilke, Altaye, & Holland, 2017) with spatially adaptive non-local means and hidden Markov random field filtering for increased homogeneity over the whole age range. Affine regularization was performed to the standard ICBM template for European brains with parameters downscaled by a factor 10 for better local fitting. This was done since some brains showed unreasonable inflation without regularization. Thus, tissue distributions are age-adjusted but the localization of the regions approximately in standard space. The original voxel size was used for reslicing (Mueller, Lepsien, Möller, & Lohmann, 2017). The BrainWavelet toolbox (Patel et al., 2014) was employed for non-linear artifact correction with the parameters "threshold" set to "15" due to the application to unsmoothed data and "chsearch" to "harsh" to be more sensitive towards slow artifacts. The data was finally smoothed with a Gaussian kernel with full width at half maximum (FWHM) of 3 times the voxel size.

FMRI modelling

An adapted CompCor approach (Behzadi, Restom, Liau, & Liu, 2007; Klöbl et al., 2020) and the Fristion-24 model (Friston, Williams, Howard, Frackowiak, & Turner, 1996) were utilized for reduction of any residual physiological or movement-related artifacts in the transfer run and RS data. The latter was further band-limited to 0.01-0.10 Hz using frequency regressors (Hallquist, Hwang, & Luna, 2013). Based on the filtered RS time series, three voxelwise models were set up: First, the percent amplitude of fluctuation (PerAF) model was used for assessing resting brain activity (Jia et al., 2020). This measure was chosen rather than (f)ALFF since it is inherently amplitude-normalized and does not rely on frequencies outside the band of interest. Second, average brain-wide and long-distance connectivity were assessed via global functional connectivity (GFC; a continuous alternative to degree centrality) (Tagliazucchi et al., 2016). Third, ReHo was assessed (Jiang & Zuo, 2016; Zang, Jiang, Lu, He, & Tian, 2004), placing itself conceptually between PerAF and GFC. Since the calculation of ReHo leads to spatial smoothing, this model was applied to the unsmoothed and filtered data. Afterwards,

the ReHo maps were smoothed with a FWHM of 2 times the voxel size achieving smoothness similar to PerAF and GFC. Finally, ReHo and GFC were Fisher z-transformed before group analysis.

The transfer runs were modeled using the 1st-level module in SPM12. The conditions ("Up" / "Down") were used as regressors and the CompCor and Friston-24 time series as nuisance signals. In addition, the equivalent of SCP differentiation between changes in negativity and positivity was calculated as the difference between the "Up" and "Down" conditions (henceforth "fMRI differentiation"). The autocorrelation model was set to "FAST" (Olszowy, Aston, Rua, & Williams, 2019).

Statistical inference

Questionnaire data was analyzed using linear mixed effects models (LMEs). Interactions between "treatment group", "time" (factors) and "ADHD baseline score" (covariate) were analyzed and dropped if non-significant. The LMEs also included participants as random intercepts.

Whole-brain inference was conducted using the 2nd-level module in SPM12. "Treatment group" by "time" and "time" effects were analyzed in one model per RS measure. Contrasts for within-group effects were further estimated in case of non-significant interactions. Correlations of regulation direction-specific activation and fMRI differentiation during the transfer run with the ERC and GEM score changes were of interest for the transfer run. Family-wise error-corrected (FWE) results are reported at the cluster- (primary threshold $p \leq 0.001$) or peak-level. Influences of ADHD were controlled for with the pre-training ADHD scores as covariate.

Associations between the RS findings and questionnaire score changes over time were investigated on an exploratory basis using partial correlation (Pearson or Spearman, depending on a visual check of the distributions, corrected for treatment group; median cluster values were extracted using the MarsBaR toolbox 0.44).

In case of significant correlations between regulation activation in the transfer run and questionnaire score changes, the latter were subsequently correlated with the average amount of SCP regulation corresponding to the activation contrast achieved during the third and last quarter (six days each) of NF training. These periods showed the strongest regulation or were closest to the second MRI session (Konicar et al., 2021). This way, we checked whether NF training success was related to the score changes (c.f., Heinrich, Gevensleben, Becker, and Rothenberger (2020)).

All tests (including neuroimaging models (Chen et al., 2019)) were two-sided and multiplicity-corrected to $p \leq 0.05$. To correct for the number of questionnaire scales / RS models / transfer run correlations, we employed an in-house developed algorithm based on the dependency-adjusted D/AP approach (Sankoh, Huque, & Dubey, 1997) (see supplement for implementation).

Results

Of the 41 participants recruited, one member of the TAU group had to be excluded due to missing compliance. Complete RS data was available from 36 and transfer run data from 20 participants (Table 1; three subjects did not participate in the second session for personal reasons, two had to be excluded due to compromised data quality).

The results of the SCP regulation training and ASD core symptoms longitudinally quantified via the Social Responsiveness Scale (Bölte & Poustka, 2008) are described in Konicar et al. (2021). A detailed analysis of the comorbid ADHD symptoms and their influence on attention and expectancy measures via the contingent negative variation (CNV) in the EEG signal is presented in Prillinger et al. (2022).

Psychometric and behavioral analysis

No significant interaction effects were found for any psychometric scale. The ER subscale (higher is better) of the ERC showed a significant symptom improvement over both groups, whereas the LN (lower is better) and GEM CE (higher is better) subscales barely fell short of statistical significance. The LN subscale positively correlated with the pre-training ADHD score.

The mean of the total displacement output of the SLOMOCO step of RS preprocessing was additionally analyzed via LME as the PerAF results indicated a potential relationship with in-scanner movement (see results on the RS models). There was a significant time by group interaction where the SCP NF group showed significantly higher pre-training movement. Total displacement did not correlate with the ADHD score over both scans (Spearman ρ = -0.05, p = 0.6790). No multiplicity correction was applied in the total displacement analysis.

Score	Neurofeedback pre	Neurofeedback post	Treatment as usual pre	Treatment as usual post	p-value
N	21	scores: 21 resting-state: 19 transfer run: 20	20	scores: 20 resting-state: 17	
Age [years] (at first magnetic resonance imaging session)	14.44 ± 1.90		15.01 ± 1.59		0.2779 [1]
Emotion Regulation Checklist: emotion regulation subscale	21.19 ± 3.98	23.05 ± 3.23	20.45 ± 5.00	21.15 ± 3.59	time: 0.0419 [2]
Emotion Regulation Checklist: lability / negativity subscale	33.71 ± 5.83	30.76 ± 5.34	32.00 ± 7.06	31.65 ± 6.43	time: 0.0506 [2] ADHD score: 0.0014 [2]
Griffith Empathy Measure: Cognitive empathy subscale	-1.00 ± 9.85	5.33 ± 9.80	0.60 ± 9.20	1.20 ± 8.89	time: 0.0601 [2]
mean total displacement	2.49 ± 0.88	2.12 ± 0.46	2.01 ± 0.41	2.16 ± 0.42	interaction: 0.0024 [2] group: 0.0145 [2]

Table 1: Overview of the psychometric and behavioral data. Pre- and post-treatment scores are given as mean ± standard deviation. ADHD: attention-deficit / hyperactivity disorder. [1] two-sample t-test, [2] linear mixed effects model.

Resting-state models

Over both groups, an increase in RS activity was found in the ventral anterior cingulate cortex (ACC). Within the TAU group, a further increase was detected in the right precentral gyrus. Another cluster of increased resting activity over both groups reaching into the medial and inferior temporal gyrus (ITG) did not survive correction for the number of models. Since, upon visual inspection (see Figure 1), especially the result in the precentral gyrus might be biased by baseline differences, LMEs of the median values extracted from the clusters were run (treatment group, time and their interaction as factors, the ADHD score as covariate, random intercept per participant). These confirmed the time effect for the ACC (p = 0.0006) and temporal gyrus (p = 5.4E-6) and indicated a baseline difference (p = 0.0326), time (p = 3.5E-5) and interaction effects (p = 0.0020; all uncorrected) for the precentral gyrus. The ADHD score had no significant influence in either model. Details are presented in the upper section of Table 2 and left column of Figure 1.

Since precentral gyrus activation might be indicative of movement, the PerAF values of each region were Pearson partially correlated (corrected for measurement, treatment group and their interaction) with the mean total displacement (thresholded at 1.5 times the interquartile range due to potential outliers) and the pre- and post-training ADHD scores on an exploratory basis. Indeed, significant relationships with total displacement were found for all results (r = [0.24, 0.49], p = [0.0479, 1.6E-5]; all p-values for the ADHD score > 0.45). The thresholded mean total displacement values were then used as additional covariate in a repeated analysis. All results survived this control for in-scanner movement with one further cluster for the TAU group in the triangular gyrus (not significant after multiplicity adjustment; see Table 2, upper section). Given the homogeneous effect on all investigated regions, the influence of total displacement was also investigated on a whole-brain level, resulting in weak but widespread patterns of correlations and anti-correlations (Figure S1).

Exploratory analyses revealed correlations between changes in the LN subscale and the middle / ITL cluster from both groups (Spearman ρ = 0.41, p = 0.0133). The GFC and ReHo models yielded no significant effects.

In-scanner transfer run

During the "Up" and "Down" conditions (corresponding to the induction of negative and positive SCP shifts, respectively), primary and secondary visual areas were activated. No significant difference was found between the two conditions (i.e., no significant fMRI differentiation). Rolandic operculum activation (stretching into the anterior insula and the Heschl gyrus) during the "Down" condition significantly correlated with increases in ER after training. FMRI differentiation in the ITL showed a significant correlation with changes in CE. A further cluster barely missing corrected significance was detected for the correlation of higher fMRI differentiation in the ventral ACC and improvements in LN. A last finding in the right insula did not survive correction for the number of correlations. The baseline ADHD score had no significant influence on any model. For details, see the lower section of Table 2 and right column of Figure 1.

The only notable correlation was found for the change in CE and the average differentiation achieved during the last quarter of the NF training (Spearman: ρ = 0.35, p = 0.1184; all other |ρ| < 0.15, p > 0.53) but was not significant.

Model / condition	Contrast	Coordinates			Cluster size	p	p$_{corr}$	Region
		x	y	z				
PerAF	post > pre	63	-34	-16	56	0.0481	0.1356	middle / inferior temporal gyrus
PerAF	post > pre	12	29	32	140	0.0003	0.0008	anterior cingulate cortex
PerAF	TAU: post > pre	60	-31	-19	peak	0.0433	0.1225	middle / inferior temporal gyrus
PerAF	TAU: post > pre	57	-10	47	96	0.0033	0.0098	precentral gyrus
PerAF - total displacement	post > pre	63	-34	-16	peak	0.0300	0.0860	middle / inferior temporal gyrus
PerAF - total displacement	post > pre	63	-34	-16	56	0.0446	0.1261	middle / inferior temporal gyrus
PerAF - total displacement	post > pre	12	29	32	133	0.0003	0.0010	anterior cingulate cortex
PerAF - total displacement	TAU: post > pre	57	-10	47	82	0.0073	0.0214	precentral gyrus
PerAF - total displacement	TAU: post > pre	54	23	29	58	0.0385	0.1095	triangular gyrus
Down	mean > 0	-18	-82	-16	543	1.8E-9	4.8E-9	primary / secondary visual cortex
Down	mean > 0	-18	-82	-16	peak	5.7E-5	0.0001	
Up	mean > 0	6	-79	-10	816	2.3E-12	5.9E-12	
Up	mean > 0	6	-79	-10	peak	0.0021	0.0056	
Down	ERC ΔER > 0	-39	-13	14	141	0.0026	0.0104	Rolandic operculum, insula, superior temporal gyrus
Down	ERC ΔER > 0	45	-4	-13	peak	0.0381	0.2683	insula
Up-Down	ERC ΔLN < 0	-12	35	20	112	0.0066	0.0515	anterior cingulate / medial prefrontal cortex
Up	GEM ΔCE > 0	54	-49	-22	72	0.0215	0.2974	inferior temporal gyrus
Up-Down	GEM ΔCE > 0	51	-46	-16	157	4.2E-4	0.0068	inferior temporal gyrus

Rows labeled "Resting-state" span the PerAF entries; rows labeled "Transfer run" span the Down/Up entries.

Table 2: Whole-brain resting-state and transfer run results. Resting-state: The original results (PerAF) and those additionally corrected for the total displacement (PerAF - total displacement) are provided. Transfer run: The mean activation per condition (mean > 0) and the correlation with psychometric score changes (ERC ΔER > 0, ER ΔLN < 0, GEM ΔCE > 0) are shown. All models were corrected for the baseline attention-deficit / hyperactivity disorder (ADHD) score. The coordinates need to be seen as rough orientation only due to a study-specific normalization. The p-values were corrected for the investigated models / contrasts. PerAF: percent amplitude of fluctuation; TAU: treatment as usual; SCP: slow cortical potentials neurofeedback; pre / post: measurement before / after intervention, Up: up-regulation, Down: down-regulation, Up-Down: difference between up- and down-regulation; ERC ΔER: Emotion Regulation Checklist emotion regulation subscale change; ERC ΔLN: Emotion Regulation Checklist lability / negativity subscale change; GEM ΔCE: Griffith Empathy Measure cognitive empathy subscale.

Figure 1: Results of the resting brain activity and transfer run analyses. The increase in the percent amplitude of fluctuations (PerAF) of the anterior cingulate cortex over both, the slow cortical potentials (SCP) neurofeedback and treatment as usual (TAU) group, is visualized in the top-left diagram. The middle-left diagram shows a PerAF increase in the precentral gyrus of the TAU and generally higher values in the SCP group. The bottom-left plot shows the PerAF changes over both groups in the temporal gyrus with stronger increases in the TAU group. The right scatter plots show the correlation of the transfer run conditions (Pearson's r corresponding to the parametric analysis as implemented in Statistical Parametric Mapping and Spearman's ρ due to potential outliers): The difference between the "Up" and "Down" conditions (i.e., fMRI differentiation) was anticorrelated with changes on the lability / negativity (ERC ΔLN) subscale of the Emotion Regulation Checklist. The SCP positivity condition correlated with changes on the emotion regulation (ERC ΔER) subscale. Changes on the Griffith Empathy Measure cognitive empathy (GEM ΔCE) subscale again significantly correlated with fMRI differentiation. The whole-brain 3D models were created using BrainNet Viewer 1.7 (Xia, Wang, & He, 2013).

Discussion

In this work, we used fMRI to assess the resting brain activity of adolescents with ASD and collected parent reports of their affective functioning before and after receiving SCP NF or TAU. In addition, the SCP NF group performed an adapted transfer run in the scanner after undergoing NF training. Using the fMRI transfer data, we found relationships between the activation during regulation and improvements in different affective domains.

Affective symptom improvements and the potential influence of regulation strategies

Changes in resting activity and affective improvements over both groups without any interaction effects indicate unspecific positive effects of SCP NF and TAU. Furthermore, while localized correlations between brain activation during regulation and improvements on all investigated affective scales point towards an influence of NF, these improvements did not significantly correlate with SCP regulation (i.e., the SCP shifts produced in the same direction for which the correlations with fMRI activation were detected). Heinrich et al. (2020) likewise concluded universal improvements of emotional and behavioral self-regulation after successfully applying SCP NF in ADHD without finding correlations of training outcomes and symptom improvements. A possible interpretation of these findings is to ascribe the affective improvements not to the achieved SCP regulation but the application of the regulation strategies. In line with this speculation, Hasslinger, D'Agostini Souto, Folkesson Hellstadius, and Bölte (2020) identified emotional strategies as a common regulation approach in SCP NF when treating ADHD. Using the same classifications, strategies from the emotional domain were also frequently reported in Konicar et al. (2021) (see supplement). Furthermore, ER is known to often be a key factor in the development of successful regulation strategies (Eldeeb et al., 2021; Friedrich et al., 2015). It however should be noted that any comparison to NF training outcomes strongly depends on the definition of learning and regulation success.

Regarding the origin of the correlation between fMRI differentiation in the right ITL and improvements in CE the support of the experimenters needs consideration. Empathic comments after negative performance feedback were shown to decreased negative feelings (Seehausen et al., 2016). The participants might have unintentionally related verbal positive reinforcement to their regulation strategies ultimately resulting in CE improvements. In a comparable scenario, the experimenters' empathy was concluded to be a potential driving factor of subjective improvements in the sham group of a NF study in primary insomnia (Schabus et al., 2017). While the weaker correlation of the "Up" condition alone appear to support this hypothesis, the stronger effect for the fMRI differentiation is probably the best argument against such an additional unintentional effect.

Also the TAU group might have improved due to the supportive clinical counseling received, leading to no significant therapeutic advantage of SCP NF. The positive correlation between the LN subscale and the ADHD baseline score corroborates the well-documented emotion dysregulation in ADHD (Christiansen, Hirsch, Albrecht, & Chavanon, 2019; Seymour, Chronis-Tuscano, Iwamoto, Kurdziel, & MacPherson, 2014; Shaw, Stringaris, Nigg, & Leibenluft, 2014). The stronger reduction in head movement (total displacement) in the SCP group, constitutes a potential unspecific training effect. The requirement to sit still for a longer time over repeated NF sessions could have led to the reduced in-scanner movement in the SCP group only.

The anterior cingulate cortex and emotional negativity

Similar to improvements on the psychometric level, we found an increase in ACC resting activity quantified as PerAF over both groups. Early and replicated neuroimaging evidence

for structural and functional alterations in the ACC of patients with ASD was found in smaller gray matter volume and decreased metabolic rate (Haznedar et al., 1997; Laidi et al., 2019). This structural anomaly might appear as lower resting activity following volumetric normalization. A negative correlation between the ADOS and ACC fALFF was found to be a distinctive feature of children diagnosed with "pervasive developmental disorder – not otherwise specified" (Qi et al., 2020). One may assume that symptomatically stronger subgroups lack the correlation between ACC resting activity and ASD symptoms due to a ceiling effect. A lack of activation in the ACC was also found for a stroop task in ADHD and related to the symptoms of inattention and impulsivity (Bush et al., 1999), which are likely shared between ADHD and ASD (Krakowski et al., 2020).

Beyond changes in PerAF, the ventral part of the ACC also showed a negative correlation between fMRI differentiation in the transfer run and the change in the LN subscale of the ERC extending into the medial prefrontal cortex. The amount of SCP differentiation was used as a primary parameter for the analysis of the NF training success as it indicates the participants' ability to differentiate between the regulation directions (Konicar et al., 2021; Ute Strehl et al., 2006). The PerAF model, being a measure of amplitude fluctuation and revealing partly overlapping changes in the ACC, might also be interpreted as a measure of fMRI "differentiation" at rest. However, the different sampling time in SCP and fMRI investigations as well as the BOLD signal containing influences of neural activity of all frequency bands might be the reasons for the missing correlation between the PerAF ACC cluster and the LN subscale changes.

The ACC is known to be related to repetitive behavior in ASD (Thakkar et al., 2008) but also to cognitive inflexibility in depression (Doss et al., 2021; Schwartz et al., 2019; Zheng et al., 2018). The latter is potentially also represented in the LN subscale (Loevaas et al., 2018), providing further evidence that behavioral flexibility might be reflected in neuronal flexibility of the ACC.

The inferior temporal gyrus and cognitive empathy
General as well as regionally specific alterations in temporal lobe structures related to ASD are well-known (Gendry Meresse et al., 2005; Zilbovicius et al., 2000). In a classification tree approach using white and gray matter volumes of various temporal regions, smaller gray matter volume of the right ITG was related to a higher probability of an ASD diagnosis (Neeley et al., 2007). Reduced gray matter volume of the left ITG was also detected in children with low-functioning ASD (Riva et al., 2013) and related to communication skills (Duan et al., 2022). As for the ACC, smaller gray matter volume in the ITG might be related to less activity after structural normalization, relating our finding of an increase after NF training to compensatory processes. In ASD, reduced resting activity estimated via ReHo was also recently reported in the left middle temporal gyrus (Lan et al., 2021). While no correlation with symptom severity was found by the authors, we found a positive relationship of PerAF changes in the affected cluster with changes in emotional negativity (LN subscale). Since this would imply a

worsening in emotional negativity accompanying an increase in ITL activity, this correlation is in contradiction with our other findings.

Besides the temporal lobe results found at rest, which did not survive correction for the number of models, we also identified a positive correlation between fMRI differentiation in the right ITG and improvements in CE. Support for laterality of this correlation as well as specificity for cognitive compared to affective empathy is gained from findings in unilateral mesial temporal lobe epilepsy (Toller et al., 2015). Furthermore, the right ITG was related to personal as well as impersonal emotional imagery whereas it was only impersonal for the left ITG (Preston et al., 2007). This could imply that the right ITG is involved when one tries to understand another person's emotions, i.e., displays CE. Preston et al. (2007) further related reduced neuronal recruitment in a region between the ITG and fusiform gyrus to a lack of CE.

The Rolandic operculum and emotion regulation

The correlation between the "Down" condition (SCP positivity) and the increase in the ER score mostly covers the posterior Rolandic operculum reaching into the insula and superior temporal gyrus. The Rolandic operculum is involved in processing of visceral signals (Blefari et al., 2017), language encoding (Indefrey et al., 2001) and emotion processing (Sutoko et al., 2020). Further, reduced activation was found in the left Rolandic Operculum of adults with ASD compared to neurotypical individuals regarding speech (Gebauer, Skewes, Hørlyck, & Vuust, 2014). However, the Rolandic Operculum showed increased activation during emotion induction with happy compared to sad music (Gebauer, Skewes, Westphael, Heaton, & Vuust, 2014). The same results were previously obtained for children with low-functioning autism and age-matched controls using speech recordings of their parents and each child's favorite song containing vocals (Lai, Pantazatos, Schneider, & Hirsch, 2012).

These findings suggest two possible conclusions related to NF: First, our reported activation related to improvements on the ER subscale was evoked by self-induced positive mood possibly in combination with sound imagination. Second, less severe language deficits facilitated the development of strategies involving inner speech, which had a positive impact on ER. On the role of SCP positivity in particular, we can only make an indirect assumption: Better SCP differentiation was previously associated with less relaxation when trying to produce negative SCP shifts (Brody et al., 1994), suggesting that the potentially induced positive mood is a side effect of more relaxation when trying to produce positive shifts. This could again be seen as support for the idea that the individual regulation strategies are responsible for the correlations between brain activation and affective improvements. In accordance with this idea, the self-reported frequency of cognitive reappraisal strategies for ER correlated with the within-run training progress in upregulating the left ventrolateral prefrontal cortex in a previous fMRI NF study (Keller et al., 2021).

The precentral gyrus at rest

Lastly, we found an increase in resting activity in the right motor cortex of the TAU group, which was, however, biased by a baseline difference. The unspecific correlations to the amount

of in-scanner (head) motion, the fact that motion decreased in the SCP NF but PerAF increases in the TAU group and the effect surviving a correction for motion on group-level (in addition to subject-level) speak against movement-related motor activity as sole cause. Altered connectivity of motor regions (Carper, Solders, Treiber, Fishman, & Müller, 2015; Nebel, Eloyan, Barber, & Mostofsky, 2014) as well as motor impairments (Duffield et al., 2013) are known in ASD. Higher visuomotor impairment was associated with increased ALFF in the precentral cortex, among other regions (Z. Wang et al., 2019). Another study reported increases in gray matter volume in the pre- and postcentral gyri of children with ASD only when no ADHD comorbidity was present (Mahajan, Dirlikov, Crocetti, & Mostofsky, 2016). In a large sample of children and young adults with ADHD, a reduction in gray matter volume was found in the precentral gyrus and several other regions (Bralten et al., 2016). Assuming that the baseline difference in precentral gyrus PerAF is reflecting more deficits / higher symptom severity in pre-treatment measures in the SCP NF group (i.e., increased movement, higher SRS scores), the observed increase in the TAU group cannot be seen as a positive therapeutic effect. This interpretation would also be in line with the previous report of increased ALFF (Z. Wang et al., 2019) and our decision to correct for baseline ADHD symptom severity.

Limitations

Some results barely missed statistical significance after multiplicity adjustment but were deemed relevant in relation to others and thus further discussed. Despite correcting for in-scanner head movement in multiple processing steps, residual artifacts are likely present in the data, as can be concluded from the correlation with the RS models (see Figure S1). However, since the PerAF findings survived an additional control for the quantified motion on group-level, in-scanner movements unlikely are the cause of our findings. The transfer run could only be reasonably acquired in the NF group and after the training, so all conclusions drawn from the data are necessarily purely correlational. We also kept the acquisition as short as possible expecting increasing movement and decreasing attention throughout each session. A general limitation of transfer runs is that it is usually not possible to check whether participants really apply the strategies learned. In addition, an influence of the different setting for the fMRI-adapted transfer run compared to the SCP version cannot be excluded. Concurrent EEG recordings would have been of little use in this case since differences between training and transfer were previously shown in our study population (Konicar et al., 2021).

Conclusion

Increases in resting activity in regions known to be affected in ASD as well as improvements in several affective domains over both treatment groups indicate unspecific positive effects. In addition, the affective improvements correlated with the activation corresponding to SCP regulation during a transfer run in comparable regions. These were however largely unrelated to the achieved degree of SCP regulation during NF training. Besides corroborating the role of regional alterations and affective functioning in ASD, our findings might suggest that what actually leads to affective symptom improvements is the application of distinct NF regulation

strategies, rather than SCP NF itself. Future research is needed to clarify the distinct influences of SCP positivity and negativity as well as the role of individual regulation strategies.

Acknowledgements

We are grateful towards the students of the ABC BRAIN LAB, who supported us during this study, our participants and their parents. Furthermore, we would like to thank the psychologists, psychiatrists and therapists, social workers and hospital staff of the Department of Child and Adolescent psychiatry of the Medical University of Vienna. We thank Carolin Mößnang for sharing the protocol for the fMRI sequence used.

CRediT author statement

Manfred Klöbl: methodology, software, formal analysis, investigation, data curation, writing – original draft, visualization

Karin Prillinger: investigation, data curation, writing – review & editing, project administration

Robert Diehm: investigation, writing – review & editing

Kamer Doganay: investigation, writing – review & editing

Rupert Lanzenberger: conceptualization, methodology, resources, writing – review & editing, supervision, project administration

Luise Poustka: conceptualization, writing – review & editing, supervision

Paul Plener: resources, supervision, writing – review & editing

Lilian Konicar: conceptualization, methodology, investigation, resources, data curation, writing – review & editing, supervision, project administration, funding acquisition

Funding

This project was funded by the Austrian Science Fund (FWF): KLI600B27.

Conflicts of interest

Rupert Lanzenberger received travel grants and/or conference speaker honoraria within the last three years from Bruker BioSpin MR and Heel, and has served as a consultant for Ono Pharmaceutical. He received investigator-initiated research funding from Siemens Healthcare regarding clinical research using PET/MR. He is a shareholder of the start-up company BM Health GmbH since 2019.

References

Albrecht, J. S., Bubenzer-Busch, S., Gallien, A., Knospe, E. L., Gaber, T. J., & Zepf, F. D. (2017). Effects of a structured 20-session slow-cortical-potential-based neurofeedback

program on attentional performance in children and adolescents with attention-deficit hyperactivity disorder: retrospective analysis of an open-label pilot-approach and 6-month follow-up. *Neuropsychiatr Dis Treat, 13*, 667-683. doi:10.2147/ndt.s119694

Amatachaya, A., Auvichayapat, N., Patjanasoontorn, N., Suphakunpinyo, C., Ngernyam, N., Aree-Uea, B., . . . Auvichayapat, P. (2014). Effect of anodal transcranial direct current stimulation on autism: a randomized double-blind crossover trial. *Behav Neurol, 2014*, 173073. doi:10.1155/2014/173073

Amatachaya, A., Jensen, M. P., Patjanasoontorn, N., Auvichayapat, N., Suphakunpinyo, C., Janjarasjitt, S., . . . Auvichayapat, P. (2015). The short-term effects of transcranial direct current stimulation on electroencephalography in children with autism: a randomized crossover controlled trial. *Behav Neurol, 2015*, 928631. doi:10.1155/2015/928631

Anninos, P., Chatzimichael, A., Adamopoulos, A., Kotini, A., & Tsagas, N. (2016). A combined study of MEG and pico-Tesla TMS on children with autism disorder. *J Integr Neurosci, 15*(4), 497-513. doi:10.1142/s0219635216500278

Arns, M., Heinrich, H., & Strehl, U. (2014). Evaluation of neurofeedback in ADHD: The long and winding road. *Biological Psychology, 95*, 108-115. doi:https://doi.org/10.1016/j.biopsycho.2013.11.013

Beall, E. B., & Lowe, M. J. (2007). Isolating physiologic noise sources with independently determined spatial measures. *Neuroimage, 37*(4), 1286-1300. doi:10.1016/j.neuroimage.2007.07.004

Beall, E. B., & Lowe, M. J. (2014). SimPACE: generating simulated motion corrupted BOLD data with synthetic-navigated acquisition for the development and evaluation of SLOMOCO: a new, highly effective slicewise motion correction. *Neuroimage, 101*, 21-34. doi:10.1016/j.neuroimage.2014.06.038

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage, 37*(1), 90-101. doi:10.1016/j.neuroimage.2007.04.042

Birbaumer, N., Elbert, T., Canavan, A. G., & Rockstroh, B. (1990). Slow potentials of the cerebral cortex and behavior. *Physiological Reviews, 70*(1), 1-41. doi:10.1152/physrev.1990.70.1.1

Blefari, M. L., Martuzzi, R., Salomon, R., Bello-Ruiz, J., Herbelin, B., Serino, A., & Blanke, O. (2017). Bilateral Rolandic operculum processing underlying heartbeat awareness reflects changes in bodily self-consciousness. *Eur J Neurosci, 45*(10), 1300-1312. doi:10.1111/ejn.13567

Bölte, S., & Poustka, F. (2008). Skala zur Erfassung sozialer Reaktivität (SRS). In *Deutsche Fassung der Social Responsiveness Scale (SRS) von John N. Constantino und Christian P. Gruber*: Testzentrale/Huber, Bern.

Bölte, S., Poustka, F., Rühl, D., & Schmötzer, G. (2006). Diagnostisches Interview für Autismus-Revidiert (ADI-R). In: Huber/Hogrefe Verlag, Bern.

Bralten, J., Greven, C. U., Franke, B., Mennes, M., Zwiers, M. P., Rommelse, N. N., . . . Buitelaar, J. K. (2016). Voxel-based morphometry analysis reveals frontal brain differences in participants with ADHD and their unaffected siblings. *J Psychiatry Neurosci, 41*(4), 272-279. doi:10.1503/jpn.140377

Brody, S., Rau, H., Köhler, F., Schupp, H., Lutzenberger, W., & Birbaumer, N. (1994). Slow cortical potential biofeedback and the startle reflex. *Biofeedback Self Regul, 19*(1), 1-11. doi:10.1007/bf01720666

Bush, G., Frazier, J. A., Rauch, S. L., Seidman, L. J., Whalen, P. J., Jenike, M. A., . . . Biederman, J. (1999). Anterior cingulate cortex dysfunction in attention-deficit/hyperactivity

disorder revealed by fMRI and the counting stroop. *Biol Psychiatry, 45*(12), 1542-1552. doi:https://doi.org/10.1016/S0006-3223(99)00083-9

Cai, R. Y., Richdale, A. L., Uljarević, M., Dissanayake, C., & Samson, A. C. (2018). Emotion regulation in autism spectrum disorder: Where we are and where we need to go. *Autism research : official journal of the International Society for Autism Research, 11*(7), 962-978. doi:10.1002/aur.1968

Carper, R. A., Solders, S., Treiber, J. M., Fishman, I., & Müller, R.-A. (2015). Corticospinal Tract Anatomy and Functional Connectivity of Primary Motor Cortex in Autism. *Journal of the American Academy of Child & Adolescent Psychiatry, 54*(10), 859-867. doi:https://doi.org/10.1016/j.jaac.2015.07.007

Chen, G., Cox, R. W., Glen, D. R., Rajendra, J. K., Reynolds, R. C., & Taylor, P. A. (2019). A tail of two sides: Artificially doubled false positive rates in neuroimaging due to the sidedness choice with t-tests. *Hum Brain Mapp, 40*(3), 1037-1043. doi:10.1002/hbm.24399

Christiansen, H., Hirsch, O., Albrecht, B., & Chavanon, M. L. (2019). Attention-Deficit/Hyperactivity Disorder (ADHD) and Emotion Regulation Over the Life Span. *Curr Psychiatry Rep, 21*(3), 17. doi:10.1007/s11920-019-1003-6

Correll, C. U., Cortese, S., Croatto, G., Monaco, F., Krinitski, D., Arrondo, G., . . . Solmi, M. (2021). Efficacy and acceptability of pharmacological, psychosocial, and brain stimulation interventions in children and adolescents with mental disorders: an umbrella review. *World Psychiatry, 20*(2), 244-275. doi:10.1002/wps.20881

Dadds, M. R., Hunter, K., Hawes, D. J., Frost, A. D. J., Vassallo, S., Bunn, P., . . . Masry, Y. E. (2008). A Measure of Cognitive and Affective Empathy in Children Using Parent Ratings. *Child Psychiatry and Human Development, 39*(2), 111-122. doi:10.1007/s10578-007-0075-4

Di Martino, A., Yan, C. G., Li, Q., Denio, E., Castellanos, F. X., Alaerts, K., . . . Milham, M. P. (2014). The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. *Mol Psychiatry, 19*(6), 659-667. doi:10.1038/mp.2013.78

Direito, B., Lima, J., Simões, M., Sayal, A., Sousa, T., Lührs, M., . . . Castelo-Branco, M. (2019). Targeting dynamic facial processing mechanisms in superior temporal sulcus using a novel fMRI neurofeedback target. *Neuroscience, 406*, 97-108. doi:10.1016/j.neuroscience.2019.02.024

Direito, B., Mouga, S., Sayal, A., Simões, M., Quental, H., Bernardino, I., . . . Castelo Branco, M. (2021). Training the social brain: Clinical and neural effects of an 8-week real-time functional magnetic resonance imaging neurofeedback Phase IIa Clinical Trial in Autism. *Autism*, 13623613211002052. doi:10.1177/13623613211002052

Döpfner, M., Görtz-Dorten, A., Lehmkuhl, G., Breuer, D., & Goletz, H. (2008). Diagnostik-System für psychische Störungen nach ICD-10 und DSM-IV für Kinder und Jugendliche - II (DISYPS-II). In: Testzentrale/Huber, Bern.

Doruk Camsari, D., Kirkovski, M., & Croarkin, P. E. (2018). Therapeutic Applications of Noninvasive Neuromodulation in Children and Adolescents. *The Psychiatric clinics of North America, 41*(3), 465-477. doi:10.1016/j.psc.2018.05.003

Doss, M. K., Považan, M., Rosenberg, M. D., Sepeda, N. D., Davis, A. K., Finan, P. H., . . . Barrett, F. S. (2021). Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder. *Transl Psychiatry, 11*(1), 574. doi:10.1038/s41398-021-01706-y

Drechsler, R., Straub, M., Doehnert, M., Heinrich, H., Steinhausen, H. C., & Brandeis, D. (2007). 1Controlled evaluation of a neurofeedback training of slow cortical potentials in

children with Attention Deficit/Hyperactivity Disorder (ADHD). *Behav Brain Funct, 3*, 35. doi:10.1186/1744-9081-3-35

Duan, Y., Zhao, W., Luo, C., Liu, X., Jiang, H., Tang, Y., . . . Yao, D. (2022). Identifying and Predicting Autism Spectrum Disorder Based on Multi-Site Structural MRI With Machine Learning. *Frontiers in human neuroscience, 15*. doi:10.3389/fnhum.2021.765517

Duffield, T. C., Trontel, H. G., Bigler, E. D., Froehlich, A., Prigge, M. B., Travers, B., . . . Lainhart, J. (2013). Neuropsychological investigation of motor impairments in autism. *Journal of clinical and experimental neuropsychology, 35*(8), 867-881. doi:10.1080/13803395.2013.827156

Eldeeb, S., Susam, B. T., Akcakaya, M., Conner, C. M., White, S. W., & Mazefsky, C. A. (2021). Trial by trial EEG based BCI for distress versus non distress classification in individuals with ASD. *Sci Rep, 11*(1), 6000. doi:10.1038/s41598-021-85362-8

Famitafreshi, H., & Karimian, M. (2018). Overview of the Recent Advances in Pathophysiology and Treatment for Autism. *CNS & Neurological Disorders - Drug Targets, 17*(8), 590-594. doi:http://dx.doi.org/10.2174/1871527317666180706141654

Friedrich, E. V., Sivanathan, A., Lim, T., Suttie, N., Louchart, S., Pillen, S., & Pineda, J. A. (2015). An Effective Neurofeedback Intervention to Improve Social Interactions in Children with Autism Spectrum Disorder. *J Autism Dev Disord, 45*(12), 4084-4100. doi:10.1007/s10803-015-2523-5

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement-related effects in fMRI time-series. *Magn Reson Med, 35*(3), 346-355. doi:10.1002/mrm.1910350312

Gebauer, L., Skewes, J., Hørlyck, L., & Vuust, P. (2014). Atypical perception of affective prosody in Autism Spectrum Disorder. *NeuroImage: Clinical, 6*, 370-378. doi:https://doi.org/10.1016/j.nicl.2014.08.025

Gebauer, L., Skewes, J., Westphael, G., Heaton, P., & Vuust, P. (2014). Intact brain processing of musical emotions in autism spectrum disorder, but more cognitive load and arousal in happy vs. sad music. *Frontiers in Neuroscience, 8*(192). doi:10.3389/fnins.2014.00192

Gendry Meresse, I., Zilbovicius, M., Boddaert, N., Robel, L., Philippe, A., Sfaello, I., . . . Chabane, N. (2005). Autism severity and temporal lobe functional abnormalities. *Ann Neurol, 58*(3), 466-469. doi:10.1002/ana.20597

Hallquist, M. N., Hwang, K., & Luna, B. (2013). The nuisance of nuisance regression: spectral misspecification in a common approach to resting-state fMRI preprocessing reintroduces noise and obscures functional connectivity. *Neuroimage, 82*, 208-225. doi:10.1016/j.neuroimage.2013.05.116

Hasslinger, J., D'Agostini Souto, M., Folkesson Hellstadius, L., & Bölte, S. (2020). Neurofeedback in ADHD: A qualitative study of strategy use in slow cortical potential training. *PLOS ONE, 15*(6), e0233343. doi:10.1371/journal.pone.0233343

Haznedar, M. M., Buchsbaum, M. S., Metzger, M., Solimando, A., Spiegel-Cohen, J., & Hollander, E. (1997). Anterior cingulate gyrus volume and glucose metabolism in autistic disorder. *Am J Psychiatry, 154*(8), 1047-1050. doi:10.1176/ajp.154.8.1047

He, B. J., & Raichle, M. E. (2009). The fMRI signal, slow cortical potential and consciousness. *Trends in Cognitive Sciences, 13*(7), 302-309. doi:https://doi.org/10.1016/j.tics.2009.04.004

Heinrich, H., Gevensleben, H., Becker, A., & Rothenberger, A. (2020). Effects of neurofeedback on the dysregulation profile in children with ADHD: SCP NF meets SDQ-DP - a retrospective analysis. *Psychol Med, 50*(2), 258-263. doi:10.1017/s0033291718004130

Helfer, B., Boxhoorn, S., Songa, J., Steel, C., Maltezos, S., & Asherson, P. (2021). Emotion recognition and mind wandering in adults with attention deficit hyperactivity disorder

or autism spectrum disorder. *J Psychiatr Res, 134*, 89-96. doi:10.1016/j.jpsychires.2020.12.059

Hinterberger, T., Veit, R., Strehl, U., Trevorrow, T., Erb, M., Kotchoubey, B., . . . Birbaumer, N. (2003). Brain areas activated in fMRI during self-regulation of slow cortical potentials (SCPs). *Experimental Brain Research, 152*(1), 113-122. doi:10.1007/s00221-003-1515-4

Holtmann, M., Steiner, S., Hohmann, S., Poustka, L., Banaschewski, T., & Bölte, S. (2011). Neurofeedback in autism spectrum disorders. *Dev Med Child Neurol, 53*(11), 986-993. doi:10.1111/j.1469-8749.2011.04043.x

Hurt, E., Arnold, L. E., & Lofthouse, N. (2014). Quantitative EEG Neurofeedback for the Treatment of Pediatric Attention-Deficit/Hyperactivity Disorder, Autism Spectrum Disorders, Learning Disorders, and Epilepsy. *Child and Adolescent Psychiatric Clinics of North America, 23*(3), 465-486. doi:https://doi.org/10.1016/j.chc.2014.02.001

Indefrey, P., Brown, C. M., Hellwig, F., Amunts, K., Herzog, H., Seitz, R. J., & Hagoort, P. (2001). A neural correlate of syntactic encoding during speech production. *Proceedings of the National Academy of Sciences, 98*(10), 5933-5936. doi:10.1073/pnas.101118098

Jia, X.-Z., Sun, J.-W., Ji, G.-J., Liao, W., Lv, Y.-T., Wang, J., . . . Zang, Y.-F. (2020). Percent amplitude of fluctuation: A simple measure for resting-state fMRI signal at single voxel level. *PLOS ONE, 15*(1), e0227021. doi:10.1371/journal.pone.0227021

Jiang, L., & Zuo, X.-N. (2016). Regional Homogeneity: A Multimodal, Multiscale Neuroimaging Marker of the Human Connectome. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry, 22*(5), 486-505. doi:10.1177/1073858415595004

Keller, M., Zweerings, J., Klasen, M., Zvyagintsev, M., Iglesias, J., Mendoza Quiñones, R., & Mathiak, K. (2021). fMRI Neurofeedback-Enhanced Cognitive Reappraisal Training in Depression: A Double-Blind Comparison of Left and Right vlPFC Regulation. *Front Psychiatry, 12*, 715898. doi:10.3389/fpsyt.2021.715898

Khader, P., Schicke, T., Röder, B., & Rösler, F. (2008). On the relationship between slow cortical potentials and BOLD signal changes in humans. *International Journal of Psychophysiology, 67*(3), 252-261. doi:https://doi.org/10.1016/j.ijpsycho.2007.05.018

Khaleghi, A., Zarafshan, H., Vand, S. R., & Mohammadi, M. R. (2020). Effects of Non-invasive Neurostimulation on Autism Spectrum Disorder: A Systematic Review. *Clin Psychopharmacol Neurosci, 18*(4), 527-552. doi:10.9758/cpn.2020.18.4.527

King, J. B., Prigge, M. B. D., King, C. K., Morgan, J., Weathersby, F., Fox, J. C., . . . Anderson, J. S. (2019). Generalizability and reproducibility of functional connectivity in autism. *Mol Autism, 10*, 27. doi:10.1186/s13229-019-0273-5

Klöbl, M., Michenthaler, P., Godbersen, G. M., Robinson, S., Hahn, A., & Lanzenberger, R. (2020). Reinforcement and Punishment Shape the Learning Dynamics in fMRI Neurofeedback. *Frontiers in human neuroscience, 14*, 304-304. doi:10.3389/fnhum.2020.00304

Konicar, L., Radev, S., Prillinger, K., Klöbl, M., Diehm, R., Birbaumer, N., . . . Poustka, L. (2021). Volitional modification of brain activity in adolescents with Autism Spectrum Disorder: A Bayesian analysis of Slow Cortical Potential neurofeedback. *Neuroimage Clin, 29*, 102557. doi:10.1016/j.nicl.2021.102557

Krakowski, A. D., Cost, K. T., Anagnostou, E., Lai, M.-C., Crosbie, J., Schachar, R., . . . Szatmari, P. (2020). Inattention and hyperactive/impulsive component scores do not differentiate between autism spectrum disorder and attention-deficit/hyperactivity disorder in a clinical sample. *Mol Autism, 11*(1), 28. doi:10.1186/s13229-020-00338-1

Lai, G., Pantazatos, S. P., Schneider, H., & Hirsch, J. (2012). Neural systems for speech and song in autism. *Brain, 135*(Pt 3), 961-975. doi:10.1093/brain/awr335

Laidi, C., Boisgontier, J., de Pierrefeu, A., Duchesnay, E., Hotier, S., d'Albis, M.-A., . . . Houenou, J. (2019). Decreased Cortical Thickness in the Anterior Cingulate Cortex in Adults with Autism. *J Autism Dev Disord, 49*(4), 1402-1409. doi:10.1007/s10803-018-3807-3

Lan, Z., Xu, S., Wu, Y., Xia, L., Hua, K., Li, M., . . . Wang, T. (2021). Alterations of Regional Homogeneity in Preschool Boys With Autism Spectrum Disorders. *Front Neurosci, 15*, 644543. doi:10.3389/fnins.2021.644543

Leitner, Y. (2014). The co-occurrence of autism and attention deficit hyperactivity disorder in children - what do we know? *Frontiers in human neuroscience, 8*, 268. doi:10.3389/fnhum.2014.00268

Loevaas, M. E. S., Sund, A. M., Patras, J., Martinsen, K., Hjemdal, O., Neumer, S. P., . . . Reinfjell, T. (2018). Emotion regulation and its relation to symptoms of anxiety and depression in children aged 8–12 years: does parental gender play a differentiating role? *BMC Psychology, 6*(1), 42. doi:10.1186/s40359-018-0255-y

Mahajan, R., Dirlikov, B., Crocetti, D., & Mostofsky, S. H. (2016). Motor Circuit Anatomy in Children with Autism Spectrum Disorder With or Without Attention Deficit Hyperactivity Disorder. *Autism research : official journal of the International Society for Autism Research, 9*(1), 67-81. doi:10.1002/aur.1497

Mayer, K., Blume, F., Wyckoff, S. N., Brokmeier, L. L., & Strehl, U. (2016). Neurofeedback of slow cortical potentials as a treatment for adults with Attention Deficit-/Hyperactivity Disorder. *Clin Neurophysiol, 127*(2), 1374-1386. doi:10.1016/j.clinph.2015.11.013

Mayer, K., Wyckoff, S. N., & Strehl, U. (2013). One Size Fits All? Slow Cortical Potentials Neurofeedback:A Review. *Journal of Attention Disorders, 17*(5), 393-409. doi:10.1177/1087054712468053

Mazza, M., Pino, M. C., Mariano, M., Tempesta, D., Ferrara, M., De Berardis, D., . . . Valenti, M. (2014). Affective and cognitive empathy in adolescents with autism spectrum disorder. *Frontiers in human neuroscience, 8*, 791. doi:10.3389/fnhum.2014.00791

McKenzie, K., Russell, A., Golm, D., & Fairchild, G. (2021). Empathic Accuracy and Cognitive and Affective Empathy in Young Adults With and Without Autism Spectrum Disorder. *J Autism Dev Disord*. doi:10.1007/s10803-021-05093-7

McPartland, J., & Volkmar, F. R. (2012). Chapter 23 - Autism and related disorders. In M. J. Aminoff, F. Boller, & D. F. Swaab (Eds.), *Handbook of Clinical Neurology* (Vol. 106, pp. 407-418): Elsevier.

Moessnang, C., Schäfer, A., Bilek, E., Roux, P., Otto, K., Baumeister, S., . . . Tost, H. (2016). Specificity, reliability and sensitivity of social brain responses during spontaneous mentalizing. *Soc Cogn Affect Neurosci, 11*(11), 1687-1697. doi:10.1093/scan/nsw098

Mohammad-Rezazadeh, I., Frohlich, J., Loo, S. K., & Jeste, S. S. (2016). Brain connectivity in autism spectrum disorder. *Curr Opin Neurol, 29*(2), 137-147. doi:10.1097/wco.0000000000000301

Mueller, K., Lepsien, J., Möller, H. E., & Lohmann, G. (2017). Commentary: Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Frontiers in human neuroscience, 11*, 345-345. doi:10.3389/fnhum.2017.00345

Nebel, M. B., Eloyan, A., Barber, A. D., & Mostofsky, S. H. (2014). Precentral gyrus functional connectivity signatures of autism. *Front Syst Neurosci, 8*, 80. doi:10.3389/fnsys.2014.00080

Neeley, E. S., Bigler, E. D., Krasny, L., Ozonoff, S., McMahon, W., & Lainhart, J. E. (2007). Quantitative temporal lobe differences: autism distinguished from controls using classification and regression tree analysis. *Brain Dev, 29*(7), 389-399. doi:10.1016/j.braindev.2006.11.006

Olszowy, W., Aston, J., Rua, C., & Williams, G. B. (2019). Accurate autocorrelation modeling substantially improves fMRI reliability. *Nature communications, 10*(1), 1220-1220. doi:10.1038/s41467-019-09230-w

Panerai, S., Tasca, D., Lanuzza, B., Trubia, G., Ferri, R., Musso, S., . . . Elia, M. (2014). Effects of repetitive transcranial magnetic stimulation in performing eye-hand integration tasks: four preliminary studies with children showing low-functioning autism. *Autism, 18*(6), 638-650. doi:10.1177/1362361313495717

Patel, A. X., Kundu, P., Rubinov, M., Jones, P. S., Vértes, P. E., Ersche, K. D., . . . Bullmore, E. T. (2014). A wavelet method for modeling and despiking motion artifacts from resting-state fMRI time series. *Neuroimage, 95*(100), 287-304. doi:10.1016/j.neuroimage.2014.03.012

Poustka, L., Rühl, D., Feineis-Matthews, S., Poustka, F., Hartung, M., & Bölte, S. (2015). ADOS-2. Diagnostische Beobachtungsskala für Autistische Störungen. In *2. Deutschsprachige Fassung der Autism Diagnostic Observation Schedule*: Testzentrale/Huber, Mannheim.

Preston, S. D., Bechara, A., Damasio, H., Grabowski, T. J., Stansfield, R. B., Mehta, S., & Damasio, A. R. (2007). The neural substrates of cognitive empathy. *Soc Neurosci, 2*(3-4), 254-275. doi:10.1080/17470910701376902

Prillinger, K., Radev, S. T., Amador de Lara, G., Klobl, M., Lanzenberger, R., Plener, P. L., . . . Konicar, L. (2021). Repeated Sessions of Transcranial Direct Current Stimulation on Adolescents With Autism Spectrum Disorder: Study Protocol for a Randomized, Double-Blind, and Sham-Controlled Clinical Trial. *Front Psychiatry, 12*, 680525. doi:10.3389/fpsyt.2021.680525

Prillinger, K., Radev, S. T., Doganay, K., Poustka, L., & Konicar, L. (2022). Impulsivity Moderates the Effect of Neurofeedback Training on the Contingent Negative Variation in Autism Spectrum Disorder. *Frontiers in human neuroscience*.

Qi, S., Morris, R., Turner, J. A., Fu, Z., Jiang, R., Deramus, T. P., . . . Sui, J. (2020). Common and unique multimodal covarying patterns in autism spectrum disorder subtypes. *Mol Autism, 11*(1), 90. doi:10.1186/s13229-020-00397-4

Reyes, N. M., Pickard, K., & Reaven, J. (2019). Emotion regulation: A treatment target for autism spectrum disorder. *Bull Menninger Clin, 83*(3), 205-234. doi:10.1521/bumc.2019.83.3.205

Riva, D., Annunziata, S., Contarino, V., Erbetta, A., Aquino, D., & Bulgheroni, S. (2013). Gray Matter Reduction in the Vermis and CRUS-II Is Associated with Social and Interaction Deficits in Low-Functioning Children with Autistic Spectrum Disorders: a VBM-DARTEL Study. *The Cerebellum, 12*(5), 676-685. doi:10.1007/s12311-013-0469-8

Ros, R., & Graziano, P. A. (2020). A Transdiagnostic Examination of Self-Regulation: Comparisons Across Preschoolers with ASD, ADHD, and Typically Developing Children. *J Clin Child Adolesc Psychol, 49*(4), 493-508. doi:10.1080/15374416.2019.1591280

Sankoh, A. J., Huque, M. F., & Dubey, S. D. (1997). Some comments on frequently used multiple endpoint adjustment methods in clinical trials. *Statistics in Medicine, 16*(22), 2529-2542. doi:https://doi.org/10.1002/(SICI)1097-0258(19971130)16:22<2529::AID-SIM692>3.0.CO;2-J

Schabus, M., Griessenberger, H., Gnjezda, M. T., Heib, D. P. J., Wislowska, M., & Hoedlmoser, K. (2017). Better than sham? A double-blind placebo-controlled neurofeedback study in primary insomnia. *Brain, 140*(4), 1041-1052. doi:10.1093/brain/awx011

Schwartz, J., Ordaz, S. J., Kircanski, K., Ho, T. C., Davis, E. G., Camacho, M. C., & Gotlib, I. H. (2019). Resting-state functional connectivity and inflexibility of daily emotions in major depression. *J Affect Disord, 249*, 26-34. doi:10.1016/j.jad.2019.01.040

Seehausen, M., Kazzer, P., Bajbouj, M., Heekeren, H. R., Jacobs, A. M., Klann-Delius, G., . . . Prehn, K. (2016). Effects of empathic social responses on the emotions of the recipient. *Brain Cogn, 103*, 50-61. doi:10.1016/j.bandc.2015.11.004

Seymour, K. E., Chronis-Tuscano, A., Iwamoto, D. K., Kurdziel, G., & MacPherson, L. (2014). Emotion Regulation Mediates the Association Between ADHD and Depressive Symptoms in a Community Sample of Youth. *Journal of Abnormal Child Psychology, 42*(4), 611-621. doi:10.1007/s10802-013-9799-8

Shaw, P., Stringaris, A., Nigg, J., & Leibenluft, E. (2014). Emotion dysregulation in attention deficit hyperactivity disorder. *Am J Psychiatry, 171*(3), 276-293. doi:10.1176/appi.ajp.2013.13070966

Shields, A., & Cicchetti, D. (1997). Emotion regulation among school-age children: the development and validation of a new criterion Q-sort scale. *Dev Psychol, 33*(6), 906-916. doi:10.1037//0012-1649.33.6.906

Sonuga-Barke, E. J., Brandeis, D., Cortese, S., Daley, D., Ferrin, M., Holtmann, M., . . . Sergeant, J. (2013). Nonpharmacological interventions for ADHD: systematic review and meta-analyses of randomized controlled trials of dietary and psychological treatments. *Am J Psychiatry, 170*(3), 275-289. doi:10.1176/appi.ajp.2012.12070991

Strehl, U., Aggensteiner, P., Wachtlin, D., Brandeis, D., Albrecht, B., Arana, M., . . . Holtmann, M. (2017). Neurofeedback of Slow Cortical Potentials in Children with Attention-Deficit/Hyperactivity Disorder: A Multicenter Randomized Trial Controlling for Unspecific Effects. *Frontiers in human neuroscience, 11*(135). doi:10.3389/fnhum.2017.00135

Strehl, U., Leins, U., Goth, G., Klinger, C., Hinterberger, T., & Birbaumer, N. (2006). Self-regulation of Slow Cortical Potentials: A New Treatment for Children With Attention-Deficit/Hyperactivity Disorder. *Pediatrics, 118*(5), e1530-e1540. doi:10.1542/peds.2005-2478

Strehl, U., Trevorrow, T., Veit, R., Hinterberger, T., Kotchoubey, B., Erb, M., & Birbaumer, N. (2006). Deactivation of brain areas during self-regulation of slow cortical potentials in seizure patients. *Appl Psychophysiol Biofeedback, 31*(1), 85-94. doi:10.1007/s10484-006-9006-6

Sutoko, S., Atsumori, H., Obata, A., Funane, T., Kandori, A., Shimonaga, K., . . . Tsuji, T. (2020). Lesions in the right Rolandic operculum are associated with self-rating affective and apathetic depressive symptoms for post-stroke patients. *Sci Rep, 10*(1), 20264. doi:10.1038/s41598-020-77136-5

Tagliazucchi, E., Roseman, L., Kaelen, M., Orban, C., Muthukumaraswamy, S. D., Murphy, K., . . . Carhart-Harris, R. (2016). Increased Global Functional Connectivity Correlates with LSD-Induced Ego Dissolution. *Curr Biol, 26*(8), 1043-1050. doi:10.1016/j.cub.2016.02.010

Taurines, R., Schwenck, C., Westerwald, E., Sachse, M., Siniatchkin, M., & Freitag, C. (2012). ADHD and autism: differential diagnosis or overlapping traits? A selective review. *ADHD Attention Deficit and Hyperactivity Disorders, 4*(3), 115-139. doi:10.1007/s12402-012-0086-2

Thakkar, K. N., Polli, F. E., Joseph, R. M., Tuch, D. S., Hadjikhani, N., Barton, J. J., & Manoach, D. S. (2008). Response monitoring, repetitive behaviour and anterior cingulate abnormalities in autism spectrum disorders (ASD). *Brain, 131*(Pt 9), 2464-2478. doi:10.1093/brain/awn099

Toller, G., Adhimoolam, B., Rankin, K. P., Huppertz, H. J., Kurthen, M., & Jokeit, H. (2015). Right fronto-limbic atrophy is associated with reduced empathy in refractory unilateral mesial temporal lobe epilepsy. *Neuropsychologia, 78,* 80-87. doi:10.1016/j.neuropsychologia.2015.09.010

Vaidya, C. J., You, X., Mostofsky, S., Pereira, F., Berl, M. M., & Kenworthy, L. (2020). Data-driven identification of subtypes of executive function across typical development, attention deficit hyperactivity disorder, and autism spectrum disorders. *J Child Psychol Psychiatry, 61*(1), 51-61. doi:10.1111/jcpp.13114

Waddington, F., Franke, B., Hartman, C., Buitelaar, J. K., Rommelse, N., & Mota, N. R. (2021). A polygenic risk score analysis of ASD and ADHD across emotion recognition subtypes. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 186*(7), 401-411. doi:https://doi.org/10.1002/ajmg.b.32818

Waddington, F., Hartman, C., de Bruijn, Y., Lappenschaar, M., Oerlemans, A., Buitelaar, J., . . . Rommelse, N. (2018). An emotion recognition subtyping approach to studying the heterogeneity and comorbidity of autism spectrum disorders and attention-deficit/hyperactivity disorder. *Journal of Neurodevelopmental Disorders, 10*(1), 31. doi:10.1186/s11689-018-9249-6

Wang, W., Liu, J., Shi, S., Liu, T., Ma, L., Ma, X., . . . Wang, M. (2018). Altered Resting-State Functional Activity in Patients With Autism Spectrum Disorder: A Quantitative Meta-Analysis. *Front Neurol, 9,* 556. doi:10.3389/fneur.2018.00556

Wang, Z., Wang, Y., Sweeney, J. A., Gong, Q., Lui, S., & Mosconi, M. W. (2019). Resting-State Brain Network Dysfunctions Associated With Visuomotor Impairments in Autism Spectrum Disorder. *Frontiers in Integrative Neuroscience, 13*(17). doi:10.3389/fnint.2019.00017

Wechsler, D., & Petermann, F. (2012). Wechsler Adult Intelligence Scale – Fourth Edition. In *Deutschsprachige Adaption der WAIS-IV von D. Wechsler*: Pearson/Testzentrale.

Wechsler, D., Petermann, F., & Petermann, U. (2010). Hamburg-Wechsler-Intelligenztest für Kinder – IV (HAWIK-IV). In: Pearson/Testzentrale.

Wilke, M., Altaye, M., & Holland, S. K. (2017). CerebroMatic: A Versatile Toolbox for Spline-Based MRI Template Creation. *Front Comput Neurosci, 11,* 5. doi:10.3389/fncom.2017.00005

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A Network Visualization Tool for Human Brain Connectomics. *PLOS ONE, 8*(7), e68910. doi:10.1371/journal.pone.0068910

Yerys, B. E., Herrington, J. D., Satterthwaite, T. D., Guy, L., Schultz, R. T., & Bassett, D. S. (2017). Globally weaker and topologically different: resting-state connectivity in youth with autism. *Mol Autism, 8,* 39-39. doi:10.1186/s13229-017-0156-6

Zang, Y., Jiang, T., Lu, Y., He, Y., & Tian, L. (2004). Regional homogeneity approach to fMRI data analysis. *Neuroimage, 22*(1), 394-400. doi:https://doi.org/10.1016/j.neuroimage.2003.12.030

Zheng, H., Li, F., Bo, Q., Li, X., Yao, L., Yao, Z., . . . Wu, X. (2018). The dynamic characteristics of the anterior cingulate cortex in resting-state fMRI of patients with depression. *J Affect Disord, 227,* 391-397. doi:10.1016/j.jad.2017.11.026

Ziermans, T., de Bruijn, Y., Dijkhuis, R., Staal, W., & Swaab, H. (2019). Impairments in cognitive empathy and alexithymia occur independently of executive functioning in college students with autism. *Autism, 23*(6), 1519-1530. doi:10.1177/1362361318817716

Zilbovicius, M., Boddaert, N., Belin, P., Poline, J. B., Remy, P., Mangin, J. F., . . . Samson, Y. (2000). Temporal lobe dysfunction in childhood autism: a PET study. Positron emission tomography. *Am J Psychiatry, 157*(12), 1988-1993. doi:10.1176/appi.ajp.157.12.1988