

## 1 Full title

2 Effects of virtual reality-based aerobic endurance training on the functional fitness of  
3 healthy older adults: A systematic review

## 4 Short title

5 Aerobic endurance training through the use of virtual reality and functional fitness in  
6 healthy older adults.

## 7 Authors

8 Daniela Ramírez Restrepo,<sup>1,3\*</sup> Julialba Castellanos Ruiz,<sup>1¶</sup> Lina María Montealegre  
9 Mesa,<sup>1¶</sup> Carolina Márquez Narváez,<sup>2&</sup> Santiago Murillo Rendón<sup>2&</sup>

- 10  
11  
12  
13 1. Department of "Cuerpo Movimiento," Universidad Autónoma de Manizales,  
14 Caldas, Colombia.  
15 2. Department of "Ciencias Computacionales," Universidad Autónoma de  
16 Manizales, Caldas, Colombia.  
17 3. Department of "Neuroaprendizaje," Universidad Autónoma de Manizales,  
18 Manizales, Caldas, Colombia.

19  
20 \*Corresponding author

21 E-mail correspondence: [daniela.ramirezr@autonoma.edu.co](mailto:daniela.ramirezr@autonoma.edu.co)

22 ORCID ID:<https://orcid.org/0000-0001-7391-9331>

23  
24  
25 ¶ These authors contributed equally to this work.

26 & These authors also contributed equally to this work.

## 27 Abstract

28  
29 **Objective:** to analytically and systematically review and integrate the available  
30 evidence of the effects of virtual reality-based aerobic endurance training on  
31 functional fitness of healthy older adults. **Materials and Methods** This systematic  
32 review of randomized controlled trials was conducted through searches in Pubmed,  
33 Science Direct, Scopus, PEDro, Web of Science, Lilacs, Scielo, Dialnet, and IEEE  
34 Xplore between February and May 2021. The evaluation of bias and methodological  
35 quality was performed following the parameters proposed by the Cochrane Manual  
36 of Systematic Reviews of Interventions and the PEDro Scale. Review Manager  
37 Software (Version 5.4.1) served for a summary of bias. **Results:** Three clinical trials

39 were selected. None of the selected studies employed any exclusively aerobic  
40 intervention with virtual reality; however, statistically significant findings for this type  
41 of intervention were found in functional fitness such as aerobic endurance, upper  
42 and lower limb strength, agility, and dynamic balance. **Conclusions:** aerobic  
43 endurance training through the use of virtual reality may provide positive effects on  
44 the functional fitness of the healthy elderly. This has implications for clinical practice,  
45 as it fosters the development of more beneficial, attractive, motivational, and quality  
46 interventions, the prevention of common conditions, and the use of technological  
47 tools by the elderly population. However, as for research implications, an increase  
48 of knowledge on the effects of technologies in the elderly functional capacity, and an  
49 implementation of scientific models or theories on physical activity and exercise are  
50 recommended with the purpose of explaining the phenomena due to the use of  
51 technologies within the processes of functional rehabilitation.

52

53 **Keywords:** Exercise, Virtual Reality, Physical Fitness, Aged, Elderly.

54

55

## 56 Introduction

57 The demographic and epidemiological transitions during the last decades have  
58 brought about a significant global increase in the number of people aged 60 years  
59 and older.(1) While in 2000 the percentage of elderly population represented 10% of  
60 the total population, it is estimated that by the year 2030 it will represent 16.6% of  
61 the total population and 22% by the year 2050.(2) A longer life expectancy makes  
62 individuals, particularly the elderly, prone to bodily impairments, functional limitations  
63 and restrictions in social participation. This is due to the fact that, in old age, there  
64 may be a greater number of health, environmental, and socioeconomic risks, as a  
65 result of a longer life trajectory.(3) Additionally, determinants such as unhealthy  
66 lifestyles, along with the physiological changes during the aging process may  
67 increase the probability of different health conditions that involve their physical  
68 activity such as aerobic endurance.(4–6)

69

70 Aerobic endurance is a remarkable component of functional fitness throughout the  
71 lifetime, as it allows for the independent performance of daily life, instrumental, and  
72 social activities with a lower level of fatigue.(7) This physical activity tends to  
73 decrease in older adults due to the cardiovascular and pulmonary diseases during  
74 the aging process.(8,9) Therefore, reduced cardiac contractility, conduction velocity  
75 and baroreceptor sensitivity, decreased cardiac output and blood flow, valvular  
76 insufficiency, increased arterial wall stiffness, decreased thoracic, alveolar and  
77 pulmonary distensibility, loss of valvular and respiratory muscle strength, reduced

78 cilia and cough reflex, among others, can reduce pulse rate during exercise.  
79 Additionally they can reduce cardiac reserve, and increase blood pressure. Reduce  
80 loss of strength of valvular and respiratory muscles, reduction of cilia and cough  
81 reflex, among others, can decrease pulse rate during exercise, reduce cardiac  
82 reserve, increase blood pressure, decrease gas exchange surface area and  
83 increase the probability of heart failure, arrhythmias, other cardiovascular diseases,  
84 infections or respiratory diseases in old age.(10–12) Also, these changes may  
85 significantly affect functional independence in the aforementioned daily life  
86 activities.(7)

87

88 Therefore, aerobic fitness is one of the most recommended strategies to maintain or  
89 improve functional fitness in this population.(13,14) Leirós et al., for example,  
90 demonstrated that aerobic training for the elderly markedly improved functional  
91 fitness such as strength and flexibility of upper and lower limbs, arm strength, and  
92 weight loss and reduced BMI.(15) Similarly, Ribeiro et al. found that water-based  
93 aerobic training showed positive findings in functionality, as well as in some aspects  
94 related to quality of life in women over 65 years of age.(16)

95

96 Great technological advances have allowed for an improvement in the effects and  
97 quality of health care, diagnosis, treatment, and intervention strategies.(17,18) For  
98 functional rehabilitation in the elderly, the incorporation of high technologies for  
99 functional fitness training, as virtual reality, has brought a lot of benefits. Performing  
100 activities in virtual environments, that resemble the real ones, and the visual and  
101 auditory feedback provided by these tools, increase individuals' motivation to  
102 exercise.(19–21) According to the Ecological Model of Physical Activity, it has been  
103 established that, beyond the influence of biological factors such as fitness level,  
104 health status, or body composition, there are psychological and environmental  
105 factors that influence and enhance the physiological effects of exercise.(22–24)

106

107 Although some benefits of conventional aerobic training and some advantages of  
108 virtual reality are already known, few studies have been conducted on the effects of  
109 this type of intervention on the functional fitness of older people, especially those  
110 with stable or healthy conditions. Van der Kolk et al., for example, evaluated the  
111 effectiveness of home and remote training in people between 30 and 75 years of  
112 age with Parkinson's Disease, which included the use of serious games (exergames)  
113 through Virtual Reality software. As results, the authors found improvements in the  
114 motor status of the participants in the experimental group, in contrast to the control  
115 group.(25) Similarly, Karssemeijer et al. evaluated the efficacy of training with  
116 exergames and conventional aerobic training in frail older people with dementia.  
117 There, it was found that the intervention with these serious games, compared to the

118 people in the control group, yielded positive results in reducing the levels of fragility  
119 in the participants of the experimental group.<sup>(26)</sup>

120

121 Thus, the objective of the present study was to analytically and systematically review  
122 and integrate the available evidence of the effects of virtual reality-based aerobic  
123 endurance training on functional fitness of healthy older adults. It is expected that  
124 the collection of information recorded in this research will serve as a point of  
125 reference for future studies in gerontological physiotherapy, especially to promote  
126 the implementation of technological devices in favor of the improvement of  
127 conventional interventions, as well as functionality, quality of life, and social  
128 participation in old age. This research aims to respond to these realities such as  
129 digital evolution, global demographic aging and stereotypes associated with the  
130 inability of older people regarding the use of technologies <sup>(1,27,28)</sup> by encouraging the  
131 transition from traditional schemes of functional rehabilitation in old age toward  
132 schemes of higher quality and demystifying existing social stigmas regarding aging,  
133 old age, and the use of technological tools.

134

135

## 136 **Material and Methods**

### 137 **Type of Study**

138 This systematic review of Randomized Controlled Trials (RCTs) was conducted on  
139 the effects of aerobic resistance training through virtual reality on the functional  
140 physical condition of healthy older people. The PICO strategy was used to guide the  
141 formulation of the research question, where P (patient or target population) = Healthy  
142 Older People; I (intervention) = Aerobic Endurance Training through Virtual Reality,  
143 either exclusively or combined with the training of other physical capacities; C  
144 (comparison or control) = Conventional Aerobic Endurance Training or No  
145 Intervention; OR (outcomes) = Aerobic Endurance (primary outcome measure),  
146 Muscular Strength, Flexibility and Balance (secondary outcome measures).  
147 Subsequently, the following research question established: What are the effects of  
148 aerobic endurance training through virtual reality on the functional physical condition  
149 of healthy older adults?

150

### 151 **Search Strategy**

152 Searches were conducted in PubMed, Science Direct, Scopus, PEDro, Web of  
153 Science, Lilacs, Scielo, Dialnet, and IEEE Xplore databases. The search was  
154 executed up to the month of May 2021 and search equations were applied that  
155 combined the following MeSH terms to identify relevant studies: Aged, Virtual  
156 Reality, Aerobic Exercise, Physical Fitness. Likewise, related terms such as Elderly,

157 Virtual Reality Exposure Therapy, Aerobic Exercises, Aerobic Training were used.  
158 For databases such as Lilacs and Dialnet, searches were also carried out with  
159 equations in the Spanish language, using DeCS terms such as Virtual Reality,  
160 Aerobic Training, Functional Physical Performance, and Elderly. All terms described  
161 were combined with the Boolean operators "AND" and "OR" to perform multiple  
162 combinations and generate several search equations. Finally, the references of  
163 some scientific articles were reviewed to identify if there were relevant studies for  
164 the present research.

165

## 166 **Study Inclusion and Exclusion Criteria**

167 The study selection comprised RCTs in any language, of healthy or with any  
168 controlled chronic health condition people of 60 years or older and that were  
169 conducted during the last 20 years (2001-2021). Other types of studies were  
170 excluded, as well as articles of people with neurological, psychiatric, or mental health  
171 conditions within their target population.

172

## 173 **Procedures**

### 174 **Study Selection**

175 Searches mentioned above in databases were carried out by one researcher, using  
176 the established search equations. First, articles were reviewed according to their  
177 titles and abstracts; then, the full text analysis of the articles that met the inclusion  
178 and exclusion criteria of this research was carried out.

179

### 180 **Extraction and Management Data**

181 The required information was extracted by a researcher. To register the articles of  
182 interest, tables were created in the Review Manager Software (Version 5.4.1)<sup>(29)</sup> that  
183 included data such as the name of the study, authors, year of publication, methods  
184 (type of study, method of intervention allocation, groups, loss to follow-up),  
185 participants (location, time frame of the study or intervention, inclusion and exclusion  
186 criteria, sociodemographic characteristics of the population), prescription  
187 parameters of the interventions of the control groups and experimental (exercise  
188 type/mode, frequency, duration, intensity), and results.

189

### 190 **Assessment of the Bias Risk and Methodological Quality of the** 191 **Articles Included**

192 Evaluation of bias and methodological quality were performed following the  
193 parameters proposed by the Cochrane Manual of Systematic Reviews of  
194 Interventions and the PEDro Scale.<sup>(30,31)</sup> First, selection, performance, detection,  
195 attrition, and reporting biases were considered; this in order to determine,

196 respectively, the systematic differences between: 1) the characteristics of the  
197 comparison groups at the beginning of the study; 2) experimental and control groups  
198 for the intervention of interest; 3) the way in which the results between the  
199 experimental group and the control group were determined; 4) groups regarding  
200 study dropouts; and 5) the findings of the study, both presented and not presented.  
201 Similarly, other biases associated with problems that were not considered in the  
202 previous types of biases were taken into account.<sup>(30)</sup>

203

204 Based on the biases described above, each type of bias in the included studies was  
205 assessed, classifying each as low bias, high bias, or unclear bias<sup>(30)</sup> Second, two  
206 researchers independently evaluated the methodological quality of the studies  
207 included through 11 items of the PEDro Scale, which presented dichotomous options  
208 as alternatives response (If: 1 point. No: 0 points) These items were<sup>(31)</sup>:

209

- 210 1. The selection criteria were specified
- 211 2. Subjects were randomly assigned to groups (in a crossover study, subjects were  
212 randomly distributed as they received treatments).
- 213 3. It was a blinded allocation
- 214 4. The groups were similar at baseline in relation to the most important prognostic  
215 indicators.
- 216 5. All individuals were blinded for treatment.
- 217 6. All therapists were blinded for treatment administration.
- 218 7. All evaluators were blinded for measuring at least one key outcome.
- 219 8. Measures of at least one of the key outcomes were obtained from more than 85%  
220 of the subjects who were initially assigned to the groups.
- 221 9. Results were presented for all subjects who received treatment or were assigned  
222 to the control group; on the contrary case, data for at least one key outcome were  
223 analyzed on an "intention-to-treat" basis.
- 224 10. Results of statistical comparisons between groups were reported for at least one  
225 key outcome.
- 226 11. This study shows point and variability measures for at least one key outcome.

227

228 Finally, these evaluations were socialized together to reach a consensus and, in  
229 cases of disagreements, these were discussed and resolved with a third evaluator.  
230 Similarly, the Cochrane's Review Manager (Version 5.4.1) was used for the creation  
231 of the summary bias analysis.<sup>(25)</sup>

232

233

## 234 **Results:**

### 235 **Study Selection**

236 A total of 241 articles from databases were selected. Then, 74 duplicate articles were  
237 removed, for a total of 167 screening articles. Out of the 167, 164 were excluded  
238 based on title, abstract and inclusion and exclusion criteria, resulting in a final  
239 number of three articles for full-text review and qualitative assessment. The details  
240 of the study selection are depicted in the PRISMA flow chart (Fig. 1)




Figure 1. PRISMA flow diagram.

## Characteristics of the Selected Studies

277 Table 1 shows the characteristics of selected studies. This systematic review  
278 comprised randomized controlled trials that measure the effects of aerobic  
279 endurance training through the use of virtual reality on the functional fitness in the  
280 healthy elderly in comparison with traditional training or absence of therapeutic  
281 intervention. Two of the studies made exclusive use of virtual reality and one of  
282 augmented reality to intervene with the experimental group; although, none of them  
283 intervened with the control group through the use of conventional therapies or  
284 exercises. Similarly, none of the selected studies involved a unique intervention for  
285 aerobic capacity in the target populations. Most of the studies were conducted on  
286 the Asian continent and their samples included a larger number of women than men.

287

## 288 **Analysis and Synthesis of Results**

289 Due to the diversity of the selected studies in terms of sample size, intervention  
290 procedure, and outcome measures, performing a quantitative analysis (meta-  
291 analysis) was not possible. However, a qualitative analysis showed positive results  
292 regarding the effects of aerobic training through the use of virtual reality on the  
293 functional fitness of the healthy elderly.

294

295 Regarding aerobic endurance, only one of the studies showed statistically significant  
296 improvements, after an augmented reality intervention, in the two-minute walk test  
297 of the Senior Fitness Test for the experimental group ( $p=0.020$ ).<sup>(26)</sup> However, for the  
298 other activities that make up functional fitness, this study also showed positive  
299 findings in muscle function ( $p=0.003$ ) and lower limb muscle strength ( $p=0.033$ ), as  
300 well as in agility and dynamic balance ( $p=0.032$ ) of the intervention group.<sup>(26)</sup>

301

302 Similarly, another selected study exhibited a statistically significant improvement in  
303 upper limb strength, agility and dynamic balance for the experimental group after the  
304 virtual reality intervention ( $p\leq 0.05$ ).<sup>(27)</sup> Also, the third selected showed significant  
305 improvements in the gait speed test ( $p\leq 0.05$ ) and in the test of getting up and sitting  
306 in a chair ( $p\leq 0.05$ ), after the intervention of the experimental group by using virtual  
307 environments.<sup>(34)</sup> With respect to this last study, although the walking speed and the  
308 sitting and standing up from a chair test are specific tasks, these require functional  
309 fitness such as muscular strength, mainly in the lower limbs, aerobic capacity and  
310 static and dynamic balance.<sup>(28,29)</sup>



Author/s	Participants (EG) – (CG)	Mean age (years)	Country	EG Intervention (System Used, Type and form of intervention, Intensity, Frequency, Duration)	CG Intervention	Outcome measures
Barsasella et al., 2021	EG = 29 (4 male and 25 female)  CG = 31 (10 male and 21 female)	70 – 74*	Taiwan	<p><b>System:</b> HTC Vive</p> <ul style="list-style-type: none"> <li>•<b>High-intensity movement:</b> engaging in some sport (Apps: Sparc, Final Soccer VR);</li> <li>•<b>Low-intensity movement:</b> climbing Mount Everest (Everest VR), moving with super human abilities (To the Top), walking around in nature (Found), or exploring a place (Google Earth)</li> <li>•<b>Very-low-intensity movement:</b> acting like a magician (Waltz of the Wizard) and exploring the body cells (Body VR).</li> </ul> <p>15 min, 2 times per week, 6 weeks</p>	Daily Routine	<p><b>Health-Related Quality of Life</b> (The EQ-5D-3L questionnaire)</p> <p><b>Happiness</b> (Mini version of the Chinese Happiness Inventory)</p> <p><b>Functional Fitness</b> (Functional Fitness Tests by Rikli and Jones)</p>
Sagwan & Jiyoun, 2020	EG = 13 (Female)  CG = 14 (Female)	72,74 ± 3,64	Korea	<p><b>System:</b> UINCARE-HEALTH TM</p> <p>•<b>Warm Up:</b> Stretching, 7–9 RPE /5 min</p> <p><b>Workout</b></p> <ul style="list-style-type: none"> <li>• <b>Upper Body Resistance Exercise (9 -11 RPE/ 5 min):</b> Shoulder Abduction 180°, Shoulder Flexion 180°, Shoulder External Rotation A 90°, Trunk, Lateral Bending 35°, Trunk Flexion 80°</li> <li>• <b>Lower Body Resistance Exercise (9 -11 RPE/ 5 min):</b> Hip Abduction 45°, Hip Flexion 90°, Knee Flexion 90°, Sitting Knee Extension 0°</li> <li>•<b>Aerobic Exercise (11-13 RPE/ 5 min):</b> Walking in place, forward, sideways, backward; Heel raises, Knee lifts, Leg curls, Front lunges</li> </ul>	Daily Routine	<p><b>Muscle Mass and Muscle Function</b></p> <p>(Standing Height: Stadiometer</p> <p>Body Mass, Lean Muscle Mass and Appendicular Skeletal Muscle Mass: - Bioelectrical Impedance Analysis</p> <p>- Skeletal Muscle Index: ASM/height<sup>2</sup></p> <p>Gait Speed and Hand Grip Strength: Walk a 6m and Hand Dynamometer)</p> <p><b>Physical Performance</b> (Senior Fitness Test)</p>

				<p>•<b>Flexibility Exercise (9-11 RPE/ 5 min):</b> Neck, Trunk, Whole Body</p> <p>•<b>Cool-Down (7-9 RPE/ 5 min):</b> Upper Body Stretching, Lower Body Stretching</p> <p>30 min, 5 times per week, 12 weeks</p>		<p><b>Exercise Self-Efficacy</b> (Exercise Self-Efficacy Scale)</p>
Valiani et al., 2017	EG = 12 (10 Female and 2 Male)	80,5 ± 4,2	United States of America	<p><b>System:</b> Jintronix Technology</p> <p>•<b>Aerobic Exercises:</b> walking in place, sideway walking, arm circles, two side Steps with arm forward, toe pointed with arms up and marching in place with Arms.</p> <p>•<b>Resistance and Strength Exercises:</b> lateral weight shift and squatting, Controlled movements with right and left arm, bilateral coordination and foot eye Coordination with leg flexion.</p> <p>•<b>Balance Exercises:</b> walking forward and backward, walking left and right, Trunk flexion and stepping.</p> <p>30 min, 2 times per week, 8 sessions</p>	Does not apply	<p><b>Feasibility and Acceptability</b></p> <p>(A quality of movements' total average score &gt; 80%: algorithms</p> <p>An appreciation total average score &gt; 80% or &gt; 2.4 points: 9-item feasibility questionnaire</p> <p>An average total time in performing the exercises &gt; 80% of expected exercise time or &gt; 192 min out of 240 min: calculated automatically by the Jintronix Software.</p> <p>A difficulty total average score &lt; 40% or &lt; 1.2 points: 9-item feasibility and acceptability questionnaire)</p> <p><b>Physical Performance</b> (Short Physical Performance Battery test)</p> <p><b>Physical Activity Levels</b> (Rapid Assessment of Physical Activity questionnaire)</p>

EG= Experimental Group; CG = Control Group; RPE: Rating of Perceived Exertion.

\* Age in this study was taken as a categorical variable.

**Table 1. Characteristics of the Selected Studies**

312 **Risk of Bias in Included Studies**

313 The summary on the risk of bias assessment is in Figure 2. One of the studies was  
 314 judged to have an unclear risk of selection bias since, although an experimental  
 315 group was established, detailed information on the sequence generation process  
 316 and concealment is not provided.(30) Additionally, an unclear risk of performance  
 317 bias was determined in the same study as information on blinding of participants and  
 318 personnel was insufficient.(30)

319  
 320 Regarding detection bias, all included studies were rated at unclear risk, since none  
 321 of them addressed the results about blinding of assessors to their research  
 322 findings.(26,27,30) However, all studies were judged to be at low risk of attrition and  
 323 reporting bias as missing outcome data were not presented and the protocol and  
 324 results for each study were explicitly stated.(26,27,30) Finally, in relation to other types  
 325 of bias, two studies were considered high risk since the method of assigning the  
 326 experimental group was not specified in any of them(30) and the type of study was  
 327 not described in the methodology of the remaining one.(26)

328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Barsasella 2020	+	+	-	?	+	+	+
Sangwan 2020	+	+	-	?	+	+	-
Valiani 2017	?	?	?	?	+	+	-

**Figure 2. Summary Risk of Bias in Included Studies**

## 352 Discussion

353 This systematic review investigated the effects of aerobic endurance training through  
354 virtual reality on the functional physical condition of healthy older people based on  
355 the findings of three RCTs. The results suggest that the interventions, although they  
356 were not exclusively aerobic, have a positive influence on physical capacities such  
357 as aerobic endurance, muscular strength, agility and dynamic balance in people  
358 aged 60 or 65 and older.

359

360 First, although few studies were included, the findings of this review are consistent  
361 with the results of certain research available within the scientific evidence. The  
362 positive findings of the Barsasella study regarding aerobic capacity(27) are consistent  
363 with the results of another investigation, which found that its experimental group,  
364 after the implementation of an exercise program with virtual reality, reached  
365 maximum heart rate and metabolic cost in a shorter period of time and significantly.  
366 This is attributed, according to this research, to the fact that immersion experiences,  
367 simulated environments, life-like experiences and visual feedback could possibly  
368 make it easier for participants to tolerate higher intensities of exercise and perceive  
369 less sensation of actual physical discomfort.(31)

370

371 Similarly, some of the findings of the study by Valiani et al., as well as the study(30)  
372 by Barsasella,(27) coincide with the results of other studies,(32,33) in which  
373 improvements in capacities such as aerobic endurance, appendicular muscle  
374 strength, flexibility, and dynamic balance were also found after interventions with  
375 reality virtual. The results of these investigations, according to their authors, were  
376 probably presented thanks to the level of attraction and security generated by virtual  
377 environments, since these can facilitate adherence to exercise and continuity in its  
378 execution by the target populations.(32,33)

379

380 Secondly, from the perspective offered by the Ecological Model of Physical Activity,  
381 one may also think that the findings of the studies included in this review are due to  
382 the benefits offered by virtual reality. In this sense, it is likely that extrinsic factors  
383 such as feedback, the ecological tasks provided to the user and the simulation of  
384 real environments offered by this type of technology, stimulate a series of  
385 psychological factors such as motivation, pleasure, fun, revitalization, and a sense  
386 of tranquility, achieving that the physiological effects of the types of exercises (which  
387 are purely biological factors), are enhanced and produce more significant  
388 improvements in the functional physical condition of the elderly than conventional  
389 training programs.(23,31,33,34) Moreover, extrinsic and psychological factors that are  
390 stimulated by the use of technologies such as virtual reality can provide multiple  
391 ways to perform activities and facilitate the adherence and establishment of exercise

392 as a habit in this population. These advantages, in the end, are often more difficult  
393 to achieve or realize when working in the traditional way.(33)

394

395 Third, although it was found that the virtual reality interventions employed in the  
396 included studies were not designed only for aerobic endurance work, several  
397 investigations have shown that this type or mode of exercise is one of the most  
398 recommended for improving functional status in old age, as well as for treating or  
399 managing chronic conditions(35–39); this is because increased tissue oxygen,  
400 increased stroke volume, improvements in perfusion, angiogenesis and increased  
401 blood vessel caliber, reduced blood pressure, and increased gas exchange surface  
402 area not only promote the function of the cardiovascular and pulmonary systems in  
403 the elderly, but also stimulate the synthesis of mitochondria and muscle proteins,  
404 improve oxygenation at the cerebral level and stimulate the release of neurotrophic  
405 factors.(4,40–42) All of this, ultimately, can translate into an improvement in other  
406 physical abilities such as strength, balance and flexibility, as well as cognition.  
407 Likewise, in terms of functionality, the maintenance or improvement of functional  
408 physical condition can ensure the development of basic, instrumental, and social  
409 activities, as well as improve well-being and quality of life during old age.(9)

410

411 Finally, this review has strengths and limitations that should be mentioned.  
412 Regarding the former, it is worth noting that in this systematic review an exhaustive  
413 bibliographic search was carried out in several databases and included studies in all  
414 languages, developed over a considerable period of time, to try to collect the most  
415 scientific information available regarding the effectiveness of aerobic resistance  
416 training by means of virtual reality on the functional physical condition of healthy  
417 elderly people. These aspects give relevance and novelty to this research since most  
418 of the studies currently developed in the field of physical therapy, aging and old age  
419 have investigated the effects of conventional exercise programs in an older adult  
420 population with one or more health conditions. However, with respect to the latter,  
421 aspects such as the limited evidence available on the topic in question, the absence  
422 of a search within the gray literature, the diversity and quality of the included studies,  
423 the lack of studies that include exclusive aerobic interventions and the small sample  
424 sizes of the RCTs reviewed are highlighted.

425

426

## 427 **Conclusions**

428 The findings of this review indicate that aerobic endurance training via virtual reality  
429 can generate positive effects on functional fitness in healthy older adults. These  
430 results have important future implications: for clinical practice, this review  
431 demonstrates that interventions that are accompanied by technologies such as

432 virtual reality, whether unimodal or multimodal, can become more beneficial and  
433 motivational within the functional rehabilitation processes of individuals aged 60 or  
434 65 years and older, while promoting adherence to exercise and primary or secondary  
435 prevention of prevalent health conditions in old age, which represent enormous  
436 health costs. In addition, the use of such tools proves to be an attractive, quality and  
437 supportive complement, quite useful for health professionals or for interdisciplinary  
438 teams that intervene in this population and that will be confronted with the  
439 consequences of population aging and the great advances of the technological era.

440

441 Similarly, for the elderly, the use of these tools can increase levels of motivation and  
442 therapeutic adherence, as well as break down social stereotypes related to their  
443 inability to use technology. This is ultimately vital to achieve exercise-related  
444 physiological adaptations, produce improvements in function and functionality,  
445 achieve a transformation of lifestyle behaviors, increase social participation,  
446 intergenerational relationships, and improve well-being and health-related quality of  
447 life in old age.

448

449 Finally, as implications for research, the results of this study should be carefully  
450 evaluated due to the limitations already described. Therefore, it is recommended to  
451 increase the generation of knowledge on these topics through experimental studies,  
452 which have greater methodological rigor, reduce the risk of bias and allow the  
453 generalization of results and recommendations to the population in general. It also  
454 reiterates the need to increase research on healthy populations, since prevention  
455 strategies during the aging and old age processes are vital for reducing morbidity  
456 and mortality rates, the demand for health services, and health costs, among others.  
457 Likewise, the review of the theoretical models or theories on physical activity and  
458 exercise is suggested, as well as using those that are relevant to explanation with  
459 greater scientific rigor, including the phenomena given in light of the research  
460 findings.

461

462

## 463 **Acknowledgements**

464 The authors would like to express their sincere gratitude to the Universidad  
465 Autónoma de Manizales and the Ministry of Science, Technology and Innovation  
466 (Minciencias) for the support and funding granted to the research project: "Desarrollo  
467 de un Sistema de Realidad Virtual para la Implementación de los protocolos de  
468 Equilibrio y Fuerza Muscular y de Estabilidad en Adultos Mayores con y sin  
469 Problemas de Estabilidad" (Development of a Virtual Reality System for the  
470 Implementation of Balance and Muscle Strength and Stability protocols in Older  
471 Adults with and without Stability Problems), within which this scientific article arises

472 within the framework of the call 831 of 2021. Likewise, the authors express their  
473 gratitude to their research team for their valuable disposition and total admiration for  
474 the elderly for being our source of inspiration in our professional endeavors. The  
475 authors would like to thank the academic and scientific communities working in the  
476 field of health, technology and aging, because without their valuable contributions,  
477 continually generating new knowledge that can increasingly benefit the elderly  
478 population would not be possible. Finally, they thank Alexandra Suaza Restrepo,  
479 who works at the translation center at UAM, for translating this final manuscript, and  
480 Gregory Wallace Amos for reviewing it.

481

482

## 483 **Supporting Information**

484 S1 Table. PRISMA checklist.

485

486

487

## 488 **Author Contributions**

- 489 • **Conceptualization:** Daniela Ramírez.
- 490 • **Data curation:** Daniela Ramírez.
- 491 • **Formal analysis:** Daniela Ramírez.
- 492 • **Methodology:** Daniela Ramírez.
- 493 • **Supervision:** Julialba Castellanos, Lina María Montealegre, Carolina Márquez,  
494 Santiago Murillo.
- 495 • **Validation:** Julialba Castellanos, Lina María Montealegre, Carolina Márquez,  
496 Santiago Murillo.
- 497 • **Writing – original draft:** Daniela Ramírez.
- 498 • **Writing – review & editing:** Julialba Castellanos, Lina María Montealegre,  
499 Carolina Márquez, Santiago Murillo.

## 500 **References**

- 501 1. McCracken K, Phillips DR. Demographic and Epidemiological Transition.  
502 International Encyclopedia of Geography: People, the Earth, Environment and  
503 Technology. 2017;1–8.
- 504 2. World Health Organization. Aging and Health [Internet]. 2021 [cited 2021 Nov 30].  
505 Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
- 506 3. Macia E, Chevé D, Montepare JM. Demographic aging and biopower. Journal of  
507 Aging Studies. 2019;51(April):100820.
- 508 4. Kozakai R, Nishita Y, Otsuka R, Ando F, Shimokata H. Age-Related Changes in  
509 Physical Fitness Among Community-Living Middle-Aged and Older Japanese: A 12-  
510 Year Longitudinal Study. Research Quarterly for Exercise and Sport.  
511 2020;91(4):662–75.
- 512 5. Tomás MT, Galán-Mercant A, Carnero EA, Fernandes B. Functional capacity and  
513 levels of physical activity in aging: A 3-year follow-up. Frontiers in Medicine.  
514 2017;4(JAN):1–8.
- 515 6. Fleg JL, Morrell CH, Bos AG, Brant LJ, Talbot LA, Wright JG, et al. Accelerated  
516 longitudinal decline of aerobic capacity in healthy older adults. Circulation.  
517 2005;112(5):674–82.
- 518 7. Fleg JL. Aerobic exercise in the elderly: a key to successful aging. Discov Med.  
519 2012;13(70):223–8.
- 520 8. Alul LAU, Gomez-Campos R, Almonacid-Fierro A, Morales-Mora L, Rojas-Mancilla  
521 E, Palomo I, et al. Aerobic capacity of chilean adults and elderly: Proposal of  
522 classification by regional percentiles. Revista Brasileira de Medicina do Esporte.  
523 2019;25(5):390–4.
- 524 9. Gómez JF, Curcio CL. Salud del anciano. In Manizales: Editorial Universidad de  
525 Caldas; 2020.
- 526 10. Karavidas A, Lazaros G, Tsiachris D, Pyrgakis V. Aging and the Cardiovascular  
527 System. Hellenic J Cardio. 2010;51:421–7.
- 528 11. Triposkiadis F, Xanthopoulos A, Butler J. Cardiovascular Aging and Heart Failure:  
529 JACC Review Topic of the Week. Journal of the American College of Cardiology.  
530 2019;74(6):804–13.
- 531 12. Sharma G, Goodwin J. Effect of aging on respiratory system physiology and  
532 immunology. Clinical interventions in aging. 2006;1(3):253–60.
- 533 13. Marriott CFS, Petrella AFM, Marriott ECS, Boa Sorte Silva NC, Petrella RJ. High-  
534 Intensity Interval Training in Older Adults: a Scoping Review. Sports Medicine - Open.  
535 2021;7(1).
- 536 14. Hwang C-L, Lim J, Yoo J-K, Kim H-K, Hwang M-H, Handberg EM, et al. Effect of All-  
537 Extremity High-Intensity Interval Training vs. Moderate-Intensity Continuous Training



- 538 on Aerobic Fitness in Middle-Aged and Older Adults with Type 2 Diabetes: A  
539 Randomized Controlled Trial. *Exp Gerontol.* 2019;116:46–53.
- 540 15. Leirós-Rodríguez R, Soto-Rodríguez A, Pérez-Ribao I, García-Soidán JL.  
541 Comparisons of the Health Benefits of Strength Training, Aqua-Fitness, and Aerobic  
542 Exercise for the Elderly. *Rehabilitation Research and Practice.* 2018;1–8.
- 543 16. Ribeiro Silva M, Lima Alberton C, Portella EG, Neves Nunes G, Gomez Martin D,  
544 Santana Pinto S. Water-based aerobic and combined training in elderly women:  
545 Effects on functional capacity and quality of life. *Experimental Gerontology.*  
546 2018;106(2017):54–60.
- 547 17. Schwartz B. Technology and the future of ophthalmology. *Survey of Ophthalmology.*  
548 1981;25(6):402–3.
- 549 18. Martinez-Martin E, Cazorla M. Rehabilitation technology: Assistance from hospital to  
550 home. *Computational Intelligence and Neuroscience.* 2019;2019(1).
- 551 19. Cano de la Cuerda R, Collado Vásquez S. *Neurorrehabilitación. Editorial Médica*  
552 *Panamericana;* 2012.
- 553 20. Cano de la Cuerda R, Martínez Piédrola RM, Miangolarra JC. Control y aprendizaje  
554 motor. *Editorial Médica Panamericana;* 2017.
- 555 21. Ulas K, Semin I. The Biological and Motivational Effects of Aerobic Exercise With  
556 Virtual Reality. *Research Quarterly for Exercise and Sport.* 2021;92(3):321–6.
- 557 22. Haese, SaHaese, S. D. ', Cardon, G., & Deforche B (2015). E y actividad física. TEEb  
558 on C and AObesity ra D', Cardon G, Deforche B. Entorno y actividad física. The  
559 ECOG's eBook on Child and Adolescent Obesity. 2015;
- 560 23. Moral Moreno L. Teorías y modelos que explican y promueven la práctica de  
561 actividad física en niños y adolescentes. *Educación y futuro: revista de investigación*  
562 *aplicada y experiencias educativas.* 2017;36(36):177–208.
- 563 24. Corporación Universitaria del Caribe. Enfoques, Teorías y Perspectivas de las  
564 Ciencias del Deporte y sus Programas Académico. Enfoques, Teorías y Perspectivas  
565 de las Ciencias del Deporte y sus Programas Académico. Sincelejo: Editorial  
566 CECAR; 2018. 146.
- 567 25. RevMan | Cochrane Training [Internet]. [cited 2021 Dec 1]. Available from:  
568 [https://training.cochrane.org/online-learning/core-software-cochrane-](https://training.cochrane.org/online-learning/core-software-cochrane-reviews/revman)  
569 [reviews/revman](https://training.cochrane.org/online-learning/core-software-cochrane-reviews/revman)
- 570 26. Jeon S, Kim J. Effects of augmented-reality-based exercise on muscle parameters,  
571 physical performance, and exercise self-efficacy for older adults. *International Journal*  
572 *of Environmental Research and Public Health.* 2020;17.
- 573 27. Barsasella D, Liu MF, Malwade S, Galvin CJ, Dhar E, Chang CC, et al. Effects of  
574 Virtual Reality Sessions on the Quality of Life, Happiness, and Functional Fitness  
575 among the Older People: A Randomized Controlled Trial from Taiwan. *Computer*  
576 *Methods and Programs in Biomedicine.* 2021;200.

- 577 28. Rybertt C, Cuevas S, Winkler X, Lavados P, Martínez S. Parámetros funcionales y  
578 su relación con la velocidad de marcha en adultos mayores chilenos residentes en la  
579 comunidad. *Biomédica*. 2015;35(212–8).
- 580 29. Río X, Guerra-Balic M, González-Pérez A, Larrinaga-Undabarrena A, Coca A.  
581 Valores de referencia del SPPB en personas mayores de 60 años en el País Vasco.  
582 *Atencion Primaria*. 2021;53(8).
- 583 30. Valiani V, Lauzé M, Martel D, Pahor M, Manini T, Anton S, et al. A New Adaptive  
584 Home-Based Exercise Technology Among Older Adults Living in Nursing Home : a  
585 Pilot Study on Feasibility , Acceptability and Physical Performance. *J Nutr Health*  
586 *Aging*. 2017;21(7):819–24.
- 587 31. Chuang TY, Sung WH, Chang HA, Wang RY. Effect of a virtual reality-enhanced  
588 exercise protocol after coronary artery bypass grafting. *Physical Therapy*.  
589 2006;86(10):1369–77.
- 590 32. Rutkowski S, Rutkowska A, Kiper P, Jastrzebski D, Rachenjuk H, Turolla A, et al.  
591 Virtual reality rehabilitation in patients with chronic obstructive pulmonary disease: A  
592 randomized controlled trial. *International Journal of COPD*. 2020;15:117–24.
- 593 33. Jo G, Rossow-Kimball B, Park G, Lee Y. Effects of virtual reality exercise for Korean  
594 adults with schizophrenia in a closed ward. *Journal of Exercise Rehabilitation*.  
595 2018;14(1):39–48.
- 596 34. Lee CC, Hsiao KL, Chen CC. Exploring the Benefit and Sacrifice Factors of Virtual  
597 Reality Gameplay. *Frontiers in Psychology*. 2020;11(March):1–8.
- 598 35. Ávila-Funes JA, García-Mayo EJ. Beneficios de la práctica del ejercicio en los  
599 ancianos. *Gaceta Medica de Mexico*. 2004;140(4):431–6.
- 600 36. Vidarte JA, Quintero MV, Herazo-Beltrán Y. Efectos Del Ejercicio Físico En La  
601 Condición Física Funcional Y La Estabilidad En Adultos Mayores. *Hacia la Promoción*  
602 *de la Salud*. 2012;17(2):79–90.
- 603 37. Bakken RC, Carey JR, Di Fabio RP, Erlandson TJ, Hake JL, Intihar TW. Effect of  
604 aerobic exercise on tracking performance in elderly people: A pilot study. *Physical*  
605 *Therapy*. 2001;81(12):1870–9.
- 606 38. Short KR, Vittone JL, Bigelow ML, Proctor DN, Nair KS. Age and aerobic exercise  
607 training effects on whole body and muscle protein metabolism. *American Journal of*  
608 *Physiology - Endocrinology and Metabolism*. 2004;286(1 49-1):92–101.
- 609 39. Varela-Gutiérrez JP, Rojas-Quirós J. Efectos De Un Programa De Ejercicio De  
610 Fuerza Y Resistencia Aeróbica En Un Adulto Mayor Pluripatológico: Estudio De  
611 Caso. *Revista digital: Actividad Física y Deporte*. 2021;7(2).
- 612 40. Patel H, Alkhawam H, Madanieh R, Shah N, Kosmas CE, Vittorio TJ. Aerobic vs  
613 anaerobic exercise training effects on the cardiovascular system . *World Journal of*  
614 *Cardiology*. 2017;9(2):134.

- 615 41. Ferguson-Stegall L, McCleave E, Ding Z, Doerner PG, Liu Y, Wang B, et al. Aerobic  
616 exercise training adaptations are increased by postexercise carbohydrate-protein  
617 supplementation. *Journal of Nutrition and Metabolism*. 2011;2011.
- 618 42. Mendez Colmenares A, Voss MW, Fanning J, Salerno EA, Gothe NP, Thomas ML,  
619 et al. White matter plasticity in healthy older adults: The effects of aerobic exercise.  
620 *NeuroImage*. 2021;239(February):118305.
- 621




Figure 1. PRISMA flow diagram.

medRxiv preprint doi: <https://doi.org/10.1101/2022.05.24.22275497>; this version posted May 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Barsasella 2020	+	+	-	?	+	+	+
Sangwan 2020	+	+	-	?	+	+	-
Valliani 2017	?	?	?	?	+	+	-

**Figure 2. Summary Risk of Bias in Included Studies.**