Concurrent Plasmodium falciparum and acute dengue virus infection in

(sub-Saharan) Africa: a systematic review and meta-analysis

Tewelde T. Gebremariam[1,2¶*], Henk D.F.H. Schallig[2¶], Zeleke Mekonnen[3&], Asrat Hailu[1&], Jonas

B. Danquah[4&], Woldaregay E. Abegaz[1&]

[1] Department of Microbiology, School of Medicine, Addis Ababa University, Addis Ababa,

Ethiopia.

[2] Department of Medical Microbiology, Academic Medical Centre, University of Amsterdam,

Amsterdam, Netherlands.

[3] School of Medical Laboratory Sciences, Jimma University, Jimma, Ethiopia.

[4] Animal Research Institute, Animal Health Division Council for Scientific & Industrial

Research, Accra, Ghana.

Corresponding author

E-mail: t.t.gebremariam@amsterdamumc.nl (TTG)

[¶] These authors contributed equally to this work.

[&] These authors also contributed equally to this work.

Abstract

Malaria and dengue fever are leading causes of acute undifferentiated febrile illness. In Africa, misdiagnoses of dengue fever as malaria is a common scenario. This systematic review was aimed to determine the prevalence of concurrent malaria and dengue fever, aetiologies, and the association between the two infections in Africa. We assessed evidence from previous studies in Africa that were available in MEDLINE and EMBASE databases between January 01, 2005, and March 30, 2022. A total of 3942 studies were identified from the databases of which 30 studies from 13 African countries, including 25,241 febrile patients, were included in the meta-analysis. The result of the meta-analysis showed a 4.0% pooled prevalence of concurrent malaria and dengue fever in (sub-Saharan) Africa, and the highest rate was recorded in Central Africa (5.7%) followed by East Africa (4.7%) and West Africa (2.9%). Children <18 years old (57%) and females (52.8%) were more affected by the co-infection than adults and females, respectively. The dengue virus serotypes identified were DENV-2, DENV-3, DENV-1, and DENV-4 in descending order with a proportion of 39%, 31%, 27%, and 3%, respectively. Moreover, *Plasmodium falciparum* was the only specified malaria parasite in the co-infection among the included studies. Significantly higher odds of malaria infection were documented due to dengue fever when compared with malaria mono-infection. However, no significant odds of acute dengue co-infection were recorded due to malaria in contrast to dengue mono-infection. This study showed a high prevalence of concurrent malaria and dengue fever in Africa. Healthcare workers should bear in mind the possibility of dengue infection as differential diagnoses for acute febrile illness as well as the possibility of co-existent malaria and dengue in endemic areas. Also, high-quality multi-centre studies are required to verify the above conclusions.

Protocol registration number: CRD42022311301.

Keywords: Prevalence, Co-infection, Malaria, Dengue fever, Dengue virus, DENV,

Plasmodium falciparum.

Introduction

Acute undifferentiated febrile illness (AUFI) is one of the most frequent reasons for healthcare seeking in Africa [1]. AUFI usually begins with non-specific symptoms such as the sudden onset of fever which fewer progresses to prolonged duration, headache, chills, and myalgia which may later involve specific organs. It can range from mild and self-limiting illness to advancing, deadly disease [2]. Malaria and dengue fever are leading causes of AUFI [3].

Africa carries the highest global malaria burden, with 2000 million cases (92%) in 2017 alone [4]. Human malaria is mainly caused by four *Plasmodium* species, namely *P. falciparum*, *P. vivax*, *P. malariae*, and *P. ovale* with variable geographic distribution. *P. falciparum* accounts for nearly all malaria deaths in sub-Saharan Africa, which bears over 90% of the global malaria burden [5]. Likewise, the prevalence of dengue in the region has dramatically increased over the past few decades although this specific infection is neither systematically investigated nor generally considered by clinicians [6]. In 2013, about 16 million apparent and over 48 million inapparent dengue were estimated to have occurred, and most countries in the continent reported recurrent outbreaks [7]. Similarly, dengue fever is caused by four genetically distinct dengue viruses (serotypes 1-4) [8].

Though malaria or dengue virus mono-infection can be severe, concomitant infections could be

even more fatal [9, 10]. The two mosquito-borne diseases have an overlapping epidemic pattern

in Africa [11]. Due to their similar clinical presentation, possible concurrent malaria-dengue fever

is often neglected [12] and generally misdiagnosed as malaria only [6, 13]. Misdiagnosis is more

probable during co-infection than mono-infection and this may result in slow identification of

dengue fever outbreaks with potentially high morbidity and mortality [6, 14].

This review aimed to gather evidence to answer the question: how common is *Plasmodium* and

dengue virus co-infection in Africa could be? The specific review objectives were (1) to determine

the prevalence of concurrent malaria and acute dengue in Africa (by region and demographic

characteristics), and (2) to describe the dengue virus serotypes in the concurrent infection.

Results

Literature retrieval and article characteristics

A total of 3942 records were identified during literature retrieval from the databases. A total of 30

studies from (sub-Saharan) Africa involving 25,241 febrile patients were included in this meta-

analysis [15-45] (Figure 10 , Table 1). The quality score of the included studies ranged from 4 to

7 on a maximum scale of quality scores of 8 for case reports, 9 for cross-sectional studies, and 11

for case-control studies using the JBI criteria (Table 1).

Table 1. Basic Characteristics and quality score of included studies.

Studies	Country	Study design	Sample size	Quality score according to JBI criteria
Stoler 2015	Ghana	Cross-sectional	218	5
Ahmed 2021	Sudan	Cross-sectional	76	4
Charrel 2005	Guinea	Case report	1	6

Sow 2016	Senegal	Cross-sectional	13845	6
Abdulaziz 2020	Nigeria	Cross-sectional	440	4
Ayorinde 2016	Nigeria	Cross-sectional	60	4
Baba 2013	Nigeria	Cross-sectional	310	4
Dariano 2017	Sierra Leone	Cross-sectional	1260	4
Miri 2021	Nigeria	Cross-sectional	94	4
Oyero 2014	Nigeria	Cross-sectional	188	4
Onyedibe 2018	Nigeria	Cross-sectional	529	5
Nassar 2019	Nigeria	Cross-sectional	170	5
Kolawole 2017	Nigeria	Cross-sectional	176	5
Vu 2016	Kenya	Cross-sectional	579	5
Ali 2020	Tanzania	Cross-sectional	149	4
Shah 2020	Kenya	Cross-sectional	862	4
Marks 2021	Burkina Faso, Madagascar, Sudan	Cross-sectional	615	6
Akelew 2022	Ethiopia	Cross-sectional	200	4
Boillat-Blanco 2018	Tanzania	Case-control	431	6
Chipwaza 2014	Tanzania	Cross-sectional	364	5
Obonyo 2018	Kenya	Cross-sectional	1332	6
Tchetgna 2021	Cameroon	Cross-sectional	320	5
Galani 2020	Cameroon	Cross-sectional	174	4
Mouiche 2020	Cameroon	Cross-sectional	629	5
Nkenfou 2021	Cameroon	Cross-sectional	349	4
Proesmans 2019	DRC	Cross-sectional	342	4
Tchuandom 2018	Cameroon	Cross-sectional	961	7
Yousseu 2018	Cameroon	Cross-sectional	114	4
Monamele 2018	Cameroon	Cross-sectional	33	4
Otu 2019	Nigeria	Cross-sectional	420	4

JBI, Joanna Briggs Institute; DRC, Democratic Republic of Congo.

Prevalence of concurrent falciparum malaria and dengue fever

The random-effect (DerSimonian-Laird) model estimator and raw proportion and/or

Freeman-Tukey double arcsine transformed proportion were used in the meta-analysis. The

prevalence of malaria and dengue co-infection in (sub-Saharan) Africa per 1000 febrile cases was

39.8 (95% CI, 32 – 48, SE = 0.004) with significant heterogeneity (Tau= 0.018, $Tau^2 = 0.00$, $I^2 =$

95.52%, and Q = 625.28, p < 0.001) (Figure 1). The prevalence significantly varies across the three

African regions ($\chi^2 = 537.13$, p < 0.00001 and ranges from 28.5 per 1000 febrile cases (95% CI,

18 – 39, SE = 0.0052) in West Africa, 47 (95% CI, 19 – 75, SE = .0142) in East Africa to 57.7

(95% CI, 30 – 86, SE = 0.0143) in Central Africa (Table 2, Figure 1 – Figure 4). A case report

[15] was excluded from this analysis.

____

Table 2. Meta-analysis of concurrent malaria and acute dengue prevalence among febrile patients in three

African regions.

| Region | No. of studies | No. of cases | **Heterogeneity** | | | | | Model | **Meta-analysis** | |
			P	T	T²	Q	I² (%)		E (95% CI) [×1000]	*P*
Central Africa	8	157	<.001	0.037	0.00	140.95	95.0	Random	57.7 (30, 86)	<.001
East Africa	8	248	<.001	0.039	0.00	214.42	96.7	Random	47 (19, 75)	<.001
West Africa	12	145	<.001	0.015	0.00	159.55	93.1	Random	28.5 (18, 39)	<.001
T, tau; Q, cochrane Q-test, p, p-value.										

99   Figure 1. Forest plot delineating prevalence of malaria and acute dengue co-infection in (sub-Saharan)

100   Africa.

Figure 2. Forest plot showing the prevalence of malaria and acute dengue co-infection in West Africa.

Figure 3. Prevalence of malaria and dengue co-infection in East Africa.

Studies	Estimate (95% CI)	Ev/Trt
Tchetgna 2021	0.072 (0.044, 0.100)	23/320
Galani 2020	0.069 (0.031, 0.107)	12/174
Mouiche 2020	0.003 (0.000, 0.008)	2/629
Nkenfou 2021	0.195 (0.153, 0.236)	68/349
Proesmans 2019	0.018 (0.004, 0.031)	6/342
Tchuandom 2018	0.042 (0.029, 0.054)	40/961
Yousseu 2018	0.026 (0.000, 0.056)	3/114
Monamele 2018	0.091 (0.000, 0.189)	3/33
Overall (I^2=95.03 % , P< 0.001)	0.058 (0.030, 0.086)	157/2922

Figure 4. Forest plot depicting prevalence of concurrent malaria and acute dengue in Central Africa.

The interaction of *Plasmodium* and dengue viruses

Twenty-two studies examined the association between acute dengue virus infection with malaria. The peto's odds ratio [46] was used to pool the odds ratio of individual studies. In our meta-analysis (Figure 5), cross-sectional studies in Cameroon [30, 45], and Kenya [22] revealed significantly lower odds of co-infection in contrast with the absence of acute dengue infections (OR: 0.60; 95% CI, 0.42 – 0.87). Cross-sectional studies in Nigeria [16, 18, 20, 25, 32, 33, 35], Tanzania [17], Sierra Leone [23], Democratic Republic of Congo (DRC) [34], Cameroon [31], Sudan [40, 42], and Senegal [21] indicated no significant odds of co-infection in comparison with acute dengue negatives (OR = 1.07; 95% CI = 0.58, 1.94). Studies in Ethiopia [44], Nigeria [29], Tanzania [36], Cameroon [43], and Kenya [39], on the other hand, indicated significantly higher odds of concurrent infection to acute dengue negatives (OR = 4.56; 95% CI = 3.29, 6.33). There was no indication of a potential outlier (studentized residuals >±3.0654) and overlay influence (cook's distance) among the studies. Above all, the summary estimate derived from 22 studies showed significantly higher odds of malaria infection among patients in the absence of acute dengue than acute dengue virus co-infection (OR = 1.51; 95% CI = 1.31, 1.75) with significant

126 heterogeneity ($I^2 = 89$; $p < 0.001$). The 95% prediction interval for the true outcomes (log peto's

127 odds ratio) was -1.41 to 2.46.

Study	Coinfection Events	Total	Malaria only Events	Total		Weight, Peto, Random, 95% CI	Odds Ratio
Abdulaziz 2020	67	332	14	108		6.80%	1.62 [0.92, 2.83]
Akelew 2022	4	15	17	185		0.73%	6.36 [1.15, 35.26]
Ayorinde 2016	1	1	34	59		0.13%	2.03 [0.12, 34.43]
Chipwaza 2014	18	76	65	288		5.88%	1.07 [0.58, 1.94]
Dariano 2017	7	47	291	1213		4.53%	0.60 [0.30, 1.20]
Miri 2021	5	52	6	42		1.34%	0.64 [0.18, 2.25]
Nassar 2019	1	3	71	167		0.40%	0.69 [0.07, 6.89]
Onyedibe 2018	7	12	110	517		1.12%	8.56 [2.16, 33.95]
Otu 2019	7	15	211	405		2.01%	0.80 [0.29, 2.25]
Oyero 2014	19	67	25	121		4.31%	1.53 [0.76, 3.10]
Proesmans 2019	9	28	146	314		3.54%	0.56 [0.26, 1.22]
Shah 2020	150	183	299	679		19.95%	4.56 [3.29, 6.33]
Tchuandom 2018	40	138	339	823		15.65%	0.60 [0.42, 0.87]
Vu 2016	33	73	300	506		8.65%	0.56 [0.34, 0.92]
Yousseu 2018	3	8	20	106		0.67%	3.15 [0.53, 18.72]
Baba 2013	18	76	53	290		5.23%	1.41 [0.75, 2.67]
Ali 2020	2	5	5	144		0.12%	3306.77 [49.61, 220423.75]
Marks 2021	19	57	149	558		5.70%	1.40 [0.76, 2.57]
Tchetgna 2021	23	41	204	279		4.10%	0.44 [0.21, 0.90]
Ahmed 2021	5	8	27	68		0.98%	2.52 [0.58, 10.99]
Sow 2016	1	3	7386	13842		0.41%	0.45 [0.05, 4.32]
Nkenfou 2021	68	70	115	279		7.75%	9.35 [5.54, 15.80]
Total (95% CI)		1310		20993		100.00%	1.51 [1.31, 1.75]

Heterogeneity: $\tau^2 = 0$, $\chi^2 = 183.73$, df= 21 (P=0) $I^2 = 89$
Test for overall effect: Z=5.54 (P=0)

0.01 0.1 1 10 100
Favours [Coinfection] Favours [Malaria only]

Figure 5. Forest plot: the difference in the prevalence of malaria due to acute dengue virus infection and

malaria mono-infection.

Similarly, the odds of acute dengue virus infection were also compared with the presence and

absence of malaria infection (Figure 6). Cross-sectional studies in Kenya [22, 39] and Cameroon

[30, 45] reported significantly low odds of acute dengue in the presence of malaria than in its

absence (OR = 0.57, 95% CI, 0.40 − 0.82). Similar studies Nigeria [18, 20, 25, 32, 33, 35],

Tanzania [17], Sierra Leone [23], DRC [34], Cameroon [31], Sudan [40, 42], and Senegal [21] ;

however, showed no notable odds of acute dengue infection in the presence of malaria when

compared to the absence of malaria (OR = 1.07, 95% CI, 0.58 –1.94). On the contrary, cross-

sectional studies in Ethiopia [44], Nigeria [16, 29], Tanzania [36], and Cameroon [43] indicated

higher odds of acute dengue infection in the presence of malaria than in its absence (OR = 6.98,

95% CI, 2.89 – 16.85). The overall estimate based on 22 studies showed no significant odds of

acute dengue co-infection in contrast to malaria mono-infection (OR = 1.00, 95% CI, 0.87 – 1.16)

with significant heterogeneity (I^2 = 86, P <0.0001). The 95% prediction interval for the true

outcomes (log peto's odds ratio) was -1.45 – 2.127.

Study	Coinfection Events	Coinfection Total	dengue only Events	dengue only Total		Odds Ratio Weight, Peto, Random, 95% CI
Abdulaziz 2020	67	81	265	359		6.40% 1.62 [0.92, 2.83]
Akelew 2022	4	22	11	178		0.71% 5.59 [1.04, 29.93]
Ayorinde 2016	1	32	0	28		0.13% 2.43 [0.15, 40.04]
Chipwaza 2014	18	83	58	281		5.54% 1.07 [0.58, 1.94]
Dariano 2017	7	298	40	962		4.27% 0.60 [0.30, 1.20]
Miri 2021	5	11	47	83		1.27% 0.64 [0.18, 2.25]
Nassar 2019	1	72	2	98		0.38% 0.69 [0.07, 6.89]
Onyedibe 2018	7	117	5	412		1.06% 8.56 [2.16, 33.95]
Otu 2019	7	218	8	202		1.89% 0.80 [0.29, 2.25]
Oyero 2014	19	44	48	144		4.06% 1.53 [0.76, 3.10]
Proesmans 2019	9	155	19	187		3.34% 0.56 [0.26, 1.22]
Shah 2020	150	449	183	413		26.65% 0.63 [0.48, 0.83]
Tchuandom 2018	40	390	98	571		14.89% 0.57 [0.40, 0.82]
Vu 2016	33	73	300	506		8.15% 0.56 [0.34, 0.92]
Yousseu 2018	3	23	5	91		0.63% 3.15 [0.53, 18.72]
Baba 2013	18	23	95	287		2.58% 6.98 [2.89, 16.85]
Ali 2020	2	7	7	142		0.20% 62.66 [2.62, 1498.59]
Marks 2021	19	168	38	447		5.37% 1.40 [0.76, 2.57]
Tchetgna 2021	23	227	18	93		3.86% 0.44 [0.21, 0.90]
Ahmed 2021	5	32	3	44		0.92% 2.52 [0.58, 10.99]
Sow 2016	1	7387	2	6458		0.39% 0.45 [0.05, 4.32]
Nkenfou 2021	68	183	2	166		7.31% 9.35 [5.54, 15.80]
Total (95% CI)		10095		12152		100.00% 1.00 [0.87, 1.16]

Heterogeneity: τ^2=0, χ^2=153.32, df=21 (P=0) I^2=86
Test for overall effect: Z=0.06 (P=0.948)

0.01 0.1 1 10 100

Favours [Coinfection] Favours [dengue only]

148   Figure 6. Forest plot showing the difference in the prevalence of acute dengue co-infection due to malaria

149   and acute dengue mono-infection.

The demographic characteristics of patients coinfected with *Plasmodium* and dengue viruses

The gender distribution among malaria and acute dengue virus coinfection showed that the proportion of

females was 52.8% (95% CI, 44.1 – 61.5, SE = 0.044) without evidence of heterogeneity (I^2 = 0%, p =

0.62), while the proportion of males was 47.2% (95% CI, 38.5 – 55.9, SE = 0.044) with no evidence of

heterogeneity (I^2 = 0%, p = 0.62) (Table 3). Furthermore, the proportion age category of the coinfection

was 57.0% (40.9-73.1, SE = 0.082) among children < 18 years old and 40.9% (95% CI, 24.5 – 57.0, SE =

0.082) in adults with significant heterogeneity (p <0.001).

Table 3. Meta-analysis of demographic characteristics of patients with concurrent malaria and dengue

virus infection in (sub-Saharan) Africa.

| Demography | No. of studies | No. of patients | Heterogeneity | | | | | Model | Meta-analysis | |
			P	T	T²	Q	I² (%)		E (95% CI) [×100]	*P*
Gender										
Female	7	62	0.62	0.00	0	6.29	0	Random	52.8 (44.1, 61.5)	<.001
Male	7	54	0.62	0.00	0	6.29	0	Random	47.2 (38.5, 55.9)	<.001
Age										
Children	8	158	<.001	0.26	0.07	641.78	98.13	Random	57 (40.9, 73.1)	<.001
Adults	7	49	<.001	0.26	0.07	637.02	98.12	Random	40.9 (24.8, 57.0)	<.001

I^2, a measure of heterogeneity; p, significant level. Abbreviations: CI, confidence interval, E, estimate; children, <18 years old; adults, ≥18 years old; T, tau; Q, cochran's Q-test.

The aetiologies of concurrent malaria and dengue fever

The meta-analysis on the proportion of aetiologies of concomitant malaria and acute dengue is

shown in Table 4. All the four dengue virus serotypes were identified, where DENV-2, DENV-3,

DENV-1, and DENV-4 were documented in descending order with a proportion of 39.5% (95%

CI, 16.8-62.2, SE = 0.116), 31.6% (95% CI, 16.0 – 47.2, SE = 0.080), 27.3% (95% CI, 14.6 – 40.1,

SE = 0.065), and 3.7% (95% CI, 1.0 – 6.3, SE = 0.014), respectively. The aetiology of malaria

among the co-infection was specified in 50.7% (95% CI, 29.6 – 71.9, SE = 0.108) of the patients,

and all were due to *P. falciparum*.

Table 4. Meta-analysis of the proportion of aetiologies of concomitant malaria and acute dengue in (sub-

Saharan) Africa.

| | No. of studies | No. of patients | Heterogeneity | | | | | Model | Meta-analysis | |
			P	T	T ²	Q	I² (%)		E (95% CI) [×100]	*P*
DENV serotypes										
DENV-2	7	97	<.001	.39	.15	331.3	96.4	Random	39.5 (16.8, 62.2)	<.001
DENV-3	7	36	<.001	.25	.06	216.2	94.5	Random	31.6 (16.0, 47.2)	<.001
DENV-1	6	33	<.001	.20	.04	118.1	89.8	Random	27.3 (14.6, 40.1)	<.001
DENV-4	3	7	0.81	.00	0	7.7	0	Random	3.7 (1.0, 6.3)	.01
***Plasmodium* species**										
Pf	16	305	<.001	.57	.33	15549.5	99.8	Random	50.7 (29.6, 71.9)	<.001
NS	14	251	<.001	.58	.33	15549.5	99.8	Random	49.3 (28.1, 70.4)	<.001

I², a measure of heterogeneity; p, significant level. Abbreviations: CI, confidence interval; E, estimate; Pf, *Plasmodium falciparum*; NS, not specified; T, tau; Q, cochran's Q-test.

Publication bias

The publication bias of included studies was evaluated using the funnel plot asymmetry tests: the

Egger's regression test and rank correlation test based on the Freeman-Tukey double arcsine

transformed proportion effect size model measures for prevalence/proportion estimates and log

odds ratio peto's method for the difference in the prevalence rates (Table 5, Figure 7 – Figure 9).

Both tests showed no significant publication bias (p >0.05) among the studies included in all the

analyses.

Table 5. Evaluation of publication bias using Egger's Regression test, and Rank Correlation test.

Publication bias assessment title	Regression test		Rank Correlation Test	
	Z	P	Kendall's Tau	P
The prevalence of malaria and dengue coinfections in Africa.	1.094	0.274	-0.025	.867
The prevalence of malaria and dengue co-infections in East Africa.	-.571	0.568	0.214	.548
The prevalence of malaria and dengue co-infections in West Africa.	1.270	.204	0.061	.841
The prevalence of malaria and acute dengue co-infections in Central Africa.	0.622	.508	0.071	.905
The difference in the prevalence of *Plasmodium* and dengue coinfection with malaria mono-infection.	1.060	2.89	1.69	.288
The difference in the prevalence of malaria and dengue coinfection with dengue mono-infection.	1.788	.074	2.73	.080
The proportion of males with malaria and acute dengue co-infection in Africa.	.222	.825	-0.088	.748
The proportion of females with malaria and acute dengue co-infection in Africa.	-.222	.825	.088	.748
The proportion of children with malaria and acute dengue co-infection in Africa.	-1.543	.123	-0.242	0.265
The proportion of adults with malaria and acute dengue co-infection in Africa.	.976	.329	0.212	0.324
The proportion of DENV-1 in malaria and acute dengue co-infection in Africa.	1.499	.134	0.340	0.111
The proportion of DENV-2 in malaria and acute dengue co-infection in Africa.	-1.62	.106	0.026	0.903
The proportion of DENV-3 in malaria and acute dengue co-infection in Africa.	.827	.408	0.261	0.220
The proportion of DENV-4 in malaria and acute dengue co-infection in Africa.	1.123	.262	0.444	0.037

Figure 7. Funnel plot showing assessment of publication bias based on the prevalence of concurrent

malaria and acute dengue in (sub-Saharan) Africa.

Figure 8. Funnel Plot depicting evaluation of publication bias based on the difference in the prevalence of

concurrent malaria and acute dengue with malaria mono-infection in (sub-Saharan) Africa.

Figure 9. Funnel Plot: evaluation of publication bias based on the difference in the prevalence of

concurrent malaria and acute dengue with acute dengue mono-infection in (sub-Saharan) Africa.

Discussion

The first report of malaria and dengue virus co-infection in Africa was documented in 2005 in a

37-years-old French woman who had recent travel history to three African countries: Guinea,

Senegal, and Sierra Leone [15]. Our meta-analysis included 25,241 febrile patients from 30 studies

conducted in 13 African countries (Ghana, Sudan, Senegal, Nigeria, Guinea, Sierra Leone, Kenya,

Tanzania, Ethiopia, Cameroon, DRC, Burkina Faso, and Madagascar) for over around 17 years.

As far as we know, this is the first description and analysis of the prevalence, demographic

characteristics, and dengue virus serotype profiles of concurrent malaria and dengue infection in

(sub-Saharan) Africa.

Based on our analysis, the pooled prevalence of concurrent malaria and dengue was 4.0%, and the

highest rate was recorded in Central Africa (5.7%), followed by East Africa (4.8%) and West

Africa (2.9%). This result is lower than the finding of a study [47] on a meta-analysis of severe

malaria and dengue co-infection which estimated a prevalence of 32%. The variation could be due

to the differences in the study population, model estimator employed, and/or geography of the

primary studies included, where our analysis was focused on studies from African continent only

while the other study included studies from all over the globe. Besides, uncomplicated febrile cases

were included in our analysis, unlike above the study which estimated severe malaria prevalence

among the co-infection.

In our study, females and children were more affected groups by the concomitant infection than

males and adults, respectively. Women and children are more susceptible to mosquito-borne

illnesses because they are exposed to mosquito bites for longer periods during dangerous hours

[48]. Moreover, malaria [49] and dengue [50] mainly affect children due to underdeveloped

specific immunity to infection [49].

In our analysis, the most prevalent dengue virus serotypes present in the concurrent infection were

serotypes 2 and 3. The DENV-2 serotype is divided into six genotypes: Asian I, Asian II,

American, Asian/American, Cosmopolitan, and Sylvatic [51]. The cosmopolitan genotype of

DENV-2 is the most widely distributed dengue virus worldwide. The DENV-3 serotype of

genotype III is found in Africa [52]. Furthermore, *Plasmodium falciparum* was the only malaria

parasite specified in the coinfection among the included studies. It is known that nearly all malaria

cases in (sub-Saharan) Africa are caused by *P. falciparum* [5].

Our study also reported significantly higher odds of malaria infection among patients with malaria

mono-infection than acute dengue virus co-infection. This could be due to the inter-species

cross-protection of dengue virus infection against malaria [11]. Our finding does not corroborate

the reports of Kotepui et al. 2019 [47] and Kotepui et al. 2020 [53]. Kotepui et al. 2019 showed

significantly lower odds of co-infection while Kotepui et al. 2020 reported no notable difference

in co-infection due to acute dengue compared to malaria mono-infection. Our study further

elucidated no significant odds of acute dengue virus co-infection infection due to malaria than

acute dengue virus mono-infection. Our report is in disagreement with the findings of Kotepui et

al. 2020 [53] which indicated higher odds of acute dengue due to malaria co-infection in contrast

to dengue virus mono-infection. The discordant findings could be attributed to the variation in

model estimator/effect size model measures and/or quality/geography of included studies.

In general, our study found a high prevalence of concurrent malaria and dengue fever among

febrile patients in (sub-Saharan) Africa with significant variation across regions. Women and

children were more affected population groups by the coinfection than men and adults,

respectively. *P. falciparum* and dengue virus serotypes 2 and 3 were the predominant aetiologies

of this coinfection in the region. Healthcare workers should bear in mind the possibility of dengue

infection as one of the differential diagnoses for acute febrile illness as well as the possibility of

co-existent malaria and dengue in endemic areas. Our study also showed significantly higher odds

of malaria infection were documented due to acute dengue co-infection in contrast to malaria

mono-infection. Furthermore, no notable odds of acute dengue infection were reported due to

malaria co-infection as compared to dengue mono-infection. Besides, high-quality multi-centre

prospective studies are required to verify the above conclusions and to gain more insights into

malaria and dengue virus co-infection in the continent.

Materials and Methods

The protocol of the review was registered in the International Prospective Register of Systematic Reviews, PROSPERO (CRD42022311301) and follows the Joanna Briggs Institute (JBI) Checklist for Prevalence Studies [54].

Databases and search strategy

The CoCoPop mnemonic (condition, context, and population) [54] was used to formulate the review question and systematically search all relevant studies published between January 01, 2005, to March 30, 2022, from MEDLINE and EMBASE databases. The search strategy used was: **(exp Malaria, Falciparum/ or Malaria/ or ((*Plasmodium falciparum* or *Plasmodium* infect* or malignant tertian or (malaria or marsh or remittent)) adj3 fever).ti,ab.) and (exp Dengue/ or exp Dengue Virus/ or ((dengue or Break?bone) adj3 fever).ti,ab.).**

Inclusion and exclusion criteria

Case report, cross-sectional and case-control studies that reported concurrent malaria and dengue fever among uncomplicated febrile cases in African regions were included. According to the United Nations, Africa is divided into five regions: Northern Africa, Central or Middle Africa, Southern Africa, East Africa, and Western Africa [55]. Similarly, the World Bank lists a total of 48 countries in the sub-Saharan African region [56].

Malaria might be diagnosed by malaria rapid diagnostic tests or microscopy and/or polymerase chain reaction while dengue fever might be identified through antigen/antibody test and/or reverse

transcriptase-polymerase chain reaction. Acute dengue or dengue fever was defined as positive for

dengue IgM or NS1 antigen test or RT-PCR.

Articles of which full texts could not be obtained or were not available in English were excluded.

The primary outcome measure was the prevalence of concurrent malaria and dengue infection in

Africa. The secondary outcomes were: (1) the proportion of males and females with concurrent

malaria and acute dengue, and (2) the proportion of dengue virus serotypes in the co-infection. The

literature screening and selection process is depicted in Fig Fig .

Figure 10. Flowchart showing identification and selection process of articles for inclusion in the review.
279

Study quality appraisal and data extraction

The Joanna Briggs Institute System for the Unified Management, Assessment and Review of

Information (JBI SUMARI) tool [57] was used to screen each article and extract relevant data for

the review. Two of the authors (TT, JD) independently screened each article at an abstract and

full-text level. The discrepancy between the two reviewers was resolved through discussion.

Articles endorsed in the full-text screening were subjected to JBI critical appraisal tool. Those with

good quality scores were undergone data extraction. Data extraction included the first author's last

name, publication year, country/region of study, sample size, number of concurrent malaria and

dengue infections, demographic characteristics (age, gender) of patients with concurrent

infections, and dengue serotypes in the concurrent infections. The JBI criteria were used to score

the quality of each study. Studies with a score greater than or equal to four were seen as good

quality to be included in the meta-analysis.

Statistical analysis

The meta-analysis was performed using Jamovi software (version 2.2.5) [58], OpenMeta-analyst

[59], and the JBI SUMARI tool [57]. The random-effect model (DerSimonian-Laid) and the effect

size model measures of raw proportion/Freeman-Tukey double arcsine transformed proportion

were used to determine the prevalence/proportion estimates. The heterogeneity of included studies

was assessed using Tau, Tau^2, I^2 statistic, and Cochran's Q-test. Publication bias was evaluated

using Egger's regression test and Rank Correlation Test. P-value <0.05 was considered statistically

significant.

Funding: Not applicable.

Acknowledgement: Not applicable.

Conflict of interest: The authors have no conflict of interest to declare.

References

1. Escadafal C, Geis S, Siqueira AM, Agnandji ST, Shimelis T, Tadesse BT, et al. Bacterial versus non-
bacterial infections: a methodology to support use-case-driven product development of diagnostics. BMJ
Glob Health. 2020;5(10):e003141.

2. Bhargava A RR, Chatterjee B, Bottieau E. Assessment and initial management of acute
undifferentiated fever in tropical and subtropical regions. BMJ. 2018;363:k4766.

3. Prasad N, Murdoch DR, Reyburn H, Crump JA. Etiology of severe febrile Illness in low- and middle-
income countries: a systematic review. PLoS One. 2015;10(6):e0127962.

4. World Health Organization. World Malaria Report. Geneva, Switzerland: WHO; 2018 [Available from:
https://www.afro.who.int/sites/default/files/halth_topics_infographics/WHO_infographics_malaria.pdf.

5. Snow RW, Omumbo JA. Disease and Mortality in Sub-Saharan Africa. 2nd edition. Chapter 14 Malaria.
Washington (DC): The International Bank for Reconstruction and Development / The World Bank; 2006
[Available from: https://www.ncbi.nlm.nih.gov/books/NBK2286/.

6. Gainor EM, Harris E, LaBeaud AD. Uncovering the burden of dengue in Africa: considerations on
magnitude, misdiagnosis, and ancestry. Viruses. 2022;14(2):233.

7. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and
burden of dengue. Nature. 2013;496(7446):504-7.

8. World Health Organization. Dengue and severe dengue.: WHO; 2022 [Available from:
https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue.

9. Epelboin L, Hanf M, Dussart P, Ouar-Epelboin S, Djossou F, Nacher M, et al. Is dengue and malaria
co-infection more severe than single infections? A retrospective matched-pair study in French Guiana.
Malar J. 2012;11:142.

10. Ward DI. A case of fatal Plasmodium falciparum malaria complicated by acute dengue fever in East
Timor. Am J Trop Med Hyg. 2006;75(1):182-5.

11. Bygbjerg IC, Simonsen L, Schiøler KL. Elimination of falciparum malaria and emergence of severe
dengue: an independent or interdependent phenomenon? Front Microbiol. 2018;9:1120.

12. Wiwanitkit V. Concurrent malaria and dengue infection: a brief summary and comment. Asian Pac J
Trop Biomed. 2011;1(4):326-7.

13. Abdul-Ghani R, Mahdy MAK, Alkubati S, Al-Mikhlafy AA, Alhariri A, Das M, et al. Malaria and
dengue in Hodeidah city, Yemen: High proportion of febrile outpatients with dengue or malaria, but low
proportion co-infected. PLoS One. 2021;16(6):e0253556.

14. Charrel RN, Brouqui P, Foucault C, de Lamballerie X. Concurrent dengue and malaria. Emerg Infect
Dis. 2005;11(7):1153-4.

15. Charrel RN, Brouqui P, Foucault C, de Lamballerie X. Concurrent dengue and malaria. Emerg Infect
Dis. 2005;11(7):1153-4.

16. Baba M, Logue CH, Oderinde B, Abdulmaleek H, Williams J, Lewis J, et al. Evidence of arbovirus
co-infection in suspected febrile malaria and typhoid patients in Nigeria. J Infect Dev Ctries.
2013;7(1):51-9.

17. Chipwaza B, Mugasa JP, Selemani M, Amuri M, Mosha F, Ngatunga SD, et al. Dengue and
chikungunya fever among viral diseases in outpatient febrile children in Kilosa District Hospital,
Tanzania. PLoS Negl Trop Dis. 2014;8(11).

18. Oyero OG, Ayukekbong JA. High dengue NS1 antigenemia in febrile patients in Ibadan, Nigeria.
Virus Res. 2014;191:59-61.

19. Stoler J, Delimini RK, Bonney JHK, Awandare GA. Evidence of recent dengue exposure among
malaria parasite-positive children in three urban centers in Ghana. Am J Trop Med Hyg. 2014;91(5
SUPPL. 1):427.

20. Ayorinde AF, Oyeyiga AM, Nosegbe NO, Folarin OA. A survey of malaria and some arboviral
infections among suspected febrile patients visiting a health centre in Simawa, Ogun State, Nigeria.
Journal of Infection and Public Health. 2016;9(1):52-9.

21. Sow A, Loucoubar C, Diallo D, Ndiaye Y, Senghor CS, Dia AT, et al. Concurrent malaria and
arbovirus infections in Kedougou, southeastern Senegal. Malar J. 2016;15(1):47.

22. Vu DM, Ripp K, Mutai N, Ndenga BA, Heath C, LaBeaud AD. Plasmodium falciparum co-infection
modulates dengue disease severity. Am J Trop Med Hyg. 2016;95(5 Supplement 1):32.

23. Dariano DF, Taitt CR, Jacobsen KH, Bangura U, Bockarie AS, Bockarie MJ, et al. Surveillance of
vector-borne infections (chikungunya, dengue, and malaria) in Bo, Sierra Leone, 2012-2013. Am J Trop
Med Hyg. 2017;97(4):1151-4.

24. Kolawole OM, Seriki AA, Irekeola AA, Bello KE, Adeyemi OO. Dengue virus and malaria
concurrent infection among febrile subjects within Ilorin metropolis, Nigeria. J Med Virol.
2017;89(8):1347-53.

25. Miri MF, Mawak JD, Chukwu CO, Chuwang NJ, Acheng SY, Ezekiel T. Detection of IgM and IgG
dengue antibodies in febrile patients suspected of malaria attending health centre in Jos, Nigeria Annals of
Medical Laboratory Science. 2017;1(1):27-35.

26. Boillat-Blanco N, Klaassen B, Mbarack Z, Samaka J, Mlaganile T, Masimba J, et al. Dengue fever in
Dar es Salaam, Tanzania: clinical features and outcome in populations of black and non-black racial
category. BMC Infect Dis. 2018;18(1):644.

27. Monamele GC, Demanou M. First documented evidence of dengue and malaria co-infection in
children attending two health centers in yaounde, cameroon. Pan Afr Med J. 2018;29:227.

28. Obonyo M, Fidhow A, Ofula V. Investigation of laboratory confirmed dengue outbreak in north-
eastern Kenya, 2011. PLoS One. 2018;13(6):e0198556.

29. Onyedibe K, Dawurung J, Iroezindu M, Shehu N, Okolo M, Shobowale E, et al. A cross sectional
study of dengue virus infection in febrile patients presumptively diagnosed of malaria in Maiduguri and
Jos plateau, Nigeria. Malawi medical journal : the journal of Medical Association of Malawi.
2018;30(4):276-82.

30. Tchuandom SB, Tchouangueu TF, Antonio-Nkondjio C, Lissom A, Djang JON, Atabonkeng EP, et
al. Seroprevalence of dengue virus among children presenting with febrile illness in some public health
facilities in Cameroon. Pan Afr Med J. 2018;31:177.

31. Youssou FBS, Nemg FBS, Ngouanet SA, Mekanda FMO, Demanou M. Detection and serotyping of

dengue viruses in febrile patients consulting at the New-Bell District Hospital in Douala, Cameroon.

PLoS One. 2018;13(10):e0204143.

32. Nassar SA, Olayiwola JO, Bakarey AS, Enyhowero SO. Investigations of dengue virus and

Plasmodium falciparum among febrile patients receiving care at a tertiary health facility in Osogbo,

south-west Nigeria. Nigerian Journal of Parasitology. 2019;40(1):18-24.

33. Otu AA, Udoh UA, Ita OI, Hicks JP, Egbe WO, Walley J. A cross-sectional survey on the

seroprevalence of dengue fever in febrile patients attending health facilities in Cross River State, Nigeria.

PLoS One. 2019;14(4):e0215143.

34. Proesmans S, Katshongo F, Milambu J, Fungula B, Mavoko HM, Ahuka-Mundeke S, et al. Dengue

and chikungunya among outpatients with acute undifferentiated fever in Kinshasa, Democratic Republic

of Congo: A crosssectional study. PLoS Negl Trop Dis. 2019;13(9):e0007047.

35. Abdulaziz MM, Ibrahim A, Ado M, Ameh C, Umeokonkwo C, Sufyan MB, et al. Prevalence and

factors associated with dengue fever among febrile patients attending secondary health facilities in Kano

metropolis, Nigeria. African J Clin Exp Microbiol. 2020;21(4):340-8.

36. Ali MA, James OC, Mohamed AA, Joachim A, Mubi M, Omodior O. Etiologic Agents of Fever of

Unknown Origin Among Patients Attending Mnazi Mmoja Hospital, Zanzibar. J Community Health.

2020;45(5):1073-80.

37. Ihahi JW, Oyugi E, Owiny M, Ogutu J, Serech H. Investigation of outbreak of malaria and other

febrile illnesses in a remote Semi-Arid County, Kenya, 2019. Am J Trop Med Hyg. 2020;103(5

SUPPL):69.

38. Mouiche MMM, Ntumvi NF, Maptue VT, Tamoufe U, Albert B, Ngum Ndze V, et al. Evidence of

Low-Level Dengue Virus Circulation in the South Region of Cameroon in 2018. Vector-Borne and

Zoonotic Diseases. 2020;20(4):314-7.

39. Shah MM, Ndenga BA, Mutuku FM, Vu DM, Grossi-Soyster EN, Okuta V, et al. High dengue burden

and circulation of 4 virus serotypes among children with undifferentiated fever, Kenya, 2014-2017.

Emerg Infect Dis. 2020;26(11):2638-50.

40. Ahmed A, Eldigail M, Elduma A, Breima T, Dietrich I, Ali Y, et al. First report of epidemic dengue

fever and malaria co-infections among internally displaced persons in humanitarian camps of North

Darfur, Sudan. Int J Infect Dis. 2021;108:513-6.

41. Galani BRT, Mapouokam DW, Simo FBN, Mohamadou H, Chuisseu PDD, Njintang NY, et al.

Investigation of dengue-malaria coinfection among febrile patients consulting at Ngaoundere Regional

Hospital, Cameroon. J Med Virol. 2021;93(6):3350-61.

42. Marks F, Liu J, Soura AB, Gasmelseed N, Operario DJ, Grundy B, et al. Pathogens that cause acute

febrile illness among children and adolescents in Burkina Faso, Madagascar, and Sudan. Clin Infect Dis

2021;73(8):1338-45.

43. Nkenfou CN, Fainguem N, Dongmo-Nguefack F, Yatchou LG, Kameni JJK, Elong EL, et al.

Enhanced passive surveillance dengue infection among febrile children: Prevalence, co-infections and

associated factors in Cameroon. PLoS Negl Trop Dis. 2021;15(4):e0009316.

44. Akelew Y, Pareyn M, Lemma M, Negash M, Bewket G, Derbew A, et al. Aetiologies of acute

undifferentiated febrile illness at the emergency ward of the University of Gondar Hospital, Ethiopia.

Trop Med Int Health. 2022;27(3):271-9.

45. Tchetgna HDS, Yousseu FS, Kamgang B, Tedjou A, McCall PJ, Wondji CS. Concurrent circulation

of dengue serotype 1, 2 and 3 among acute febrile patients in Cameroon. Int J Infect Dis.

2022;116(Supplement):S125.

46. Brockhaus AC, Bender R, Skipka G. The Peto odds ratio viewed as a new effect measure. Stat Med.

2014;33(28):4861-74.

47. Kotepui M, Kotepui KU. Prevalence and laboratory analysis of malaria and dengue co-infection: a

systematic review and meta-analysis. BMC Public Health. 2019;19(1):1148.

48. Quaresima V, Agbenyega T, Oppong B, Awunyo JADA, Adomah PA, Enty E, et al. Are malaria risk

factors based on gender? A mixed-methods survey in an urban setting in Ghana. Trop Med Infect Dis.

2021;6(3).

49. Schumacher RF, Spinelli E. Malaria in children. Mediterr J Hematol Infect Dis. 2012;4(1):e2012073.

50. Tantawichien T. Dengue fever and dengue haemorrhagic fever in adolescents and adults. Paediatr Int

Child Health. 2012;32 Suppl 1(s1):22-7.

51. Waman VP, Kolekar P, Ramtirthkar MR, Kale MM, Kulkarni-Kale U. Analysis of genotype diversity

and evolution of Dengue virus serotype 2 using complete genomes. PeerJ. 2016;4:e2326.

52. Dorji T, Yoon IK, Holmes EC, Wangchuk S, Tobgay T, Nisalak A, et al. Diversity and origin of

dengue virus serotypes 1, 2, and 3, Bhutan. Emerg Infect Dis. 2009;15(10):1630-2.

53. Kotepui M, Kotepui KU, Milanez GJ, Masangkay FR. Prevalence of and risk factors for severe

malaria caused by Plasmodium and dengue virus co-infection: a systematic review and meta-analysis.

Infect Dis Poverty. 2020;9(1):134.

54. Munn Z, Moola S, Lisy K, Riitano D, Tufanaru C. Methodological guidance for systematic reviews of

observational epidemiological studies reporting prevalence and cumulative incidence data. Int J Evid

Based Healthc. 2015;13(3):147-53.

55. Pitt. African Studies and African Country Resources: University of Pittsbergh; 2022 [Available from:

https://pitt.libguides.com/.

56. World B. All countries of Sub-Saharan Africa. : World Bank; 2022 [Available from:

https://openknowledge.worldbank.org/pages/focus-sub-saharan-africa.

57. SUMARI J. End to end support for developing systematic reviews. 2022 [Available from:

https://sumari.jbi.global/.

58. Project TJ. Jamovi (version 2.2), computer software. 2021 [Available from: https://www.jamovi.org.

59. Wallace BC, Dahabre IJ, Trikalinos TA, Lau J, Trow P, Schmid CH. Closing the gap between

methodologists and end-users: R as a computational back-end. J Stat Softw. 2012;49(5):1-15.

Supporting information captions

Fig 1. Forest plot delineating prevalence of malaria and acute dengue co-infection in (sub-Saharan)

Africa.

Fig 2. Forest plot showing the prevalence of malaria and acute dengue co-infection in West Africa.

Fig 3. Prevalence of malaria and dengue co-infection in East Africa.

Fig 4. Forest plot depicting prevalence of concurrent malaria and acute dengue in Central Africa.

Fig 5. Forest plot: the difference in the prevalence of malaria and dengue virus concurrent infection with

malaria mono-infection.

Fig 6. Forest plot showing the difference in the prevalence of acute dengue with presence versus absence

of malaria infection.

Fig 7. Funnel plot showing assessment of publication bias based on the prevalence of concurrent malaria

and acute dengue in (sub-Saharan) Africa.

Fig 8. Funnel Plot depicting evaluation of publication bias based on the difference in the prevalence of

concurrent malaria and acute dengue with malaria mono-infection in (sub-Saharan) Africa.

Fig 9. Funnel Plot: evaluation of publication bias based on the difference in the prevalence of concurrent

malaria and acute dengue with acute dengue mono-infection in (sub-Saharan) Africa.

Fig 10. Flowchart showing identification and selection process of articles for inclusion in the

review.