Population-weighted greenspace exposure tied to lower COVID-19 mortality rates: A nationwide dose-response study

Yuwen Yang[a, b], Yi Lu[c], Bin Jiang[a, b], *

Affiliations:

[a]Urban Environments and Human Health Lab, HKUrbanLabs, Faculty of Architecture, The University of Hong Kong, HK.

[b]Division of Landscape Architecture, Department of Architecture, The University of Hong Kong, HK.

[c]Department of Architecture and Civil Engineering, College of Engineering, City University of Hong Kong, HK.

*Corresponding author: Bin Jiang, RM614, 6/F, Knowles Building, The University of Hong Kong, Pok fu lam Road, Hong Kong; email: jiangbin@hku.hk

Population-weighted greenspace exposure tied to lower COVID-19 mortality rates: A nationwide dose-response study

Abstract

The COVID-19 outbreak has caused enormous deaths and profound social and economic disruption globally. Accumulating evidence suggests exposure to greenspace may reduce the risk of COVID-19 mortality. Greenspace exposure enhances immune functioning, reduces inflammation, and replenishes gut microbiota may protect against the risk of mortality among those with COVID-19. However, previous studies often fail to distinguish the health effect of different types of greenspace, explore the dose-response association and optimal buffer distance, and consider the spatial dynamics of population distribution and geographic locations of greenspace.

This study examined the associations among ratio of different types of greenspaces, population-weighted exposure to different types of greenspaces, and COVID-19 mortality rates using a negative binomial generalized linear mixed effects model across 3,025 counties, adjusted for socioeconomic, demographic, pre-existing chronic disease, policy and regulation, behavioral, and environmental factors. The population-weighted measure gave proportionally greater weight to greenspace near areas of higher population density.

Exposure to forest and pasture was negatively associated with COVID-19 mortality rates, while developed open space has insignificant or positive associations with mortality rates. *Forest outside park* has the largest effect size across all buffer distances, followed by *forest inside park*. The optimal exposure buffer distance is 1km for *forest outside park*, with 1 unit of increase in exposure associated with a 9.9% decrease in mortality rates (95% confidence interval: 6.9% -12.8%). The optimal exposure buffer distance of *forest inside park* is 400m, with 1 unit of increase in exposure, associated with a 4.7% decrease in mortality rates (95% confidence interval: 2.4% - 6.9%).

Greenspaces, especially nearby forest, may be effective at lowering the mortality risk of COVID-19 patients. Our findings suggest that policymakers and planners should prioritize forestry within walking distance of residential clusters to mitigate mortality rates during current and future respiratory pandemics.

Keywords: *COVID-19 mortality rate; Greenspace; Forest; Dose-response; Optimal buffer distance*

1. Introduction

Since its outbreak in 2019, COVID-19 has spread rapidly throughout the world, leading to numerous infections and deaths. In the United States, COVID-19 is largely responsible for the substantial 17.7% increase in total deaths from 2019 to 2020 and became the third leading cause of death following heart disease and cancer (Ahmad, Cisewski, Miniño, & Anderson, 2021). By the end of 2020, COVID-19 deaths were estimated at 348,600; by February 2020, they had reached over 933,000 (Johns Hopkins University & Medicine, 2021; Viglione, 2020). In the US, COVID-19 was estimated to reduce life expectancy by 1.13 years annually (Andrasfay & Goldman, 2021).

The COVID-19 pandemic also overwhelmed healthcare systems and caused substantial economic loss. COVID-19 infection and deaths cast escalating pressure on testing capacities and hospitalizations in the U.S. (Dyer, 2020; Miller, Becker, Grenfell, & Metcalf, 2020). Critically ill COVID-19 patients faced shortages in intensive care units (ICUs) compounded by other critical health conditions (Halpern & Tan, 2020). The cumulative economic costs of the COVID-19 pandemic due to premature deaths, unemployment, and business revenue decline was estimated to be US$1.4 trillion GDP by 2030 (Cutler & Summers, 2020; Chen et al. 2021).

Accumulating evidence suggests links between both nature and the built environment and COVID-19 mortality rates. Exposure to air pollution (Ali & Islam, 2020; Konstantinoudis et al., 2021; Liang et al., 2020), crowded housing (Brandén et al., 2020; Hu, Roberts, Azevedo, & Milner, 2021; van Ingen et al., 2021), and lower average temperature (Ma et al., 2020; Perone, 2021) were found to increase COVID-19 deaths. The relationship between greenspace and COVID-19 mortality rates has received far less attention (Jiang, Yang, et al., 2021; Klompmaker et al., 2021; Lu, Chen, et al., 2021), despite the numerous salutary effects of nature exposure on human health.

1.1 How might exposure to greenspace alleviate COVID-19 mortality rates?

An overwhelming amount of research has shown that exposure to greenspace can improve both

physical and mental health (Jiang, Chang, & Sullivan, 2014; Lu, Chen, et al., 2021). Particularly,

studies have shown that contact with greenspace boosts our defense capacity against viruses by

increasing Natural Killer (NK) and T cells and cytotoxic activities (Liisa Andersen, Sus Sola Sola

Corazon, & Ulrika Karlsson Karlsson Stigsdotter, 2021; Li, 2010; Roviello, Gilhen-Baker,

Vicidomini, & Roviello, 2021), reducing inflammation (Kuo, 2015; Ribeiro, Tavares, Guttentag,

& Barros, 2019), and replenishing gut microbiota (Parajuli, 2019; Parajuli et al., 2020; Marja I.

Roslund et al., 2020). Hospitalized COVID-19 patients with severe or fatal cases have consistently

shown immune interference (e.g., lower NK and T cell count, exaggerated cytotoxic activities)

(Castelli, Cimini, & Ferri, 2020; Girija, Shankar, & Larsson, 2020; Qin et al., 2020), hyper-

inflammation or 'cytokine storm'(e.g., delayed but elevated of pro-inflammatory cytokines)

(Paranjpe et al., 2020; Potempa, Rajab, Hart, Bordon, & Fernandez-Botran, 2020; Yang et al.,

2020), and decreased gut microbiota diversity (Dhar & Mohanty, 2020) compared to non-critically

ill patients. Thus, contact with nature has the potential to mitigate severe COVID-19 prognosis

and deaths.

1.2 A critical gap: The relationship between different types of greenspace and COVID-19

mortality rate.

Several studies have shown a significant association between greenness and COVID-19

mortality rates in the U.S. (Klompmaker et al., 2021; Lee et al., 2021; Russette et al., 2021;

Spotswood et al., 2021). These studies define greenspace as the total area of vegetation within a

boundary (e.g., Normalized Difference Vegetation Index and Leaf Area Index).They did not

distinguish between open space, forest, grassland/herbaceous, and hay/pasture, nor did they

consider the impacts of greenspace proximity and recreational function provision on COVID-19

health outcomes. While previous evidence suggests different types of greenspace does not have

same impacts on health outcomes (Akpinar, Barbosa-Leiker, and Brooks 2016; Ekkel & de Vries

2017; Kim & Miller 2019; Johnson et al. 2020; Ma et al. 2022). For instance, greenspace such as

greenness and park had negative association with COVID-19 infection (Russette et al. 2021;

Spotswood et al. 2021; Wang et al. 2021; Johnson et al. 2020), but park mobility and green space

with better accessibility were positively associated with COVID-19 transmission (Pan, Bardhan,

and Jin 2021; DePhillipo et al., 2021). We still do not know whether and to what extent different

types of greenspace can influence the COVID-19 mortality rates.

Second, existing studies estimate the amount of greenness in a county but ignore the spatial

distribution of greenspace in relation to population (Klompmaker et al., 2021; Russette et al., 2021).

Despite being widely used and effective, the accuracy of 'greenness' metric can be greatly

improved by considering spatial relations between location of greenspace and population

distribution (Ben et al., 2019). Further, one study used deciles of 'greenness' to assess the dose-

response association (Russette et al., 2021). Though, the dose-response associations for different

types of greenspace within various buffer distances are unclear. Many previous studies suggested

that distance matters for greenspace's impact on health outcomes, and health effect might drop

after a threshold distance (Coombes, Jones, and Hillsdon 2010; Grahn and Stigsdotter 2003;

Nielsen and Hansen 2007). Still, we do not know whether the association of nearby greenspace on

mortality risk is significantly stronger than distant ones. We do not know which distances are

optimal. Without addressing these critical gaps, policymakers and urban planners are unable to

develop evidence-based urban greening solutions and policy to promote public health for current

and future pandemics.

1.3 Research questions

In this study, we investigated the associations among the ratio of six types of greenspace,

population-weighted exposure to six types of greenspace at different buffer distances, and full-

year COVID-19 mortality rates after controlling for potential confounding covariates. We seek to

answer the following three questions: 1) What are the associations between the ratio of six types

of greenspace and COVID-19 mortality rates after controlling for confounding variables? 2) What

are the associations between population-weighted exposure to six types of greenspace and

COVID-19 mortality rates within various buffer distances after controlling for confounding

variables? (3) Which exposure distances of the significant types of greenspace have the strongest

association with COVID-19 mortality rates?

2. Methods

We combined COVID-19 mortality data, sociodemographic characteristics, healthcare and

testing data, pre-existing chronic disease data, policy and regulation data, behavior data, and

environmental factors from diverse sources for 3,025 counties. The greenspace exposures were

calculated in GEE. We first used a negative binomial generalized linear mixed effects model to

evaluate the association between the ratio of six types of greenspace and COVID-19 mortality

rates in the U.S. from January 22 to December 31, 2020, adjusted for socioeconomic, demographic,

pre-existing chronic disease, policy and regulation, behavioral, and environmental factors. Then,

we examined the associations between the population-weighted exposure to greenspace at varying

distances within 4km and COVID-19 mortality rates, adjusted for confounders.

2.1 Data

2.1.1 COVID-19 mortality data

COVID-19 mortality data are publicly available at the US Centers for Disease Control and

Prevention (CDC) and State government websites (Kolak et al., 2021). We define the COVID-19

mortality rates as the cumulative number of COVID-19 deaths per 100,000 people for each of

3,025 counties from January 22, 2020 to December 31, 2020 (Fig. 1). We limit our research period

at the end of 2020 to avoid the possible confounding effect from large-scale vaccination, which

will have significantly impact on mortality rates (see Supplementary Table 1 for descriptive

COVID-19 mortality data).

Fig. 1 | County-level COVID-19 deaths per 100,000 population in the United States (from
Jan 18 to Dec 30, 2020).
151
2.1.2 Greenspace exposure data

We assessed greenspace exposure using two metrics. First, we quantified the ratios of six types

of greenspaces within a county, which measures the area of each greenspace over the county area.

The ratio of *forest inside park*, *forest outside park*, grassland/herbaceous, hay/pasture, *open space*

inside park, and *open space outside park* were calculated use the National Land Cover Database

2016 (NLCD, 2016) (Fig. 2). We distinguished open space and forest within park from open

space and forest outside park use the boundary derived from the USA parks from Esri (Esri,

2021).

Fig. 2 | Ratio of greenspace at the county level in the United States. (**A**) forest inside park (**B**) forest outside park;
(**C**) hay/pasture; (**D**) grassland/herbaceous; (**E**) developed open space inside park; (**F**) developed open space outside
park.

Second, we quantified the population-weighted exposure to greenspaces within different

distances from human settlements using two datasets: the National Land Cover Datasets in 2016

(Yang et al., 2018) and the 2020 WorldPop Global Project Population Data (Sorichetta et al.,

2015). The 30-meter resolution NLCD 2016 Landsat imagery was re-projected to match the 100-

168 meter spatial resolution of WorldPop Dataset. We estimated population-weighted greenspace

169 exposures within 4km in GEE (Gorelick et al., 2017), because past studies suggest few walking

170 activities occur beyond 4km (Yang & Diez-Roux, 2012). These measures considered population

171 spatial distributions and gave proportionally greater weight to greenspace near areas of higher

172 population density, which previous studies fail to address (Chen et al., 2018). Considering

173 population distribution in greenspace exposure measurement can reduce bias caused by a

174 mismatch between population and greenspaces within an area. The buffer interval is set as 200m

175 within 2km and 500m between 2km to 4m. The population-weighted exposure to greenspaces

176 with varying buffer sizes in each county is defined by Equation 1 (Chen et al., 2018),

177          $$FE = \frac{\sum_{i=1}^{N} P_i \times F_i^b}{\sum_{i=1}^{N} P_i} \text{ (Equation 1)}$$

178 where P_i represents the population of the i^{th} grid, F_i^b represents the land cover of the i^{th} grid at a

179 buffer size of b meters, N denotes the total number of grids for a given county, and FE is the

180 estimated greenspace exposure level for the given county (see Supplementary Table 2 for

181 descriptive greenspace exposure data).

182 *2.1.3. Predictors of COVID-19 mortality*

183  We considered a number of predictors of COVID-19 mortality as potential covariates in analyses.

184 Many studies have found sociodemographic, chronic disease, behavioral, healthcare, and

185 environmental factors linked to COVID-19 mortality. The county-level sociodemographic,

186 healthcare, and testing data from the US Census Bureau (US Census Bureau, 2019) and US

187 COVID Atlas of the Center for Spatial Data Science (Kolak et al., 2021). These variables included

188 population density, the ratio of female household, non-Hispanic black, white, and Hispanic, the

189 proportion of residents older than 65, median household income, Gini index, poverty rate, median

190 housing value, unemployment rate, Gini index, the ratio of residents without a high school diploma

and without a college degree, percent without health insurance coverage, and COVID-19 testing

rates.

The pre-existing chronic diseases were shown to affect COVID-19 mortality risk, included rates

of hypertension, heart failure, stroke mortality, diabetes, and obesity (CDC, 2021; Prevention.,

2022). The policy and regulation factors included stay-at-home order intensity, public mask

mandates, and bar and restaurant closing and reopening orders (Chernozhukov et al., 2021;

VoPham et al., 2020). The behavior risk factors included the proportion of current smokers and

the proportion of essential workers, the proportion of workers who commuted to work by public

transportation, walking, and private cars, and the proportion of leisure-time physical inactivity,

median max-distance traveled, and foot traffic to different out-of-home activities. The

environmental risk factors included particulate matter (e.g., $PM_{2.5}$ and PM_{10}), temperature, relative

humidity, precipitation, wind speed, and transportation density. All descriptive statistics of

covariates and data sources are available in Supplementary Table 1.

2.2 Statistical analysis

We used a negative binomial generalized linear mixed effects model to evaluate the associations

between the ratios of six types of greenspace and COVID-19 mortality rates, and state was used

in analyses as a random effect to account for state-level variability and non-independence in our

data. The analyses were adjusted for a range of covariates. We applied restricted maximum

likelihood (REML) with a negative binomial link function. The negative binomial mixed effect

model and state as random effect accounts for our over-dispersed count data and partially

accounts for the presence of spatial autocorrelation. The variance inflation factor (VIF) test was

used to identify multi-collinearity between the independent variables. Variables with a VIF ≥ 4

were excluded from our models (O'Brien, 2007).

To identify the optimal exposure distance for significant greenspaces, we used a negative

binomial generalized linear mixed effects model to evaluate associations between population-

weighted exposures to six types of greenspace with COVID-19 mortality rates. The analyses use

the same sets of covariates as previous analysis and state was used as random effect. All

explanatory variables were centered and scaled.

We used Moran's I test to assess spatial autocorrelation of COVID-19 mortality residuals. We

confirmed the presence of spatial autocorrelation with Moran's I=0.21, p < 0.0001. The Moran's

I value equal to 0 indicates a lack of spatial autocorrelation, and positive values indicate

clustering of similar values. The analyses were performed in R v.4.1.2 (Team, 2015), and Moran'

I test was performed using the package 'spdep' (Bivand & Wong, 2018). The negative binomial

mixed effects models were performed using the package lme4 (Bates, Mächler, Bolker, &

Walker, 2014).

2.3 Model validation

The negative binomial generalized linear mixed effects model provides an appropriate error

structure for the overdispersed COVID-19 mortality count data. Due to the presence of spatial

autocorrelation (Moran's I=0.21, p < 0.0001), we built additional spatial autoregressive models

(SAR) to validate the results of the negative binomial mixed effects model. The queen's criteria

were used to build the neighbors matrix. We used the Akaike information criterion (AIC) value

to compare the spatial error model, spatial lag model, and spatial Durbin model. The spatial error

model has the lowest AIC values, which suggests that spatial dependence occurs in the error

term. The model validation confirms the negative associations between *forest inside park*, *forest*

outside park, pasture, and COVID-19 mortality rates (See results of SAR models in

Supplementary Table 5). Given the structure of our data, model coefficients and the magnitude

of effects, we chose to interpret our result using the negative binomial generalized linear mixed

effects model.

3. Results

3.1 Associations among ratio of six types of greenspaces and COVID-19 mortality rates.

We found *forest inside park* and *forest outside park* are significantly negatively associated with

COVID-19 mortality rates ($p < 0.0001$); *open space outside park* is significantly positively

associated with COVID-19 mortality rates ($p < 0.01$); grassland/herbaceous, hay/pasture and *open*

space inside park are not significantly associated with COVID-19 mortality rates, after controlling

for all covariates (Fig. 3). Among the six types of greenspace, *forest outside park* has the greatest

effect size ($\beta = -0.097$), which is slightly larger than that of *forest inside park* ($\beta = -0.082$). We

found 1 unit of increase in *forest outside park* is associated with a 9.2% decrease in COVID-19

mortality rates (MRR 95% CI: 6.3 - 12.1%), whereas 1 unit of increase in *forest inside park* is

associated with a 7.8% decrease (MRR 95% CI: 4.3 - 11.2%). In contrast, 1 unit of increase in

developed open space outside park is associated with a 5.8% increase in COVID-19 mortality rates

(MRR 95% CI: 2.3 - 9.5%) (Table 1).

252
Fig. 3 | Exposure to forest is associated with lower COVID-19 mortality rates adjusted for covariates. Coefficient
values represent effect sizes for the associations between mortality rates of COVID-19 (cases per 100,000 people) and
ratio of grassland/herbaceous, hay/pasture, *open space in park*, *open space outside park*, *forest inside park*, *forest*
outside park, and all covariates. Coefficient values are represented as dots, bars represent 95% CI, and significant
variables are shown in color: grey = $p \geq 0.01$; yellow = $p < 0.01$; red = $p < 0.0001$.

Table 1| Mortality rate ratios (MRR), 95% confidence intervals (CI) of all variables in the model
for COVID-19 mortality rates.

Variables		MRR (95% CI)
Socioeconomic and demographic factors	Population density	1.073 (1.041, 1.105)
	Black non-Hispanic	1.057 (1.021, 1.095)
	Population aged 65+	1.162 (1.129, 1.196)
	Gini Index	1.019 (0.993, 1.045)
	Median home value	0.878 (0.842, 0.915)
	Unemployment rate	0.970 (0.941, 1.000)
	Population without high school diploma	1.139 (1.095, 1.185)
Healthcare and testing factors	Population without insurance	0.973 (0.920, 1.029)
	Testing rate	1.092 (1.060, 1.126)
Pre-existing chronic disease factors	Diabetes rate	1.018 (0.993, 1.045)
	Obesity rate	0.988 (0.961, 1.015)
	Stroke mortality	1.033 (1.004, 1.062)
	Hypertension mortality	1.013 (0.986, 1.040)
	Heart disease mortality	1.024 (0.996, 1.052)
Behavioral factors	Smoker	0.993 (0.951, 1.037)
	Essential worker	1.031 (0.996, 1.067)
	POI visits	1.022 (0.997, 1.047)
	Commute to work by walking or bicycle	0.987 (0.962, 1.014)
	Leisure time physical inactivity	1.042 (1.011, 1.074)
	Mobility	0.981 (0.962,1.000)
	Mobility 50index	0.988 (0.966, 1.011)
Policy and regulation factors	Stay-at-home orders	1.027 (0.987, 1.068)
	Public mask mandates	0.950 (0.831, 1.085)
	Bar	1.097 (0.956, 1.258)
	Restaurant	0.95 (0.808, 1.118)
Environmental factors	Crowded Housing	1.073 (1.042, 1.105)
	Proximity to highway	1.013 (0.990, 1.038)
	Airport density	0.968 (0.947, 0.989)
	Railway density	0.964 (0.938, 0.990)
	Highway and secondary road density	1.001 (0.964, 1.038)
	$PM_{2.5}$	1.015 (0.978, 1.052)
	PM_{10}	0.968 (0.924, 1.014)
	NO_2	1.012 (0.976, 1.050)
	Maximum temperature	0.975 (0.915, 1.038)
	Humidity	0.886 (0.842, 0.933)
	Wind speed	1.010 (0.980, 1.042)

Green space factors	Grassland / Herbaceous	1.016 (0.986, 1.046)
	Hay / Pasture	0.970(0.945, 0.995)
	Open space in park	1.024 (0.988, 1.060)
	Open space outside park	1.058 (1.023, 1.095)
	Forest in park	0.922 (0.888, 0.957)
	Forest outside park	0.908 (0.879, 0.937)

3.2 Associations of population-weighted exposures to greenspace with COVID-19 mortality

rates at various buffer distances.

We also found population-weighted exposure to *forest inside park*, *forest outside park* and

pasture are significantly and negatively associated with COVID-19 mortality rates. Population-

weighted exposure to *forest inside park* is significantly and negatively associated with mortality

rates between 100 to 400m and 1,800m to 4km, and the distance for largest effect size is 4km ($\beta =$

-0.050). The effect size increases as buffer distance gets larger, though the increase remains limited

(400m β= -0.048 vs 4km β = -0.050, 4% of increase) (Fig. 4). With 1 unit of increase in *forest*

exposure in park at 4km, there is a 4.9% decrease in COVID-19 mortality rates (MRR 95% CI:

2.7-7.0%) (Table 2).

The population-weighted to *forest outside park* is consistently and significantly negatively

associated with COVID-19 mortality rates across all buffer distances, and the greatest reduction

occurs at 1km (β = -0.104). The effect size increases as the buffer increases from 100m to 1km

and decrease beyond 1km (Fig. 5). We found a 9.9% decrease in COVID-19 mortality rates per

unit increase in *forest exposure outside park* at 1km (MRR 95% CI: 6.9–12.8%) (Table 2).

The population-weighted exposure to *pasture* is significantly negatively associated with

mortality rates from 2,500m to 4km with increasing effect size, and reaches optimal effect at 4km

(β = -0.036). With 1 unit of increase in pasture exposure at 4km, the associated mortality rates

decrease by 3.5% (MRR 95% CI: 5.9 - 10.0%) (Table 2).

Fig. 4 | The effect size of population-weighted exposure to *forest inside park* within 4km on COVID-19 mortality rates. Coefficient values represent effect sizes from a negative binomial mixed effects model for the relationship between mortality of COVID-19 mortality rates (death count per 100,000 people) and population-weighted exposure to *forest inside park*. Coefficient values are represented as dots, grey = $p > 0.05$; red = $p < 0.0001$.

Fig. 5 | The effect size of population weighted exposure to *forest outside park* within 4km on COVID-19 mortality rates. Coefficient values represent effect sizes from a negative binomial mixed effects model for the relationship between mortality of COVID-19 mortality rates (death count per 100,000 people) and population-weighted exposure to *forest outside park*. Coefficient values are represented as dots, red = $p < 0.0001$.

294 **Table 2 |** Mortality Rate Ratio in models of population-weighted exposure to greenspaces with
295 COVID-19 mortality rates at optimal distances.
296

Variables	COVID-19 MRR (95% CI)
Forest inside park 400m	0.953 (0.931, 0.976)
Forest inside park 4km	0.951 (0.930, 0.973)
Forest outside park 1km	0.901 (0.872, 0.931)
Hay/pasture 4km	0.965 (0.941, 0.990)

4. Discussion

We found exposure to forest and pasture to be negatively associated with COVID-19 mortality rates, while exposure to developed open space has mixed association with mortality rates. Exposure to *forest outside park* has the largest effect size on reduced COVID-19 mortality rates across all buffer distances, followed by *forest inside park*. Further, the effect size of exposure to *forest outside park* increases until the distance reaches 1km, then declines beyond 1km. The effect size of population-weighted exposure to *forest inside park* increase with larger buffer size and is greatest at 4km, though similar to that at 400m.

While this cross-sectional study cannot infer any causal relationships, previous findings suggest multiple mechanisms that might explain the observed associations. We proposed a framework of potential mechanisms that may contribute to the observed associations. We consider why exposure to *forest outside park* may have a larger effect size than *forest inside park*. We provide explanations for optimal exposure buffer size for significant greenspace types. Last, we discuss the contributions of our findings and identify questions for future research.

4.1. Potential mechanisms for observed associations.

4.1.1 How might forest and pasture alleviate COVID-19 mortality risk?

We found that forest and pasture exposure were significantly and negatively associated with COVID-19 mortality rates in the US, after control for all covariates. This finding aligns with previous studies (Klompmaker et al., 2021; Lee et al., 2021; Russette et al., 2021; Spotswood et al., 2021). Greenspace may lower COVID-19 mortality risk if it boosts biological processes that fight against the prognosis of COVID-19 (Andersen et al., 2021; Roslund et al., 2020; Roviello et al., 2021), making affected patients less vulnerable to death (Roviello & Roviello, 2021). We suggest that contact with greenspace may reduce mortality rates through increased biogenic

volatile organic compound (VOC) exposure, increased environmental microbiota exposure,

reduced psychological stress and air pollution, and increased physical activity (Fig. 6).

Fig. 6 | The proposed theoretical model for negative associations between greenspace exposure and COVID-19
mortality rates.

Exposure to the forest environment increases exposure to biogenic volatile organic

compounds (VOCs). Forests are abundant with phytoncides (e.g., terpenes, limonene and pinene),

a group of biogenic VOCs given off by forest trees that have been found to enhance immune

capacity and reduce inflammation (Andersen, Corazon, & Stigsdotter, 2021; Cho et al., 2017;

Huang et al., 2020; Kim, Song, Cho, & Lee, 2020). Studies suggest forest bathing increases natural

killer (NK) cell counts and activity, and the effect can last for more than a month (Li, 2010; Li et

al., 2008; Li et al., 2007; Tsao et al., 2018). NK cells activate receptors to recognize virus-infected

cells and trigger cytotoxicity (phagocytosis or apoptosis) (Market et al., 2020; Yokoyama, 2005).

A recent study found COVID-19 mortality rates are lower in areas with high ratios of hectares of

Mediterranean forest per capita, where biogenic VOCs are abundant (Roviello & Roviello, 2021).

In addition, studies have shown that nearby greenspaces are associated with a lower level of a

biomarker of inflammation - high sensitivity C-reactive protein (Del Valle et al., 2020; Mandel,

Harari, Gurevich, & Achiron, 2020; Ribeiro et al., 2019). Forests can increase exposure to VOCs

that might boost the NK defenses, compensate for reduced NK cell counts, and modulate excess

inflammatory responses in severely affected patients (Market et al., 2020; Osman et al., 2020);

Exposure to forest and pasture can diversify gut microbiota profile. Though our gut

microbial composition is shaped by the interplay of multiple factors, such as diet and genetics

(Dhar & Mohanty, 2020), many studies have found microbiome from surrounding green

environments can transfer to humans (Grönroos et al., 2019; Parajuli et al., 2018; Parajuli et al.,

2020). A 28-day intervention study in Finland found that daily contact with backyard forest and

grass areas within 500m can diversify children's gut microbiota profile and enhance immune

capacity (e.g., increases in plasma TGF-β1 levels and the proportion of regulatory T cells)

(Roslund et al., 2020). If exposed to greenspaces, the disturbed gut microbe condition and immune

function may improve for COVID-19 patients (Claesson et al., 2012; Donati Zeppa, Agostini,

Piccoli, Stocchi, & Sestili, 2020; Roslund et al., 2020; Yeoh et al., 2021; Zuo et al., 2020).

Greenspaces can decrease patients' exposure to air pollutants. Many studies have noted an

inverse association between air pollution and COVID-19 mortality rates (Jaafari, Shabani,

Moeinaddini, Danehkar, & Sakieh, 2020; Martelletti & Martelletti, 2020; Shen & Lung, 2017).

Chronic exposure to air pollutants was associated with delays in recovery of COVID-19 patients

and led to more fatal conditions (Domingo & Rovira, 2020). This may be attributed to modified

host respiratory immune responses, perturbed anti-microbial responses, and triggered

inflammatory cytokine release from air pollution exposure (Bauer, Diaz-Sanchez, & Jaspers, 2012;

Ciencewicki & Jaspers, 2007; Glencross, Ho, Camiña, Hawrylowicz, & Pfeffer, 2020). Forests can

reduce air pollutants by intercepting particulate matter on plant surfaces and absorbing gaseous

pollutants (Nowak, Hirabayashi, Bodine, & Greenfield, 2014; Nowak, Hirabayashi, Doyle,

McGovern, & Pasher, 2018). Trees and forests in the U.S. remove an estimated 17.4 million tonnes

(t) of air pollution annually (Nowak et al., 2014). It is thus reasonable to speculate that forests can

reduce COVID-19 mortality rates by removing air pollutants.

Forests can reduce patients' psychological stress. Patients infected with COVID-19 show a

high prevalence of mental problems (Kong et al., 2020; Wang et al., 2021). Mental stress has been

linked to dysregulation of the immune system and increased pro-inflammatory cytokines (Gouin

et al., 2012; Morey, Boggero et al., 2015; Steptoe et al, 2007). Patients with increased contact with

nature or nature views from home during the pandemic were found to have decreased depression

and anxiety (Soga et al., 2021). Previous theoretical and empirical studies support nature's stress-

reducing effect (Gidlow et al., 2016; Jiang et al., 2014; Lee et al., 2011; Mancus et al., 2020; Ulrich

et al., 1991). It is possible that patients can have strengthened immune function and healthier

inflammation level benefit from the stress-reduction effect of contact with forest.

Green space can promote physical activities (PA) during the pandemic. Since the COVID-

19 outbreak, people have escaped to nature. Recreational activities, such as walking or cycling in

parks and trails, have spiked globally during the pandemic (Geng et al., 2021; Lu et al., 2021;

Venter et al., 2020; Venter et al, 2021). Physical activity can boost COVID-19 patients' immune

response, modulate inflammation levels, and lower the risk of obesity. Exercise increases NK & T

cells, enhances recirculation, and increases lymphocyte concentration and cytotoxic activity

(Amatriain-Fernández et al., 2020; Fernandez et al., 2018; Nieman & Wentz, 2019). People who

exercise in forested areas can benefit from the synergistic effect from physical activity and

exposure to forest (Pretty et al, 2005). Moreover, physical activity has a direct anti-inflammatory

effect and can dampen systemic inflammation (Biddle et al., 2019; DeSantis et al., 2012; Nieman

& Wentz, 2019). Exercise lowers the risk of obesity, a risk factor for COVID-19. Obesity is the

precursor of a range of chronic diseases that have been found to increase COVID-19 mortality

(Bastien et al., 2014; Calle & Thun, 2004; Chan et al., 1994; Hussain et al., 2020; Klang et al.,

2020; Krauss et al., 1998). Thus, the benefits of greenspace on reduced obesity through physical

activity may contribute to the negative association between greenspace and COVID-19 mortality

rates (Coombes et al., 2010; De la Fuente et al., 2021; Ghimire et al., 2017; Jia et al., 2020).

4.1.2 How might open space exacerbates COVID-19 mortality risk?

We found the ratio of *open space outside park* is significantly and positively associated with

COVID-19 mortality rates. This finding contradicts previous studies that have reported health

benefits from exposure to open spaces. Our results suggest that exposure to open space may not

be effective or may even be detrimental to COVID-19 mortality rate. Open space in our study is

defined as "large-lot single-family housing units or vegetation planted in developed settings"

(NLCD, 2016). On one hand, open space can provide health benefits by promoting physical

activity, social interaction, and reduced air pollutants (Lu, Chen, et al., 2021), though the effect

may not be as strong as it is for forests (Reid, Clougherty, Shmool, & Kubzansky, 2017). On the

other hand, the lower supply of open space per capita in urban areas makes it hard to comply with

safe social distancing in non-park open spaces (e.g., streets and backyards), and may exacerbate

mortality risk. Though outdoor transmission of COVID-19 is rare (Bulfone et al. 2021), people

who participate in outdoor social activities such as talking or partying were at higher risk of

spreading the disease (Domènech-Montoliu et al. 2021; Peng et al. 2022). Presumably, the higher

infection risk might lead to higher COVID-19 mortality rates. More evidence is needed to

understand the mechanisms of the mixed effects of open space on COVID-19 mortality rates.

4.1.3 Why might forest outside park have a stronger effect than forest inside park?

We found *forest outside park* to have a larger effect size than *forest inside park* on COVID-19

mortality rates after accounting for other covariates. This finding aligns with previous studies of

stronger health-promoting effect outside park areas (Reid et al., 2017; Allard-Poesi et al., 2022).

The difference in forest exposure inside and outside park may explain the stronger effect of *forest*

outside park. The US population has a ten times greater opportunity to be exposed to *forest inside*

park than *forest outside park* within walking distance (Fig. 7). Second, social activities in parks

might increase risk of close contact and inhale droplets from infected people (DePhillipo et al.,

2021; Praharaj & Han, 2021), thus increase infection risk. Thus, it is reasonable to argue that the

increased infection risk caused by the social interactions in parks may offset the other health

benefits of *forests inside park* on mortality. Further, the health effect of *forest inside park* may be

weakened in part due to shutdown policies in some states, which closed parks due to COVID-19

spread risk (Volenec et al., 2021; Smith et al., 2021).

4.1.4 Optimal exposure buffer distance: what and why?

We found the effect of population-weighted exposure to *forest outside park* increases with a

larger buffer distance and reaches an optimal effect at 1km. This suggests that exposure to nearby

forest outside park within 1km is more effective than exposure to forests that are at a greater

walking distance. This may be because nearby forests are visited more often than forests located

further away. Studies suggest that the frequency of visits to greenspaces declines as distance

increases (Coombes et al., 2010). A distance of 1,120 m (0.7 miles) is the mean walking distance

in the U.S. (Yang & Diez-Roux, 2012).

We also found the effect size of *forest inside park* is optimal at 4km, though the effect size is

close to that at 400m (2% increase). This suggests the effect of *forest inside park* is less sensitive

to buffer size within walking distance. Studies suggest people walk much longer for recreation

purposes as opposed to other purposes (Yang & Diez-Roux, 2012). Considering the effect size and

previous literature on walking behaviors, we suggest the optimal exposure buffer distance for

forest outside park to be 1km and the optimal exposure buffer distance for *forest inside park* to be

400m.

Fig. 7 | Mean population-weighted exposure to *forest inside park* and *forest outside park* within 4km. The bar
represents the average population-weighted forest exposure at county level within each buffer distance.

4.2 Contributions and implications

To our knowledge, this is the first nationwide study distinguish the impact of different types of

greenspace on COVID-19 mortality rates in the U.S. In this study, the greenspace exposure

measure integrated the dynamics of population distribution and geographic location of greenspaces

into exposure assessment, providing a more precise and reasonable exposure estimates. The dose-

response associations between different types of green spaces at various buffer distances and

COVID-19 mortality rates were examined, allowing us to identify an optimal effect distance,

which was previously lacking in the literature.

Evidence from this study suggests that planners and policymakers should prioritize the supply

of nearby forests. Specifically, we recommend ensuring that forests outside parks be located within

1000m from residents, and forests inside parks be located within 400m from residents. These

forested areas will be especially beneficial in highly urbanized, low-SES, and minority-dominated

areas where COVID-19 mortality rates are disproportionally high (Lu et al., 2021). Many

greenspaces were temporarily closed during the pandemic to reduce the spread of disease (Ugolini

et al., 2020). The findings in this study advocate for keeping nearby greenspace open, especially

forested areas. Cities with accessible forested areas can promote health and resilience during the

current and future pandemics.

4.3 Limitations and future research opportunities

This study has several limitations, which pose opportunities for future research. This is an

ecological study using aggregated data at the county level. It is subjected to ecological fallacy.

Future studies can use individual level data or experimental studies to confirm the causal relations

and the potential underlying mechanisms (Jiang et al., 2021).

Second, the unit of analysis is the county due to the availability of COVID-19 mortality data

and other confounding variable. Though county data are widely used in nationwide studies, future

studies should use finer-grained data (i.e., census tract level data). Different scales of analyses may

reveal different associations between neighborhood greenspace and health outcomes (Richardson

et al., 2012).

Third, our research investigated associations using data from 2020, but the situation has

continued to evolve with the emergence of vaccines and COVID-19 variants (e.g., Delta and

Omicron). Future studies should consider the new situations accordingly.

5. Conclusion

Our findings suggest that during 2020, exposure to more green spaces, especially forests, was

significantly associated with a lower level of COVID-19 mortality rates, while exposure to more

developed open space has insignificant or positive associations with the COVID-19 mortality rates.

Forest outside park is more beneficial than *forest inside park,* with the optimal buffer distance

being 1km for *forest outside park*, and within 400m for *forest inside park*. These findings imply

that policymakers and planners should prioritize urban greening within optimal distances of

residential clusters and keep beneficial greenspaces accessible for COVID-19 and future

pandemics.

Acknowledgements
We thank to Long Chen, Xueming Liu, and Xueying Wu for data collection; Bin Chen for
methodological help; and Linda Larsen for proof reading the article.

Funding
This work was supported by the University Grants Committee [grant numbers
102010054.088616.01100.302.01] and the Research Grants Council of the Hong Kong SAR
[grant number CityU11207520].

References

1. Ahmad, F. B., Cisewski, J. A., Miniño, A., & Anderson, R. N. (2021). Provisional mortality data—united states, 2020. *Morbidity and Mortality Weekly Report,* 70 (14), 519.

2. Akpinar, Abdullah, Celestina Barbosa-Leiker, and Kerry R Brooks. 2016. "Does green space matter? Exploring relationships between green space type and health indicators." *Urban Forestry & Urban Greening* 20: 407-418.

3. Ali, N., & Islam, F. (2020). The Effects of Air Pollution on COVID-19 Infection and Mortality-A Review on Recent Evidence. *Frontiers in public health,* 8, 580057-580057.

4. Allard-Poesi, F., Matos, L. B. S., & Massu, J. (2022). Not all types of nature have an equal effect on urban residents' well-being: A structural equation model approach. *Health & place*, 74, 102759.

5. Amatriain-Fernández, S., Gronwald, T., Murillo-Rodríguez, E., Imperatori, C., Solano, A. F., Latini, A., & Budde, H. (2020). Physical exercise potentials against viral diseases like COVID-19 in the elderly. *Frontiers in medicine, 7*, 379.

6. Andersen, L., Corazon, S. S. S., & Stigsdotter, U. K. K. (2021). Nature exposure and its effects on immune system functioning: a systematic review. *International Journal of Environmental Research and Public Health, 18*(4), 1416.

7. Andrasfay, T., & Goldman, N. (2021). Reductions in 2020 US life expectancy due to COVID-19 and the disproportionate impact on the Black and Latino populations. *Proceedings of the National Academy of Sciences, 118*(5), e2014746118.

8. Bastien, M., Poirier, P., Lemieux, I., & Després, J.-P. (2014). Overview of epidemiology and contribution of obesity to cardiovascular disease. *Progress in cardiovascular diseases, 56*(4), 369-381.

9. Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.

10. Bauer, R. N., Diaz-Sanchez, D., & Jaspers, I. (2012). Effects of air pollutants on innate immunity: the role of Toll-like receptors and nucleotide-binding oligomerization domain-like receptors. *The Journal of allergy and clinical immunology, 129*(1), 14-26.

11. Ben, Y., Ma, F., Wang, H., Hassan, M. A., Yevheniia, R., Fan, W., . . . Dong, Z. (2019). A spatio-temporally weighted hybrid model to improve estimates of personal PM2. 5 exposure: Incorporating big data from multiple data sources. *Environmental pollution, 253*, 403-411.

12. Biddle, S. J., Ciaccioni, S., Thomas, G., & Vergeer, I. (2019). Physical activity and mental health in children and adolescents: An updated review of reviews and an analysis of causality. *Psychology of Sport and Exercise, 42*, 146-155.

13. Bivand, R. S., & Wong, D. W. (2018). Comparing implementations of global and local indicators of spatial association. *Test, 27*(3), 716-748.

14. Brandén, M., Aradhya, S., Kolk, M., Härkönen, J., Drefahl, S., Malmberg, B., . . . Mussino, E. (2020). Residential context and COVID-19 mortality among adults aged 70 years and older in Stockholm: a population-based, observational study using individual-level data. *The Lancet Healthy Longevity, 1*(2), e80-e88.

15. Bulfone, Tommaso Celeste, Mohsen Malekinejad, George W Rutherford, and Nooshin Razani. 2021. "Outdoor transmission of SARS-CoV-2 and other respiratory viruses: a systematic review." *The Journal of infectious diseases* 223 (4): 550-561.

16. Bureau, U. S. C. (2019). *Interactive Atlas of Heart Disease and Stroke.* Retrieved from https://nccd.cdc.gov/DHDSPAtlas/Default.aspx

17. Calle, E. E., & Thun, M. J. (2004). Obesity and cancer. *Oncogene, 23*(38), 6365-6378.

18. Castelli, V., Cimini, A., & Ferri, C. (2020). Cytokine storm in COVID-19:"when you come
out of the storm, you won't be the same person who walked in". *Frontiers in
Immunology, 11*.

19. CDC. (2021). Interactive Atlas of Heart Disease and Stroke. Retrieved from
http://nccd.cdc.gov/DHDSPAtlas.   http://nccd.cdc.gov/DHDSPAtlas.

20. Chan, J. M., Rimm, E. B., Colditz, G. A., Stampfer, M. J., & Willett, W. C. (1994). Obesity,
fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes
care, 17*(9), 961-969.

21. Chen, B., Song, Y., Jiang, T., Chen, Z., Huang, B., & Xu, B. (2018). Real-time estimation of
population exposure to PM2. 5 using mobile-and station-based big data. *International
Journal of Environmental Research and Public Health, 15*(4), 573.

22. Chen, B., Song, Y., Kwan, M.-P., Huang, B., & Xu, B. (2018). How do people in different
places experience different levels of air pollution? Using worldwide Chinese as a lens.
Environmental pollution, 238, 874-883.

23. Chen, Simiao, Klaus Prettner, Michael Kuhn, and David E Bloom. 2021. "The economic
burden of COVID-19 in the United States: Estimates and projections under an infection-
based herd immunity approach." *The Journal of the Economics of Ageing* 20: 100328.

24. Chernozhukov, V., Kasahara, H., & Schrimpf, P. (2021). Causal impact of masks, policies,
behavior on early covid-19 pandemic in the U.S. *Journal of Econometrics, 220*(1), 23-62.

25. Cho, K. S., Lim, Y.-r., Lee, K., Lee, J., Lee, J. H., & Lee, I.-S. (2017). Terpenes from forests
and human health. *Toxicological research, 33*(2), 97-106.

26. Ciencewicki, J., & Jaspers, I. (2007). Air pollution and respiratory viral infection. *Inhalation
toxicology, 19*(14), 1135-1146.

27. Claesson, M. J., Jeffery, I. B., Conde, S., Power, S. E., O'Connor, E. M., Cusack, S., . . .
O'Toole, P. W. (2012). Gut microbiota composition correlates with diet and health in the
elderly. *Nature, 488*(7410), 178-184.

28. Coombes, E., Jones, A. P., & Hillsdon, M. (2010). The relationship of physical activity and
overweight to objectively measured green space accessibility and use. *Social Science &
Medicine, 70*(6), 816-822.

29. Coombes, Emma, Andrew P. Jones, and Melvyn Hillsdon. 2010. "The relationship of
physical activity and overweight to objectively measured green space accessibility and
use." *Social Science & Medicine* 70 (6): 816-822.

30. Cutler, D. M., & Summers, L. H. (2020). The COVID-19 Pandemic and the $16 Trillion
Virus. *JAMA, 324*(15), 1495-1496. doi:10.1001/jama.2020.19759

31. De la Fuente, F., Saldías, M. A., Cubillos, C., Mery, G., Carvajal, D., Bowen, M., &
Bertoglia, M. P. (2021). Green space exposure association with type 2 diabetes mellitus,
physical activity, and obesity: a systematic review. *International Journal of
Environmental Research and Public Health, 18*(1), 97.

32. Del Valle, D. M., Kim-Schulze, S., Huang, H.-H., Beckmann, N. D., Nirenberg, S., Wang,
B., . . . Gnjatic, S. (2020). An inflammatory cytokine signature predicts COVID-19
severity and survival. *Nature Medicine, 26*(10), 1636-1643.

33. DePhillipo, N. N., Chahla, J., Busler, M., & LaPrade, R. F. (2021). Mobile phone gps data
and prevalence of covid-19 infections: Quantifying parameters of social distancing in the
us. *Archives of Bone and Joint Surgery*, *9*(2), 217.

34. DeSantis, A., DiezRoux, A., Hajat, A., Aiello, A., Golden, S., Jenny, N., . . . Shea, S. (2012). Associations of salivary cortisol levels with inflammatory markers: the Multi-Ethnic Study of Atherosclerosis. *Psychoneuroendocrinology, 37*(7), 1009-1018.

35. Dhar, D., & Mohanty, A. (2020). Gut microbiota and Covid-19- possible link and implications. *Virus Research, 285*, 198018.

36. Domènech-Montoliu, Salvador, Maria Rosario Pac-Sa, Paula Vidal-Utrillas, Marta Latorre-Poveda, Alba Del Rio-González, Sara Ferrando-Rubert, Gema Ferrer-Abad, Manuel Sánchez-Urbano, Laura Aparisi-Esteve, and Gema Badenes-Marques. 2021. "Mass gathering events and COVID-19 transmission in Borriana (Spain): A retrospective cohort study." *PloS one* 16 (8): e0256747.

37. Domingo, J. L., & Rovira, J. (2020). Effects of air pollutants on the transmission and severity of respiratory viral infections. *Environmental research, 187*, 109650.

38. Donati Zeppa, S., Agostini, D., Piccoli, G., Stocchi, V., & Sestili, P. (2020). Gut Microbiota Status in COVID-19: An Unrecognized Player? *Frontiers in Cellular and Infection Microbiology, 10*(742).

39. Donovan, G. H., Butry, D. T., Michael, Y. L., Prestemon, J. P., Liebhold, A. M., Gatziolis, D., & Mao, M. Y. (2013). The Relationship Between Trees and Human Health: Evidence from the Spread of the Emerald Ash Borer. *American journal of preventive medicine, 44*(2), 139-145.

40. Dyer, O. (2020). Covid-19: US testing ramps up as early response draws harsh criticism. In: British Medical Journal Publishing Group.

41. Ekkel, E. Dinand, and Sjerp de Vries. 2017. "Nearby green space and human health: Evaluating accessibility metrics." *Landscape and Urban Planning* 157: 214-220.

42. Esri. (2021). *USA Parks*. Retrieved from: https://www.arcgis.com/home/item.html?id=578968f975774d3fab79fe56c8c90941

43. Fernandez, D. M., Clemente, J. C., & Giannarelli, C. (2018). Physical Activity, Immune System, and the Microbiome in Cardiovascular Disease. *Frontiers in physiology, 9*(763).

44. Geng, D., Innes, J., Wu, W., & Wang, G. (2021). Impacts of COVID-19 pandemic on urban park visitation: a global analysis. *Journal of Forestry Research, 32*(2), 553-567.

45. Ghimire, R., Ferreira, S., Green, G. T., Poudyal, N. C., Cordell, H. K., & Thapa, J. R. (2017). Green space and adult obesity in the United States. *Ecological Economics, 136*, 201-212.

46. Gidlow, C. J., Randall, J., Gillman, J., Smith, G. R., & Jones, M. V. (2016). Natural environments and chronic stress measured by hair cortisol. *Landscape and Urban Planning, 148*, 61-67.

47. Girija, A., Shankar, E. M., & Larsson, M. (2020). Could SARS-CoV-2-induced hyperinflammation magnify the severity of coronavirus disease (CoViD-19) leading to acute respiratory distress syndrome? *Frontiers in Immunology, 11*, 1206.

48. Glencross, D. A., Ho, T.-R., Camiña, N., Hawrylowicz, C. M., & Pfeffer, P. E. (2020). Air pollution and its effects on the immune system. *Free Radical Biology and Medicine, 151*, 56-68.

49. Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote sensing of Environment, 202*, 18-27.

50. Gouin, J.-P., Glaser, R., Malarkey, W. B., Beversdorf, D., & Kiecolt-Glaser, J. (2012). Chronic stress, daily stressors, and circulating inflammatory markers. *Health Psychology, 31*(2), 264.

51. Grahn, Patrik, and Ulrika A. Stigsdotter. 2003. "Landscape planning and stress." *Urban*
Forestry & Urban Greening 2 (1): 1-18.
52. Grönroos, M., Parajuli, A., Laitinen, O. H., Roslund, M. I., Vari, H. K., Hyöty, H., . . .
Sinkkonen, A. (2019). Short-term direct contact with soil and plant materials leads to an
immediate increase in diversity of skin microbiota. *Microbiologyopen, 8*(3), e00645.
doi:10.1002/mbo3.645
53. Halpern, N., & Tan, K. (2020). United states resource availability for COVID-19. Society of
Critical Care Medicine. *Updated May, 12*.
54. Hu, M., Roberts, J. D., Azevedo, G. P., & Milner, D. (2021). The role of built and social
environmental factors in Covid-19 transmission: A look at America's capital city.
Sustainable Cities and Society, 65, 102580.
55. Huang, Q., Wu, X., Zheng, X., Luo, S., Xu, S., & Weng, J. (2020). Targeting inflammation
and cytokine storm in COVID-19. *Pharmacological Research, 159*, 105051.
56. Hussain, A., Mahawar, K., Xia, Z., Yang, W., & El-Hasani, S. (2020). Obesity and mortality
of COVID-19. Meta-analysis. *Obesity research & clinical practice, 14*(4), 295-300.
57. Jaafari, S., Shabani, A. A., Moeinaddini, M., Danehkar, A., & Sakieh, Y. (2020). Applying
landscape metrics and structural equation modeling to predict the effect of urban green
space on air pollution and respiratory mortality in Tehran. *Environmental Monitoring and*
Assessment, 192(7), 412-412.
58. Jia, P., Cao, X., Yang, H., Dai, S., He, P., Huang, G., . . . Wang, Y. (2020). Green space
access in the neighbourhood and childhood obesity. *Obesity Reviews*.
59. Jiang, B. (2014). Establishing dose-response curves for the impact of urban forests on
recovery from acute stress and landscape preference. University of Illinois at Urbana-
Champaign.
60. Jiang, B., Chang, C.-Y., & Sullivan, W. C. (2014). A dose of nature: Tree cover, stress
reduction, and gender differences. *Landscape and Urban Planning, 132*, 26-36.
61. Jiang, B., Shen, K., Sullivan, W. C., Yang, Y., Liu, X., & Lu, Y. (2021). A natural
experiment reveals impacts of built environment on suicide rate: Developing an
environmental theory of suicide. *Science of the total environment, 776*, 145750.
62. Jiang, B., Yang, Y., Chen, L., Liu, X., Wu, X., Chen, B., . . . Lu, Y. (2021). Green spaces,
especially forest, linked to lower SARS-CoV-2 infection rates: A one-year nationwide
study. *medRxiv*.
63. Johnson, Thomas F, Lisbeth A Hordley, Matthew P Greenwell, and Luke C Evans. 2020.
"Effect of park use and landscape structure on COVID-19 transmission rates." *MedRxiv*:
2020.10. 20.20215731.
64. Kim, Gunwoo, and Patrick A Miller. 2019. "The impact of green infrastructure on human
health and well-being: The example of the Huckleberry Trail and the Heritage
Community Park and Natural Area in Blacksburg, Virginia." *Sustainable Cities and*
Society 48: 101562.
65. Kim, T., Song, B., Cho, K. S., & Lee, I.-S. (2020). Therapeutic potential of volatile terpenes
and terpenoids from forests for inflammatory diseases. *International journal of molecular*
sciences, 21(6), 2187.
66. Klang, E., Kassim, G., Soffer, S., Freeman, R., Levin, M. A., & Reich, D. L. (2020). Severe
obesity as an independent risk factor for COVID-19 mortality in hospitalized patients
younger than 50. *Obesity, 28*(9), 1595-1599.

67. Klompmaker, J. O., Hart, J. E., Holland, I., Sabath, M. B., Wu, X., Laden, F., . . . James, P.
(2021). County-level exposures to greenness and associations with COVID-19 incidence
and mortality in the United States. *Environmental Research, 199*, 111331.
68. Kolak, M., Lin, Q., Halpern, D., Paykin, S., Martinez-Cardoso, A., & Li, X. (2021). The US
Covid Atlas. Retrieved from https://www.uscovidatlas.org
69. Kong, X., Zheng, K., Tang, M., Kong, F., Zhou, J., Diao, L., . . . Dong, Y. (2020). Prevalence
and factors associated with depression and anxiety of hospitalized patients with COVID-
19. *medRxiv*.
70. Konstantinoudis, G., Padellini, T., Bennett, J., Davies, B., Ezzati, M., & Blangiardo, M.
(2021). Long-term exposure to air-pollution and COVID-19 mortality in England: A
hierarchical spatial analysis. *Environment international, 146*, 106316.
71. Krauss, R. M., Winston, M., Fletcher, B. J., & Grundy, S. M. (1998). Obesity: impact on
cardiovascular disease. *Circulation, 98*(14), 1472-1476.
72. Kuo, M. (2015). How might contact with nature promote human health? Promising
mechanisms and a possible central pathway. *Frontiers in psychology, 6*, 1093.
73. Lee, J., Park, B. J., Tsunetsugu, Y., Ohira, T., Kagawa, T., & Miyazaki, Y. (2011). Effect of
forest bathing on physiological and psychological responses in young Japanese male
subjects. *Public health, 125*(2), 93-100.
74. Lee, W., Kim, H., Choi, H. M., Heo, S., Fong, K. C., Yang, J., . . . Bell, M. L. (2021). Urban
environments and COVID-19 in three Eastern states of the United States. *Science of the
total environment, 779*, 146334.
75. Li, Q. (2010). Effect of forest bathing trips on human immune function. *Environmental
health and preventive medicine, 15*(1), 9-17.
76. Li, Q., Morimoto, K., Kobayashi, M., Inagaki, H., Katsumata, M., Hirata, Y., . . . Miyazaki,
Y. (2008). A forest bathing trip increases human natural killer activity and expression of
anti-cancer proteins in female subjects. *J Biol Regul Homeost Agents, 22*(1), 45-55.
77. Li, Q., Morimoto, K., Nakadai, A., Inagaki, H., Katsumata, M., Shimizu, T., . . . Miyazaki,
Y. (2007). Forest bathing enhances human natural killer activity and expression of anti-
cancer proteins. *International journal of immunopathology and pharmacology, 20*, 3-8.
78. Liang, D., Shi, L., Zhao, J., Liu, P., Sarnat, J. A., Gao, S., . . . Chang, H. H. (2020). Urban
Air Pollution May Enhance COVID-19 Case-Fatality and Mortality Rates in the United
States. *The Innovation, 1*(3), 100047.
79. Lu, Y., Chen, L., Liu, X., Yang, Y., Sullivan, W. C., Xu, W., . . . Jiang, B. (2021). Green
spaces mitigate racial disparity of health: A higher ratio of green spaces indicates a lower
racial disparity in SARS-CoV-2 infection rates in the USA. *Environment international,
152*, 106465.
80. Lu, Y., Sarkar, C., & Xiao, Y. (2018). The effect of street-level greenery on walking
behavior: Evidence from Hong Kong. *Social Science & Medicine, 208*, 41-49.
81. Lu, Y., Zhao, J., Wu, X., & Lo, S. M. (2021). Escaping to nature during a pandemic: A
natural experiment in Asian cities during the COVID-19 pandemic with big social media
data. *Science of the total environment, 777*, 146092.
82. Ma, Jianfang, Haihong Zhu, Peng Li, Chengcheng Liu, Feng Li, Zhenwei Luo, Meihui
Zhang, and Lin Li. 2022. "Spatial Patterns of the Spread of COVID-19 in Singapore and
the Influencing Factors." *ISPRS International Journal of Geo-Information* 11 (3): 152.

83. Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu, S., . . . Luo, B. (2020). Effects of
temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Science*
of the total environment, 724, 138226.
84. Mancus, G., Cimino, A. N., Hasan, M. Z., Campbell, J. C., Winch, P. J., Sharps, P., . . .
Stockman, J. K. (2020). Residential Greenness Positively Associated with the Cortisol to
DHEA Ratio among Urban-Dwelling African American Women at Risk for HIV. *Journal*
of Urban Health, 1-9.
85. Mandel, M., Harari, G., Gurevich, M., & Achiron, A. (2020). Cytokine prediction of
mortality in COVID19 patients. *Cytokine, 134*, 155190.
86. Market, M., Angka, L., Martel, A. B., Bastin, D., Olanubi, O., Tennakoon, G., . . . Auer, R.
C. (2020). Flattening the COVID-19 Curve With Natural Killer Cell Based
Immunotherapies. *Frontiers in Immunology, 11*(1512).
87. Martelletti, L., & Martelletti, P. (2020). Air Pollution and the Novel Covid-19 Disease: a
Putative Disease Risk Factor. *SN comprehensive clinical medicine*, 1-5.
88. Medicine, J. H. U. (2021). COVID-19 United States Deaths by County. Retrieved from
https://coronavirus.jhu.edu/us-map
89. Miller, I. F., Becker, A. D., Grenfell, B. T., & Metcalf, C. J. E. (2020). Disease and
healthcare burden of COVID-19 in the United States. *Nature Medicine, 26*(8), 1212-
1217.
90. Morey, J. N., Boggero, I. A., Scott, A. B., & Segerstrom, S. C. (2015). Current Directions in
Stress and Human Immune Function. *Current opinion in psychology, 5*, 13-17.
91. Nielsen, Thomas Sick, and Karsten Bruun Hansen. 2007. "Do green areas affect health?
Results from a Danish survey on the use of green areas and health indicators." *Health &*
Place 13 (4): 839-850.
92. Nieman, D. C., & Wentz, L. M. (2019). The compelling link between physical activity and
the body's defense system. *Journal of sport and health science, 8*(3), 201-217.
93. NLCD. (2016). National Land Cover Database Class Legend and Description Retrieved from
https://www.mrlc.gov/data/legends/national-land-cover-database-class-legend-and-
description
94. Nowak, D. J., Hirabayashi, S., Bodine, A., & Greenfield, E. (2014). Tree and forest effects
on air quality and human health in the United States. *Environmental pollution, 193*, 119-
129.
95. Nowak, D. J., Hirabayashi, S., Doyle, M., McGovern, M., & Pasher, J. (2018). Air pollution
removal by urban forests in Canada and its effect on air quality and human health. *Urban*
Forestry & Urban Greening, 29, 40-48.
96. Osman, M., Faridi, R. M., Sligl, W., Shabani-Rad, M.-T., Dharmani-Khan, P., Parker, A., . . .
Cohen Tervaert, J. W. (2020). Impaired natural killer cell counts and cytolytic activity in
patients with severe COVID-19. *Blood advances, 4*(20), 5035-5039.
97. Pan, Jiayu, Ronita Bardhan, and Ying Jin. 2021. "Spatial distributive effects of public green
space and COVID-19 infection in London." *Urban Forestry & Urban Greening* 62:
127182.
98. Parajuli, A. (2019). The effect of living environment and environmental exposure on the
composition of microbial community in soil, on human skin and in the gut.
99. Parajuli, A., Grönroos, M., Siter, N., Puhakka, R., Vari, H. K., Roslund, M. I., . . . Hyöty, H.
(2018). Urbanization reduces transfer of diverse environmental microbiota indoors.
Frontiers in microbiology, 9, 84.

100. Parajuli, A., Hui, N., Puhakka, R., Oikarinen, S., Grönroos, M., Selonen, V. A., . . . Vari, H. K. (2020). Yard vegetation is associated with gut microbiota composition. *Science of the total environment, 713*, 136707.

101. Paranjpe, I., Russak, A., De Freitas, J. K., Lala, A., Miotto, R., Vaid, A., . . . Meyer, D. (2020). Clinical characteristics of hospitalized Covid-19 patients in New York City. *medRxiv*.

102. Peng, Zhe, AL Pineda Rojas, E Kropff, W Bahnfleth, Giorgio Buonanno, SJ Dancer, Jarek Kurnitski, Yuguo Li, Marcel GLC Loomans, and Linsey C Marr. 2022. "Practical indicators for risk of airborne transmission in shared indoor environments and their application to covid-19 outbreaks." *Environmental science & technology*.

103. Perone, G. (2021). The determinants of COVID-19 case fatality rate (CFR) in the Italian regions and provinces: An analysis of environmental, demographic, and healthcare factors. *Science of the total environment, 755*, 142523.

104. Potempa, L. A., Rajab, I. M., Hart, P. C., Bordon, J., & Fernandez-Botran, R. (2020). Insights into the use of C-reactive protein as a diagnostic index of disease severity in COVID-19 infections. *The American journal of tropical medicine and hygiene, 103*(2), 561.

105. Pretty, J., Peacock, J., Sellens, M., & Griffin, M. (2005). The mental and physical health outcomes of green exercise. *International Journal of Environmental Health Research, 15*(5), 319-337.

106. Prevention., C. f. D. C. a. (2022). Interactive Atlas of Heart Disease and Stroke. Retrieved from http://nccd.cdc.gov/DHDSPAtlas.

107. Qin, C., Zhou, L., Hu, Z., Zhang, S., Yang, S., Tao, Y., . . . Tian, D. S. (2020). Dysregulation of Immune Response in Patients With Coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis, 71*(15), 762-768.

108. Qiu, Y., Zuo, S., Yu, Z., Zhan, Y., & Ren, Y. (2021). Discovering the effects of integrated green space air regulation on human health: A bibliometric and meta-analysis. *Ecological Indicators, 132*, 108292.

109. Reid, C. E., Clougherty, J. E., Shmool, J. L., & Kubzansky, L. D. (2017). Is all urban green space the same? A comparison of the health benefits of trees and grass in New York City. *International Journal of Environmental Research and Public Health, 14*(11), 1411.

110. Ribeiro, A. I., Tavares, C., Guttentag, A., & Barros, H. (2019). Association between neighbourhood green space and biological markers in school-aged children. Findings from the Generation XXI birth cohort. *Environment international, 132*, 105070.

111. Richardson, E. A., Mitchell, R., Hartig, T., De Vries, S., Astell-Burt, T., & Frumkin, H. (2012). Green cities and health: a question of scale? *J Epidemiol Community Health, 66*(2), 160-165.

112. Roslund, M. I., Puhakka, R., Grönroos, M., Nurminen, N., Oikarinen, S., Gazali, A. M., . . . Vari, H. K. (2020). Biodiversity intervention enhances immune regulation and health-associated commensal microbiota among daycare children. *Science Advances, 6*(42), eaba2578.

113. Roviello, V., & Roviello, G. N. (2021). Less COVID-19 deaths in southern and insular Italy explained by forest bathing, Mediterranean environment, and antiviral plant volatile organic compounds. *Environmental Chemistry Letters*, 1-11.

114. Roviello, V., Gilhen-Baker, M., Vicidomini, C., & Roviello, G. N. (2021). Forest-bathing
and physical activity as weapons against COVID-19: a review. *Environmental Chemistry*
Letters, 1-10.

115. Russette, Helen, Jon Graham, Zachary Holden, Erin O Semmens, Elizabeth Williams, and
Erin L Landguth. 2021. "Greenspace exposure and COVID-19 mortality in the United
States: January–July 2020." *Environmental research* 198: 111195.

116. Shen, Y.-S., & Lung, S.-C. C. (2017). Mediation pathways and effects of green structures
on respiratory mortality via reducing air pollution. *Scientific reports, 7*(1), 1-9.

117. Smith, M Kyle S, Izak PJ Smit, Louise K Swemmer, Mohlamatsane M Mokhatla, Stefanie
Freitag, Dirk J Roux, and Luthando Dziba. 2021. "Sustainability of protected areas:
Vulnerabilities and opportunities as revealed by COVID-19 in a national park
management agency." *Biological Conservation* 255: 108985.

118. Soga, M., Evans, M. J., Tsuchiya, K., & Fukano, Y. (2021). A room with a green view: the
importance of nearby nature for mental health during the COVID-19 pandemic.
Ecological Applications, 31(2), e2248.

119. Sorichetta, A., Hornby, G. M., Stevens, F. R., Gaughan, A. E., Linard, C., & Tatem, A. J.
(2015). High-resolution gridded population datasets for Latin America and the Caribbean
in 2010, 2015, and 2020. *Scientific data, 2*(1), 1-12.

120. Spotswood, E. N., Benjamin, M., Stoneburner, L., Wheeler, M. M., Beller, E. E., Balk,
D., . . . McDonald, R. I. (2021). Nature inequity and higher COVID-19 case rates in less-
green neighbourhoods in the United States. *Nature Sustainability, 4*(12), 1092-1098.

121. Steptoe, A., Hamer, M., & Chida, Y. (2007). The effects of acute psychological stress on
circulating inflammatory factors in humans: a review and meta-analysis. *Brain, behavior,*
and immunity, 21(7), 901-912.

122. Team, R. C. (2015). R: A language and environment for statistical computing. Vienna,
Austria; 2014. In.

123. Tsao, T.-M., Tsai, M.-J., Hwang, J.-S., Cheng, W.-F., Wu, C.-F., Chou, C.-C. K., & Su, T.-
C. (2018). Health effects of a forest environment on natural killer cells in humans: an
observational pilot study. *Oncotarget, 9*(23), 16501-16511.

124. Ugolini, F., Massetti, L., Calaza-Martínez, P., Cariñanos, P., Dobbs, C., Ostoic, S. K., . . .
Šaulienė, I. (2020). Effects of the COVID-19 pandemic on the use and perceptions of
urban green space: an international exploratory study. *Urban Forestry & Urban*
Greening, 126888.

125. Ulrich, R. S., Simons, R. F., Losito, B. D., Fiorito, E., Miles, M. A., & Zelson, M. (1991).
Stress recovery during exposure to natural and urban environments. *Journal of*
environmental psychology, 11(3), 201-230.

126. Van Ingen, T., Brown, K. A., Buchan, S. A., Akingbola, S., Daneman, N., & Smith, B. T.
(2021). Neighbourhood-level risk factors of COVID-19 incidence and mortality.
medRxiv.

127. Venter, Z. S., Barton, D. N., Gundersen, V., Figari, H., & Nowell, M. S. (2021). Back to
nature: Norwegians sustain increased recreational use of urban green space months after
the COVID-19 outbreak. *Landscape and Urban Planning, 214*, 104175.

128. Venter, Z., Barton, D., Gundersen, V., Figari, H., & Nowell, M. (2020). Urban nature in a
time of crisis: recreational use of green space increases during the COVID-19 outbreak in
Oslo, Norway. *Environmental Research Letters*.

129. Viglione, G. (2020). How many people has the coronavirus killed? Retrieved from
867   https://www.nature.com/articles/d41586-020-02497-w
868 130. Villeneuve, P. J., Jerrett, M., G. Su, J., Burnett, R. T., Chen, H., Wheeler, A. J., &
869   Goldberg, M. S. (2012). A cohort study relating urban green space with mortality in
870   Ontario, Canada. *Environmental Research, 115*, 51-58.
871 131. Volenec, Z. M., Abraham, J. O., Becker, A. D., & Dobson, A. P. (2021). Public parks and
872   the pandemic: How park usage has been affected by COVID-19 policies. *PloS one, 16*(5),
873   e0251799.
874 132. VoPham, T., Weaver, M. D., Hart, J. E., Ton, M., White, E., & Newcomb, P. A. (2020).
875   Effect of social distancing on COVID-19 incidence and mortality in the US. *medRxiv*.
876 133. Wang, Jingjing, Xueying Wu, Ruoyu Wang, Dongsheng He, Dongying Li, Linchuan Yang,
877   Yiyang Yang, and Yi Lu. 2021. "Review of associations between built environment
878   characteristics and severe acute respiratory syndrome coronavirus 2 infection risk."
879   *International Journal of Environmental Research and Public Health* 18 (14): 7561.
880 134. Wang, M., Hu, C., Zhao, Q., Feng, R., Wang, Q., Cai, H., . . . Guo, C. (2021). Acute
881   psychological impact on COVID-19 patients in Hubei: a multicenter observational study.
882   *Translational psychiatry, 11*(1), 1-9.
883 135. Yang, B.-Y., Hu, L.-W., Jalaludin, B., Knibbs, L. D., Markevych, I., Heinrich, J., . . . Dong,
884   G.-H. (2020). Association Between Residential Greenness, Cardiometabolic Disorders,
885   and Cardiovascular Disease Among Adults in China. *JAMA Network Open, 3*(9),
886   e2017507-e2017507.
887 136. Yang, L., Jin, S., Danielson, P., Homer, C., Gass, L., Bender, S. M., . . . Fry, J. (2018). A
888   new generation of the United States National Land Cover Database: Requirements,
889   research priorities, design, and implementation strategies. *ISPRS Journal of
890   Photogrammetry and Remote Sensing, 146*, 108-123.
891 137. Yang, Y., & Diez-Roux, V. A. (2012). Walking Distance by Trip Purpose and Population
892   Subgroups. *American journal of preventive medicine, 43*(1), 11-19.
893 138. Yeoh, Y. K., Zuo, T., Lui, G. C.-Y., Zhang, F., Liu, Q., Li, A. Y., . . . Fung, K. S. (2021).
894   Gut microbiota composition reflects disease severity and dysfunctional immune
895   responses in patients with COVID-19. *Gut, 70*(4), 698-706.
896 139. Yokoyama, W. M. (2005). Natural killer cell immune responses. *Immunol Res, 32*(1-3),
897   317-325.
898 140. Zuo, T., Zhang, F., Lui, G. C. Y., Yeoh, Y. K., Li, A. Y. L., Zhan, H., . . . Ng, S. C. (2020).
899   Alterations in Gut Microbiota of Patients With COVID-19 During Time of
900   Hospitalization. *Gastroenterology, 159*(3), 944-955.e948.