Altered brain activity and functional connectivity after MDMA-assisted therapy for post-

traumatic stress disorder

S. Parker Singleton[1], Julie B. Wang, M.P.H, Ph.D.[2], Michael Mithoefer, M.D. [2,3], Colleen

Hanlon, Ph.D.[4], Mark S. George, M.D.[3,5], Annie Mithoefer[2], Oliver Mithoefer, M.D.[3], Allison R.

Coker, Ph.D. [2,6], Berra Yazar-Klosinski, Ph.D.[7], Amy Emerson [2], Rick Doblin, Ph.D.[7], Amy

Kuceyeski, Ph.D.[1,8]

[1] Department of Computational Biology, Cornell University, Ithaca, NY 14850, USA

[2] MAPS Public Benefit Corporations, San Jose, CA 95117, USA.

[3] Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina,

Charleston, SC 29425-0742, USA.

[4] Wake Forest School of Medicine, Winston-Salem, NC 27101 USA

[5] Ralph H. Johnson VA Medical Center, Charleston, SC 29425 USA

[6] Department of Neurology, University of California, San Francisco, San Francisco, CA, USA

[7] Multidisciplinary Association for Psychedelic Studies, San Jose, CA 95117, USA.

[8] Department of Radiology, Weill Cornell Medicine, New York, New York 10021, USA

Corresponding Author: S. Parker Singleton; Email: sps253@cornell.edu

Role of Funding Sources: The clinical trial was sponsored by the Multidisciplinary Association

for Psychedelic Studies (MAPS), a 501(c)(3) nonprofit organization. MAPS provided the

MDMA and fully funded this study from private and foundation donations. MAPS Public

Benefit Corporation (MAPS PBC), wholly owned by MAPS, was the trial organizer.

Contributors: SPS and AK had full access to all the data in the study and take responsibility for

the integrity of the data and for the accuracy of the data analysis. BYK, AE, MM, RD conceived

and designed the clinical trial part of the study. MM & AM carried out experimental treatments

and recorded participant scripts. AE carried out sponsor oversight of data collection. CH and

MSG designed the imaging tasks and acquisition. CH, MSG, and OM oversaw the scanning and

data collection. SPS, AK, JW, AC drafted the paper. SPS carried out preprocessing, analysis, and

interpretation of data. RD obtained funding for the study. All authors contributed to the critical

review and final version of the manuscript.

Conflict of Interest: MM was paid as a contractor by MAPS PBC. AE, CP, AC, and JW

received salary support for full-time employment with MAPS PBC. BYK and RD received

salary support for full-time employment with MAPS. MM is on the Clinical Advisory Board of

Awakn Life Sciences.

Acknowledgements: The authors would like to thank the candidates who were willing to be

screened for eligibility; the participants in the trial who contributed their data; S. Sadler for her

dedication as the Study Coordinator; S. Braswell for serving as night attendant, M. Wagner and

J. Wymer for Independent Rater assessments. The authors would also like to thank: J. Holland

who supported this trial as a Medical Monitor; L. Jerome for her numerous and varied

contributions to this Clinical Development Program since prior to its inception with global

systematic literature reviews and medical coding; C. Hennigan for data management; R.

Matthews and B. Shechet for clinical operations and monitoring; J. Sonstroem and A. Seltzer for

randomization support and system programming; E. Sola, Y. Gelfand, and B. Cohen for

conducting adherence ratings to facilitate standardization of therapy; I. Gorman for development

of adherence ratings; A. Wilens for supporting video recording. The authors would like to thank

Dr. Edmund Higgins for catalyzing the first discussions about acquiring imaging data in the

MAPS trial.

ABSTRACT

3,4-methylenedioxymethamphetamine-assisted therapy (MDMA-AT) for post-traumatic stress

disorder (PTSD) has demonstrated promise in multiple clinical trials. MDMA is hypothesized to

facilitate the therapeutic process, in part, by decreasing fear response during fear memory

processing while increasing extinction learning. The acute administration of MDMA in healthy

controls modifies recruitment of brain regions involved in the hyperactive fear response in PTSD

such as the amygdala, hippocampus, and insula. However, to date there have been no

neuroimaging studies aimed at directly elucidating the neural impact of MDMA-AT in PTSD

patients. We analyzed brain activity and connectivity via functional MRI during both rest and

autobiographical memory (trauma and neutral) response before and two-months after MDMA-

AT in nine veterans and first-responders with chronic PTSD of 6 months or more. We

hypothesized that MDMA-AT would increase amygdala-hippocampus resting-state functional

connectivity, however we only found evidence of a trend in the left amygdala – left hippocampus

(t = -2.91, uncorrected p = 0.0225, corrected p = 0.0901). We also found reduced activation

contrast (trauma > neutral) after MDMA-AT in the cuneus. Finally, the amount of recovery from

PTSD after MDMA-AT correlated with changes in four functional connections during

autobiographical memory recall: the left amygdala – left posterior cingulate cortex (PCC), left

amygdala – right PCC, left amygdala – left insula, and left isthmus cingulate – left posterior

hippocampus. Amygdala – insular functional connectivity is reliably implicated in PTSD and

anxiety, and both regions are impacted by MDMA administration. These findings compliment

previous research indicating that amygdala, hippocampus, and insula functional connectivity is a

potential target of MDMA-AT, and highlights other regions of interest related to memory

processes. More research is necessary to determine if these findings are specific to MDMA-AT

compared to other types of treatment for PTSD.

This study: NCT02102802   Parent-study: NCT01211405

Keywords: PTSD, MDMA, amygdala, hippocampus, precuneus, symptom provocation, insula,

fMRI, functional connectivity, resting-state, autobiographical memory.

1. INTRODUCTION

Post-traumatic stress disorder (PTSD), which can arise following exposure to a traumatic event

or repeated stressful events and is a debilitating social and economic burden on individuals and

their families (Davidson 2000; Gradus 2017). PTSD is associated with an increased fear response

(VanElzakker, Staples-Bradley, and Shin 2018) and distressing and intrusive re-experiencing of

traumatic memories (Ehlers 2010) that often serves as a barrier to the therapeutic process. This

may explain why individuals with the most severe PTSD symptoms after trauma are more likely

to end up with chronic PTSD durations (Shalev et al. 2019; van der Mei et al. 2020). Lifetime

occurrence of PTSD in the general population is around 8%, and, prevalence is significantly

higher in military personnel (17.1%) and first responders (10-32%), individuals on the front lines

of societal trauma (Hoge et al. 2004; Javidi and Yadollahie 2012). A meta-analysis of trials for

military-related PTSD found that cognitive behavioral therapy and prolonged exposure therapy

delivered clinically meaningful symptom improvements in 49-70% of patients, however 60-72%

of veterans receiving these therapies still retained their PTSD diagnosis (Steenkamp et al. 2015).

Adverse outcomes such as increased symptoms and disengagement from treatment cause many

current psychological therapies for PTSD to have high dropout rates (Goetter et al. 2015),

especially trauma focused therapies (Lewis et al. 2020; Mott et al. 2014).

One approach to developing more effective psychotherapies for PTSD is to administer a drug

alongside psychotherapy to aid the therapeutic process (Feduccia et al. 2018). 3,4-

methylenedioxymethamphetamine-assisted therapy (MDMA-AT) is hypothesized to reduce the

fear response associated with re-experiencing traumatic memories, and therefore may facilitate

tolerable processing of traumatic content in patients with PTSD (Mithoefer et al. 2011). Phase 2

and 3 trials have demonstrated promise for MDMA-AT as a viable treatment for PTSD (Mitchell

et al. 2021; Mithoefer et al. 2019; Wang et al. 2021; Jerome et al. 2020). MDMA, particularly

the *R*-enantiomer, increases pro-social behavior and enhances fear extinction in mice and this

effect appears to be mediated by serotonergic mechanisms (Curry et al. 2018; M. B. Young et al.

2017). In healthy humans, acute administration of MDMA has been shown to enhance positive

and reduce negative affect during the recollection of autobiographical memories, while

preserving vividness and emotional intensity (Carhart-Harris et al. 2014). In another study,

MDMA was found to preserve the memory accuracy when administered during both encoding

and retrieval phases, while attenuating the recollection of salient details for both positive and

negative memories, suggesting that MDMA alters emotional memory representations (Doss et al.

2018). Again in healthy controls, MDMA was found to enhance fear extinction

learning/retention rates compared to placebo when administered during extinction training

phases (Vizeli et al. 2022; Maples-Keller et al. 2022). These findings suggest that MDMA may

aid the therapeutic process, in part, by enabling patient access to emotionally challenging

material and facilitating memory reconsolidation/fear extinction processes (Feduccia and

Mithoefer 2018).

Functional magnetic resonance imaging (fMRI) measures changes in regional blood oxygenation

over time and is thus used as a proxy for fluctuating neuronal activity. In-scanner environments

can be absent of stimuli (resting-state fMRI – thought to measure intrinsic brain activity), or

tasks or stimuli may be presented to study the regional brain dynamics underlying specific

cognitive processes (Glover 2011). In addition to the study of isolated regional activation

changes, functional connectivity (FC) can be assessed to infer interaction between two or more

brain regions. FC is defined as the statistical relationship (Pearson correlation in the case of the

present study) between two brain regions' activity over time. These tools have been used to study

functioning of brain regions and their networks in a wide range of neuropathology and

psychiatric disorders (Zhou et al. 2017; Lucassen et al. 2014; Woodward and Cascio 2015).

PTSD patients have shown altered functioning of the precuneus, posterior cingulate cortex

(PCC), anterior cingulate cortex (ACC), insula, prefrontal and frontoparietal regions, as well as

the hippocampus and amygdala (Sartory et al. 2013; Harnett, Goodman, and Knight 2020;

Abdallah et al. 2017; Lazarov et al. 2017; Malivoire et al. 2018; Brown et al. 2014; Pitman et al.

2012; Gilboa et al. 2004; Fonzo et al. 2021; Rabinak et al. 2011; Belleau et al. 2020; Thome et

al. 2020; R. A. Lanius et al. 2006; Sripada et al. 2012), suggesting augmented recruitment of

brain regions involved in self-referential processing (Cavanna and Trimble 2006), salient

autobiographical memory (Svoboda, McKinnon, and Levine 2006; Spreng, Mar, and Kim 2009;

Sestieri et al. 2011; Maddock 1999; Spreng and Grady 2010), and fear and emotion (LeDoux

2003).

The specific effects of MDMA-AT on brain function in individuals with PTSD have not been

characterized, but several studies suggest the amygdala and hippocampus may play an important

role. The amygdala is broadly associated with fear response, and the hippocampus, associated

with learning and memory, may provide contextual information necessary for cognitive-affect

during memory recall (LeDoux 2003; Harnett, Goodman, and Knight 2020; Pitman et al. 2012).

Sripada and colleagues (2012) found combat veterans with PTSD have decreased amygdala-

hippocampal resting-state functional connectivity (RSFC) compared to combat veterans without

PTSD, which the authors speculate may represent an inability to contextualize affective

information in PTSD. Increased amygdala-hippocampal RSFC following stress/trauma exposure

has been shown to correlate with recovery from stress or trauma symptoms (Ben-Zion, Keynan,

et al. 2020; Vaisvaser et al. 2013; Fan et al. 2015), suggesting a possible adaptive mechanism to

threat exposure. In healthy volunteers, the acute administration of MDMA increases amygdala-

hippocampal RSFC (Carhart-Harris, 2015). Intranasal oxytocin also increases amygdala-

hippocampal RSFC (Fan et al. 2015), and this effect appears to be mediated by serotonin system

signaling (Lan et al. 2022) – two neuromodulators that MDMA significantly increases in the

extracellular/plasma concentration of, and that play a crucial role in its effects on pro-social

behavior and fear extinction (Dumont et al. 2009; Feduccia and Duvauchelle 2008; de la Torre et

al. 2004; Green et al. 2003; Thompson et al. 2007; Nardou et al. 2019; Young et al. 2017).

Despite evidence that RSFC between the amygdala and hippocampus is implicated in PTSD and

that this connection may be modulated by MDMA, no study to date has shown relationships

between changes in these regions' functional connectivity and the therapeutic effects of MDMA-

AT.

Herein, we describe results from a study of combat veterans and first-responders undergoing

MDMA-AT for PTSD in a randomized, double-blind, dose-response phase 2 clinical trial

(Mithoefer et al. 2018). The Clinician-Administered PTSD Scale (CAPS-IV) (an hour-long,

semi-structured interview centered around an index trauma) (Blake et al. 1990) was assessed

throughout the study to track PTSD severity. Enrolled individuals had moderate-to-severe PTSD

with a chronic PTSD duration of 6 months or more. Both resting-state and task-fMRI data,

acquired while individuals listened to trauma-related and neutral audio scripts, were collected

before and two months after MDMA-AT (follow-up scans were collected after the blind was

broken).

Prior to analysis, we hypothesized that MDMA-AT would increase RSFC between the amygdala

and hippocampus (Sripada et al. 2012; Carhart-Harris et al. 2015; Ben-Zion, Zeevi, et al. 2020;

Vaisvaser et al. 2013; Fan et al. 2015; Lan et al. 2022). We further hypothesized that, at baseline,

brain activity would be higher during the trauma-related listening task compared with the neutral

listening task in regions associated with autobiographical memory, fear, and emotion, and that

this effect would be reduced post-treatment (Sartory et al. 2013). In a final set of analyses, we

assessed pre-to-post treatment change in the FC of several regions of interest contained within

the limbic, salience, and default mode networks known to be hyperactive in PTSD (Etkin and

Wager 2007) during the trauma and neutral autobiographical memory task-fMRI scans. FC

changes were then correlated with the pre-to-post treatment recovery in overall PTSD

symptomatology – as measured by decreases in CAPS-IV total severity scores.

2. METHODS

Figure 1: Simplified study design. Subjects were assessed and imaged at the start of the study (baseline). All subjects (low dose (LD; 30 mg MDMA), medium dose (MD; 75 mg MDMA), and high dose (HD; 125 mg MDMA)) underwent three non-drug preparatory therapy sessions prior to their first MDMA dosing session. Each MDMA session was followed by three non-drug integration therapy sessions. After MDMA Session 2 and the subsequent integration sessions, subjects were assessed and the dosing blind was broken. HD subjects completed their final set of drug and non-drug therapy sessions unblinded, and LD/MD subjects crossed over into the HD arm where they completed three sets of drug and non-drug sessions, now with the higher dose and unblinded. All subjects were assessed and underwent MRI approximately two months following their last HD MDMA session. See the Methods section for a full description of study design and scanning protocols.

2.1 Trial design

The present study analyzed data from a sub-study (NCT02102802) of a Phase 2 randomized,

double-blind, dose-response trial of MDMA-AT in veterans and first responders with severe and

chronic PTSD (NCT01211405) (Mithoefer et al. 2018). A detailed study description of the

parent study can be found in (Mithoefer et al. 2018), and we summarize the study design in the

SI. Here, we provide a description of the MRI-based sub-study design.

Participants in the parent study were able to opt into the MRI-based sub-study after which they

provided written informed consent approved by the Medical University of South Carolina

Institutional Review Board. They were screened for additional neuroimaging related eligibility

criteria and were excluded for any conditions that could render MRI unsafe. A script-driven

autobiographical memory paradigm was used to assess brain activity during symptom

provocation (Ruth A Lanius et al. 2002; Ruth A. Lanius et al. 2004; Mertens et al. 2022).

Following baseline CAPS-IV assessment in the parent study, sub-study participants worked with

investigators to create two scripts: one describing a personally traumatic event and one reflecting

their typical morning routine at home. Two audio recordings, each six minutes in length, were

created from the participant's reading of each script. Each audio recording was divided into two

3-minute blocks for the task-fMRI. All participants, in all arms, were imaged at baseline, prior to

therapy, and again at the follow-up visit two months after their final dosing session. LD (N = 2)

and MD (N = 2) participants were additionally imaged after the primary endpoint visit in Stage 1

(one month following their second dosing session), however the small sample sizes prevented

any meaningful analysis with these scans. The present analysis focuses on the pre- and post-

therapy effects of MDMA-AT on fMRI biomarkers, and thus uses the scans collected at pre-

treatment (baseline) and at least 2 months after the largest dose of MDMA (follow-up).

2.2 MRI acquisition

At each scanning session, participants underwent MRI on a 32 channel 3T Siemens system. T1

anatomical scans with TR/TE=1900/2.34 ms and 0.9x0.9x1.0 mm voxel size were collected,

followed by two identical task fMRI (design described below) (TR/TE = 2200/35ms, 3.0 mm

isotropic voxel size, length of each scan = 14:25 min) and one resting state fMRI (TR/TE =

2000/30ms, 3.3x3.3x3.0 mm voxel size, length = 5:00 min).

Participants' 6-minute trauma and neutral audio scripts were divided into two three-minute

trauma and neutral blocks each (See 2.1 Trial design for description of audio recordings). During

fMRI, participants were presented with the visual cue "allow" and instructed to allow themselves

to experience the scripts as their audio recordings were played for both neutral and trauma

blocks. Each task scan had an alternating block design (neutral 1, trauma 1, neutral 2, trauma 2)

with an 18 second 'rest' period at the start of the scan and between each block, and about a

minute of rest at the end of the scan. The precise length of each audio block was 2.95 min.

2.3 Image preprocessing

FreeSurfer (Dale, Fischl, and Sereno 1999) was applied to the T1s to create white matter (WM),

gray matter (GM) and cerebrospinal fluid (CSF) segmentations. FMRIB Software Library (FSL)

(Smith et al. 2004) was used for 1) brain extraction, 2) registration between T1s and fMRI's

(brain-boundary registration, non-linear, full-search), 3) high-pass filtering 4) slice-time and 5)

motion correction.

2.4 Motion

High-motion frames (defined as > 0.9 mm relative framewise displacement (FD); CONN

Toolbox standard parameter (Whitfield-Gabrieli and Nieto-Castanon 2012)) were counted as

outlier volumes and scrubbed from functional connectivity and activation analyses. The

percentage of scrubbed volumes for resting-state scans ranged from 0 to 6.8% (mean = 2.1%) of

total volumes. The percentage of scrubbed volumes for task scans ranged from 0 to 11% (mean =

1.7%) of total volumes. The mean composite FD (Power et al. 2012) for each condition (rest and

task) was calculated and compared across time-points using two-sided, paired t-tests. Mean FD

during resting-state fMRI scans at baseline and follow-up were 0.12($\pm$ 0.06 s.d.) mm and 0.12($\pm$

0.07 s.d.) mm, respectively, and were not significantly different from one another (*t*-statistic = -

0.02; p = 0.99). Mean FD during task fMRI scans at baseline and follow-up were 0.22($\pm$ 0.16

s.d.) mm and 0.17($\pm$ 0.09 s.d.) mm, respectively, and were not significantly different from one

another (*t*-statistic = 1.23, p = 0.25).

2.5 Activation analysis: brain response to trauma versus neutral audio listening

FSL's fMRI Expert Analysis Tool (FEAT) (Woolrich et al. 2004) was used for fitting a general

linear model (GLM) to the voxelwise timeseries for each task scan after spatial smoothing using

a Gaussian kernel function (6 mm full width at half maximum (FWHM)). For 1st- level analysis,

models were generated for the neutral block, the trauma block, and a contrast of the two (trauma

> neutral). Confound explanatory variables (EVs) included the temporal derivative of each block,

5 nuisance regressors each for WM and CSF signal, outlier volumes (spikes in global signal (> 5

standard deviations) and motion (>0.9 mm FD); CONN Toolbox standard parameters (Whitfield-

Gabrieli and Nieto-Castanon 2012)), and 24 motion confounds (Friston et al. 1996). Second-

level analysis averaged the models from each of the two task scans performed at each time point.

Third-level analyses, using a two-sided, one-sample t-test (FSL randomize; Winkler et al. 2014)

identified group-level response for the contrast model (i) at baseline, and (ii) at the two-month

follow-up. A final third-level analysis (iii) compared the group-level responses to the contrast

model at baseline and follow-up using a two-sample, two-sided, paired t-test (FSL randomize;

Winkler et al. 2014). Third-level results were corrected for multiple comparisons using

threshold-free cluster enhancement (TFCE; alpha = 0.05) (Smith and Nichols 2009) which

identifies significant clusters based on the extent of local support from surrounding voxels.

2.6 RSFC analysis

Prior to extraction of RSFC, in addition to the preprocessing steps taken in 2.3, fMRI data were

further denoised using an in-house pipeline (https://github.com/kjamison/fmriclean). FMRIs

were bandpass filtered and regressed for 24 motion confounds (Friston et al. 1996), 5 nuisance

regressors each for WM and CSF, and one for global GM signal. The first five frames (scanner

start-up noise) and confound frames (spikes in global signal and motion) were discarded. RSFC

(Fisher Z-transformed Pearson correlation values) between the right and left amygdala and right

and left hippocampus (Desikan et al. 2006) was calculated for each resting-state scan. For the

supplemental RSFC analysis, each hippocampus was further segmented into head and tail

portions using FreeSurfer's hippocampal subregion segmentation tool (Iglesias et al. 2015).

Two-tailed, paired *t*-tests were used to compare the 4 RSFC measures before and after MDMA-

AT. Baseline to follow-up changes in these 4 measures were also correlated (Pearson's) with

individual level reductions in CAPS-IV using participant's age and mean FD changes (follow-up

– baseline) as covariates of non-interest. All statistical tests were performed at an alpha level of

0.05. P-values were corrected for multiple comparisons using the Benjamini-Hochberg algorithm

(Benjamini and Hochberg 1995) where indicated (pFDR).

2.7 Task FC analysis

Prior to extraction of task FC, the preprocessed residuals from the task activation analysis (2.5)

were further denoised with bandpass filtering and regressed for global GM signal. The first five

frames (scanner start-up noise) and confound frames (spikes in global signal (> 5 standard

deviations) and motion (>0.9 mm FD); CONN Toolbox standard parameters (Whitfield-Gabrieli

and Nieto-Castanon 2012)) were ignored. Functional connectivity (Fisher Z-transformed Pearson

correlation values) during each task fMRI scan (task FC) was calculated between 18 regions of

interest (ROIs): the right and left hippocampus head, hippocampus tail, amygdala, precuneus,

caudal anterior cingulate cortex (ACC), rostral ACC, posterior cingulate cortex (PCC), isthmus

cingulate, and insula. All ROIs were extracted from the Disikan-Killiany cortical atlas (see SI

Figures 7 and 8 for ROI definitions) (Desikan et al. 2006) , and head and tail portions of the

hippocampus were created using FreeSurfer's hippocampal subregion segmentation tool (Iglesias

et al. 2015). Two identical task scans were collected at each time point, thus FC values obtained

from both scans were averaged to give a single value for each connection per subject. Group-

level changes from pre- to post-therapy in the strength of functional connections were assessed

using two-tailed, paired *t*-tests. Pearson correlations were calculated between individuals'

changes in functional connection strength and change in CAPS-IV total severity scores (follow-

up – baseline) using participant age and mean FD changes between baseline and follow-up as

covariates of non-interest. All statistical tests were performed at an alpha level of 0.05. P-values

were corrected for multiple comparisons using the Benjamini-Hochberg algorithm (Benjamini

and Hochberg 1995) where indicated (pFDR).

3.   RESULTS

3.1 CAPS-IV total severity scores significantly decreased after HD MDMA-AT

Ten participants enrolled in the sub-study, and one withdrew consent after baseline due to

anxiety in the MRI scanner, leaving nine participants with MRI data at both time points (6 male,

3 female, aged 41.3; standard deviation (SD) = ± 10.9 years; 8 veterans and 1 first-responder; see

SI Table 1 for additional demographic information). All participants had chronic PTSD (mean

duration = 84 (± 45) months). One participant's baseline resting-state fMRI was truncated due to

technical issues, leaving eight participants for resting-state analysis and nine for the task fMRI

analysis. One participant began the trial with moderate PTSD (CAPS-IV > 39), while the

remaining eight presented with severe PTSD (CAPS-IV >59). Mean (SD) CAPS-IV total

severity scores of the nine individuals pre- and post-MDMA-AT were 86 (± 16) and 39 (± 25),

respectively, representing a significant decrease in PTSD symptom severity between the two

time points (Figure 2; N=9, t = 6.36, p = 0.00022). The average percent decrease in CAPS was

57 (± 26)%. Results on all participants enrolled in the Phase 2 parent trial have been previously

reported (Mithoefer et al. 2018).

Figure 2: Patient CAPS-IV total severity scores at the baseline (pre-therapy) and two-month

follow-up (post-therapy) scanning sessions. Black solid and dashed lines indicate group means

and medians, respectively. Blue marker = male. Red marker = female. A significant reduction is

PTSD severity was observed after MDMA-AT (baseline > follow-up; N=9, $t = 6.36$, p =

0.00022).

3.2 Baseline versus two-month follow-up amygdala-hippocampal RSFC

The RSFC was assessed between the amygdala and hippocampus before and after MDMA-AT

and the strengths of these connections are illustrated in Figure 3. All connections trended

towards increased RSFC after therapy compared to before therapy (using a two-sided paired t-

test), with left amygdala to left hippocampus having a significant increase prior to (but not after)

corrections for multiple comparisons (N=8, $t = -2.91$, uncorrected p = 0.0225, pFDR = 0.0901).

Figure 3: RSFC between the amygdala and hippocampus before and after MDMA-AT. P-values from a two-sided, paired t-test. Blue = male. Red = female. Black solid and dashed lines indicate group means and medians, respectively. (N = 8; t-statistics indicate baseline > follow-up; * uncorrected p < 0.05).

Individual-level pre-to-post-therapy changes between the strength of these functional connections were then correlated (two-sided Pearson's) with changes in CAPS scores (SI Figure 1). Only one of these correlations (right amygdala to left hippocampus FC versus CAPS) was significant before correction (N = 8; R = -0.820, uncorrected p = 0.0460, pFDR = 0.183).

3.4 Brain activation during symptom provocation

A script-driven autobiographical memory paradigm was used to assess brain activity during

symptom provocation (Ruth A Lanius et al. 2002; Ruth A. Lanius et al. 2004; Mertens et al.

2022). We compared whole-brain, script-driven activations (trauma > neutral) at baseline and

follow-up (Figure 4). Before therapy (baseline), there tended to be larger magnitude activation in

response to the trauma script versus the neutral script, as evidenced by the generally positive t-

statistics (Figure 4A-D). After correction using threshold free cluster enhancement (TFCE), there

was significantly greater activation during the trauma scripts compared to the neutral scripts in

four separate areas (see Figure 4 caption for details of each). After therapy, there were smaller

magnitude differences between brain activity in response to the two scripts, with no significant

clusters (Figure 4E). Finally, we assessed the differences in the contrast before and after therapy

(follow-up > baseline). There was generally greater contrast between the trauma and neutral

scripts at baseline compared to at follow-up, with one cluster in the bilateral cuneus and lingual

gyrus demonstrating significance after correction using TFCE (Figure 4F).

Figure 4: Group-level activation contrasts for trauma versus neutral script listening tasks (N = 9).

All panels show t-statistics for corresponding contrasts. Analyses were performed in 3 mm MNI

standard space, however results here are interpolated into 1 mm MNI standard space and clipped

to only show *t*-statistics greater than +/-1 for visualization purposes. Statistics reported below

(voxels, volume, t-statistic, corrected p-value) were calculated from original 3mm results. P-

values were corrected using threshold-free cluster enhancement (TFCE; see methods). For panels

A-E, positive *t*-statistics indicate greater activation to trauma scripts compared to neutral. For

panel F, sign indicates the direction of change in trauma>neutral contrast from baseline (i.e.

negative *t*-statistics indicate the contrast between trauma and neutral scripts was decreased at the

two-month follow-up compared to baseline). Crosshairs are located on the center of gravity

(c.o.g.) of significant clusters. (A) Cluster 1 for the baseline contrast is located primarily in the

right and left isthmus cingulate, with some overlap into the right and left precuneus (c.o.g.

MNI152 [0, -48, 24]; 6 voxels (162 mm^3); c.o.g. *t* = 9.61, p(TFCE) = 0.0293). (B) Cluster 2 for

the baseline contrast is located in the left caudal middle-frontal gyrus (c.o.g. MNI152 [-36, 21,

51]; 3 voxels (81 mm³); c.o.g. $t = 9.01$, p(TFCE) = 0.0234). (C) Cluster 3 for the baseline

contrast is located in the right medial prefrontal cortex (c.o.g. MNI152 [6, 57, 30]; 2 voxels (54

mm³); c.o.g. $t = 7.41$, p(TFCE) = 0.0488). (D) Cluster 4 for the baseline contrast is located in the

left rostral middle frontal gyrus (c.o.g. MNI152 [-21 54 15]; 1 voxel (27 mm³); c.o.g. $t = 9.45$,

p(TFCE) = 0.0312). (E) There were no significant activation contrasts at the two-month follow-

up (crosshairs shown at MNI152 [0, 0, 0]). (F) Comparing the group-level contrasts between

time points (follow-up > baseline), there exists one significant cluster located primarily in the

right and left cuneus, with some overlap into the right and left lingual gyrus (c.o.g. MNI152 [3, -

90, 3]; 47 voxels (1,269 mm³); c.o.g $t = -9.31$, p(TFCE) = 0.0391).

3.4 Baseline versus two-month follow-up changes in task FC

We compared the pre- and post-therapy FC strength between 18 brain regions of interest (ROIs)

during the task fMRI scans involving neutral and traumatic autobiographical audio recordings

(Figure 5A). The ROIs are as follows: the right and left hippocampus head, hippocampus tail,

amygdala, precuneus, caudal anterior cingulate cortex (ACC), rostral ACC, posterior cingulate

cortex (PCC), isthmus cingulate, and the insula. Only one functional connection was

significantly modified at follow-up compared to baseline: the right amygdala to left caudal ACC

(N=9; t-statistic = 3.04, p = 0.0148). This finding was no longer significant after corrections for

multiple comparisons, however (pFDR = 0.9875).

Individual-level pre-to-post therapy changes in these functional connections were then correlated

(two-sided, Pearson's) with the individual-level reductions in CAPS-IV scores (Figure 5B). Most

correlations were positive, meaning that larger reductions in connectivity from pre- to post-

therapy corresponded to larger improvements in PTSD symptoms. Four correlations between FC

and CAPS-IV changes were significant following multiple comparisons correction: the left

amygdala and left PCC (N=9; Pearson's R = 0.951, pFDR = 0.0462), the left amygdala and right

PCC (N=9; Pearson's R = 0.972, pFDR = 0.0197), the left amygdala and left insula (N=9;

Pearson's R = 0.977, pFDR = 0.0197), and the left isthmus cingulate and left hippocampal tail

(N=9; Pearson's R = 0.947, pFDR = 0.0462) (Figure 5C).

Figure 5: (A) Paired t-statistics shown for differences in functional connectivity between all brain regions of interest during the task fMRI scan involving neutral and trauma memory audio listening (N = 9; baseline > follow-up; * two-sided uncorrected p < 0.05; ** pFDR < 0.05, corrected). (B) Pearson correlation values between changes in ROI functional connectivity and reduction in CAPS scores. Changes were calculated as follow-up values minus baseline values. (N=9; * uncorrected p < 0.05; ** pFDR < 0.05) Age and mean FD difference between baseline

and follow-up were included as covariates of non-interest. (C) Scatter plots of the three

correlations that remained significant after corrections for multiple comparisons (i.e. pFDR <

0.05). Red marker = female. Blue marker = male.

3.5 **Supplemental analyses**

Although not the primary focus of our analysis, we also repeated the previous correlations using

other secondary outcome measures in place of CAPS-IV total severity scores. Namely, changes

between baseline and follow-up in the BDI-II (depression symptoms), the PSQI (sleep quality),

the PTGI (perceived growth following trauma), the DES-II (symptoms of dissociation), and the

GAF (general psychological function) were used (SI Figure 2). Following correction for

multiple comparisons, none of these correlations were significant.

We also replicated our *a priori* analyses of amygdala-hippocampal RSFC using head (anterior)

and tail (posterior) sub-regions of the hippocampus (SI Figure 3). The left hippocampal head to

left amygdala RSFC was increased at follow-up compared to baseline (N=8; t = -2.593,

uncorrected p = 0.0358), as was the RSFC between the left hippocampal tail and right amygdala

(N=8; t = -3.00, uncorrected p = 0.0199). Neither of these effects were significant after

corrections for multiple comparisons.

Lastly, we replicate our main functional connectivity analyses without the use of global signal

regression and find that these results largely show the same trends, however there is less

significance in some cases (SI Figures 4-6). The correlation between the left amygdala and left

insula task functional connectivity change and CAPS reductions was significant in both with and

without the use of global signal regression (SI Figure 6; N=9; R = 0.971, pFDR = 0.0229).

DISCUSSION

We report signatures of brain response during rest and audio listening task in eight veterans and

one first-responder with clinically diagnosed chronic and severe PTSD before and two-months

after MDMA-assisted therapy. We found a significant reduction in CAPS-IV total severity

scores after therapy (Figure 2), indicating our sub-study participants mirrored the results from the

parent study (Mithoefer et al. 2018). We found a trend suggesting that RSFC between the

amygdala and hippocampus was strengthened post-therapy, particularly in the left hemisphere

(Figure 3). Prior work suggests that modulation of amygdalae-hippocampal RSFC may be an

important component of MDMA-AT for PTSD (Sripada et al. 2012; Ben-Zion, Keynan, et al.

2020; Vaisvaser et al. 2013; Fan et al. 2015; Carhart-Harris et al. 2015; Lan et al. 2022), thus

investigating this connection in future studies is warranted. We also found participants had

increased activation in areas involved with self-referential processing and autobiographical

memory while listening to traumatic versus neutral memory narrations pre-therapy (Figure 4A-

D), and that no significant contrast existed after MDMA-AT (Figure 4E). Comparing trauma

versus neutral contrasts between baseline and follow-up revealed a significant decrease in cuneus

contrast after MDMA-AT (Figure 4F). Finally, the pre- to post-therapy reductions in FC during

the script listening task between the left amygdala and right PCC, left PCC, and left insula, as

well as FC between the left isthmus cingulate and left hippocampal tail strongly and significantly

correlated with PTSD symptom improvement (Figure 5C). These results begin to shed light on

the neurological mechanisms that may drive MDMA-AT for PTSD.

Previous work quantifying functional connectivity in PTSD and in stress exposure (Sripada et al.

2012; Ben-Zion, Keynan, et al. 2020; Vaisvaser et al. 2013; Fan et al. 2015) and acute MDMA

administration in controls (Carhart-Harris et al. 2015) suggests one mechanism of MDMA-AT

may be to increase pathologically lowered amygdala-hippocampal RSFC (Feduccia and

Mithoefer 2018). The amygdala is associated with fear expression, threat recognition, and

heightened response to emotional memories and is often dysregulated in patients with PTSD

(LeDoux 2003; Pitman et al. 2012; Liberzon et al. 1999; Etkin and Wager 2007; Bremner et al.

2005; Harnett, Goodman, and Knight 2020). The hippocampus also plays a central role in PTSD

as it is thought to provide contextual information important for cognitive-affect during memory

recollection (Harnett, Goodman, and Knight 2020; Pitman et al. 2012). Sripada and colleagues

(2012) found combat veterans with PTSD had reduced amygdala-hippocampal RSFC compared

to combat-exposed controls, leading them to speculate that this may relate to an inability to

contextualize affective information in PTSD. Carhart-Harris and colleagues (2015) demonstrated

that amygdala-hippocampal RSFC is increased acutely in MDMA administration compared to

placebo and this increase occurred in a manner that correlated with the drug's subjective effects

at a near-significant level, leading these researchers to propose that this functional connection

was a primary target of MDMA-AT. Increased amygdala-hippocampal RSFC has also been

linked to intranasal oxytocin administration after stress exposure (Fan et al. 2015) and this effect

was mediated by serotonin signaling (Lan et al. 2022) – two neuromodulators that play a

significant role in the pro-social and fear extinction effects of MDMA (Dumont et al. 2009; de la

Torre et al. 2004; Feduccia and Duvauchelle 2008; Green et al. 2003; Thompson et al. 2007;

Nardou et al. 2019; Young et al. 2017). Prior to our analysis (although after the study was

designed and the data collected), we hypothesized that the RSFC between the amygdala and

hippocampus would be higher after MDMA-AT compared to pre-therapy levels. Only the RSFC

between the left amygdala and left hippocampus was significantly increased (Figure 3) however,

this finding no longer met thresholds for significance after multiple comparisons correction

(pFDR = 0.09). We also found that the amount of increased right amygdala – left hippocampal

RSFC after MDMA-AT positively correlated with PTSD symptom improvement at a near-

significant level (SI Figure 1; R = -0.820, uncorrected p = 0.046, pFDR = 0.183). Our current

findings, though inconclusive, are suggestive of a trend towards moderate increases in amygdala

– hippocampal RSFC two-months after MDMA-AT. It is possible that more significant changes

would have been observed with a larger sample size, longer resting-state scans, or imaging

performed closer to MDMA administration. These findings justify the continued investigation of

amygdala-hippocampal RSFC in the therapeutic mechanisms of MDMA-AT in future studies.

We next sought to study brain response during autobiographical memory listening to draw

additional conclusions about MDMA-AT's effects in individuals with PTSD. Before therapy,

participants had larger activation in four areas during an individualized trauma script listening

task compared to neutral script listening: the right and left isthmus cingulate and precuneus, the

left caudal middle frontal gyrus, the right medial prefrontal cortex, and the left rostral middle

frontal gyrus (Figure 4A-D). These regions are broadly involved in self-processing operations

(e.g. first-person perspective taking), episodic memory retrieval, visual-spatial imagery, auto-

biographical memory recollection, and are included in or interact with the default mode network

(Cavanna and Trimble 2006; Cauda et al. 2010; Spreng, Mar, and Kim 2009; Spreng and Grady

2010; Kalenzaga et al. 2015; Demblon, Bahri, and D'Argembeau 2016; Raichle 2015). The

retrosplenial cortex – located within the isthmus cingulate - is also found to be consistently

activated by emotionally salient stimuli, and has been proposed to play a role in the interaction

between emotion and memory (Maddock 1999). We conjecture that increased activation in these

regions during traumatic compared to neutral audio listening (Figure 4A-D) could be related to

an increased intensity of the recollection or re-experiencing of traumatic memories compared to

neutral ones for patients before therapy. At 2-month follow-up to MDMA-AT, there was no

significant difference in the trauma vs neutral script activation (Figure 4E). The longitudinal

comparison of these two time points indicated that the contrast between trauma and neutral was

larger at baseline, particularly in a significant cluster in the right and left cuneus/lingual gyrus

(Figure 4F). Cuneus activity during autobiographical memory tasks often coincides with activity

in the frontal regions highlighted by the baseline contrast, and has been found to correlate with

memory recall accuracy (Spreng and Grady 2010; Demblon, Bahri, and D'Argembeau 2016;

Kalenzaga et al. 2015). Cuneus activity is thought to enhance the visual imagery of

autobiographical memory recollection (Cabeza and St Jacques 2007), therefore decreased

contrast in this area at follow-up suggests that intensity of visual imagery contrast between

trauma and neutral memories may be decreased after MDMA-AT. Larger studies may allow

more statistical power to identify additional longitudinal differences. Other longitudinal studies

of individuals with PTSD have found that decreases in precuneus, isthmus cingulate, and middle

frontal gyrus activation during symptom provocation is correlated with reductions in PTSD

symptom severity (Garrett et al. 2019; Ke et al. 2016).

PTSD is often associated with hyperactivity in the amygdala (Pitman et al. 2012); the acute

administration of MDMA in healthy volunteers decreases blood flow to the amygdala during rest

(Carhart-Harris et al. 2015) and attenuates its response to angry faces (Bedi et al. 2009). We had

hypothesized that we would observe hyperactivity of the amygdala to trauma versus neutral

scripts at baseline and that MDMA-AT would attenuate this response, however we observed

neither. It is important to note inconsistencies in the literature here. Amygdala hyperactivity in

PTSD is not always observed, possibly due to differences in subtypes, sex, cultural

representations, or choice of paradigm (van Huijstee and Vermetten 2018; R. A. Lanius et al.

2001; Ruth A Lanius et al. 2002; Helpman et al. 2021; Chiao et al. 2008; Liddell and Jobson

2016). Additionally, while MDMA did suppress amygdala activity during rest and in response to

angry faces as previously mentioned, there was no observed impact on its response to

autobiographical memories (Carhart-Harris et al. 2014). While activation-based analyses deserve

continued attention in future studies to rectify these inconsistencies, functional connectivity is a

complimentary approach we can use to extract additional information from fMRI.

Functional connectivity analyses in individuals with PTSD have revealed aberrant connectivity

between several regions within default mode, limbic, and salience networks, and more broadly,

regions involved in emotional and self-referential processing (Abdallah et al. 2017; Lazarov et al.

2017; Malivoire et al. 2018; Brown et al. 2014; Gilboa et al. 2004; Fonzo et al. 2021; Rabinak et

al. 2011; Belleau et al. 2020; Thome et al. 2020; Ruth A. Lanius et al. 2004; Sripada et al. 2012),

and, further, the administration of MDMA in healthy volunteers has been shown to disrupt the

functional integrity of these networks (Walpola et al. 2017; Carhart-Harris et al. 2015; Müller et

al. 2021; Dipasquale et al. 2019; Avram et al. 2022). Here, we measured functional connectivity

during script-driven autobiographical memory recall between the right and left hippocampus head,

hippocampus tail, amygdala, precuneus, caudal ACC, rostral ACC, PCC, isthmus cingulate, and

the insula. Our ROIs were defined and labeled based on the Desikan-Killiany (DK) brain atlas (SI

Figures 7 and 8) (Desikan et al. 2006). We chose to segment the hippocampus into anterior (head)

and posterior (tail) ROIs based on recent work showing that the two portions' FC are differentially

effected by PTSD (Malivoire et al. 2018). We assessed group-level changes in the strength of these

functional connections and found no significant differences between baseline and follow-up after

corrections for multiple comparisons (Figure 5A). However, we did find that greater recovery

(larger decreases in CAPS-IV at follow-up) was associated with reductions in FC between the left

amygdala and the right and left PCC, as well as the left insula (Figure 5C). The acute effects of

MDMA in healthy volunteers has been shown to decrease the FC of the PCC (Müller et al. 2021;

Dipasquale et al. 2019) and insula (Walpola et al. 2017), and alter amygdala and hippocampus FC

(Carhart-Harris et al. 2015), highlighting the potential relevance of our current findings. Amygdala

to posterior and mid-cingulate cortex FC has been shown to be associated with PTSD severity at

different stages of disease progression, although differing patient populations and assessment time-

lines lead to conflicting results (Belleau et al. 2020; R. A. Lanius et al. 2010; Bluhm et al. 2009;

Cisler et al. 2016; Shou et al. 2017). One finding in healthy adults shows increased amygdala -

PCC FC following the acute exposure to stress (Veer et al. 2011), thus the association between

recovery and reduced amygdala – PCC task FC at follow-up possibly relates to reduced stress

response to trauma memories (although the finding by Veer and colleagues is more posterior to

the ROI used here). Amygdala and insula RSFC is increased in PTSD (Sripada et al. 2012; Rabinak

et al. 2011) (except in one study which finds the opposite (Fonzo et al. 2021)), and reduced

amygdala-insula FC during negative image reappraisal is associated with larger improvements in

PTSD symptoms (Cisler et al. 2016). The strength of left amygdala-insula FC also positively

correlates with the amount of acute anxiety measured in participants just before scanning (Baur et

al. 2013). Attenuated functional connectivity of these two regions at follow-up in the present study

possibly suggests a decreased intensity of recalled events, less 're-experiencing', or reduced

anxiety during the script-driven memory recall (Etkin and Wager 2007). Lastly, we found that

reductions in CAPS-IV at follow-up were associated with reduced FC between the left isthmus

cingulate and left hippocampal tail (Figure 5C). The isthmus cingulate labeled here consists of the

most posterior potions of the PCC (SI Figure 7). Increased FC between these two regions has

previously been reported in PTSD patients compared to trauma-exposed health controls (Malivoire

et al. 2018; Lazarov et al. 2017), and the present finding is possibly indicative of changes in

memory contextualization and reduced threat sensitivity at two-month follow-up to MDMA-AT

compared to baseline (Bluhm et al. 2009).

PTSD is characterized by decreased fear extinction in response to trauma-related stimuli. One

possible mechanism through which MDMA-AT operates is enhanced reconsolidation and/or fear

extinction processes (Feduccia and Mithoefer 2018). Several studies with MDMA implicate

reconsolidation or fear extinction processes, and while it is currently unclear whether MDMA

acts on only one or both, it is important to note that the two interact (Suzuki et al. 2004). Rodent

models have demonstrated that the administration of MDMA prior to extinction learning

enhances extinction retention (tested 48 hours after learning) and this effect is blocked by acute

and chronic treatment with a serotonin transporter inhibitor (M. B. Young et al. 2015; 2017).

Hake et al. (2019) found that MDMA administered during extinction learning phases did not

enhance fear extinction memory, while MDMA administration during reconsolidation phases

resulted in prolonged reductions in conditioned fear. In addition, MDMA administered prior to

trauma-cue exposure (reconsolidation phase) in rodents resulted in reduced stress-related

behavioral responses 7 days later (Arluk et al. 2022). Two recent trials in healthy humans found

that MDMA (100 mg and 125 mg, respectively) administered prior to extinction learning

resulted in improved extinction learning at extinction recall phases (48hr and 24hr later,

respectively) compared to the placebo group (Maples-Keller et al. 2022; Vizeli et al. 2022).

Doss and colleagues (2018) found that 1 mg/kg of MDMA in healthy humans attenuated the

encoding and retrieval of salient details from positive and negative stimuli (but not neutral

stimuli), suggesting an ability for MDMA to alter emotional memory representation.

Interestingly, a fMRI study in healthy humans found decreased activation in the precuneus/PCC

during fear extinction learning (Ridderbusch et al. 2021), regions highlighted by our present

study and others in PTSD (Garrett et al. 2019; Ke et al. 2016).

LIMITATIONS

The small sample size of the present study and the lack of a control population (e.g. trauma-

exposed healthy controls) may decrease the generalizability of these findings. The trial design

was placebo-controlled for dose-response (low, medium, and high), however, the follow-up

scans used in this study were after the breaking of the blind and dose cross-over (low/medium to

high) had occurred. For neuroimaging studies, comparisons with control populations are helpful

for contextualizing longitudinal changes in brain response and provide information about

whether changes in patient populations represent an abnormal response being restored to

normality or a compensatory mechanism. In addition, multi-point imaging of healthy control or

non-treatment (placebo) groups allow for the quantification of test-retest variability. Lastly, had

we imaged a cohort that received therapy without MDMA, we would have been able to assess

longitudinal brain changes that were unique to or enhanced by MDMA.

Here it must be discussed that PTSD is a disorder exhibiting at least two major sub-types

(dissociative and non-dissociative) with characteristically opposing phenomenological and

physiological responses to symptom provocation, which may explain inconsistencies in the

634 PTSD neurobiology literature (van Huijstee and Vermetten 2018). PTSD sub-type information

635 was not collected in the present study. In addition to sub-type heterogeneity, males and females

636 may also differ in their adaptive neural responses to trauma (Helpman et al. 2021). Limited by

637 our sample size, we did not investigate differences between males and females in this study.

639  The accepted standard for assessing PTSD severity is the Clinician-Administered PTSD Scale

640 (Blake et al. 1990). Specifically, CAPS-IV was used in this study. CAPS-IV involves an hour-

641 long semi-structured interview with a clinician and, though comprehensive, faces limitations. In

642 their baseline CAPS-IV assessment, and subsequently thereafter, patients were asked to refer to

643 an index trauma that was measured throughout the study. This may present an issue in accurately

644 assessing global PTSD severity if an adjacent or un-related trauma surfaces during therapy and

645 becomes the prominent driver of their symptoms. These issues, combined with difficulty in

646 blinding and expectancy effects, present additional challenges in accurately mapping fMRI

647 metrics to clinical outcomes.

649 The task design used in this study examined differences in brain response to personalized audio

650 scripts generated from narrations of traumatic and neutral memories. Many different stimuli have

651 been used in fMRI studies of PTSD (R. A. Lanius et al. 2006; Sartory et al. 2013; Patel et al.

652 2012), each providing its own unique advantages and disadvantages. Our present design

653 optimizes personal relevance of the stimuli; however, this has the consequence of presenting

654 each subject with a different set of stimuli, whereby brain responses within each block are not

655 time-locked across participants. Also, it has previously been shown that PTSD survivors take

656 longer to retrieve unrelated autobiographical information when listening to taped imagery scripts

of their traumatic memories (Kleim, Wallott, and Ehlers 2008). This suggests the possibility that

those with the most severe PTSD will take the longest to cognitively transition to the neutral

block from the trauma block. If this is true, then there would perhaps be an inverse-"U"

relationship between PTSD severity and contrast between the trauma and neutral conditions, if

the blocks are not spaced far enough apart to allow adequate time for patients to return to a

baseline level of cognitive functioning. Additionally, because our repeated task fMRI scans were

identical (rather than counterbalanced for condition), there could be primacy effects in the

neutral condition (which was always first) and/or fatigue effects in the trauma condition (which

was always last).

We did not aim to characterize lateralization in our findings, and though most of our significant

FC results were found in the left amygdala, we did not test for statistical interaction effects

between hemispheres. While lateralization of the amygdala remains debated, it has been

suggested by early work that the left amygdala is more strongly related to conscious (versus

unconscious) perception and emotional regulation (Morris, Öhman, and Dolan 1998).

Finally, the pre-specified aim of this study was to estimate longitudinal (baseline to 2-months

after final MDMA session) changes in ROI response to traumatic audio scripts. Between the start

of data collection and analysis, new literature emerged (Carhart-Harris et al. 2015; Lan et al.

2022) implicating amygdala-hippocampus RSFC as a potential target of MDMA-AT, compelling

us to expand our analysis beyond the pre-specified aims. Functional connectivity estimates from

shorter scans (e.g. five minutes in the case of our resting-state data) can have lower reliability

(Birn et al. 2013) and therefore the trends in increased RSFC between amygdala and

hippocampal regions reported here should be considered preliminary.

CONCLUSION

We report functional brain changes associated with MDMA-AT in veterans and first responders

with moderate-to-severe and chronic PTSD. We had hypothesized that MDMA-AT may act

through strengthening the RSFC between the amygdala and hippocampus, a connection which is

weaker in PTSD populations (Sripada et al. 2012) and increased acutely by MDMA in healthy

volunteers (Carhart-Harris et al. 2015). The trends found here are suggestive of such in the left

amygdala – left hippocampus, however larger studies are needed. We also provide preliminary

evidence that MDMA-AT alters brain response during symptom provocation in regions

associated with fear response, anxiety, self-referential processing, and salient autobiographical

memory, and are commonly found to be hyperactive in PTSD patients (Sartory et al. 2013; Patel

et al. 2012). Finally, the reduction of several functional connections during autobiographical

memory audio co-varied with symptom reduction in PTSD. Of these connections, the left

amygdala – left insula is perhaps the most interesting, due to the role of amygdala-insular FC in

anxiety and PTSD symptomatology (Baur et al. 2013; Cisler et al. 2016; Sripada et al. 2012;

Rabinak et al. 2011). More research is necessary to confirm these results and to disentangle

effects specific to MDMA and its combination with psychotherapy.

----------------------- *END OF MANUSCRIPT* ------------------------

REFERENCES

Abdallah, C. G., K. M. Wrocklage, C. L. Averill, T. Akiki, B. Schweinsburg, A. Roy, B. Martini, S. M. Southwick, J. H. Krystal, and J. C. Scott. 2017. "Anterior Hippocampal Dysconnectivity in Posttraumatic Stress Disorder: A Dimensional and Multimodal Approach." *Translational Psychiatry* 7 (2): e1045–e1045. https://doi.org/10.1038/tp.2017.12.

Arluk, Shira, Michael A. Matar, Lior Carmi, Oded Arbel, Joseph Zohar, Doron Todder, and Hagit Cohen. 2022. "MDMA Treatment Paired with a Trauma-Cue Promotes Adaptive Stress Responses in a Translational Model of PTSD in Rats." *Translational Psychiatry* 12 (1): 181. https://doi.org/10.1038/s41398-022-01952-8.

Avram, Mihai, Felix Müller, Helena Rogg, Alexandra Korda, Christina Andreou, Friederike Holze, Patrick Vizeli, Laura Ley, Matthias E. Liechti, and Stefan Borgwardt. 2022. "Characterizing Thalamocortical (Dys)Connectivity Following D-Amphetamine, LSD, and MDMA Administration." *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* 7 (9): 885–94. https://doi.org/10.1016/j.bpsc.2022.04.003.

Baur, Volker, Jürgen Hänggi, Nicolas Langer, and Lutz Jäncke. 2013. "Resting-State Functional and Structural Connectivity within an Insula-Amygdala Route Specifically Index State and Trait Anxiety." *Biological Psychiatry* 73 (1): 85–92. https://doi.org/10.1016/j.biopsych.2012.06.003.

Bedi, Gillinder, K. Luan Phan, Mike Angstadt, and Harriet de Wit. 2009. "Effects of MDMA on Sociability and Neural Response to Social Threat and Social Reward." *Psychopharmacology* 207 (1): 73–83. https://doi.org/10.1007/s00213-009-1635-z.

Belleau, Emily L., Lauren E. Ehret, Jessica L. Hanson, Karen J. Brasel, Christine L. Larson, and Terri A. deRoon-Cassini. 2020. "Amygdala Functional Connectivity in the Acute Aftermath of Trauma Prospectively Predicts Severity of Posttraumatic Stress Symptoms." *Neurobiology of Stress* 12 (May): 100217. https://doi.org/10.1016/j.ynstr.2020.100217.

Benjamini, Yoav, and Yosef Hochberg. 1995. "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing." *Journal of the Royal Statistical Society. Series B (Methodological)* 57 (1): 289–300.

Ben-Zion, Ziv, Nimrod Jackob Keynan, Roee Admon, Haggai Sharon, Pinchas Halpern, Israel Liberzon, Arieh Shalev, and Talma Henlder. 2020. "Hippocampal-Amygdala Resting State Functional Connectivity Serves as Resilience Factor for Short- and Long-Term Stress Exposure." *Biological Psychiatry*, 75 Annual Scientific Convention and Meeting, 87 (9, Supplement): S88–89. https://doi.org/10.1016/j.biopsych.2020.02.248.

Ben-Zion, Ziv, Yoav Zeevi, Nimrod Jackob Keynan, Roee Admon, Tal Kozlovski, Haggai Sharon, Pinchas Halpern, et al. 2020. "Multi-Domain Potential Biomarkers for Post-Traumatic Stress Disorder (PTSD) Severity in Recent Trauma Survivors." *Translational Psychiatry* 10 (1): 208. https://doi.org/10.1038/s41398-020-00898-z.

Birn, Rasmus M., Erin K. Molloy, Rémi Patriat, Taurean Parker, Timothy B. Meier, Gregory R. Kirk, Veena A. Nair, M. Elizabeth Meyerand, and Vivek Prabhakaran. 2013. "The Effect of Scan Length on the Reliability of Resting-State FMRI Connectivity Estimates." *NeuroImage* 83 (December): 550–58. https://doi.org/10.1016/j.neuroimage.2013.05.099.

Blake, Dudley, Frank Weathers, LN Nagy, D. Kaloupek, G. Klauminzer, Dennis Charney, and Terence Keane. 1990. "A Clinician Rating Scale for Assessing Current and Lifetime PTSD: The CAPS-1." *Behav. Ther.* 18 (January).

Bluhm, Robyn L, Peter C Williamson, Elizabeth A Osuch, Paul A Frewen, Todd K Stevens, Kristine Boksman, Richard W J Neufeld, Jean Théberge, and Ruth A Lanius. 2009. "Alterations in Default Network Connectivity in Posttraumatic Stress Disorder Related to Early-Life Trauma." *Journal of Psychiatry & Neuroscience : JPN* 34 (3): 187–94.

Bremner, J. Douglas, Eric Vermetten, Christian Schmahl, Viola Vaccarino, Meena Vythilingam, Nadeem Afzal, Christian Grillon, and Dennis S. Charney. 2005. "Positron Emission Tomographic Imaging of Neural Correlates of a Fear Acquisition and Extinction Paradigm in Women with Childhood Sexual-Abuse-Related Post-Traumatic Stress Disorder." *Psychological Medicine* 35 (6): 791–806. https://doi.org/10.1017/S0033291704003290.

Brown, Vanessa M, Kevin S LaBar, Courtney C Haswell, Andrea L Gold, Gregory McCarthy, and Rajendra A Morey. 2014. "Altered Resting-State Functional Connectivity of Basolateral and Centromedial Amygdala Complexes in Posttraumatic Stress Disorder." *Neuropsychopharmacology* 39 (2): 351–59. https://doi.org/10.1038/npp.2013.197.

Cabeza, Roberto, and Peggy St Jacques. 2007. "Functional Neuroimaging of Autobiographical Memory." *Trends in Cognitive Sciences* 11 (5): 219–27. https://doi.org/10.1016/j.tics.2007.02.005.

Carhart-Harris, Murphy K, Leech R, Erritzoe D, Wall MB, Ferguson B, Williams LT, et al. 2015. "The Effects of Acutely Administered 3,4-Methylenedioxymethamphetamine on Spontaneous Brain Function in Healthy Volunteers Measured With Arterial Spin Labeling and Blood Oxygen Level-Dependent Resting State Functional Connectivity." *Biological Psychiatry* 78 (8). https://doi.org/10.1016/J.BIOPSYCH.2013.12.015.

Carhart-Harris, R L, M B Wall, D Erritzoe, M Kaelen, B Ferguson, I De Meer, M Tanner, et al. 2014. "The Effect of Acutely Administered MDMA on Subjective and BOLD-FMRI Responses to Favourite and Worst Autobiographical Memories." *The International Journal of Neuropsychopharmacology* 17 (4): 527–40. https://doi.org/10.1017/S1461145713001405.

Cauda, Franco, Giuliano Geminiani, Federico D'Agata, Katiuscia Sacco, Sergio Duca, Andrew P. Bagshaw, and Andrea E. Cavanna. 2010. "Functional Connectivity of the Posteromedial Cortex." *PLoS ONE* 5 (9): e13107. https://doi.org/10.1371/journal.pone.0013107.

Cavanna, Andrea E., and Michael R. Trimble. 2006. "The Precuneus: A Review of Its Functional Anatomy and Behavioural Correlates." *Brain: A Journal of Neurology* 129 (Pt 3): 564–83. https://doi.org/10.1093/brain/awl004.

Chiao, Joan Y., Tetsuya Iidaka, Heather L. Gordon, Junpei Nogawa, Moshe Bar, Elissa Aminoff, Norihiro Sadato, and Nalini Ambady. 2008. "Cultural Specificity in Amygdala Response to Fear Faces." *Journal of Cognitive Neuroscience* 20 (12): 2167–74. https://doi.org/10.1162/jocn.2008.20151.

Cisler, J. M., B. A. Sigel, J. S. Steele, S. Smitherman, K. Vanderzee, J. Pemberton, T. L. Kramer, and C. D. Kilts. 2016. "Changes in Functional Connectivity of the Amygdala during Cognitive Reappraisal Predict Symptom Reduction during Trauma-Focused Cognitive-Behavioral Therapy among Adolescent Girls with Post-Traumatic Stress Disorder." *Psychological Medicine* 46 (14): 3013–23. https://doi.org/10.1017/S0033291716001847.

Curry, Daniel W., Matthew B. Young, Andrew N. Tran, Georges E. Daoud, and Leonard L. Howell. 2018. "Separating the Agony from Ecstasy: R(–)-3,4-Methylenedioxymethamphetamine Has Prosocial and Therapeutic-like Effects without

Signs of Neurotoxicity in Mice." *Neuropharmacology* 128 (January): 196–206. https://doi.org/10.1016/j.neuropharm.2017.10.003.

Dale, A. M., B. Fischl, and M. I. Sereno. 1999. "Cortical Surface-Based Analysis. I. Segmentation and Surface Reconstruction." *NeuroImage* 9 (2): 179–94. https://doi.org/10.1006/nimg.1998.0395.

Davidson, J. R. T. 2000. "Trauma: The Impact of Post-Traumatic Stress Disorder." *Journal of Psychopharmacology* 14 (2_suppl1): S5–12. https://doi.org/10.1177/02698811000142S102.

Demblon, Julie, Mohamed Ali Bahri, and Arnaud D'Argembeau. 2016. "Neural Correlates of Event Clusters in Past and Future Thoughts: How the Brain Integrates Specific Episodes with Autobiographical Knowledge." *NeuroImage* 127 (February): 257–66. https://doi.org/10.1016/j.neuroimage.2015.11.062.

Desikan, Rahul S., Florent Ségonne, Bruce Fischl, Brian T. Quinn, Bradford C. Dickerson, Deborah Blacker, Randy L. Buckner, et al. 2006. "An Automated Labeling System for Subdividing the Human Cerebral Cortex on MRI Scans into Gyral Based Regions of Interest." *NeuroImage* 31 (3): 968–80. https://doi.org/10.1016/j.neuroimage.2006.01.021.

Dipasquale, Ottavia, Pierluigi Selvaggi, Mattia Veronese, Anthony S. Gabay, Federico Turkheimer, and Mitul A. Mehta. 2019. "Receptor-Enriched Analysis of Functional Connectivity by Targets (REACT): A Novel, Multimodal Analytical Approach Informed by PET to Study the Pharmacodynamic Response of the Brain under MDMA." *NeuroImage* 195 (July): 252–60. https://doi.org/10.1016/j.neuroimage.2019.04.007.

Doss, Manoj K., Jessica Weafer, David A. Gallo, and Harriet de Wit. 2018. "MDMA Impairs Both the Encoding and Retrieval of Emotional Recollections." *Neuropsychopharmacology* 43 (4): 791–800. https://doi.org/10.1038/npp.2017.171.

Dumont, G. J. H., F. C. G. J. Sweep, R. van der Steen, R. Hermsen, A. R. T. Donders, D. J. Touw, J. M. A. van Gerven, J. K. Buitelaar, and R. J. Verkes. 2009. "Increased Oxytocin Concentrations and Prosocial Feelings in Humans after Ecstasy (3,4-Methylenedioxymethamphetamine) Administration." *Social Neuroscience* 4 (4): 359–66. https://doi.org/10.1080/17470910802649470.

Ehlers, Anke. 2010. "Understanding and Treating Unwanted Trauma Memories in Posttraumatic Stress Disorder." *Zeitschrift Für Psychologie / Journal of Psychology* 218 (2): 141–45. https://doi.org/10.1027/0044-3409/a000021.

Etkin, Amit, and Tor D. Wager. 2007. "Functional Neuroimaging of Anxiety: A Meta-Analysis of Emotional Processing in PTSD, Social Anxiety Disorder, and Specific Phobia." *American Journal of Psychiatry* 164 (10): 1476–88. https://doi.org/10.1176/appi.ajp.2007.07030504.

Fan, Yan, Karin Pestke, Melanie Feeser, Sabine Aust, Jens C. Pruessner, Heinz Böker, Malek Bajbouj, and Simone Grimm. 2015. "Amygdala–Hippocampal Connectivity Changes During Acute Psychosocial Stress: Joint Effect of Early Life Stress and Oxytocin." *Neuropsychopharmacology* 40 (12): 2736–44. https://doi.org/10.1038/npp.2015.123.

Feduccia, Allison A., and Christine L. Duvauchelle. 2008. "Auditory Stimuli Enhance MDMA-Conditioned Reward and MDMA-Induced Nucleus Accumbens Dopamine, Serotonin and Locomotor Responses." *Brain Research Bulletin* 77 (4): 189–96. https://doi.org/10.1016/j.brainresbull.2008.07.007.

Feduccia, Allison A., and Michael C. Mithoefer. 2018. "MDMA-Assisted Psychotherapy for PTSD: Are Memory Reconsolidation and Fear Extinction Underlying Mechanisms?"

Progress in Neuro-Psychopharmacology and Biological Psychiatry 84 (June): 221–28. https://doi.org/10.1016/J.PNPBP.2018.03.003.

Feduccia, Allison A., Michael C. Mithoefer, Lisa Jerome, Julie Holland, Amy Emerson, and Rick Doblin. 2018. "Response to the Consensus Statement of the PTSD Psychopharmacology Working Group." *Biological Psychiatry* 84 (2): e21–22. https://doi.org/10.1016/j.biopsych.2017.11.023.

Fonzo, Gregory A., Madeleine S. Goodkind, Desmond J. Oathes, Yevgeniya V. Zaiko, Meredith Harvey, Kathy K. Peng, M. Elizabeth Weiss, et al. 2021. "Amygdala and Insula Connectivity Changes Following Psychotherapy for Posttraumatic Stress Disorder: A Randomized Clinical Trial." *Biological Psychiatry* 89 (9): 857–67. https://doi.org/10.1016/j.biopsych.2020.11.021.

Friston, Karl J., Steven Williams, Robert Howard, Richard S. J. Frackowiak, and Robert Turner. 1996. "Movement-Related Effects in FMRI Time-Series." *Magnetic Resonance in Medicine* 35 (3): 346–55. https://doi.org/10.1002/mrm.1910350312.

Garrett, Amy, Judith A. Cohen, Sanno Zack, Victor Carrion, Booil Jo, Joseph Blader, Alexis Rodriguez, Thomas J. Vanasse, Allan L. Reiss, and W. Stewart Agras. 2019. "Longitudinal Changes in Brain Function Associated with Symptom Improvement in Youth with PTSD." *Journal of Psychiatric Research* 114 (July): 161–69. https://doi.org/10.1016/j.jpsychires.2019.04.021.

Gilboa, Asaf, Arieh Y Shalev, Lucian Laor, Hava Lester, Yoram Louzoun, Roland Chisin, and Omer Bonne. 2004. "Functional Connectivity of the Prefrontal Cortex and the Amygdala in Posttraumatic Stress Disorder." *Biological Psychiatry* 55 (3): 263–72. https://doi.org/10.1016/j.biopsych.2003.08.004.

Glover, Gary H. 2011. "Overview of Functional Magnetic Resonance Imaging." *Neurosurgery Clinics of North America*, Functional Imaging, 22 (2): 133–39. https://doi.org/10.1016/j.nec.2010.11.001.

Goetter, Elizabeth M., Eric Bui, Rebecca A. Ojserkis, Rebecca J. Zakarian, Rebecca Weintraub Brendel, and Naomi M. Simon. 2015. "A Systematic Review of Dropout From Psychotherapy for Posttraumatic Stress Disorder Among Iraq and Afghanistan Combat Veterans." *Journal of Traumatic Stress* 28 (5): 401–9. https://doi.org/10.1002/jts.22038.

Gradus, Jaimie L. 2017. "Prevalence and Prognosis of Stress Disorders: A Review of the Epidemiologic Literature." *Clinical Epidemiology* 9 (May): 251–60. https://doi.org/10.2147/CLEP.S106250.

Green, A. Richard, Annis O. Mechan, J. Martin Elliott, Esther O'Shea, and M. Isabel Colado. 2003. "The Pharmacology and Clinical Pharmacology of 3,4-Methylenedioxymethamphetamine (MDMA, 'Ecstasy')." *Pharmacological Reviews* 55 (3): 463–508. https://doi.org/10.1124/pr.55.3.3.

Hake, Holly S., Jazmyne K. P. Davis, River R. Wood, Margaret K. Tanner, Esteban C. Loetz, Anais Sanchez, Mykola Ostrovskyy, et al. 2019. "3,4-Methylenedioxymethamphetamine (MDMA) Impairs the Extinction and Reconsolidation of Fear Memory in Rats." *Physiology & Behavior* 199 (February): 343–50. https://doi.org/10.1016/j.physbeh.2018.12.007.

Harnett, Nathaniel G., Adam M. Goodman, and David C. Knight. 2020. "PTSD-Related Neuroimaging Abnormalities in Brain Function, Structure, and Biochemistry." *Experimental Neurology* 330 (August): 113331. https://doi.org/10.1016/j.expneurol.2020.113331.

Helpman, Liat, Xi Zhu, Sigal Zilcha-Mano, Benjamin Suarez-Jimenez, Amit Lazarov, Bret Rutherford, and Yuval Neria. 2021. "Reversed Patterns of Resting State Functional Connectivity for Females vs. Males in Posttraumatic Stress Disorder." *Neurobiology of Stress*, September, 100389. https://doi.org/10.1016/J.YNSTR.2021.100389.

Hoge, Charles W., Carl A. Castro, Stephen C. Messer, Dennis McGurk, Dave I. Cotting, and Robert L. Koffman. 2004. "Combat Duty in Iraq and Afghanistan, Mental Health Problems, and Barriers to Care." *The New England Journal of Medicine* 351 (1): 13–22. https://doi.org/10.1056/NEJMoa040603.

Huijstee, Jytte van, and Eric Vermetten. 2018. "The Dissociative Subtype of Post-Traumatic Stress Disorder: Research Update on Clinical and Neurobiological Features." *Current Topics in Behavioral Neurosciences* 38: 229–48. https://doi.org/10.1007/7854_2017_33.

Iglesias, Juan Eugenio, Jean C. Augustinack, Khoa Nguyen, Christopher M. Player, Allison Player, Michelle Wright, Nicole Roy, et al. 2015. "A Computational Atlas of the Hippocampal Formation Using Ex Vivo, Ultra-High Resolution MRI: Application to Adaptive Segmentation of in Vivo MRI." *NeuroImage* 115 (July): 117–37. https://doi.org/10.1016/j.neuroimage.2015.04.042.

Javidi, H., and M. Yadollahie. 2012. "Post-Traumatic Stress Disorder." *The International Journal of Occupational and Environmental Medicine* 3 (1): 2–9.

Jerome, Lisa, Allison A. Feduccia, Julie B. Wang, Scott Hamilton, Berra Yazar-Klosinski, Amy Emerson, Michael C. Mithoefer, and Rick Doblin. 2020. "Long-Term Follow-up Outcomes of MDMA-Assisted Psychotherapy for Treatment of PTSD: A Longitudinal Pooled Analysis of Six Phase 2 Trials." *Psychopharmacology* 237 (8): 2485–97. https://doi.org/10.1007/s00213-020-05548-2.

Kalenzaga, Sandrine, Marco Sperduti, Adèle Anssens, Penelope Martinelli, Anne-Dominique Devauchelle, Thierry Gallarda, Marion Delhommeau, et al. 2015. "Episodic Memory and Self-Reference via Semantic Autobiographical Memory: Insights from an FMRI Study in Younger and Older Adults." *Frontiers in Behavioral Neuroscience* 8. https://www.frontiersin.org/articles/10.3389/fnbeh.2014.00449.

Ke, Jun, Li Zhang, Rongfeng Qi, Weihui Li, Cailan Hou, Yuan Zhong, Zhong He, Lingjiang Li, and Guangming Lu. 2016. "A Longitudinal FMRI Investigation in Acute Post-Traumatic Stress Disorder (PTSD)." *Acta Radiologica* 57 (11): 1387–95. https://doi.org/10.1177/0284185115585848.

Kleim, Birgit, Franziska Wallott, and Anke Ehlers. 2008. "Are Trauma Memories Disjointed from Other Autobiographical Memories in   Posttraumatic Stress Disorder? An Experimental Investigation." *Behavioural and Cognitive Psychotherapy* 36 (2): 221–34. https://doi.org/10.1017/S1352465807004080.

Lan, Chunmei, Congcong Liu, Keshuang Li, Zhiying Zhao, Jiaxin Yang, Yina Ma, Dirk Scheele, Shuxia Yao, Keith M. Kendrick, and Benjamin Becker. 2022. "Oxytocinergic Modulation of Stress-Associated Amygdala-Hippocampus Pathways in Humans Is Mediated by Serotonergic Mechanisms." *The International Journal of Neuropsychopharmacology* 25 (10): 807–17. https://doi.org/10.1093/ijnp/pyac037.

Lanius, R. A., R. L. Bluhm, N. J. Coupland, K. M. Hegadoren, B. Rowe, J. Théberge, R. W. J. Neufeld, P. C. Williamson, and M. Brimson. 2010. "Default Mode Network Connectivity as a Predictor of Post-Traumatic Stress Disorder Symptom Severity in Acutely Traumatized Subjects." *Acta Psychiatrica Scandinavica* 121 (1): 33–40. https://doi.org/10.1111/j.1600-0447.2009.01391.x.

Lanius, R. A., R. Bluhm, U. Lanius, and C. Pain. 2006. "A Review of Neuroimaging Studies in PTSD: Heterogeneity of Response to Symptom Provocation." *Journal of Psychiatric Research* 40 (8): 709–29. https://doi.org/10.1016/j.jpsychires.2005.07.007.

Lanius, R. A., P. C. Williamson, M. Densmore, K. Boksman, M. A. Gupta, R. W. Neufeld, J. S. Gati, and R. S. Menon. 2001. "Neural Correlates of Traumatic Memories in Posttraumatic Stress Disorder: A Functional MRI Investigation." *The American Journal of Psychiatry* 158 (11): 1920–22. https://doi.org/10.1176/appi.ajp.158.11.1920.

Lanius, Ruth A, Peter C Williamson, Kristine Boksman, Maria Densmore, Madhulika Gupta, Richard W.J Neufeld, Joseph S Gati, and Ravi S Menon. 2002. "Brain Activation during Script-Driven Imagery Induced Dissociative Responses in PTSD: A Functional Magnetic Resonance Imaging Investigation." *Biological Psychiatry* 52 (4): 305–11. https://doi.org/10.1016/S0006-3223(02)01367-7.

Lanius, Ruth A., Peter C. Williamson, Maria Densmore, Kristine Boksman, R. W. Neufeld, Joseph S. Gati, and Ravi S. Menon. 2004. "The Nature of Traumatic Memories: A 4-T FMRI Functional Connectivity Analysis." *American Journal of Psychiatry* 161 (1): 36–44. https://doi.org/10.1176/appi.ajp.161.1.36.

Lazarov, Amit, Xi Zhu, Benjamin Suarez-Jimenez, Bret R. Rutherford, and Yuval Neria. 2017. "Resting-State Functional Connectivity of Anterior and Posterior Hippocampus in Posttraumatic Stress Disorder." *Journal of Psychiatric Research* 94 (November): 15–22. https://doi.org/10.1016/j.jpsychires.2017.06.003.

LeDoux, Joseph. 2003. "The Emotional Brain, Fear, and the Amygdala." *Cellular and Molecular Neurobiology* 23 (4): 727–38. https://doi.org/10.1023/A:1025048802629.

Lewis, Catrin, Neil P. Roberts, Samuel Gibson, and Jonathan I. Bisson. 2020. "Dropout from Psychological Therapies for Post-Traumatic Stress Disorder (PTSD) in Adults: Systematic Review and Meta-Analysis." *European Journal of Psychotraumatology* 11 (1): 1709709. https://doi.org/10.1080/20008198.2019.1709709.

Liberzon, Israel, Stephan F. Taylor, Richard Amdur, Tara D. Jung, Kenneth R. Chamberlain, Satoshi Minoshima, Robert A. Koeppe, and Lorraine M. Fig. 1999. "Brain Activation in PTSD in Response to Trauma-Related Stimuli." *Biological Psychiatry* 45 (7): 817–26. https://doi.org/10.1016/S0006-3223(98)00246-7.

Liddell, Belinda J., and Laura Jobson. 2016. "The Impact of Cultural Differences in Self-Representation on the Neural Substrates of Posttraumatic Stress Disorder." *European Journal of Psychotraumatology* 7 (1): 30464. https://doi.org/10.3402/ejpt.v7.30464.

Lucassen, Paul J., Jens Pruessner, Nuno Sousa, Osborne F. X. Almeida, Anne Marie Van Dam, Grazyna Rajkowska, Dick F. Swaab, and Boldizsár Czéh. 2014. "Neuropathology of Stress." *Acta Neuropathologica* 127 (1): 109–35. https://doi.org/10.1007/s00401-013-1223-5.

Maddock, Richard J. 1999. "The Retrosplenial Cortex and Emotion: New Insights from Functional Neuroimaging of the Human Brain." *Trends in Neurosciences* 22 (7): 310–16. https://doi.org/10.1016/S0166-2236(98)01374-5.

Malivoire, Bailee L., Todd A. Girard, Ronak Patel, and Candice M. Monson. 2018. "Functional Connectivity of Hippocampal Subregions in PTSD: Relations with Symptoms." *BMC Psychiatry* 18 (1): 129. https://doi.org/10.1186/s12888-018-1716-9.

Maples-Keller, Jessica L, Seth D Norrholm, Mark Burton, Collin Reiff, Callan Coghlan, Tanja Jovanovic, Carly Yasinski, et al. 2022. "A Randomized Controlled Trial of 3,4-Methylenedioxymethamphetamine (MDMA) and Fear Extinction Retention in Healthy

Adults." *Journal of Psychopharmacology (Oxford, England)* 36 (3): 368–77.
https://doi.org/10.1177/02698811211069124.

Mei, Willem F. van der, Anna C. Barbano, Andrew Ratanatharathorn, Richard A. Bryant,
Douglas L. Delahanty, Terri A. deRoon-Cassini, Betty S. Lai, et al. 2020. "Evaluating a
Screener to Quantify PTSD Risk Using Emergency Care Information: A Proof of
Concept Study." *BMC Emergency Medicine* 20 (March): 16.
https://doi.org/10.1186/s12873-020-00308-z.

Mertens, Yoki L., Antje Manthey, Anika Sierk, Henrik Walter, and Judith K. Daniels. 2022.
"Neural Correlates of Acute Post-Traumatic Dissociation: A Functional Neuroimaging
Script-Driven Imagery Study." *BJPsych Open* 8 (4): e109.
https://doi.org/10.1192/bjo.2022.65.

Mitchell, Jennifer M., Michael Bogenschutz, Alia Lilienstein, Charlotte Harrison, Sarah
Kleiman, Kelly Parker-Guilbert, Marcela Ot'alora G., et al. 2021. "MDMA-Assisted
Therapy for Severe PTSD: A Randomized, Double-Blind, Placebo-Controlled Phase 3
Study." *Nature Medicine* 27 (6): 1025–33. https://doi.org/10.1038/s41591-021-01336-3.

Mithoefer, Michael C., Allison A. Feduccia, Lisa Jerome, Anne Mithoefer, Mark Wagner, Zach
Walsh, Scott Hamilton, Berra Yazar-Klosinski, Amy Emerson, and Rick Doblin. 2019.
"MDMA-Assisted Psychotherapy for Treatment of PTSD: Study Design and Rationale
for Phase 3 Trials Based on Pooled Analysis of Six Phase 2 Randomized Controlled
Trials." *Psychopharmacology* 236 (9): 2735–45. https://doi.org/10.1007/s00213-019-
05249-5.

Mithoefer, Michael C., Ann T. Mithoefer, Allison A. Feduccia, Lisa Jerome, Mark Wagner, Joy
Wymer, Julie Holland, et al. 2018. "3,4-Methylenedioxymethamphetamine (MDMA)-
Assisted Psychotherapy for Post-Traumatic Stress Disorder in Military Veterans,
Firefighters, and Police Officers: A Randomised, Double-Blind, Dose-Response, Phase 2
Clinical Trial." *The Lancet Psychiatry* 5 (6): 486–97. https://doi.org/10.1016/S2215-
0366(18)30135-4.

Mithoefer, Michael C., Mark T. Wagner, Ann T. Mithoefer, Lisa Jerome, and Rick Doblin. 2011.
"The Safety and Efficacy of {+/-}3,4-Methylenedioxymethamphetamine-Assisted
Psychotherapy in Subjects with Chronic, Treatment-Resistant Posttraumatic Stress
Disorder: The First Randomized Controlled Pilot Study." *Journal of
Psychopharmacology (Oxford, England)* 25 (4): 439–52.
https://doi.org/10.1177/0269881110378371.

Morris, J. S., A. Öhman, and R. J. Dolan. 1998. "Conscious and Unconscious Emotional
Learning in the Human Amygdala." *Nature* 393 (6684): 467–70.
https://doi.org/10.1038/30976.

Mott, Juliette M., Sasha Mondragon, Natalie E. Hundt, Melissa Beason-Smith, Rebecca H.
Grady, and Ellen J. Teng. 2014. "Characteristics of U.S. Veterans Who Begin and
Complete Prolonged Exposure and Cognitive Processing Therapy for PTSD." *Journal of
Traumatic Stress* 27 (3): 265–73. https://doi.org/10.1002/jts.21927.

Müller, Felix, Friederike Holze, Patrick Dolder, Laura Ley, Patrick Vizeli, Alain Soltermann,
Matthias E. Liechti, and Stefan Borgwardt. 2021. "MDMA-Induced Changes in within-
Network Connectivity Contradict the Specificity of These Alterations for the Effects of
Serotonergic Hallucinogens." *Neuropsychopharmacology* 46 (3): 545–53.
https://doi.org/10.1038/s41386-020-00906-2.

Nardou, Romain, Eastman M. Lewis, Rebecca Rothhaas, Ran Xu, Aimei Yang, Edward Boyden, and Gül Dölen. 2019. "Oxytocin-Dependent Reopening of a Social Reward Learning Critical Period with MDMA." *Nature* 569 (7754): 116–20. https://doi.org/10.1038/s41586-019-1075-9.

Patel, Ronak, R. Nathan Spreng, Lisa M. Shin, and Todd A. Girard. 2012. "Neurocircuitry Models of Posttraumatic Stress Disorder and beyond: A Meta-Analysis of Functional Neuroimaging Studies." *Neuroscience & Biobehavioral Reviews* 36 (9): 2130–42. https://doi.org/10.1016/j.neubiorev.2012.06.003.

Pitman, Roger K., Ann M. Rasmusson, Karestan C. Koenen, Lisa M. Shin, Scott P. Orr, Mark W. Gilbertson, Mohammed R. Milad, and Israel Liberzon. 2012. "Biological Studies of Post-Traumatic Stress Disorder." *Nature Reviews Neuroscience* 13 (11): 769–87. https://doi.org/10.1038/nrn3339.

Power, Jonathan D., Kelly A. Barnes, Abraham Z. Snyder, Bradley L. Schlaggar, and Steven E. Petersen. 2012. "Spurious but Systematic Correlations in Functional Connectivity MRI Networks Arise from Subject Motion." *NeuroImage* 59 (3): 2142–54. https://doi.org/10.1016/j.neuroimage.2011.10.018.

Rabinak, Christine, Mike Angstadt, Robert Welsh, Amy Kennedy, Mark Lyubkin, Brian Martis, and K. Luan Phan. 2011. "Altered Amygdala Resting-State Functional Connectivity in Post-Traumatic Stress Disorder." *Frontiers in Psychiatry* 2. https://www.frontiersin.org/articles/10.3389/fpsyt.2011.00062.

Raichle, Marcus E. 2015. "The Brain's Default Mode Network." *Annual Review of Neuroscience* 38 (1): 433–47. https://doi.org/10.1146/annurev-neuro-071013-014030.

Ridderbusch, Isabelle C., Adrian Wroblewski, Yunbo Yang, Jan Richter, Maike Hollandt, Alfons O. Hamm, Hans-Ulrich Wittchen, et al. 2021. "Neural Adaptation of Cingulate and Insular Activity during Delayed Fear Extinction: A Replicable Pattern across Assessment Sites and Repeated Measurements." *NeuroImage* 237 (August): 118157. https://doi.org/10.1016/j.neuroimage.2021.118157.

Sartory, Gudrun, Jan Cwik, Helge Knuppertz, Benjamin Schürholt, Morena Lebens, Rüdiger J. Seitz, and Ralf Schulze. 2013. "In Search of the Trauma Memory: A Meta-Analysis of Functional Neuroimaging Studies of Symptom Provocation in Posttraumatic Stress Disorder (PTSD)." *PloS One* 8 (3): e58150. https://doi.org/10.1371/journal.pone.0058150.

Sestieri, Carlo, Maurizio Corbetta, Gian Luca Romani, and Gordon L. Shulman. 2011. "Episodic Memory Retrieval, Parietal Cortex, and the Default Mode Network: Functional and Topographic Analyses." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 31 (12): 4407–20. https://doi.org/10.1523/JNEUROSCI.3335-10.2011.

Shalev, Arieh Y., Martin Gevonden, Andrew Ratanatharathorn, Eugene Laska, Willem F. van der Mei, Wei Qi, Sarah Lowe, et al. 2019. "Estimating the Risk of PTSD in Recent Trauma Survivors: Results of the International Consortium to Predict PTSD (ICPP)." *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)* 18 (1): 77–87. https://doi.org/10.1002/wps.20608.

Shou, Haochang, Zhen Yang, Theodore D. Satterthwaite, Philip A Cook, Steven E. Bruce, Russell T. Shinohara, Benjamin Rosenberg, and Yvette I. Sheline. 2017. "Cognitive Behavioral Therapy Increases Amygdala Connectivity with the Cognitive Control Network in Both MDD and PTSD." *NeuroImage: Clinical* 14 (January): 464–70. https://doi.org/10.1016/J.NICL.2017.01.030.

Smith, Stephen M., Mark Jenkinson, Mark W. Woolrich, Christian F. Beckmann, Timothy E. J. Behrens, Heidi Johansen-Berg, Peter R. Bannister, et al. 2004. "Advances in Functional and Structural MR Image Analysis and Implementation as FSL." *NeuroImage* 23 Suppl 1: S208-219. https://doi.org/10.1016/j.neuroimage.2004.07.051.

Smith, Stephen M., and Thomas E. Nichols. 2009. "Threshold-Free Cluster Enhancement: Addressing Problems of Smoothing, Threshold Dependence and Localisation in Cluster Inference." *NeuroImage* 44 (1): 83–98. https://doi.org/10.1016/j.neuroimage.2008.03.061.

Spreng, R. Nathan, and Cheryl L. Grady. 2010. "Patterns of Brain Activity Supporting Autobiographical Memory, Prospection, and Theory of Mind, and Their Relationship to the Default Mode Network." *Journal of Cognitive Neuroscience* 22 (6): 1112–23. https://doi.org/10.1162/jocn.2009.21282.

Spreng, R. Nathan, Raymond A. Mar, and Alice S. N. Kim. 2009. "The Common Neural Basis of Autobiographical Memory, Prospection, Navigation, Theory of Mind, and the Default Mode: A Quantitative Meta-Analysis." *Journal of Cognitive Neuroscience* 21 (3): 489–510. https://doi.org/10.1162/jocn.2008.21029.

Sripada, Rebecca K., Anthony P. King, Sarah N. Garfinkel, Xin Wang, Chandra S. Sripada, Robert C. Welsh, and Israel Liberzon. 2012. "Altered Resting-State Amygdala Functional Connectivity in Men with Posttraumatic Stress Disorder." *Journal of Psychiatry & Neuroscience : JPN* 37 (4): 241. https://doi.org/10.1503/JPN.110069.

Steenkamp, Maria M., Brett T. Litz, Charles W. Hoge, and Charles R. Marmar. 2015. "Psychotherapy for Military-Related PTSD: A Review of Randomized Clinical Trials." *JAMA* 314 (5): 489–500. https://doi.org/10.1001/jama.2015.8370.

Suzuki, Akinobu, Sheena A. Josselyn, Paul W. Frankland, Shoichi Masushige, Alcino J. Silva, and Satoshi Kida. 2004. "Memory Reconsolidation and Extinction Have Distinct Temporal and Biochemical Signatures." *Journal of Neuroscience* 24 (20): 4787–95. https://doi.org/10.1523/JNEUROSCI.5491-03.2004.

Svoboda, Eva, Margaret C. McKinnon, and Brian Levine. 2006. "The Functional Neuroanatomy of Autobiographical Memory: A Meta-Analysis." *Neuropsychologia* 44 (12): 2189–2208. https://doi.org/10.1016/j.neuropsychologia.2006.05.023.

Thome, Janine, Braeden A. Terpou, Margaret C. McKinnon, and Ruth A. Lanius. 2020. "The Neural Correlates of Trauma-Related Autobiographical Memory in Posttraumatic Stress Disorder: A Meta-Analysis." *Depression and Anxiety* 37 (4): 321–45. https://doi.org/10.1002/da.22977.

Thompson, M. R., P. D. Callaghan, G. E. Hunt, J. L. Cornish, and I. S. McGregor. 2007. "A Role for Oxytocin and 5-HT1A Receptors in the Prosocial Effects of 3,4 Methylenedioxymethamphetamine ('Ecstasy')." *Neuroscience* 146 (2): 509–14. https://doi.org/10.1016/j.neuroscience.2007.02.032.

Torre, Rafael de la, Magí Farré, Pere N. Roset, Neus Pizarro, Sergio Abanades, Mireia Segura, Jordi Segura, and Jordi Camí. 2004. "Human Pharmacology of MDMA: Pharmacokinetics, Metabolism, and Disposition." *Therapeutic Drug Monitoring* 26 (2): 137–44.

Vaisvaser, Sharon, Tamar Lin, Roee Admon, Ilana Podlipsky, Yona Greenman, Naftali Stern, Eyal Fruchter, et al. 2013. "Neural Traces of Stress: Cortisol Related Sustained Enhancement of Amygdala-Hippocampal Functional Connectivity." *Frontiers in Human Neuroscience* 7 (July): 313. https://doi.org/10.3389/fnhum.2013.00313.

VanElzakker, Michael B., Lindsay K. Staples-Bradley, and Lisa M. Shin. 2018. "The Neurocircuitry of Fear and PTSD." In *Sleep and Combat-Related Post Traumatic Stress Disorder*, edited by Eric Vermetten, Anne Germain, and Thomas C. Neylan, 111–25. New York, NY: Springer. https://doi.org/10.1007/978-1-4939-7148-0_10.

Veer, Ilya M., Nicole Y. L. Oei, Philip Spinhoven, Mark A. van Buchem, Bernet M. Elzinga, and Serge A. R. B. Rombouts. 2011. "Beyond Acute Social Stress: Increased Functional Connectivity between Amygdala and Cortical Midline Structures." *NeuroImage* 57 (4): 1534–41. https://doi.org/10.1016/j.neuroimage.2011.05.074.

Vizeli, Patrick, Isabelle Straumann, Urs Duthaler, Nimmy Varghese, Anne Eckert, Martin P. Paulus, Victoria Risbrough, and Matthias E. Liechti. 2022. "Effects of 3,4-Methylenedioxymethamphetamine on Conditioned Fear Extinction and Retention in a Crossover Study in Healthy Subjects." *Frontiers in Pharmacology* 13 (July): 906639. https://doi.org/10.3389/fphar.2022.906639.

Walpola, Ishan C, Timothy Nest, Leor Roseman, David Erritzoe, Amanda Feilding, David J Nutt, and Robin L Carhart-Harris. 2017. "Altered Insula Connectivity under MDMA." *Neuropsychopharmacology* 42 (11): 2152–62. https://doi.org/10.1038/npp.2017.35.

Wang, Julie B., Jessica Lin, Leah Bedrosian, Allison Coker, Ilsa Jerome, Allison Feduccia, Alia Lilienstein, et al. 2021. "Scaling Up: Multisite Open-Label Clinical Trials of MDMA-Assisted Therapy for Severe Posttraumatic Stress Disorder." *Journal of Humanistic Psychology*, June, 00221678211023663. https://doi.org/10.1177/00221678211023663.

Whitfield-Gabrieli, Susan, and Alfonso Nieto-Castanon. 2012. "Conn: A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks." *Brain Connectivity* 2 (3): 125–41. https://doi.org/10.1089/brain.2012.0073.

Winkler, Anderson M., Gerard R. Ridgway, Matthew A. Webster, Stephen M. Smith, and Thomas E. Nichols. 2014. "Permutation Inference for the General Linear Model." *NeuroImage* 92 (May): 381–97. https://doi.org/10.1016/j.neuroimage.2014.01.060.

Woodward, Neil D., and Carissa J. Cascio. 2015. "Resting-State Functional Connectivity in Psychiatric Disorders." *JAMA Psychiatry* 72 (8): 743–44. https://doi.org/10.1001/jamapsychiatry.2015.0484.

Woolrich, Mark W., Timothy E. J. Behrens, Christian F. Beckmann, Mark Jenkinson, and Stephen M. Smith. 2004. "Multilevel Linear Modelling for FMRI Group Analysis Using Bayesian Inference." *NeuroImage* 21 (4): 1732–47. https://doi.org/10.1016/j.neuroimage.2003.12.023.

Young, M. B., R. Andero, K. J. Ressler, and L. L. Howell. 2015. "3,4-Methylenedioxymethamphetamine Facilitates Fear Extinction Learning." *Translational Psychiatry* 5 (9): e634–e634. https://doi.org/10.1038/tp.2015.138.

Young, M. B., Seth D. Norrholm, Lara M. Khoury, Tanja Jovanovic, Sheila A.M. Rauch, Collin M. Reiff, Boadie W. Dunlop, Barbara O. Rothbaum, and Leonard L. Howell. 2017. "Inhibition of Serotonin Transporters Disrupts the Enhancement of Fear Memory Extinction by 3,4-Methylenedioxymethamphetamine (MDMA)." *Psychopharmacology* 234 (19): 2883–95. https://doi.org/10.1007/s00213-017-4684-8.

Zhou, Juan, Siwei Liu, Kwun Kei Ng, and Juan Wang. 2017. "Applications of Resting-State Functional Connectivity to Neurodegenerative Disease." *Neuroimaging Clinics* 27 (4): 663–83. https://doi.org/10.1016/j.nic.2017.06.007.