**Factor structure of the 19-item Comprehensive Psychopathological Rating Scale Self-**
**rating Scale for Affective Syndromes in a Swedish nationwide clinical sample of eating**
**disorder patients**

Christopher Hübel[1,2,3], Andreas Birgegård[4], Therese Johansson[5,6], Liselotte V. Petersen[3], Rasmus
Isomaa[7,8], Moritz Herle[1,9]

[1] Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and

Neuroscience, King's College London, Denmark Hill, Camberwell, London, UK

[2] UK National Institute for Health Research (NIHR) Biomedical Research Centre for Mental

Health, South London and Maudsley Hospital, London, UK

[3] National Centre for Register-based Research, Aarhus BSS Business and Social Sciences, Aarhus

University, Aarhus, Denmark

[4] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,

Sweden

[5] Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

[6] Centre for Women's Mental Health during the Reproductive Lifespan – Womher, Uppsala

University, Uppsala, Sweden

[7] The Wellbeing Services County of Ostrobothnia, Finland

[8] Faculty of Education and Welfare Studies, Åbo Akademi University, Vasa, Finland

[9] Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology &

Neuroscience, King's College London, London, UK

**Corresponding author:** Dr. Christopher Hübel, MD, PhD, (christopher.1.huebel@kcl.ac.uk),
Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology &
Neuroscience, King's College London, London, SE5 8AF, UK.

**Running title:** CPRS-S-A in eating disorders

**Key words.** Depression; compulsion; anxiety; disinterest; hypochondria; psychometrics;
questionnaire

**Abstract (250/250)**

**Objective:**

Individuals with eating disorders have anxiety and depression symptoms. To study these

comorbidities, we need high-quality self-report measures appropriate for patients and

population samples covering a wide range of symptoms. One potential tool is the 19-item self-

rated version of the Comprehensive Psychopathological Rating Scale for Affective Syndromes

(CPRS-S-A), but its factor structure has not been validated in eating disorder patients. We

investigated its factor structure and tested if the derived subscales are invariant across eating

disorders and if eating disorder patients differ on these scales.

**Method:**

The sample comprises patients registered by 45 treatment units in the nationwide Stepwise

quality assurance database for specialised eating disorder care in Sweden ($n$ = 9,509). Patients

self-reported their anxiety and depression symptoms using the CPRS-S-A. Analyses included an

exploratory factor analysis in a split sample, a confirmatory factor analysis on a hold-out

sample, and invariance testing. We examined differences in subscales across different eating

disorders.

**Results:**

Results suggested a four-factor solution with the subscales: Depression, Somatic & fear

symptoms, Disinterest, and Worry. Further multigroup confirmatory factor analysis indicated

that the factor structure of the subscales held across the different eating disorders. Patients with

anorexia nervosa binge-eating/purging subtype scored the highest and patients with

unspecified feeding and eating disorders the lowest on all subscales. Additionally, we detected

factor-specific differences amongst eating disorders.

**Discussion:**

In summary, our four-factor solution of the CPRS-S-A is suitable for patients with different

eating disorders and may help to identify differences in anxiety and depression domains

amongst eating disorders.

**Public Significance Statement:**

Patients with eating disorders often have anxiety and depression symptoms that impact

treatment and recovery. Therefore, we need questionnaires that measure these symptoms so

we can study them. More than 9,000 Swedish patients with eating disorders answered an

anxiety and depression questionnaire. We tested if the questionnaire is suitable for patients

with eating disorders. We could group questions resulting in a fine-tuned tool to study anxiety

and depression symptoms.

**1. Introduction**

**1.1 Comorbidity among affective disorders and eating disorders**

Symptoms of anxiety and depression, such as disinterest, low mood, and suicidality are

commonly found amongst patients with eating disorders from population and clinical samples

(Dolan et al., 2021; Eckert et al., 1982; Kaye et al., 2004; Martín et al., 2019; Puccio et al., 2016;

Udo & Grilo, 2019). Overall, the prevalence of comorbid depression among individuals with any

eating disorders is estimated at 75% (Godart et al., 2015). Specifically, ~60% of adolescents

with anorexia nervosa report depressive symptoms (Blinder et al., 2006; Bühren et al., 2014)

and patients with anorexia nervosa often have a comorbid clinical diagnosis of depression (Jaite

et al., 2013; Ulfvebrand et al., 2015). This high level of comorbidity with  depression is also

observed in individuals with bulimia nervosa,  in population (Hudson et al., 2007; Swanson et

al., 2011) and clinical samples (Fischer & le Grange, 2007; Kaye et al., 2004; Ulfvebrand et al.,

2015). Similarly to depression, anxiety and eating disorders also co-occur (Garcia et al., 2020;

Godart et al., 2002; Kaye et al., 2004; Kerr-Gaffney et al., 2018; Steinhausen et al., 2015;

Swinbourne et al., 2012; Ulfvebrand et al., 2015). However, prevalence estimates show

considerable heterogeneity depending on measure or assessment type (e.g. self-report versus

clinical interview) (Godart et al., 2007; Meier et al., 2015). Overall, the co-occurrence of

depressive or anxiety symptoms with eating disorder symptoms complicates treatment (Brand-

Gothelf et al., 2014; Martín et al., 2019; Thornton et al., 2011). Therefore, accurate assessment of

anxiety and depression symptoms in eating disorders may be beneficial for treatment planning.

**1.2 Measurement issues**

To study comorbidity among eating disorders, anxiety, and depression, we need high-quality

measures of symptoms to delineate differences in eating disorder presentation in clinical and

population samples. Measures like the Patient Health Questionnaire-9 (PHQ-9; Kroenke et al.,

2001) and the Generalised Anxiety Disorder Assessment (GAD-7; Spitzer et al., 2006) are widely

used in research. The PHQ-9 and GAD-7 factor structure and scores have been validated in

samples of eating disorder patients and the general population, indicating that both

questionnaires are suitable (Wisting et al., 2021). However, as they are strictly based on

diagnostic criteria, these questionnaires only cover a limited range of anxiety and depression

symptoms. Hence, in order to explore the whole spectrum of symptoms, broader assessment

tools to better understand the heterogeneity in eating disorder presentations, evading the cost-

and time-related limitations of diagnostic interviews, are urgently needed. One potential scale of

interest is the Comprehensive Psychopathological Rating Scale (Asberg & Schalling, 1979)

which consists of 65 items and was originally developed to evaluate treatment outcomes in

psychological interventions. The scale includes items covering symptoms of psychiatric disorders, such as schizophrenia but also anxiety and depression. The scale was originally developed in Sweden, and has been translated into most other European languages. The complete version of the CPRS is rarely used, but shorter subscales have been deemed to be more useful, such as the Montgomery Åsberg Depression Rating Scale (MADRS; Montgomery & Asberg, 1979) and the Self-rating Scale for Affective Syndromes (CPRS-S-A; Svanborg & Asberg, 1994). The latter is in focus here, and is designed to contain subscales for depression, anxiety, and compulsivity.

A previous analysis of the CPRS-S-A questionnaire in a subsample of the data available for our investigation showed that patients with an unspecified feeding or eating disorder reported fewer problems than patients with other eating disorders. Additionally, patients with the anorexia nervosa binge-eating/purging subtype reported more problems compared with atypical anorexia nervosa patients (Ekeroth et al., 2013). One issue of the questionnaire lies in the construction of its three subscales. When calculating the subscales, it is advised to include the same item in several subscales. Therefore, the subscales are highly correlated. In our sample, these correlations ranged from 0.78 to 0.86, and are likely to be inflated, potentially rendering the original subscales unreliable. Therefore, in this study, we explored and confirmed the factor structure of the Self-rating Scale for Affective Syndromes (CPRS-S-A); a short form of the CPRS (Svanborg & Asberg, 1994) in a large clinical sample of individuals with eating disorders in Sweden. Moreover, we investigated our newly derived CPRS-S-A subscales further by studying their differences across eating disorder subtypes.

## 2. Methods
### 2.1 Sample
The sample comprises inpatients and outpatients registered by 45 treatment units in the Stepwise quality assurance database for specialised eating disorder care in Sweden aged 18 years and older (Birgegård et al., 2010). Stepwise is a nationwide internet-based data collection system, which includes individuals through medical or self-referral, if intention to treat has been established, and if the individual received a formal eating disorder diagnosis. The database has been used since 2005 and our data were extracted on 23[rd] of November 2017. At data extraction, approximately 10,470 adult patients had been registered.

**2.2 Eating disorder diagnosis**

Clinicians registered patients' eating disorders diagnosis based on Diagnostic and Statistical

Manual of Mental Disorders, Fourth Edition (DSM-IV) (American Psychiatric Association, 2013)

In our analysis, we translated DSM-IV to DSM-5 eating disorders to reflect the current

understanding of eating disorders. Depending on the patient's endorsement of binge eating or

purging in either the Eating Disorder Examination questionnaire (EDE-Q) (Luce & Crowther,

1999) or the Structured Eating Disorder Interview (SEDI) (de Man Lapidoth & Birgegård, 2010),

we re-assigned DSM-5 diagnoses. We used 18.5 $kg/m^2$ as the cutoff value for underweight in

anorexia nervosa. Anorexia nervosa without weight criterion ($n = 50$) or without amenorrhea ($n$

$= 186$) that had a BMI lower than 18.5 $kg/m^2$ who endorsed any binge eating or purging were

assigned anorexia nervosa binge-eating/purging. If none endorsed, they were assigned an

anorexia nervosa restricting diagnosis ($n_{without weight criterion} = 84$ or $n_{without amenorrhea} = 144$). If their

BMI was above 18.5 $kg/m^2$, we assigned an atypical anorexia nervosa diagnosis ($n_{without weight}$

$_{criterion} = 441$ or $n_{without amenorrhea} = 402$). Eating Disorder Not Otherwise Specified (EDNOS) type 3

or bulimia nervosa without sufficient duration/frequency criteria ($n = 833$) was assigned as

bulimia nervosa diagnosis because those criteria are relaxed in DSM-5. Further, EDNOS example

4 was kept as "purging disorder". The remaining unspecified eating disorders that were not

classified into either of these categories were termed "Unspecified feeding or eating disorder"

(UFED), consisting of patients with "chewing and spitting", bulimia nervosa/binge-eating

disorder with low frequency/duration, or other residual types that did not fit any of the main

categories.

**2.3 Exclusion**

We excluded 801 duplicated entries of repeated registrations of the same individual.

Subsequently, we iteratively excluded 2 individuals with missing age, 16 not assigned a

treatment centre, 120 without a clinical eating disorder diagnosis, and 22 because they had not

answered the CPRS questionnaire. The final sample comprised 9,509 eating disorder patients.

**2.4 Ethics**

When patients were entered into the database, clinicians recorded consent for general research

use of their data and 3% declined participation. This study is approved by the Stockholm

Regional Ethics Board (Reg. no. 2009/196-31/4).

## 2.5 Comprehensive Psychopathological Rating Scale, Self-rated Version for Affective Syndromes

At registration, the patients answered 19 items of the CPRS-S-A. We present the instrument as **Supplementary Material**. The answer options are different for each question, but they are on a scale from 0 to 3, rated in 0.5-point increments. We recoded these values to 0–6. We renamed item 19, titled "Zest for life" in the MADRS-S to "Suicidal thoughts" to represent its content better.

*Figure 1.* Pairwise Pearson's correlations among the Self-rating Scale for Affective Syndromes (CPRS-S-A) items. We calculated the correlations in 9,509 participants registered in Stepwise, the Swedish clinical eating disorder database. We estimated the number of independent traits in the matrix using the Galwey method and adjusted the ⍺ threshold (⍺ = 0.003) accordingly. All correlations are statistically significant at this ⍺ threshold. Saturation represents the strength of the correlation. Positive correlations are red, negative correlations are blue.

**2.6 Exploratory factor analyses**

We calculated pairwise Pearson correlations among all items (**Figure 1**) in the full sample ($n$ =

9,509). We inspected the matrix visually for singularity, multicollinearity, and redundancy of

items (i.e., values <0.30 and >0.90). We calculated the determinant of the matrix (Dziuban &

Shirkey, 1974), the Kaiser-Meyer-Olkin (KMO) statistic (Kaiser, 1974), and performed Bartlett's

Test of Sphericity (Bartlett, 1950), to test if our data are suitable for an exploratory factor

analysis. To inform our decision on the underlying factor structure, we performed parallel

analysis (Horn, 1965), and calculated the Very Simple Structure criterion (VSS Revelle &

Rocklin, 1979), and Velicer's Minimum Average Partial (MAP) criterion (Velicer, 1976). We

performed the exploratory factor analysis on 70% ($n$ = 6,656) of the sample using the maximum

likelihood estimator in the "psych" R package (Revelle & Revelle, 2015). Given that the CPRS

items have seven answer options, we treated them as continuous. We allowed the factors to

correlate using oblimin rotation. To judge the fit of our model, we applied the criteria as

outlined in **Table 1** (Hu & Bentler, 1999). We retained the items with factor loadings of >0.30. If

multiple models showed adequate fit, we would choose the model with factors that encompass

the greatest number of items.

**Table 1.** *Criteria for a good fit (Hu & Bentler, 1999)*

| Root Mean Square Error of Approximation (RMSEA) | ≤0.05 |
|---|---|
| Tucker Lewis Index (TLI) | ≥0.95 |
| Standardised Root Mean Square Residuals (SRMR) | ≤0.05 |
| Bayesian Information Criteria (BIC) | Smaller than other models |

208
209

**2.7 Confirmatory factor analysis and factor scores**

We validated our exploratory factor analysis model with a confirmatory factor analysis (CFA) on

the remaining 30% participants using the "lavaan" R package (Rosseel, 2012). We interpreted fit

statistics (Hu & Bentler, 1999; Schreiber et al., 2006) and considered a Comparative Fit Index

(CFI) ≥0.95 as good fit. Subsequently, we computed the confirmatory factor analysis in the full

sample ($n$ = 9,509) to provide fit statistics and calculate factor scores, using the Bartlett

estimator for continuous items.

**2.8 Descriptive indices and psychometric properties**

We show responses to the individual items and distributions of the factor scores as frequency

and box plots. We also report mean and standard deviations for our generated factor scores and

report Cronbach's ⬚ (Bland & Altman, 1997; Cronbach, 1951) and McDonald's ⬚ (Hayes &

Coutts, 2020) as measures of internal consistency.

**2.9 Multigroup confirmatory factor analysis**

We performed a multigroup confirmatory factor analysis (MGCFA) to test if the questionnaire

elicits the same responses, response patterns, and has the same underlying factor structure

across eating disorder diagnostic groups. If statistical invariance in responding is found, then we

can compare scores and subscale scores across groups. Different types of measurement

invariance exist: configural, the factor structure is similar across groups; metric, factor loadings

are similar across groups; scalar, intercepts (i.e., group means) are similar; and strict, residuals

(i.e., variances) are similar across the groups. We tested for these invariance models in a

stepwise procedure from the least restricted model to the fully restricted model. Overall,

invariance indicates that different groups are from the same population.

**2.10 Group comparisons**

We judged the distribution of the factor scores by visually inspecting qq and distribution plots

(**Supplementary Figure S1**). None of the four subscales showed a normal distribution.

Therefore, we performed non-parametric Kruskal-Wallis one-way ANOVAs. If significant,

Dunn's post-hoc tests were carried out with a Benjamini Hochberg-adjusted level of significance

for the pairwise comparisons.

**3. Results**

**3.1 Descriptives**

The patients in our sample were on average 26 years ($SD$ = 8) old and the age ranged from 18 to

70 years, with 96% of the sample being female. Of the patients, 1,363 (14%) received an

anorexia nervosa restricting, 702 (7%) anorexia nervosa binge-eating/purging, 832 (9%) an

atypical anorexia nervosa, 3,807 (40%) a bulimia nervosa, 658 (7%) a binge-eating disorder,

1,711 (18%) a purging disorder diagnosis, and 436 UFED (5%).

**3.2. Descriptives of the CPRS-S-A in Stepwise**

The CPRS-S-A showed a Cronbach's ⍺ (⍺ = 0.90) and McDonald's ⍵ (⍵ = 0.92) in our sample

(**Supplementary Table S1**). The distribution of answers to the questionnaire items is displayed

in **Figure 2** for the full sample and **Supplementary Table S2** for the discovery sample.

*Figure 2. Endorsement of the Comprehensive Psychopathological Rating Scale Self-rating*

*Scale for Affective Syndromes (CPRS-S-A) 19 items version in the Stepwise sample (n =*

*9,509). The saturation of blue indicates a higher endorsement on the specific item. We display*

*percentages. Originally, each item had seven answer options ranging from 0-3, which were later*

*recorded from 0-6 for the main analyses. The answer options differed across items, with higher*

*values indicating a stronger endorsement.*

**3.3 Suitability of the data for factor analysis**

Prior to factor analyses, the suitability of the data was investigated. None of the items showed

zero or near-zero variance (**Supplementary Table S3**). Kaiser-Meyer-Olkin measure (KMO =

0.94, **Supplementary Table S4**) and significant Bartlett test of sphericity ($p < 2.22 \times 10^{-16}$)

indicated that the data were suitable for factor analyses. Pearson's correlations ranged from

0.09 to 0.68 (**Figure 1**). The exploratory factor analysis was conducted on one random split of

the sample ($n = 6,656$; 70%). As we were primarily interested in core anxiety and depression

symptoms, we excluded the items "14. Obsessions" and "15. Compulsions" from the factor

analysis. Furthermore, they loaded strongly on one factor by themselves, representing an index

of compulsion. If these items had remained in the model, they would have lowered our power to

measure meaningful underlying factors as they would have distorted the model towards their

own factor. We, furthermore, excluded the item "11. Health concerns", because its correlation

with the other items was small (*r* = 0.09 to 0.29; **Figure 1**), rendering it unsuitable for factor

analysis. Cronbach's  remained stable after these items were dropped (**Supplementary Table**

**S5**)

*Figure 3. Exploratory factor analysis of 16 items of the Comprehensive Psychopathological*

*Rating Scale Self-rating Scale for Affective Syndromes (CPRS-S-A). The path diagram shows*

*item factor loadings and between-factor correlations for the four factors of Depression, Somatic &*

*fear symptoms, Disinterest, and Worry. Paths with a factor loading of <0.3 were omitted.*

### 3.4 Exploratory factor analysis

Very simple structure (**Supplementary Table S6**) and parallel analysis (**Supplementary Table S7**) suggested a one factor solution. However, as we are interested in different anxiety and depression symptoms, a comparison of fit statistics suggested that the five factor solution fitted the data best. However, the model contained two factors on which only one item loaded (i.e., 2. Feelings of unease & 12. Worry about minor things) and therefore the model was unsuitable. Hence, we chose the four-factor solution as our final model which explained 34% of the total variance. The factor solution had a low RMSEA (0.042, 90% CI: 0.039 0.045) and low Bayesian Information Criterion (BIC = 245; for full results, see **Table 2** and **Supplementary Tables S8-12**). As factors were considered to be correlated, factors were realigned using an oblique rotation. The factor loadings for each item, after rotation, are listed in **Figure 3**. Items 3 and 4 (Irritation and anger, and Sleep, respectively) did not load on any of the factors and are therefore not included in the confirmatory factor analysis. We labelled the four factors: F1 Depression, F2 Somatic & fear symptoms, F3 Disinterest, and F4 Worry

### *Table 2. Model fit statistics for exploratory factor analysis*

*The factor analysis was performed on 16 items of the Comprehensive Psychopathological Rating Scale Self-rating Scale for Affective Syndromes (CPRS-S-A) in the Swedish quality register for eating disorder care, Stepwise (n = 6,656).*

| Number of factors | df | RMSEA (≤ 0.06) | RMSEA 90% CI | TLI (≥ 0.95) | BIC | SRMR (≤ 0.08) | Cumulative variance | Minimum item loading |
|---|---|---|---|---|---|---|---|---|
| 1 | 104 | 0.078 | [0.076, 0.080] | 0.882 | 3351 | 0.05 | 0.38 | 16 |
| 2 | 89 | 0.062 | [0.060, 0.065] | 0.923 | 1608 | 0.03 | 0.35 | 7 |
| 3 | 75 | 0.053 | [0.051 0.056] | 0.944 | 840 | 0.03 | 0.33 | 3 |
| **4** | **62** | **0.042** | **[0.039 0.045]** | **0.965** | **245** | **0.02** | **0.34** | **2** |
| 5 | 50 | 0.035 | [0.033 0.038] | 0.975 | 29 | 0.01 | 0.34 | 1 |

df: Degrees of freedom; RMSEA: Root Mean Square Error of Approximation; TLI: Tucker-Lewis Fit Index; BIC: Bayesian Information Criterion; SRMR: Standardised Root Mean Square Residuals. The cut off for each statistic to signify good fit is listed in each header (Hu & Bentler,

310     1999). The model with the lowest BIC is preferred. Cumulative variance indicates the part of the
311     total variance explained by all items comprising the factors.

**3.5 Confirmatory factor analysis**

We conducted the confirmatory factor analysis in the remaining 30% of the sample ($n$ = 2,853;
**Supplementary Table S13**). Results confirmed that the four-factor model was a good fit. The
RMSEA (0.060, 90% CI: 0.056, 0.064), the CFI (0.952), and the SRMR (0.032) indicated good fit
(Hu & Bentler, 1999). The TLI (0.939) was slightly above the threshold for good fit. We also ran
the confirmatory factor analysis in the full sample ($n$ = 9,509) which yielded the following fit
statistics: CFI = 0.953, TLI = 0.940, RMSEA = 0.060 [90% CI, 0.058, 0.062], SRMR = 0.030). We
show the resulting factor scores and their distribution in **Figure 4** and **Supplementary Figure**
**S1.**

*Figure 4. Distribution of factor scores across eating disorders.*

*Boxplots represent median and interquartile range in the whole Stepwise sample (n = 9,509).*

*Note. AN, anorexia nervosa, UFED, unspecified feeding or eating disorder*

**3.6 Multigroup confirmatory factor analysis**

Our multigroup confirmatory factor analysis resulted in full configural and metric invariance,

indicating that the factor structure and the factor loadings are comparable across eating

disorders (**Supplementary Table S14**). Furthermore, the questionnaire showed partial scalar

invariance when freeing up the intercepts of item five and eight, meaning that the means were

similar across groups apart from item five (less appetite) and eight (less motivation).

**Table 3. Median differences and results from Dunn's post-hoc tests**

*We performed Kruskal-Wallis one-way Analysis of Variance (ANOVA) and Dunn's post-hoc tests for pairwise comparisons and judged significance by adjusting the alpha level using the Benjamini Hochberg approach.*

| Factor | ANR vs. ANBP | ANR vs. AAN | ANR vs. BN | ANR vs. BED | ANR vs. PUR | ANR vs. UFED | ANBP vs. AAN | ANBP vs. BN | ANBP vs. BED | ANBP vs. PUR | ANBP vs. UFED | AAN vs. BN | AAN vs. BED | AAN vs. PUR | AAN vs. UFED | BN vs. BED | BN vs. PUR | BN vs. UFED | BED vs. PUR | BED vs. UFED | PUR vs. UFED |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| F1 | 0.6**** | 0.0 | 0.1 | 0.0 | 0.0 | -0.6**** | -0.6**** | -0.5**** | -0.6**** | -0.6**** | -1.2**** | 0.1 | 0.0 | 0.0 | -0.6**** | -0.1 | -0.1 | -0.7**** | 0.0 | -0.6**** | -0.6**** |
| F2 | 0.3**** | 0.0 | -0.1** | -0.1 | 0.1 | -0.4**** | -0.3**** | -0.4**** | -0.4**** | -0.2**** | -0.7**** | -0.1*** | -0.1** | -0.1** | -0.4**** | 0.0 | 0.2**** | -0.3**** | 0.2** | -0.3*** | -0.5**** |
| F3 | 0.3**** | -0.1 | 0.1* | -0.1 | 0.0 | -0.7**** | -0.4**** | -0.2**** | -0.4**** | -0.3**** | -1.0**** | 0.2 | 0.0 | 0.0 | -0.6**** | -0.2** | -0.1** | -0.8**** | 0.1 | -0.6**** | -0.7**** |
| F4 | 0.4**** | 0.0 | 0.0 | -0.3* | 0.0 | -0.8**** | -0.4**** | -0.4**** | -0.7**** | -0.4**** | -1.2**** | 0.0 | -0.3* | -0.3* | -0.8**** | -0.3** | 0.0 | -0.8**** | 0.3** | -0.5**** | -0.8**** |

**Note.** F1 Depression, F2 Somatic & fear symptoms, F3 Disinterest, and F4 Worry, **** < 1 x 10$^{-4}$, *** < 0.001, ** < 0.01, * < 0.05

ANR, anorexia nervosa restricting; ANBP, anorexia nervosa binge-eating/purging; AAN, atypical anorexia nervosa, BN, bulimia nervosa; BED, binge-eating disorder, PUR, purging disorder patients, UFED, unspecified feeding or eating disorder.

### 3.7 Factor scores

We calculated factor scores for each individual based on the final model (**Figure 4** & **Supplementary Table S15**). We compared the factor scores using Kruskal–Wallis one-way analysis of variance, and Dunn's post hoc test (**Table 3**). Overall, individuals with anorexia nervosa binge-eating/purging scored higher on all four subscales than all other eating disorders, including the restricting subtype of anorexia nervosa. However, there was no statistically significant difference on any of the four scores between the restricting subtype and atypical anorexia nervosa or purging disorder. Individuals with UFED scored lower on all four scales than all other eating disorders.

On factor 2 Somatic & fear symptoms, patients with anorexia nervosa or purging disorder scored higher than individuals with either bulimia nervosa or binge-eating disorder. Patients with anorexia nervosa restricting subtype or atypical anorexia nervosa reported depressive symptoms on the same median level as patients with either bulimia nervosa, binge-eating disorder, or purging disorder. However, compared with anorexia nervosa binge-eating/purging, all other eating disorders reported fewer depressive symptoms. On factor 3 Disinterest, the results showed a mixed picture: patients with either anorexia nervosa binge-eating/purging subtype or bulimia nervosa reported more disinterest than the other eating disorders.

### 4. Discussion
### 4.1 Summary
**Summary of findings.** Using a factor analytic approach, we propose and confirm a four factor structure of the Self-rating Scale for Affective Syndromes (CPRS-S-A) questionnaire in the world's largest clinical sample of individuals with eating disorders. Based on questionnaire responses of these Swedish patients, we propose four latent factors capturing specific aspects of depression and anxiety symptoms: 1. Depression (4 items), 2. Somatic & fear symptoms (5 items), 3. Disinterest (3 items) and 4. Worry (2 items).
**Reduced scale.** Our psychometric analysis suggests items can be removed due to their low correlations with other items. Our proposed reduced scale has a total of 14 questionnaire items and is hence shorter than the original CPRS-S-A with 19 items. First, we could drop item "11. Health concerns" as it barely correlated with other items on the scale. Second, we dropped two items regarding compulsiveness (i.e., item 14 & 15) as these were not deemed core to depression and anxiety, were highly correlated with each other, and hence would have distorted the factor analysis. Further, two items (i.e., 4. Sleep and 3. Irritation and anger) did not sufficiently load onto any of our four factors and were hence dropped.

**4.2 Context of existing literature**

**Difference to original scale.** Our factor structure differs substantially from the original CPRS-S-A (Svanborg & Asberg, 1994). In contrast to the original structure (**Supplementary Table 1**), our new structure splits traditional depression symptoms into separate factors: depression and disinterest. Depression mostly included indicator items of low mood, pessimism, and lack of enjoyment, whereas Disinterest revolved around cognition, such as lack of concentration and decision making. Anxiety symptoms were also split into two factors. First, the Somatic & fear symptoms factor grouped together general pain, bodily discomfort, physical panic attack symptoms, such as heart palpitations and dizziness, with phobias which can present with somatic symptoms. The factor also included an item probing differences in appetite. This item may be inappropriate in the context of eating disorders as changes in appetite can be a central symptom of eating disorders; however, the changes may be diametral depending on the eating disorder type. The appetite item loaded poorly on its factor and may be removed at the researcher's or clinician's discretion. Second, the Worry factor groups General worry about minor things and Feeling of unease together. This differs from the original CPRS-S-A that combined worry symptoms with the somatic and fear-based symptoms.

Overall, our analyses suggest a substantially different factor structure compared with the original structure. Our results highlight that we cannot readily apply questionnaires developed for one patient group (here, patients with depression and anxiety) to patients with different disorders as the factor structure may not hold in the new context.

**General differences amongst eating disorders.** We also explored differences in the new subscales amongst eating disorders. On the one hand, comparisons suggest that patients with anorexia nervosa binge-eating/purging score higher on all four subscales, consistent with the previous report based on a subsample of our analysis (Ulfvebrand et al., 2015), on the other hand, UFED patients had the lowest scores across all four subscales in line with their subsyndromal expression of eating disorders.

**Specific differences.** In addition to these overarching differences, we detected differences for specific factors. On factor 2 Somatic & fear symptoms, patients with anorexia nervosa or purging disorder scored higher than individuals with either bulimia nervosa or binge-eating disorder. These differences may indicate that the somatic complications seen in anorexia nervosa (Westmoreland et al., 2016) and purging disorder may be captured by items on this factor summarising somatic fear symptoms. Furthermore, patients with anorexia nervosa and purging disorders may perceive these somatic and fear symptoms more strongly than patients with

bulimia nervosa or binge-eating disorder. Fear has been proposed as a fundamental mechanism in the development of anorexia nervosa (Murray et al., 2018).

Depression and anxiety are risk factors for eating disorders (Meier et al., 2015; Steinhausen et al., 2015), but certain symptoms of anxiety or depression may represent somatic or psychiatric complications or sequelae of the eating disorder itself. In some cases, depressive and anxiety symptoms may be independent of the eating disorder.

### 4.3 Limitations

Our study is biased due to the following limitations. The sample consisted predominantly of women which limits the ability to identify sex differences. Eating disorders are more commonly diagnosed among women, however, men with eating disorders are consistently underrepresented in eating disorder research. This may be due to a lack of awareness and understanding for these disorders among the wider community and clinicians or may represent an underlying sex difference. Our sample included Swedish treatment seeking patients of mostly white European ancestry limiting the generalisability of our findings. Furthermore, patients in healthcare registers often represent a more severe subpopulation of individuals with eating disorders. Hence, the factor structure and our observed differences amongst eating disorders may not replicate across other ancestry or cultural groups or of individuals with a less severe presentation. Our analyses were cross-sectional.

### 4.4 Future directions

To address a few of our limitations, future studies should confirm our newly detected factor structure in community samples, samples with other psychiatric disorders, and include a healthy comparison group. Optimally, researchers would collect repeated measures of the CPRS-S-A that would further our understanding of how these constructs develop over time and how levels of depression, disinterest, fear, and worry may change with treatment. Future studies should investigate the convergent validity of the questionnaire and could investigate clinical cutoffs to measure comorbid depressive and anxiety disorders.

### 4.5 Conclusions

In summary, our four factor solution of the CPRS-S-A is suitable for adult patients with different eating disorders and may help to identify differences in anxiety and depression domains. An easily administered, reliable self-report measure for the most common forms of comorbidity in eating disorders is clinically important. The CPRS-S-A may aid the clinician in case formulation and treatment planning. It may also be relevant for the patient's own understanding of their

situation. A discussion between patient and clinician, facilitated by the individual CPRS-S-A results, of depression and anxiety symptoms in relation to eating disorder symptoms may also improve therapeutic alliance and thus treatment outcome.

**Data availability.**

Due to regional legal regulations, the data from the quality register cannot be shared.

**Conflict of Interest Statement.**

The authors report no conflict of interest.

**Acknowledgement**

Liselotte V. Petersen and Christopher Hübel acknowledge funding by Lundbeckfonden (R276-2018-4581). Moritz Herle is funded by a fellowship from the Medical Research Council UK (MR/T027843/1).

**Authors contribution statement**

Christopher Hübel: Conceptualisation, Software, Visualisation, Formal analyses, Writing - Original Draft, Writing - Review & Editing,. Andreas Birgegård: Data Curation, Supervision, Writing - Review & Editing , Funding acquisition, Resources, Therese Johansson: Software, Writing - Review & Editing. Rasmus Isomaa: Supervision, Writing - Review & Editing, Liselotte V. Petersen: Supervision, Writing - Review & Editing , Funding acquisition, Moritz Herle: Supervision, Conceptualisation, Writing - Original Draft, Writing - Review & Editing, Project administration

**References**

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). American Psychiatric Association.

Asberg, M., & Schalling, D. (1979). Construction of a new psychiatric rating instrument, the Comprehensive Psychopathological Rating Scale (CPRS). *Progress in Neuro-Psychopharmacology*, 3(4), 405–412.

Bartlett, M. S. (1950). Tests of significance in factor analysis. *The British Journal of Mathematical and Statistical Psychology*, 3(2), 77–85.

Birgegård, A., Björck, C., & Clinton, D. (2010). Quality assurance of specialised treatment of eating disorders using large-scale Internet-based collection systems: methods, results and lessons learned from designing the Stepwise database. *European Eating Disorders Review: The Journal of the Eating Disorders Association*, *18*(4), 251–259.

Bland, J. M., & Altman, D. G. (1997). Cronbach's alpha. *BMJ*, *314*(7080), 572.

Blinder, B. J., Cumella, E. J., & Sanathara, V. A. (2006). Psychiatric comorbidities of female inpatients with eating disorders. *Psychosomatic Medicine*, *68*(3), 454–462.

Brand-Gothelf, A., Leor, S., Apter, A., & Fennig, S. (2014). The impact of comorbid depressive and anxiety disorders on severity of anorexia nervosa in adolescent girls. *The Journal of Nervous and Mental Disease*, *202*(10), 759–762.

Bühren, K., Schwarte, R., Fluck, F., Timmesfeld, N., Krei, M., Egberts, K., Pfeiffer, E., Fleischhaker, C., Wewetzer, C., & Herpertz-Dahlmann, B. (2014). Comorbid Psychiatric Disorders in Female Adolescents with First-Onset Anorexia Nervosa: Comorbidity in Adolescent Anorexia Nervosa. *European Eating Disorders Review: The Journal of the Eating Disorders Association*, *22*(1), 39–44.

Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, *16*(3), 297–334.

de Man Lapidoth, J., & Birgegård, A. (2010). Validation of the structured eating disorder

interview (SEDI) against the eating disorder examination (EDE). *Stockholm: Karolinska Institutet*.

Dolan, S. C., Khindri, R., Franko, D. L., Thomas, J. J., Reilly, E. E., & Eddy, K. T. (2021). Anhedonia in eating disorders: A meta-analysis and systematic review. *The International Journal of Eating Disorders*. https://doi.org/10.1002/eat.23645

Dziuban, C. D., & Shirkey, E. C. (1974). When is a correlation matrix appropriate for factor analysis? Some decision rules. *Psychological Bulletin*, *81*(6), 358–361.

Eckert, E. D., Goldberg, S. C., Halmi, K. A., Casper, R. C., & Davis, J. M. (1982). Depression in anorexia nervosa. *Psychological Medicine*, *12*(1), 115–122.

Ekeroth, K., Clinton, D., Norring, C., & Birgegård, A. (2013). Clinical characteristics and distinctiveness of DSM-5 eating disorder diagnoses: findings from a large naturalistic clinical database. *Journal of Eating Disorders*, 1, 31.

Fischer, S., & le Grange, D. (2007). Comorbidity and high-risk behaviors in treatment-seeking adolescents with bulimia nervosa. *The International Journal of Eating Disorders*, *40*(8), 751–753.

Garcia, S. C., Mikhail, M. E., Keel, P. K., Burt, S. A., Neale, M. C., Boker, S., & Klump, K. L. (2020). Increased rates of eating disorders and their symptoms in women with major depressive disorder and anxiety disorders. *The International Journal of Eating Disorders*, *53*(11), 1844–1854.

Godart, N. T., Flament, M. F., Perdereau, F., & Jeammet, P. (2002). Comorbidity between eating disorders and anxiety disorders: a review. *The International Journal of Eating Disorders*, *32*(3), 253–270.

Godart, N. T., Perdereau, F., Rein, Z., Berthoz, S., Wallier, J., Jeammet, P., & Flament, M. F. (2007). Comorbidity studies of eating disorders and mood disorders. Critical review of the literature. *Journal of Affective Disorders*, *97*(1-3), 37–49.

Godart, N. T., Radon, L., Curt, F., Duclos, J., Perdereau, F., Lang, F., Venisse, J. L., Halfon, O.,

Bizouard, P., Loas, G., Corcos, M., Jeammet, P., & Flament, M. F. (2015). Mood disorders in

    eating disorder patients: Prevalence and chronology of ONSET. *Journal of Affective*

    *Disorders, 185*, 115–122.

Hayes, A. F., & Coutts, J. J. (2020). Use omega rather than cronbach's alpha for estimating

    reliability. But.... *Communication Methods and Measures, 14*(1), 1–24.

Horn, J. L. (1965). A RATIONALE AND TEST FOR THE NUMBER OF FACTORS IN FACTOR

    ANALYSIS. *Psychometrika, 30*, 179–185.

Hudson, J. I., Hiripi, E., Pope, H. G., Jr, & Kessler, R. C. (2007). The prevalence and correlates of

    eating disorders in the National Comorbidity Survey Replication. *Biological Psychiatry,*

    *61*(3), 348–358.

Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:

    Conventional criteria versus new alternatives. *Structural Equation Modeling: A*

    *Multidisciplinary Journal, 6*(1), 1–55.

Jaite, C., Hoffmann, F., Glaeske, G., & Bachmann, C. J. (2013). Prevalence, comorbidities and

    outpatient treatment of anorexia and bulimia nervosa in German children and adolescents.

    *Eating and Weight Disorders: EWD, 18*(2), 157–165.

Kaiser, M. O. (1974). Kaiser-Meyer-Olkin measure for identity correlation matrix. *Journal of the*

    *Royal Statistical Society, 52*(1), 296–298.

Kaye, W. H., Bulik, C. M., Thornton, L., Barbarich, N., Masters, K., & Group, the Price Foundation

    Collaborative. (2004). Comorbidity of Anxiety Disorders With Anorexia and Bulimia

    Nervosa. *The American Journal of Psychiatry, 161*(12), 2215–2221.

Kerr-Gaffney, J., Harrison, A., & Tchanturia, K. (2018). Social anxiety in the eating disorders: a

    systematic review and meta-analysis. *Psychological Medicine, 48*(15), 2477–2491.

Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: validity of a brief depression

    severity measure. *Journal of General Internal Medicine, 16*(9), 606–613.

Luce, K. H., & Crowther, J. H. (1999). The reliability of the Eating Disorder Examination-Self-

    Report Questionnaire Version (EDE-Q). *The International Journal of Eating Disorders, 25*(3),

349–351.

Martín, J., Arostegui, I., Loroño, A., Padierna, A., Najera-Zuloaga, J., & Quintana, J. M. (2019). Anxiety and depressive symptoms are related to core symptoms, general health outcome, and medical comorbidities in eating disorders. *European Eating Disorders Review: The Journal of the Eating Disorders Association.* https://doi.org/10.1002/erv.2677

Meier, S. M., Bulik, C. M., Thornton, L. M., Mattheisen, M., Mortensen, P. B., & Petersen, L. (2015). Diagnosed Anxiety Disorders and the Risk of Subsequent Anorexia Nervosa: A Danish Population Register Study. *European Eating Disorders Review: The Journal of the Eating Disorders Association*, *23*(6), 524–530.

Montgomery, S. A., & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *The British Journal of Psychiatry: The Journal of Mental Science*, *134*, 382–389.

Murray, S. B., Strober, M., Craske, M. G., Griffiths, S., Levinson, C. A., & Strigo, I. A. (2018). Fear as a translational mechanism in the psychopathology of anorexia nervosa. *Neuroscience and Biobehavioral Reviews*, *95*, 383–395.

Puccio, F., Fuller-Tyszkiewicz, M., Ong, D., & Krug, I. (2016). A systematic review and meta-analysis on the longitudinal relationship between eating pathology and depression. *The International Journal of Eating Disorders*, *49*(5), 439–454.

Revelle, W., & Revelle, M. W. (2015). Package "psych." *The Comprehensive R Archive Network*, *337*, 338.

Revelle, W., & Rocklin, T. (1979). Very Simple Structure: An Alternative Procedure For Estimating The Optimal Number Of Interpretable Factors. *Multivariate Behavioral Research*, *14*(4), 403–414.

Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling and more. Version 0.5--12 (BETA). *Journal of Statistical Software*, *48*(2), 1–36.

Schreiber, J. B., Nora, A., Stage, F. K., Barlow, E. A., & King, J. (2006). Reporting Structural Equation Modeling and Confirmatory Factor Analysis Results: A Review. *The Journal of Educational Research*, *99*(6), 323–338.

Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing

generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, *166*(10), 1092–

1097.

Steinhausen, H.-C., Jakobsen, H., Helenius, D., Munk-Jørgensen, P., & Strober, M. (2015). A nation-

wide study of the family aggregation and risk factors in anorexia nervosa over three

generations. *The International Journal of Eating Disorders*, *48*(1), 1–8.

Svanborg, P., & Asberg, M. (1994). A new self-rating scale for depression and anxiety states

based on the Comprehensive Psychopathological Rating Scale. *Acta Psychiatrica

Scandinavica*, *89*(1), 21–28.

Swanson, S. A., Crow, S. J., Le Grange, D., Swendsen, J., & Merikangas, K. R. (2011). Prevalence

and correlates of eating disorders in adolescents. Results from the national comorbidity

survey replication adolescent supplement. *Archives of General Psychiatry*, *68*(7), 714–723.

Swinbourne, J., Hunt, C., Abbott, M., Russell, J., St Clare, T., & Touyz, S. (2012). The comorbidity

between eating disorders and anxiety disorders: prevalence in an eating disorder sample

and anxiety disorder sample. *The Australian and New Zealand Journal of Psychiatry*, *46*(2),

118–131.

Thornton, L. M., Dellava, J. E., Root, T. L., Lichtenstein, P., & Bulik, C. M. (2011). Anorexia nervosa

and generalized anxiety disorder: further explorations of the relation between anxiety and

body mass index. *Journal of Anxiety Disorders*, *25*(5), 727–730.

Udo, T., & Grilo, C. M. (2019). Psychiatric and medical correlates of DSM-5 eating disorders in a

nationally representative sample of adults in the United States. *The International Journal of

Eating Disorders*, *52*(1), 42–50.

Ulfvebrand, S., Birgegård, A., Norring, C., Högdahl, L., & von Hausswolff-Juhlin, Y. (2015).

Psychiatric comorbidity in women and men with eating disorders results from a large

clinical database. *Psychiatry Research*, *230*(2), 294–299.

Velicer, W. F. (1976). Determining the number of components from the matrix of partial

correlations. *Psychometrika*, *41*(3), 321–327.

Westmoreland, P., Krantz, M. J., & Mehler, P. S. (2016). Medical Complications of Anorexia

    Nervosa and Bulimia. *The American Journal of Medicine*, *129*(1), 30–37.

Wisting, L., Johnson, S. U., Bulik, C. M., Andreassen, O. A., Rø, Ø., & Bang, L. (2021). Psychometric

    properties of the Norwegian version of the Patient Health Questionnaire-9 (PHQ-9) in a

    large female sample of adults with and without eating disorders. *BMC Psychiatry*, *21*(1), 6.