The Cultural Evolution of Vaccine Hesitancy: Modeling the Interaction between Beliefs and

Behaviors

Kerri-Ann Anderson and Nicole Creanza

Department of Biological Sciences and Evolutionary Studies Initiative, Vanderbilt University,

Nashville, TN 37212

Abstract:

In the last decade, despite the proven efficacy of vaccines, the developed world has seen a

resurgence of vaccine-preventable diseases (VPDs) such as measles, pertussis, and polio. Vaccine

hesitancy, an individual behavior influenced by historical, political, and socio-cultural factors, is

believed to be a primary factor responsible for decreasing vaccine coverage, thereby increasing

the risk and occurrence of VPD outbreaks. Society, culture, and individual motivations affect

human decisions regarding health behavior and preventative care, and health perceptions and

health-related behaviors can change at the population level as cultures evolve. In recent years,

mathematical models of disease dynamics have begun to incorporate aspects of human behavior,

however they do not address how evolving cultures influence these health behaviors. Here, using

a mathematical modeling framework, we explore the effects of cultural evolution on vaccine

hesitancy and vaccination behavior. With this model, we shed light on the facets of cultural

evolution (vertical and oblique transmission, homophily, etc.) that promote the spread of vaccine

hesitancy, ultimately affecting levels of vaccination coverage and VPD outbreak risk in a

population. In addition, we present our model as a generalizable framework for exploring cultural

evolution when beliefs influence, but do not strictly dictate, human behaviors. We show vaccine

confidence and vaccine-conferred benefits can be driving forces of vaccine coverage, and we

demonstrate that an assortative preference among vaccine-hesitant individuals can lead to

increased vaccine hesitancy and lower vaccine coverage. Further, we show that vaccine mandates

can lead to a phenomenon in which high vaccine hesitancy co-occurs with high vaccination

coverage, and that high vaccine confidence can be maintained even in areas where access to

vaccines is limited.

Introduction:

In conjunction with genetics, variations in human behaviors such as diet, hand-washing, and vaccination result in differential susceptibility to a range of diseases. Improvements in sanitation and healthcare practices have reduced the mortality and morbidity of many infectious diseases [1]. For example, the implementation of childhood vaccination policies has led to the eradication of smallpox and the elimination of poliomyelitis (polio) in the United States [2–4], and the high efficacy of the measles vaccine, combined with wide vaccine acceptance in developed countries, had resulted in measles previously being targeted for elimination by 2020 [5]. Recent infectious disease trends, however, have illustrated how easily efforts to eradicate diseases can be undermined: over the past decade, there has been a resurgence of vaccine-preventable diseases (VPDs) in developed countries despite the safety and efficacy of vaccines and high overall childhood vaccination rates [6–9]. Outbreaks of VPDs such as measles, pertussis, and polio have been linked to pockets of under-vaccinated or unvaccinated individuals, which often go relatively unnoticed by the public due to high national estimates of vaccination coverage and the assumption of associated herd immunity [8].

Human decisions regarding health-related behaviors, including preventative care, are influenced by both internal and external motivations. The term "local vaccination cultures" describes the shared beliefs among individuals within a community or region about disease etiology, prevention, and treatment, as well as the experiences with health services and vaccination settings that influence individual vaccine decisions [10]. In areas where there exists a pro-vaccination culture, there may nonetheless be individuals who are vaccine hesitant (as defined by [11]), delaying or refusing to use vaccines despite their availability. Vaccine hesitancy is a complex and context-specific individual behavior influenced by factors that shape vaccination culture such as complacency (the belief that vaccination is unnecessary when the perceived risk of VPDs is low), convenience (the accessibility and affordability of vaccines), and confidence (the level of trust in the efficacy and safety of the vaccine, and in the healthcare system) [9,11]. Vaccine hesitancy was named one of the World Health Organization's ten threats to global health in 2019 [12], as it is believed to be responsible for decreasing vaccination coverage and thus increasing the risk of vaccine-preventable disease outbreaks worldwide.

Vaccine hesitancy and vaccination opposition are, however, not recent phenomena: since the creation of vaccines in the 18th century, public opposition to vaccinations has been based in theology, politics, law, and general skepticism [13–16]. Two persistent themes of vaccine

hesitancy include the idea that vaccinations are more harmful than the diseases they intend to

prevent (usually occurring at the introduction of new vaccines), and the idea that vaccines may

not be necessary during a decline in disease (usually after compulsory vaccination mandates)

[13]. With the introduction of compulsory vaccination in the 19th century came the precursors of

contemporary vaccination exemptions [17,18]. The Anti-Vaccination League, founded in mid-19th-

century London, argued that compulsory vaccination invaded the people's liberties [19]. A 1969

review of mandatory vaccination in the United States revealed a similar objection—"infringement

on personal liberty"—along with an aversion to government intrusion on religious beliefs and a

general distrust of medical science [20]. Thus, since the advent of vaccinations, cultural context

has interacted with vaccine-related beliefs and in turn influenced vaccination behaviors,

ultimately affecting population-level immunity and public health.

Due to developments in media and communication, including widespread engagement with

social media platforms, all types of information are more accessible and faster spreading than

ever before. Unfortunately, negative information (such as anecdotes about adverse drug

reactions), misinformation (such as the discredited link between vaccines and autism), and

maladaptive practices (such as delayed vaccination schedules [21,22]) can spread more quickly

through these digital channels than by traditional word-of-mouth [23], resulting in the increased

adoption of harmful beliefs and practices. In particular, a small but very vocal community,

sometimes dubbed "Anti-Vaxxers," stoke the fears and emotions that contribute to vaccine

hesitancy, and their reach is aided by social media [9,24]. Anti-vaccine sentiments are still on the

rise despite numerous studies countering, discrediting, and debunking the spurious connection

between vaccines and autism [25] and other anti-vaccination arguments [26]. Additionally,

individuals are typically more trusting of those with whom they have a personal connection and

more skeptical of health providers and scientists [27], therefore anti-vaccine attitudes from one's

social network are more likely to be internalized than pro-vaccine information from healthcare

professionals.

Health-related behaviors are also affected by homophily—the tendency of individuals to

choose social contacts and mates who are similar to themselves [28–31]. For example, individuals

with online social contacts who share similar demographic characteristics had significantly higher

adoption of new health-related behaviors introduced in that online environment [32]. In addition,

network-based simulations suggest that individuals with similar vaccine-hesitant opinions form

groups that are more susceptible to vaccine-preventable diseases, impeding the attainment of

herd immunity and substantially increasing the likelihood of disease outbreak in these clusters

[33]. Some of the predictions of these simulations have been observed in the real world: for

example, a number of recent VPD outbreaks have been linked to tight-knit cultural groups, such

as the Ohio Amish community, which has specific teachings on health practices [34,35], and the

Minnesota Somali immigrant community, which was vulnerable to anti-vaccination rhetoric

[36,37]. If individuals have a tendency toward homophily on the basis of vaccine-related beliefs,

preferentially partnering with others who have similar beliefs and mindsets, children might be

increasingly likely to have two vaccine-hesitant or two vaccine-confident parents and to inherit

their beliefs. When vaccine-related beliefs are clustered on a social network, homophily could

further expand and reinforce these clusters as vaccine-hesitant individuals seek out interactions

with one another, exacerbating the risk of disease outbreak. Thus, a particular health culture—

collective beliefs and behaviors regarding health—could be seen as a cultural "niche" in the sense

that it influences an individual's beliefs, behaviors, and social network connections.

Niche construction is a process in which organisms modify their local environment, thus

altering selection pressures on themselves and the other organisms that share that environment

[38,39]. In *cultural* niche construction, humans modify their cultural environments—for example,

their beliefs, behaviors, preferences, and social contacts—in ways that subsequently alter

evolutionary pressures on themselves and/or their culture [39]. Within various human-

constructed cultural niches, for example in agricultural niches, there exist niche-specific

differential effects on health and health-related behaviors (for example, the link between dairy

farming and lactose tolerance in adulthood [40]. Mathematical models of niche construction have

been used to help understand biological evolution, and this type of model has been recently

expanded to explain the evolution of cultural behaviors, with applications to religion, fertility, and

the evolution of large-scale human conflict [29,30,39,41–43]. The cultural niche framing of health

cultures allows us to apply these modeling frameworks to the evolution of human disease

response. Modeling the belief-behavior interactions underlying vaccination coverage allows us to

better understand how vaccination cultures are formed and how they can be transformed to

promote public health.

Compartmental models, such as the Susceptible-Infected-Recovered (SIR) model and its

various extensions, are the most established set of epidemiological models used to predict the

spread of infectious disease. Even though more epidemiological models have begun to

incorporate aspects of human behavior (e.g [44,45]), these models do not typically incorporate

the effects of population beliefs and changing cultural landscapes on disease transmission. For

example, they do not consider social preferences or early-life influences on adult behaviors, and

how these affect the health beliefs that drive specific behaviors. SIR models with a vaccination

component (e.g. [46,47]) demonstrate that the introduction of vaccination decreases the

susceptible and infected populations proportional to vaccination rate [46]. Though these models

are very useful in determining the intensity of intervention needed to address an epidemic [46],

they do not explain fluctuations in vaccination rates or lower-than-expected rates of adoption

based on cultural factors. In wealthier countries, belief systems act as the main barrier to

vaccination, as opposed to lack of vaccine access [33,48]. The perception of vaccine importance

and disease risk is a driving factor in vaccine behavior, even for vaccinated individuals [27,49–53].

The rates of non-medical exemption from vaccines (exemption on the basis of religious,

philosophical, and personal beliefs), have been increasing in the United States, with some

believing the "negative consequences of exemption are not sufficient to justify violating parental

autonomy" [54,55]. Thus, understanding and incorporating the underlying health cultures and

their evolution, including the interplay between beliefs and behaviors, will allow us to build more

comprehensive and representative models of vaccinations dynamics and better support public

health efforts.

In this study, we model the development of vaccine hesitancy through a cultural evolution

framework, incorporating the transmission of vaccine culture both from parents and from the

community. We shed light on the situations in which vaccine hesitancy is most likely to spread

and potentially lead to an increase in vaccine-preventable diseases. By modeling the evolution of

vaccination behaviors through the lens of cultural niche construction, we aim to assess the

dynamic interactions between vaccine beliefs (shaped by social interactions) and behaviors

(influenced by these beliefs). With this framework, we can also model the effects of vaccine

mandates and of vaccine accessibility on both beliefs and behaviors. In addition, we consider that

the perception of the relative risks of a disease and its preventive vaccine can fluctuate based on

the prevalence of vaccination, such that the population's vaccination coverage can influence

individual decision-making. We further incorporate homophily, here represented as the

likelihood that people with similar vaccine-related beliefs will preferentially mate with one

another, to understand how social subcultures influence behaviors.

Beliefs influence but do not dictate an individual's behaviors, and these belief-behavior

interactions are influenced by the broader cultural landscape. We propose that scientific research

can better address how to mitigate VPD outbreaks by understanding the cultural dynamics of

vaccine hesitancy. In this manuscript, we aim to take a cultural approach to understanding the

evolution of vaccine hesitancy and its interactions with vaccination coverage and vaccine-

preventable disease using a generalizable modeling framework for belief-behavior interactions.

Methods:

To model the cultural niche construction of vaccine beliefs and behaviors, we build on the

framework of [30] to assess the effects of vaccine attitudes on vaccination behaviors and on the

resulting vaccination culture. We explore how vaccination patterns evolve in a population when a

cultural trait, such as vaccine hesitancy, can influence but not perfectly predict a behavior, such

as vaccinating one's children.

Figure 1. Workflow of a single iteration of the model: The schematic shows the processes within a single
model iteration. The model is initialized with the phenotypic frequencies (V^+A^+, V^+A^-, V^-A^+, V^-A^-) in the
population. After mate selection, individuals reproduce; they then vertically transmit vaccination and
attitude traits to their offspring. Vaccination trait frequencies are further modulated by cultural selection.
Oblique transmission (cultural transmission from non-parental adults in the population) follows, in which
offspring may alter their attitude state. (Parameters, their definitions, and default values are listed in
Table 1.)

Table 1: List of parameters, their definitions, and default or initial values

Parameter	Meaning	Parameter	Meaning
V	Vaccination state (V^+ vaccinated, V^- unvaccinated)	**A**	Vaccine attitude (A^+ confident, A^- hesitant)
m_{ij}	Mating frequencies (given in **Table S1**)	α_k	Assortative mating parameter (homophily) **Default: $\alpha_1 = 0$, $\alpha_2 = 0$**
$B_{m,n}$	Probability that parental pairs vaccinate their children, which depends upon the parents' vaccination states (b_m) and vaccine attitude (c_n) (given in **Table S2**)	C_n	Probability that parental pairs transmit vaccine confidence to their children **Default: $C_0 = 0.01$, $C_1 = C_2 = 0.5$, $C_3 = 0.99$**
b_m	Probability that parental pairs support offspring vaccination given their vaccination states **Default: $b_0 = 0.01$, $b_1 = b_2 = 0.5$, $b_3 = 0.99$**	c_n	Probability that parental pairs support offspring vaccination given their vaccine attitude **Default: $c_0 = 0.01$, $c_1 = c_2 = 0.5$, $c_3 = 0.99$**
σ	Comprehensive selection coefficient for V^+, dependent on the population-wide vaccination rate (see **Figure 2**)	σ_{max}	The highest additional benefit that can be conferred by vaccination **Default: $\sigma_{max} = 0.1$**
Initial Phenotype Frequencies		$x_1(V^+A^+) = 0.81$, $x_2(V^+A^-) = 0.1$, $x_3(V^-A^+) = 0.07$, $x_4(V^-A^-) = 0.02$	

We consider two cultural traits: **V**, a vaccination trait, and **A**, a vaccine attitude trait. Each trait has two possible states, V^+ or V^- and A^+ or A^-, respectively. V^+ individuals are vaccinated and V^- individuals are unvaccinated, and A^+ individuals are vaccine confident and A^- individuals are vaccine hesitant; thus, there are four possible phenotypes: V^+A^+ (type 1: vaccinated and confident), V^+A^- (type 2: vaccinated and hesitant), V^-A^+ (type 3: unvaccinated and confident), and V^-A^- (type 4: unvaccinated and hesitant), whose population frequencies are denoted by $x_1, x_2, x_3,$ and x_4, respectively, with $\sum_{i=1}^{4} x_i = 1$.

The attitude trait (**A**) can influence the dynamics of the vaccination trait (**V**) in two ways: by affecting the likelihood that couples vaccinate their offspring, and by determining with whom each adult will preferentially pair in assortative interactions. The state of the vaccine attitude trait (**A**) informs the value of an assortative mating parameter (α_k), which measures the departure from random mating. We define a 'choosing parent,' arbitrarily assigned as the first member of

each mating pair. The choosing parent's **A** state dictates the level of assortative mating, that is,

the degree to which an individual of a given **A** state will preferentially mate with another

individual of the same state, expressed by parameters α_k where $k = \{1, 2\}$ and $0 \leq \alpha_k \leq 1$ (**Table S1**).

If the choosing parent is A^+, this individual mates preferentially with other A^+ individuals with

probability α_1, and mates randomly with probability $1-\alpha_1$, whereas if the choosing parent is A^-,

this individual mates preferentially with other A^- individuals with probability α_2, and mates

randomly with probability $1-\alpha_2$. There are sixteen possible mating pairs from the four

phenotypes described, and we use the notation $m_{i,j}$ to indicate the frequency of a mating

between a choosing parent of type i and the second parent of type j where $i, j = \{1, 2, 3, 4\}$ (**Table**

S1); in other words, $m_{1,3}$ represents the mating frequency of V^+A^+ (x_1) and V^-A^+ (x_3).

Since the two traits in question (**A** and **V**) are transmitted vertically, for each phenotype we

must specify the probability that the mating produces an offspring of that phenotype. The

vaccine confidence trait (A^+) is transmitted with probability C_n, and the vaccine hesitancy trait (A^-)

is transmitted with probability $1-C_n$ (for $n = \{0, 1, 2, 3\}$ as shown in **Tables 2** and **S2**). If $C_0 = 0$, two

A^- parents will always produce A^- offspring, and if $C_3 = 1$, two A^+ parents will always produce A^+

offspring. However, if $C_0 > 0$, two A^- parents can produce A^+ offspring at some probability, and

similarly if $C_3 < 1$, two A^+ parents can produce A^- offspring with some probability.

Transmission of vaccination (V^+ with probability $B_{m,n}$ for $m, n = \{0, 1, 2, 3\}$; **Table 1**) is more

complex, since parents' vaccine attitudes (**A**), in addition to their own vaccination states (**V**), can

influence their behavior in vaccinating their offspring via a set of "influence parameters" that

inform vaccination probabilities. The probability that each mating pair produces an offspring with

the V^+ trait (i.e. vaccinates their offspring) is a scaled product of the influence of parental

attitudes (c_n for $n = \{0, 1, 2, 3\}$) and the influence of parental vaccination states (b_m for $m = \{0, 1, 2,$

$3\}$) (**Tables 2 and S2**). For example, for mating pair $V^+A^+ \times V^+A^-$, their combined vaccination states

($V^+ \times V^+$) will influence vaccination behavior by b_3, and their combined attitude states, ($A^+ \times A^-$),

will influence vaccination behavior by c_2. Therefore, a $V^+A^+ \times V^+A^-$ mating will produce a V^+

offspring with probability $B_{3,2} = c_2 \left(\frac{1+b_3}{2} \right)$; this pair will also produce an A^+ offspring with

probability C_2 based on their combined attitude states. Thus, according to the model, this pairing

will produce a V^+A^+ offspring with probability $B_{3,2}C_2$ and a V^+A^- offspring with probability

$B_{3,2}(1-C_2)$. We note that assortative mating ($\alpha_k > 0$) will increase the frequency of matings between

individuals that share an attitude trait, with these non-random interactions in turn skewing

vaccination outcomes in line with those of same-state couples (via c_0 and c_3).

Table 2: Presence (+) and absence (−) subscript assignments. Demonstrating the trait presence (+) and absence (−) combinations associated with m, n subscripts. For example, the + × − combinations is associated with m and n subscript value 2: an $A^+ \times A^-$ pairing transmits A^+ at probability C_2. This rule applies to parameters C_n, b_m, $B_{m,n}$, c_n, as shown in **Table S2**.

Subscript Value (m, n; e.g. b_m, C_n)	Associated Pairing (e.g. V × V, A × A)
0	− × −
1	− × +
2	+ × −
3	+ × +

Transmission and influence probabilities are constant throughout a single simulation, with values ranging from 0 to 1. At default settings, the influence parameters b_m and c_n, and the transmission parameter C_n would take the following values: C_0, b_0, $c_0 = 0.01$; C_1, C_2, b_1, b_2, c_1, $c_2 = 0.5$; and C_3, b_3, $c_3 = 0.99$. In our model, parental vaccine beliefs (c_n) have a greater influence than their own vaccination status (b_n) on their likelihood of vaccinating their offspring, so offspring vaccination is guaranteed at some probability only if $c_n > 0$.

The cultural selection pressure on vaccination is given by the parameter σ, such that the frequency of the V^+A^+ and V^+A^- phenotypes are multiplied by 1+σ after vertical cultural transmission has occurred. At the end of each timestep, the frequency of each phenotype is divided by the sum of all four frequencies, ensuring that the frequencies sum to 1. This cultural selection coefficient is implemented in the same way as a selection coefficient in a population-genetic model, but unlike the latter, it is structured to encompass both biological fitness and cultural selection pressures, including perceived risks or benefits of the vaccine itself, personal cost-benefit analyses of preventative health behaviors, and the structural or societal-level factors influencing vaccination rates [56,57]. Since the frequencies of V^+ phenotypes are multiplied by 1+σ, this parameter modulates whether there are more or fewer vaccinated individuals than expected: in other words, when σ>0, vaccinated individuals are more common in a set of offspring than would be expected strictly by parental beliefs and vaccination statuses. We calculate σ in each timestep as a function of the current vaccination coverage (frequency of V^+, i.e. $x_1 + x_2$), and in each simulation we specify σ_{max} as the maximum cultural selection pressure of

getting vaccinated ($-1 \leq \sigma_{max} \leq 1$) (see the cultural selection coefficient function in **Figure 2**). This

function was constructed by fitting a curve to pre-specified conditions: we assume that when

vaccination coverage is low, the real and perceived benefits of vaccination are highest and thus

the cultural selection pressure is near σ_{max}, however, as vaccination coverage increases, the

perceived benefits of vaccination decrease and the cultural selection pressure is reduced (**Figure**

2).

Figure 2: Cultural selection coefficient function. The cultural selection coefficient function was
constructed by fitting a curve to specified conditions, and considers both health and non-health related
effects. The selection coefficient (σ; vertical-axis) is dependent on the frequency of vaccinated individuals
(V^+) in the population (horizontal-axis). σ_{max} is the maximum cultural selection coefficient associated with
being vaccinated. Perceived vaccine benefit is reduced as vaccination coverage increases, since the
negative effects of the disease will be less apparent. Note: Of the σ_{max} values shown, only $\sigma_{max} = 0.1$ allows
the cultural selection pressure to be either positive or negative at a given timepoint depending on the
frequency of vaccination.

Thus far, we have described vertical cultural transmission from parent to offspring. The

model incorporates a second phase with oblique cultural transmission (i.e. influence from non-

parental adults), in which individuals can change their inherited vaccine attitudes (**A**) due to

influence from other adults in the population. There are two probabilities associated with

attitude modulation: the probability that an vaccine hesitant (A⁻) individual adopts the vaccine

confident (A⁺) state (A⁻ to A⁺ transition probability, given by $A_{\to Confident}$ in **Figure 3**), and the

probability that an A⁺ individual adopts the A⁻ state (A⁺ to A⁻ transition probability, given by

$A_{\to Hesitant}$ in **Figure 3**). As with the strength of cultural selection (σ) described previously, the

probability that offspring change their vaccine attitude is a function of the V⁺ frequency in the

population. As the frequency of vaccinated individuals (V⁺) increases in the population, vaccine-

confident individuals (A⁺) are more likely to become hesitant ($A_{\to Hesitant}$ probability increases)

and vaccine-hesitant individuals (A⁻) are less likely to become confident ($A_{\to Confident}$ probability

decreases). By modulating the attitude transition probabilities according to the vaccination

coverage in this manner, we assume that when vaccine coverage (V⁺ frequency, $x_1 + x_2$) is low,

disease occurrence is high and the negative effects of the disease are experienced widely, thus

the benefits of being vaccinated (and the costs of not being vaccinated) are more evident [58,59].

As vaccination coverage (V⁺) increases in the population, and thus disease occurrence is low, the

benefits to being vaccinated are less obvious, while low-probability costs such as adverse

reactions become more apparent and could be perceived as being riskier than the disease itself.

Modulating both the attitude transition probabilities and the cultural selection coefficient

according to the level of vaccination coverage in a population reflects that perceptions about the

vaccine and its associated effects on health could be meaningfully different in a population with

high vaccination coverage than in one with low coverage.

Figure 3: Attitude Transition Probability Function. Attitude transition probability functions were
constructed by fitting a curve to specified values. Attitude transition probability (vertical axis) is a function
of the vaccination frequency in the population (V$^+$; horizontal axis). The probability that a vaccine hesitant
individual adopts vaccine confidence (A$^-$ to A$^+$ transition probability, shown in black) is determined by the
function $A_{\rightarrow Confident}$, and the probability that a vaccine confident individual adopts vaccine hesitancy (A$^+$
to A$^-$ transition probability, shown with a blue dashed line) is determined by the function $A_{\rightarrow Hesitant}$.

To compute the frequency of a given phenotype in the next iteration, we sum the
probability that each mating pair produces offspring of that phenotype over each of the sixteen
possible mating pairs. Cultural selection (σ), described above, then operates on offspring with the
V$^+$ trait. The full recursions, giving x_i' phenotype frequencies in the next iteration in terms of x_i in
the current iteration, are given in **Text S1**. If x_i' is equal to x_i, the system is at equilibrium. Unless
otherwise stated, the model is initialized with phenotypic frequencies structured to represent
those of the United States: x_1 (frequency of V$^+$A$^+$) = 0.81, x_2 (V$^+$A$^-$) = 0.1, x_3 (V$^-$A$^+$) = 0.07, x_4 (V$^-$A$^-$) =
0.02. These frequencies were estimated using reports of Measles-Mumps-Rubella (MMR)
vaccination rates and estimates of vaccine attitude frequencies obtained from various sources in
the literature [60,61] and the Centers of Disease Control ChildVax database [62,63].

Results:

Parent-to-Offspring Interactions (Simulations with vertical transmission only)

We began by testing this model with only vertical transmission dynamics, i.e. only parent-

to-offspring transmission, varying parameter values in turn to test their effects on population

vaccination behavior and attitudes. In the vertical transmission phase of the model, parents

choose whether to vaccinate their offspring (i.e., transmit V^+) or to not vaccinate (V^-), and

parents also transmit a vaccine attitude (confidence, A^+, or hesitancy, A^-), each with a specified

probability given the phenotypes of the parents. The parental attitude state, vaccination status,

assortative mating levels, and cultural selection parameters interact to affect vaccination

coverage (frequency of V^+) and vaccine confidence (frequency of A^+). Couples with mixed

vaccination and/or attitude states ($V^+ \times V^-$, $A^+ \times A^-$) are assumed to be more variable in their

decision to vaccinate their offspring than parents who share the same state. Hence, we primarily

modulated probabilities associated with these mixed-phenotype pairings for the purposes of

testing our model.

First, we compared the effects of varying the confidence transmission probabilities for

mixed-attitude couples (C_1 and C_2), i.e. those with one vaccine-hesitant individual and one

vaccine-confident individual, in combination with multiple factors: 1) the maximum cultural

selection coefficient (σ_{max}) (**Figure 4A-B**), 2) the vaccination influence parameters b_1 and b_2

(**Figure 4C-D**), 3) the attitude influence parameters c_1 and c_2 (**Figure 4E-F**), and 4) the vaccination

probabilities of couples with mixed traits, $B_{1,1}$, $B_{1,2}$, $B_{2,1}$, $B_{2,2}$ (**Figure 4G-H**). In each examination,

we observed a C_n threshold: there is a mid-range value of C_n at which vaccination coverage and

vaccine confidence traits are polymorphic (i.e. both forms of each trait coexist in the population),

separating definitive high ($\gtrsim 80\%$) and low ($\lesssim 30\%$) levels of vaccination coverage and confidence.

This C_n threshold value is more sensitive to σ_{max} than to b_m, c_n, or $B_{m,n}$: the threshold value is

lowered as σ_{max} increases (diagonal line in **Figure 4A-B**). Although vaccination probability ($B_{m,n}$) is

dependent on both c_n, the influence of parental vaccine attitude, and b_m, the influence of

parental vaccination state (**Table S2**), modulating either type of influence of mixed-state parents

has little effect on the level of vaccination coverage and negligible effects on confidence levels at

each non-threshold C_n (**Figure 4C-F**).

Interestingly, direct modulation of the mixed-trait couple vaccination probability ($B_{1,1} =$

$B_{1,2} = B_{2,1} = B_{2,2}$) also has little power in affecting coverage and confidence levels at equilibrium

(**Figure 4G-H**). We hypothesize that predominantly high or predominantly low confidence

transmission within a population reduces the occurrence of "mixed-trait" pairings, i.e. if the

majority of the population becomes confident or hesitant, there are fewer confident-hesitant and

vaccinated-unvaccinated pairings. Thus, the effect of modulating mixed-trait vaccination

probabilities ($B_{1,1}$, $B_{1,2}$, $B_{2,1}$, $B_{2,2}$) is significantly minimized as these couples approach low

frequencies in the population, and confidence transmission dominates the vaccination patterns.

To further explore this relationship, we varied all vaccination probabilities ($B_{m,n}$), not just those of

mixed couples, while modulating mixed trait confidence transmission probabilities ($C_1 = C_2$) as

before. To directly alter vaccination probabilities while still accounting for the couple's vaccine

attitudes, we set ranges of values for $B_{m,n}$ that vary along the horizontal axis of **Figure 5**, with the

vaccination probability for two hesitant parents (e.g. $B_{0,0}$) on the lower end of the range and the

vaccination probability for two confident parents (e.g. $B_{3,3}$) on the higher end of the range (**Table**

S3). In these tests, we observe increasing equilibrium vaccination coverage as $B_{m,n}$ probabilities

increase, with higher coverage in high-confidence transmission environments (**Figure 5**). If we

vary both confidence transmission parameters and vaccination probability parameters by

implementing range shifts in both C_n and $B_{m,n}$, we observe an interaction between confidence

transmission and vaccination probability that determines vaccination coverage (**Figure S1**). In

both, we confirm vaccination coverage levels are determined by an interaction between

confidence transmission and vaccination probability, whereas confidence levels are dictated

primarily by levels of confidence transmission. In sum, the degree to which parents with mixed

vaccine-hesitant and vaccine-confident attitudes transmit vaccine confidence instead of vaccine

hesitancy to their offspring is a key factor in determining population trait majorities which can

drastically shift population dynamics.

Figure 4: **Vaccine Confidence Transmission Dictates Vaccination Coverage and Confidence Levels**
Heatmaps showing final vaccination coverage and vaccine confidence after 100 time-steps with no oblique
transmission, only parent-to-offspring transmission. The top row (**A, C, E, G**) shows vaccination coverage
(i.e. frequency of V^+ in the population) with low coverage in blue and high coverage in yellow; the bottom
row (**B, D, F, H**)) shows the corresponding final vaccine confidence (i.e. frequency of A^+), with low
confidence in red and high confidence in yellow. Unless varied on the horizontal or vertical axis, other
parameters are set to the default values given in **Table 1**. In our model, parents' likelihood of vaccinating
their children depends on both their vaccination state and their attitude state. This figure shows that the
strength of parental transmission of vaccine confidence (C_n) has a much stronger effect on the equilibrium
levels of both vaccine coverage (V^+) and confidence (A^+) than other parameters: the maximum cultural
selection coefficient, σ_{max} (**A,B**), the influence of parental vaccination state, b_m (**C, D**), the level of influence
of parental vaccine attitudes on their vaccination behaviors, c_n (**E,F**), and the probability that mixed-state
parents vaccinate their offspring $B_{m,n}$ (**G,H**).

Figure 5: Vaccination coverage levels are determined by an interaction between confidence transmission and vaccination probability. Heatmaps showing final vaccination coverage (**A**) and corresponding vaccine confidence (**B**) after 100 time-steps with no oblique transmission. Confidence transmission in mixed-attitude couples ($C_1 = C_2$) is varied along the vertical axis, while the vaccination probabilities ($B_{m,n}$) are set within the range indicated on the horizontal axis, with $B_{0,0}$, $B_{1,0}$, $B_{2,0}$ and $B_{3,0}$ taking the lowest value and $B_{3,3}$ taking the highest value (**Table S3**). We show increased equilibrium vaccination coverage with increasing vaccination probability ranges, while confidence levels are primarily dictated by proportion of the population transmitting confidence or hesitancy.

Next, we hold vaccine confidence transmission (C_n) at default probabilities, reminiscent of Mendelian transmission, such that two vaccine confident or two vaccine hesitant parents predictably transmit their vaccine attitude, and parents with differing vaccine attitudes each have a ~50% chance of transmitting their phenotype, e.g. C_0 near 0, C_1 and C_2 at 0.5, C_3 near 1 (**Table 1**). We then varied cultural selection in combination with vaccination-associated probabilities (b_m, c_n, $B_{m,n}$). With C_n held constant, cultural selection (σ_{max}) is the primary factor determining vaccination coverage and confidence levels (**Figure 6**. Raising the maximum cultural selection coefficient increases the equilibrium level of vaccination coverage and vaccine confidence across various levels of vaccination state influence (b_m) (**Figure 6A-B**), vaccination attitude influence (c_n) (**Figure 6C-D**), and vaccination probability ($B_{m,n}$) (**Figure 6E-F**). Unlike in **Figure 4**, vaccine confidence does not always mirror vaccination coverage across all levels of attitude influence (c_n) or vaccination probabilities. Instead, vaccine confidence levels decline with increased c_n and increased $B_{m,n}$ for $\sigma_{max} \lessgtr 0.3$ (**Figure 6D, F**), as well as for both increased c_n and increased b_m (**Figure S2**. This dynamic is interesting as these parameters influence vaccination behavior,

hinting that high vaccination rates could reduce a populations' expected vaccine confidence.

Vaccination coverage and vaccine confidence remain low when cultural selection does not favor

vaccination ($\sigma_{max} \lesseqgtr 0$), i.e. parents vaccinate their children at or below the levels expected based

on cultural transmission rates.

Figure 6: **Cultural Selection Influences Vaccination Coverage and Vaccine Confidence.** Heatmaps showing

final vaccination coverage (**A, C, E**) and final vaccination confidence (**B, D, F**) after 100 time-steps with no

oblique transmission, only parent-to-offspring transmission. As in **Figure 4**, parameters not varied here are

given in **Table 1**. Parents' likelihood of vaccinating their children depends on both their vaccination state

and their attitude state. At default probabilities of vaccine confidence transmission (C_n values in **Table 1**),

these figures show that modulating the maximum cultural selection coefficient affects the equilibrium

levels of vaccination coverage and vaccine confidence across the range of specified parameters: parental

vaccination state influence, b_m (**A, B**), parental attitude state influence, c_n (**C,D**), and offspring vaccination

probability, $B_{m,n}$ (**E,F**). Unless directly modulated (as in panels **E-F**), $B_{m,n}$ varies with b_m and c_n ($B_{m,n}$ =

$c_n((1+b_m)/2)$.

Temporal dynamics of vaccine-related beliefs and behaviors

To test whether the equilibrium phenotype frequencies were sensitive to starting

frequencies, we plotted the dynamics of each phenotype over time at default parameters (given

in **Table 1**). For each set of initial phenotype proportions tested, each phenotypic frequency

quickly adjusted to approach equilibrium values and then gradually plateaued to a stable

equilibrium (vertical transmission: **Figure 7 and Figure S3**, vertical+oblique transmission: **Figure**

S4. This demonstrates that equilibrium frequencies of vaccination coverage and vaccine

confidence are determined by the parameter conditions rather than by the initial frequencies

themselves.

437
Figure 7: Equilibrium frequencies are determined by the parameter space, not by initial frequencies. The

change in each of the four phenotype frequencies and the total V^+ and A^+ frequencies (vertical axis) over

100 iterations of the model (horizontal axis). Initial frequencies are varied, such that we begin each

simulation with a different phenotype at an initial high frequency (0.81): V^+A^+ in panel **A**, V^+A^- in panel **B**,

V^-A^+ in panel **C**, V^-A^- in panel **D**; the remaining phenotypes are set to lower frequencies (0.1, 0.07, 0.02).

See **Figure S3** for a full listing of these initial frequencies. The remaining parameters are held at default

values (**Table 1**).

When two parameters in particular are varied—maximum cultural selection (σ_{max}) or

confidence transmission ($C_1 = C_2$)—we observe a trade-off between the V^-A^- phenotype, which

dominates at lower values of these parameters, and the V^+A^+ phenotype, which dominates at

higher values (**Figures 8-9**). Interestingly, the "conflicting" phenotypes (when an individual's

attitude toward vaccinating their children does not match their own vaccination state: V^-A^+ and

V^+A^-) are present at their highest frequencies at neutral cultural selection ($\sigma_{max} = 0$, **Figure 8B**)

and/or neutral confidence transmission ($C_1 = C_2 = 0.5$, **Figure 9B**). Vaccinated individuals have the

same fitness regardless of their attitude (V^+A^+ bears the same selection pressure as V^+A^-), so it is

worth noting that at higher levels of confidence transmission and cultural selection, V^+A^+

increases in frequency but V^+A^- decreases in frequency (compare **Figure 8B-C, Figure 9B-C**). This

pattern seems to reflect their differing likelihoods of vaccinating their offspring: across all

possible partners, vaccinated but vaccine-hesitant parents (V^+A^-) are less likely to vaccinate their

offspring than vaccinated and vaccine-confident parents (V^+A^+), resulting in more V^- offspring.

Thus, when V^+ is favored by cultural selection, there is indirect selection against the V^+A^-

phenotype (**Figure 8**). Similarly, indirect selection against V^-A^+ occurs when V^- is favored by

cultural selection (**Figure 8A**): compared to **Figure 8B**, we observe an increase in V^-A^- individuals

but a decrease in V^-A^+ individuals, who because of their vaccine confidence have more V^+

offspring, which are culturally disfavored in this environment. When cultural transmission from

non-parental adults (oblique transmission) was included, described in following sections, we

observed similar patterns, but the final equilibria were more likely to be polymorphic, with

vaccinated, unvaccinated, confident, and hesitant phenotypes stabilizing at more moderate

frequencies than they would have with only vertical transmission (Compare **Figures 8-9 to Figures**

S5-S6).

Low confidence transmission ($C_1 = C_2 = 0.1$, **Figure 9A**) increases the frequency of vaccine

hesitancy (A^-) in the population over time, increasing the probability that more couples choose

not to vaccinate their offspring. However, the increase in vaccine hesitancy does not occur

equally in vaccinated and unvaccinated individuals: A^- frequency may increase overall in this

environment, but V^+A^- frequencies are lower and V^-A^- frequencies are higher (compared to

Figure 9B-C and **Figure S6**). At neutral confidence transmission probabilities (i.e. when couples

with one confident and one hesitant parent are equally likely to transmit either attitude), there is

a higher chance that the vaccinated but vaccine-hesitant (V^+A^-) phenotype is replenished.

However, if confidence is highly transmitted ($C_1 = C_2 = 0.8$), the V^+A^- frequency will be reduced, as

this phenotype is more likely to produce A^+ offspring than A^-, thus increasing V^+A^+ phenotype

frequencies in the population (**Figure 9** and **Figure S6**). If we turn to the other conflicting

phenotype, unvaccinated but vaccine-confident (V^-A^+) individuals become more common when

A^+ increases in frequency in the population as $C_1 = C_2$ increases from 0.1 to 0.5 (**Figure 9** and

Figure S6). In contrast, higher vaccine confidence transmission ($C_1 = C_2 = 0.8$) can lead to a

vaccination-promoting environment in which V^- frequencies are reduced over time; thus the V^-A^+

phenotype becomes rare and V^+A^+ predominates (**Figure 9** and **Figure S6**).

Figure 8: **Temporal Effects of Cultural Selection**. The change in equilibrium phenotype frequencies as the

maximum cultural selection coefficient (σ_{max}) is varied: **A**. $\sigma_{max} = -0.1$; **B**. $\sigma_{max} = 0$; **C**. $\sigma_{max} = 0.1$; **D**. $\sigma_{max} =$

0.5, while other parameters are held at default values (**Table 1**). Cultural selection against vaccinated

individuals increases the frequency of V^-A^-, decreasing the other frequencies (**A**), whereas increased

cultural selection favoring vaccinated individuals increases V^+A^+ frequencies while decreasing the other

frequencies (**B, C, D**).

Figure 9: **Temporal Effects of Confidence Transmission**. The change in phenotype frequencies over 50
iterations as vaccine confidence transmission in mixed couples ($C_1 = C_2$) is varied (**A**. $C_1 = C_2 = 0.1$; **B**. $C_1 = C_2$
= 0.5; **C**. $C_1 = C_2 = 0.8$) while other parameters are held at default values (**Table 1**). The population
equilibrates at over 90% A^-V^- at low confidence transmission (A). Increasing the probability of confidence
transmission results in less vaccine hesitancy and, in turn, higher vaccination frequencies (V^+A^+).

Offspring can Change their Inherited Hesitancy State (Vertical and Oblique Dynamics)

Increased exposure to the attitudes of the broader community (i.e. oblique cultural

transmission from non-parental adults in the population) could influence and change vaccination

beliefs inherited in childhood. Therefore, we next included these oblique effects in our model to

understand how they might modulate vaccine confidence and vaccination coverage levels. In the

oblique transmission phase of the model, offspring can change their vaccine attitude with some

probability based on the frequency of vaccination in the population (**Figure 3**). Thus, in addition

to the vertical transmission of attitudes and behaviors, phenotype frequencies are further

affected by the probability that adult offspring change their attitude (i.e. transition from vaccine

confident (A^+) to hesitant (A^-) and vice versa).

The addition of oblique dynamics produces a pattern of vaccination coverage and vaccine

confidence similar to that of simulations run with solely vertical transmission (**Figure 4 and Figure**

6 compared to Figure 10 and Figure 11, and Figures 7-9 compared to **Figures S4-6**)—the level of

(vertical) vaccine confidence transmission still largely determines the level of vaccination

coverage and vaccine confidence (**Figure 10**). However, oblique cultural influences expanded the

polymorphic space, resulting in a wider range of intermediate C_n in which the different

phenotypes (vaccinated, unvaccinated, confident, and hesitant) are present in the population in

roughly equal proportions. In other words, there is a wider horizontal stripe of moderate values

between the definitively high and definitively low equilibrium frequencies in **Figure 10** than in

Figure 4). Overall, the addition of oblique transmission appears to lead to less polarized results,

moving the equilibrium levels of vaccination coverage and vaccine confidence away from extreme

values.

Figure 10: Vaccine confidence transmission dictates vaccination coverage and confidence levels (with oblique transmission). Heatmaps showing final vaccination coverage (i.e. frequency of V^+ in the population, with low coverage in blue and high coverage in yellow (**A, C, E, G**)) and final vaccine confidence (i.e. frequency of A^+, with low confidence in red and high confidence in yellow (**B, D, F, H**)) after 100 time-steps in which oblique transmission of vaccine attitude can occur after parent-to-offspring transmission. The likelihood that individuals change their vaccine beliefs depends on the current vaccination coverage of the population (**Figure 2**). Unless varied on the horizontal or vertical axes, other parameters are set to the default values given in **Table 1**. Parents' likelihood of vaccinating their children depends on both their vaccination state and their attitude state; these figures show that the strength of parental transmission of vaccine confidence (C_n) has a much stronger effect on the equilibrium levels of both vaccine coverage (V^+) and confidence (A^+) than other tested parameters do: the maximum cultural selection coefficient, σ_{max}, (**A,B**), the influence of parental vaccination state, b_m, (**C, D**), the level of influence of parental attitudes on their vaccination behaviors, c_n, (**E,F**), and offspring vaccination probability, $B_{m,n}$ (**G,H**).

With neutral confidence transmission ($C_1 = C_2 = 0.5$), we also observe an expansion of the

polymorphic space when we modulate cultural selection (σ_{max}) alongside the influence and

transmission parameters (**Figure 11**). Interestingly, in the cultural environment defined by this

parameter space, we observe a pattern that deviates from the expected association between

high vaccine confidence and high vaccination coverage: as the influence of vaccine attitudes (c_n)

and vaccination probabilities ($B_{m,n}$) increase (**Figure 11**, horizontal axes), the population's

equilibrium vaccination coverage increases while its vaccine confidence decreases. This pattern

persisted across all tested levels of maximum cultural selection (σ_{max}) (**Figure 11**, vertical axes). In

 other words, we observe higher levels of confidence at low influence and transmission

 probabilities than we do at higher probabilities (**Figure 11D, F**).

Figure 11: Cultural selection influences vaccination coverage and vaccine confidence (with oblique transmission). Heatmaps showing final vaccination coverage **(A, C, E)** and final vaccination confidence **(B, D, F)** after 100 time-steps with oblique transmission. As in previous figures, parameters not varied are given in Table 1. Parents' likelihood of vaccinating their children depends on both their vaccination state and their attitude state. At default probabilities of vaccine confidence transmission (C_n), these figures show that modulating the maximum cultural selection coefficient affects the equilibrium levels of vaccination coverage and vaccine confidence across the range of specified parameters: parental vaccination state influence, b_m **(A, B)**, parental attitude state influence, c_n **(C,D)**, and offspring vaccination probability, $B_{m,n}$ **(E,F)**. Unless directly varied (as in panels **E-F**), $B_{m,n}$ varies as b_m and c_n are varied, as shown in **Table 1**.

We explored the interaction between the influence parameters, b_m and c_n, at various maximum cultural selection coefficients (σ_{max}) (**Figure 12**). Vaccination coverage and vaccine confidence equilibrate at mid-range frequencies (between 0.3 and 0.8) across the range of b_m and c_n, indicating that these trait frequencies are not particularly sensitive to either parameter. Cultural selection favoring vaccination increases the equilibrium level of vaccination coverage and vaccine confidence (**Figure 12 and Figure S7**). The most notable deviation between equilibrium

confidence and vaccination frequencies occurs at the intersection of the highest influence

parameter values (b_m and c_n), circumstances in which the parents' vaccination states and vaccine

attitudes overwhelmingly support offspring vaccination; in this top right section of the heat maps,

vaccination coverage is high while vaccine confidence is lower, indicating a behavioral pattern in

which mixed-trait couples are more inclined to vaccinate their offspring than transmit vaccine

confidence. Overall, the addition of oblique transmission to a population that would otherwise

equilibrate at high vaccination coverage (**Figure S2**) leads to increased attitude transition to

vaccine hesitancy and subsequently lower vaccine coverage.

Figure 12: **The influence of parental traits on vaccination coverage and vaccine confidence at different**

levels of cultural selection. Heatmaps showing final vaccination coverage (**A, C**) and final vaccination

confidence (**B, D**) after 100 timesteps with oblique transmission. We modulate the interaction between

vaccination state influence (b_m; vertical axis) and attitude influence (c_n; horizontal axis) at various

maximum cultural selection coefficients: σ_{max} = 0 (**A, B**) and σ_{max} = 0.1 (**C, D**). As in previous figures,

unvaried parameters are given in **Table 1**. Vaccination frequency increases as both influence probabilities

increase and vaccination confidence decreases as both influence probabilities increase.

Mating Preferences:

We hypothesized that mating preference (assortative mating) could modulate belief and

behavior dynamics and thus the vaccination coverage and confidence levels in the population. If

individuals are more likely to pair with individuals of the same vaccine attitude, such that same-

attitude couples become more common and mixed-attitude couples are less common, the

parameter values for mixed-attitude couples may have less impact on vaccination coverage and

confidence dynamics. Therefore, we analyzed the interaction between A^+ homophily (with α_1

indicating the preference of A^+ individuals for other A^+ individuals) and A^- homophily (with α_2

indicating the preference of A^- individuals for other A^- individuals) at various σ_{max} levels. When

vaccine attitudes are transmitted both from parent to offspring and between unrelated

individuals (vertical and oblique transmission) and there is neither cultural selection for nor

against being vaccinated (σ_{max} = 0), we observe a threshold region at roughly equal mating

preferences ($\alpha_1 \approx \alpha_2$; diagonal lines in **Figure 13A-B**); above this boundary (when $\alpha_1 > \alpha_2$)

vaccination coverage and confidence are much higher than below this boundary (when $\alpha_1 < \alpha_2$) .

When cultural selection favors being vaccinated (σ_{max} > 0, **Figure 13C-D, Figure S8C-D**), the

threshold between high and low equilibrium values is shifted, such that high coverage and high

confidence levels can potentially be attained even when vaccine hesitant individuals

preferentially pair with each other more than vaccine confident individuals do ($\alpha_1 < \alpha_2$). Likewise,

if cultural selection explicitly does not favor vaccination (e.g. σ_{max} = −0.1, **Figure S8**), low

vaccination coverage and confidence can occur even when there are more vaccine confident

couples in the population than hesitant couples ($\alpha_1 > \alpha_2$). We observe qualitatively similar

patterns when vaccine attitudes are only transmitted from parent to offspring (**Figure S9**); as we

have previously observed, the addition of oblique transmission leads to a broader polymorphic

region than vertical transmission alone. These patterns illustrate two overarching themes: 1)

preferential interactions between individuals with similar vaccine beliefs can dramatically shift

the equilibrium levels of vaccination coverage and confidence with all other parameters

remaining equal, and 2) the actual and perceived quality and efficacy of the vaccine are

important to determining vaccine acceptance.

Figure 13: Homophily between individuals with similar vaccine beliefs can shift equilibrium frequencies of both vaccination coverage and confidence. Heatmaps showing final vaccination coverage (**A, C**) and final vaccine confidence (**B, D**) after 100 timesteps with oblique transmission. As in previous figures, unspecified parameters are given in **Table 1**. As vaccine-hesitant individuals (A⁻) increasingly prefer to pair with one another (α_2; horizontal axis), vaccine-confident individuals (A⁺) must also preferentially interact to maintain high vaccine coverage (α_1; vertical axis); this tradeoff is modulated by the cultural selection pressures on vaccination ($\sigma_{max} = 0$ (**A, B**) and $\sigma_{max} = 0.1$ (**C, D**)).

Mandatory Vaccines and Vaccine Inaccessibility:

Compulsory Vaccination

The implementation of mandatory childhood vaccinations has resulted in almost 100% vaccination coverage in the United States [9,61]. Allowing non-medical exemptions such as those based on philosophical or religious beliefs, however, has contributed to the recent decrease in vaccination rates [64]. We suspect that parental vaccine attitudes also influence their use of exemptions and thus levels of non-vaccination under a mandated vaccination system. Therefore, we simulate the effects of varying levels of compulsory vaccine mandate strictness (i.e. the ease of obtaining an exemption) by modulating the influence of vaccine attitude on offspring vaccination for each pairing. We assume the implementation of strict mandates (difficulty in obtaining exemptions) would increase vaccination in all couple types. If vaccination exemptions are permitted, we expect that $A^- \times A^-$ couples (those with two vaccine-hesitant individuals) would be most likely to obtain exemptions, followed by mixed attitude ($A^- \times A^+$ or $A^+ \times A^-$) couples, with vaccine confident couples ($A^+ \times A^+$) being least likely. Hence, to model the effects of implementing a strict mandate, we increase attitude influence parameters from baseline values (**Table 1**) to $c_0 = 0.5$, $c_1 = c_2 = 0.9$, $c_3 = 0.99$, then examined the effect of cultural selection coefficient and confidence transmission probability (**Figure 14A-B**); we then compare the results of this strict mandate both to our baseline parameters and to a more lenient mandate represented by $c_0 = 0.3$, $c_1 = c_2 = 0.7$, $c_3 = 0.99$ (**Figure S10**).

Our previous simulations consistently showed a positive correlation between population-level frequencies of vaccination coverage and vaccine confidence (**Figures 8-10, 13**). However, modeling a cultural environment under strict mandates reveals a decoupling of vaccination coverage and vaccine confidence at low confidence transmission probabilities (**Figure 14A-B**). Even when vaccine confidence is very low (specifically at mixed-trait couple confidence transmission probabilities below 0.5; red region in **Figure 14B**), vaccination coverage is much higher than without the mandate (compare **Figure 14A-B** to **Figure 10A-B**, see also **Figure S10**). This suggests that an external pressure to vaccinate is able to overcome the opposing cultural pressure imposed by hesitancy in the population. If we lower the barrier to acquiring an exemption (using more lenient mandate parameters), vaccination coverage and vaccine confidence dynamics begin to mirror one another, as they do in simulations without vaccine mandates (**Figure S10**).

Vaccine Inaccessibility

We next modeled the dynamics of vaccination behaviors and attitudes when a population

has limited access to vaccines. To represent this scenario, we reduced the influence of parental

vaccine attitudes on vaccination behaviors for couples with at least one confident individual (i.e.

reducing c_1, c_2, c_3 from default values). In this simple representation of a vaccine-scarce

environment, we assume that parents' confidence in vaccines would have reduced influence on

their ability to vaccinate their offspring, that is, their vaccine confidence does not ensure their

ability to overcome vaccine inaccessibility. Hesitant couples are least likely to vaccinate their

offspring regardless of vaccine availability, but couples who would likely vaccinate their offspring

given the chance would have difficulty doing so due to the lack of resources. This scenario can be

contrasted with the strict vaccination mandate scenario described above, in which a couple

would often vaccinate their children despite their hesitancy. Attitude influence parameters were

set to $c_0 = 0.01$; $c_1 = c_2 = 0.1$, and $c_3 = 0.5$, and as before, we modulated the maximum cultural

selection coefficient of vaccination (σ_{max}) and confidence transmission (C_1 and C_2) **(Figure 14C-D)**.

Vaccination coverage was noticeably reduced overall, while vaccine confidence increased slightly

across the parameter space. Juxtaposed with the strict mandate **(Figure 14A-B)**, our vaccine

scarcity models produce an opposite deviation of vaccination coverage from vaccine confidence

levels: when vaccines are mandated, we observe increased vaccination coverage in low-

confidence environments, and when vaccines are inaccessible, we observe lower than expected

vaccination coverage (<50%) in a predominantly vaccine-confident environment (>90%) (**Figure**

14).

Figure 14: External factors (vaccine mandates and vaccine scarcity) disconnect levels of vaccine confidence from vaccination coverage. Equilibrium vaccine coverage and vaccine confidence levels for varying values of confidence transmission (C_n; vertical axis) and maximum selection coefficient (σ_{max}; horizontal axis). A strict vaccine mandate (**A, B**) is simulated by $c_0 = 0.5$, $c_1 = c_2 = 0.9$, $c_3 = 0.99$; Vaccine scarcity (**C, D**) is simulated by $c_0 = 0.01$; $c_1 = c_2 = 0.1$, and $c_3 = 0.5$. As in previous figures, unspecified parameters are given in **Table 1**. These simulations show an inverse correlation between vaccination coverage and vaccine confidence at $C_n < 0.5$ under a strict mandate, and $C_n > 0.5$ when vaccine access is severely limited.

Discussion:

690
In this manuscript, we present a simplified model of a complex process: the spread of vaccine
attitudes and their effects on vaccination behavior in a population. Increasing and maintaining
sufficient vaccination coverage to combat disease is more complex than simply increasing vaccine
availability or providing accurate information. A number of factors affect a person's vaccine-
related beliefs and a family's decision to vaccinate their children, including their history with
vaccinations and their perception of the disease and vaccine effects. As such, it is important that
we understand how these personal factors can shape vaccination cultures and thus affect public
health. Using a cultural niche construction framework, we modeled the spread of vaccine
attitudes and vaccination behavior in a variety of circumstances and measured the resulting
levels of vaccination coverage and vaccine confidence in the modeled population. Using this
novel approach of modeling dynamically interacting beliefs and behaviors, we are able to explore
the interplay of cultural factors that drive vaccine attitudes and vaccination behavior, providing
insight into how vaccination cultures are formed, maintained, and evolve.

Our model shows vaccine confidence transmission (C_n) to be the parameter that most
strongly determines vaccination coverage and confidence levels. That is, the probability of
transmitting vaccine-positive attitudes is a strong predictor of future vaccination coverage
(**Figures 4 and 10**). We note, however, that the vaccine confidence transmission of mixed-trait
couples (those with one hesitant parent and one confident parent, $C_1 = C_2$) has a much larger
effect on the levels of vaccination coverage than mixed-trait couple vaccination probability itself,
likely because the spread of vaccine confidence or hesitancy makes mixed-trait couples relatively
rare in the population. By modulating the vaccination probabilities of all individuals directly
(varying all $B_{m,n}$), while modulating mixed-trait confidence transmission ($C_1 = C_2$) (**Figure 5**) or all
confidence transmission parameters (C_n) (**Figure S1**), we show that vaccination probabilities work
alongside vaccine confidence transmission to determine vaccination coverage equilibria, while
confidence equilibria are dictated primarily by confidence transmission probabilities.

It is important to note that even though this model demonstrates the powerful effects of
beliefs on vaccination behaviors, it also shows that vaccine efficacy and perceived value are
important to maintaining sufficient levels of vaccination coverage, especially if vaccine confidence
is not being robustly transmitted (or maintained in adulthood). Further, we show that a culture in
which vaccine-hesitant individuals preferentially assort with one another more so than vaccine-
confident individuals ($\alpha_2 > \alpha_1$) can allow vaccine hesitancy to more easily gain a foothold,

lowering vaccination coverage unless vaccines have a high perceived value (i.e. high σ_{max}, **Figure**

13, Figure S9). Thus, our simulations suggest that a pro-vaccination health culture can be

undermined by a vaccine-hesitancy "echo chamber". This phenomenon would likely be amplified

if we modeled homophily of oblique interactions as well, for example if individuals preferentially

interacted with friend groups or news sources that shared similar beliefs and values to their own.

Further, our model demonstrates that the equilibrium frequencies of vaccination coverage

and vaccine confidence are determined by the cultural context rather than by the initial

frequencies. In other words, the current cultural landscape—beliefs, behaviors, policies, etc.—are

more predictive of future levels of vaccine coverage and confidence than current coverage and

confidence levels in the population. This result further supports the importance of considering

the cultural factors that have shaped current health-related beliefs and behaviors if health

policies aim to maintain or change the current conditions.

In the absence of external pressures, populations reach an equilibrium at which vaccination

coverage and vaccine confidence mirror each other across the tested parameters: high

vaccination coverage coincides with high levels of confidence in vaccines, and low coverage

coincides with low confidence levels. This result suggests, intuitively, that when population traits

are at or near an equilibrium, we can infer that a population with high vaccination coverage will

have low rates of vaccine hesitancy and vice versa. We then model two possible exceptions to

this pattern—vaccine mandates and a lack of access to vaccines. When there is increased

pressure to vaccinate or difficulty in acquiring vaccination exemptions, an undercurrent of

vaccine hesitancy can persist in a relatively well-vaccinated population, potentially limiting the

adoption of newly introduced vaccines. This possibly contributes to the unexpected lag in uptake

of newer vaccines, such as the COVID or HPV vaccines, in communities with otherwise high

vaccination rates [65–67]. The perceived increase in hesitancy surrounding new vaccines may

actually be existing vaccine hesitancy becoming evident. In addition, "fence sitters", those who

have not made a firm stance regarding vaccines and thus could be more influenced by targeted

campaigns [61], may develop higher levels of uncertainty about new vaccines than their parents

had about existing ones.

In contrast to the effect of vaccine mandates, by modeling vaccine inaccessibility we

illustrate another important pattern: reduced vaccination coverage in a vaccine confident culture.

In a vaccine-scarce environment, an individual's attitude regarding vaccines has less influence on

vaccination behavior due to the barrier imposed by resource availability. As a result, a population

may be undervaccinated despite holding vaccine-affirming beliefs. In addition, a health culture

previously shaped by vaccine inaccessibility could potentially ingrain specific behavioral practices

(for example, visiting the doctor only when a child is sick and not for a regular vaccine schedule)

that are not easily modified even if vaccines become more readily available. These vaccine

scarcity scenarios are most likely to exist in low- and middle-income countries in which vaccine

acquisition, storage and/or distribution resources are insufficient [68–70] whereas the opposite

scenario (low confidence/high vaccination) after vaccine mandates is most common in developed

nations [71]. In summary, we find that vaccine mandates can result in high vaccination coverage

even in a culture of hesitancy, and that lack of access to vaccines can produce the inverse: low

vaccination coverage in a culture of confidence.

Individuals consider multiple factors alongside vaccine efficacy when deciding whether to

vaccinate themselves or their children, essentially performing an internal cost-benefit analysis

based on their circumstances and interpretation of accessible information. We aimed to be

inclusive of these considerations via our comprehensive cultural selection coefficient; this

parameter allows the frequency of vaccination to deviate from the level that is expected given

the cultural transmission probabilities. Increasing positive public perception through honest and

effective communication, while reducing public concern about vaccines and increasing vaccine

safety, together could drive increased vaccination trust and acceptance. Achieving the optimal

vaccination coverage lies not only in the hands of the public by vaccinating themselves and their

children, but also in the efforts of health officials and leaders in creating an environment that

fosters confidence by assuring the public of vaccine efficacy, safety, and value, while providing

convenient avenues to attain vaccines.

As with any model, we cannot fully capture the complex reality of vaccine hesitancy and

vaccination behavior. First, though vaccination frequency data is available for numerous vaccines,

frequency data for vaccine attitudes are much less common, with the two traditionally not

surveyed together. Thus, there is no dataset that exactly estimates the phenotypes presented

here, for example, the number of vaccinated but hesitant (V^+A^-) individuals in a population. The

goals of vaccination attitude surveys have been primarily to identify themes of vaccine hesitancy,

and to a lesser degree, the themes of vaccination. However, they do not report parent

vaccination states or whether the child was actually vaccinated (on schedule). With data

presenting parent vaccination states alongside their vaccine attitudes and vaccination decisions,

we would be able to more accurately inform phenotype frequencies, possibly extending the

model to incorporate various types of hesitancy. We note, however, that our results did not
depend on the initial proportions of vaccination status or vaccine hesitancy, so these data would
primarily be for comparison to our equilibrium outcomes. We were also constrained by limited
data to inform the cultural transmission and transition probabilities. In our model, baseline
confidence transmission and influence probabilities are structured according to a simple pattern
of inheritance, such that each parent is equally likely to influence an offspring's phenotype.
However, cultural traits and vaccination attitudes may not strictly follow this pattern: one parent
might have more influence, or one variant of a trait might be more likely to be transmitted. In
addition, transmission probabilities are constant in our model, remaining unaffected by changing
cultural conditions throughout each simulation, but in reality, these probabilities may fluctuate in
response to a variety of factors including vaccine type or family structure. Future developments
of the model could include modulating the probability of transmission of vaccine confidence
dependent on aspects of the cultural environment, such as the attitude frequencies in the
population. Our cultural selection coefficient and attitude transition probabilities did vary with
the frequency of vaccination coverage, but the exact relationships could not be informed by
existing data. As a result, we generated frequency-dependent functions to fit a set of
assumptions: for example, that the transition to vaccine hesitancy might be most common when
the vaccination coverage is low and the benefits of vaccination are more apparent.
Though vaccination coverage and vaccine confidence stabilized (reached equilibria) in our
simulations, in reality vaccination rates fluctuate over time in response to changing population
dynamics, perhaps never arriving at a stable equilibrium. For example, the increasingly rapid
spread of information [23] may cause attitudes and behaviors to change frequently over short
periods of time. In our model, most of the phenotype frequency fluctuations occur in the first few
iterations before quickly adjusting to an equilibrium. Unlike many models of population
dynamics, this model has a discrete-time format and does not consider a birth-death cycle or
asynchrony in population turnover. Thus, the timescale of our model might not translate directly
to years or generations, and we avoid interpreting the number of iterations in literal terms. It is
possible that if more realistic birth and death processes were incorporated, the cultural dynamics
would occur at different timescales and would continue to fluctuate instead of approaching a
stable equilibrium. In addition, parents' vaccination decisions are also influenced by the
grandparents of the children to be vaccinated [72]. A restructuring of the timescale or the

incorporation of population asynchrony in our model could allow for consideration of these

influences.

Finally, we constructed the offspring vaccination probability to be informed primarily by

parents' vaccine attitudes and secondarily by their own vaccination status. Though it is

understood that there is an interaction between parents' beliefs and their own experiences with

vaccines regarding their decision to vaccinate their children, accurately modeling the relative

contribution of these two factors could benefit from empirical studies on parental willingness to

vaccinate based on their beliefs and vaccination status. With our current formula ($B_{m,n}$, **Table S2**),

vaccine-confident parents who did not themselves receive childhood vaccines have a reduced

likelihood of vaccinating their offspring than vaccinated parents. In reality, parental vaccine

attitudes might even further outweigh their own vaccination status in their decision making than

we model here.

Our findings, which are based on cultural evolutionary modeling in a public-health

context, suggest several avenues for policy and outreach recommendations. First, we note that

vaccine mandates with limited exemptions will increase the vaccination coverage but may mask

the spread of vaccine hesitancy. In addition to vaccine policy, our results suggest that a broad

effort to encourage and inform the public about vaccine safety and efficacy will foster higher

vaccine coverage. In this vein, we note that individuals who are skeptical about vaccines might

invest more time in seeking out information about them [73–75], thus we recommend that

accurate information about vaccines should be readily accessible through a variety of means, be

easily understood, and be supported by personal anecdotes. If misinformation is easier to

encounter and digest than accurate information, it could have an outsized impact on individuals

who are "on the fence" [61]. Relatedly, we note that research suggests transparency and trust go

hand in hand: if individuals perceive the healthcare system is concealing information to make

vaccines appear more safe, trust in that system will decrease and people will be more susceptible

to vaccine hesitancy [73–76]. Finally, dialogue between people with different beliefs and

attitudes can help to break the "echo chambers" of homophily, encouraging individuals to

communicate and empathize with one another. Therefore, to address vaccine hesitancy, our

results underscore the importance of considering the cultural beliefs that underpin health

behaviors.

References:

1. Institute of Medicine, Division of Health Care Services, Committee for the Study of the Future of
Public Health. The Future of Public Health. National Academies Press; 1988.

2. Fenner F. A successful eradication campaign. Global eradication of smallpox. Rev Infect Dis. 1982;4:
916–930.

3. Kim-Farley R, Schonberger L, Nkowane B, Kew O, Bart K, Orenstein W, et al. POLIOMYELITIS IN THE
USA: VIRTUAL ELIMINATION OF DISEASE CAUSED BY WILD VIRUS. The Lancet. 1984. pp. 1315–1317.
doi:10.1016/s0140-6736(84)90829-8

4. Salk D. Eradication of Poliomyelitis in the United States. I. Live Virus Vaccine-Associated and Wild
Poliovirus Disease. Clinical Infectious Diseases. 1980. pp. 228–242. doi:10.1093/clinids/2.2.228

5. Thompson KM, Strebel PM, Dabbagh A, Cherian T, Cochi SL. Enabling implementation of the Global
Vaccine Action Plan: developing investment cases to achieve targets for measles and rubella
prevention. Vaccine. 2013;31 Suppl 2: B149–56.

6. Atwell JE, Salmon DA. Pertussis resurgence and vaccine uptake: implications for reducing vaccine
hesitancy. Pediatrics. 2014. pp. 602–604.

7. Kubin L. Is There a Resurgence of Vaccine Preventable Diseases in the U.S.? Journal of Pediatric
Nursing. 2019. pp. 115–118. doi:10.1016/j.pedn.2018.11.011

8. Falagas ME, Zarkadoulia E. Factors associated with suboptimal compliance to vaccinations in children
in developed countries: a systematic review. Curr Med Res Opin. 2008;24: 1719–1741.

9. Dubé E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. Vaccine hesitancy: an overview. Hum
Vaccin Immunother. 2013;9: 1763–1773.

10. Streefland P, Chowdhury AMR, Ramos-Jimenez P. Patterns of vaccination acceptance. Social Science
& Medicine. 1999. pp. 1705–1716. doi:10.1016/s0277-9536(99)00239-7

11. MacDonald NE, SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: Definition, scope and
determinants. Vaccine. 2015;33: 4161–4164.

12. Scheres J, Kuszewski K. The Ten Threats to Global Health in 2018 and 2019. A welcome and
informative communication of WHO to everybody. Zdrowie Publiczne i Zarządzanie. 2019. pp. 2–8.
doi:10.4467/20842627oz.19.001.11297

13. Schwartz JL. New media, old messages: themes in the history of vaccine hesitancy and refusal. Virtual
Mentor. 2012;14: 50–55.

14. Koslap-Petraco M. Vaccine hesitancy: Not a new phenomenon, but a new threat. J Am Assoc Nurse
Pract. 2019;31: 624–626.

15. Siddiqui M, Salmon DA, Omer SB. Epidemiology of vaccine hesitancy in the United States. Hum Vaccin
Immunother. 2013;9: 2643–2648.

16.   Callender D. Vaccine hesitancy: More than a movement. Hum Vaccin Immunother. 2016;12: 2464–
2468.

17.   Durbach N. Bodily Matters: The Anti-Vaccination Movement in England, 1853–1907. Duke University
Press; 2005.

18.   Swales JD. The Leicester anti-vaccination movement. The Lancet. 1992. pp. 1019–1021.
doi:10.1016/0140-6736(92)93021-e

19.   Wolfe RM, Sharp LK. Anti-vaccinationists past and present. BMJ. 2002;325: 430–432.

20.   Jackson CL. State laws on compulsory immunization in the United States. Public Health Rep. 1969;84:
787–795.

21.   Feikin DR. Individual and Community Risks of Measles and Pertussis Associated With Personal
Exemptions to Immunization. JAMA. 2000. p. 3145. doi:10.1001/jama.284.24.3145

22.   Salmon DA, Haber M, Gangarosa EJ, Phillips L, Smith NJ, Chen RT. Health Consequences of Religious
and Philosophical Exemptions From Immunization Laws. JAMA. 1999. p. 47.
doi:10.1001/jama.282.1.47

23.   Hornik J, Satchi RS, Cesareo L, Pastore A. Information dissemination via electronic word-of-mouth:
Good news travels fast, bad news travels faster! Computers in Human Behavior. 2015. pp. 273–280.
doi:10.1016/j.chb.2014.11.008

24.   Spier RE. Perception of risk of vaccine adverse events: a historical perspective. Vaccine. 2001;20 Suppl
1: S78–84; discussion S75–7.

25.   Eggertson L. Lancet retracts 12-year-old article linking autism to MMR vaccines. CMAJ. 2010;182:
E199–200.

26.   Rao TSS, Andrade C. The MMR vaccine and autism: Sensation, refutation, retraction, and fraud. Indian
J Psychiatry. 2011;53: 95–96.

27.   Yaqub O, Castle-Clarke S, Sevdalis N, Chataway J. Attitudes to vaccination: A critical review. Social
Science & Medicine. 2014. pp. 1–11. doi:10.1016/j.socscimed.2014.04.018

28.   Burley N. The meaning of assortative mating. Ethol Sociobiol. 1983;4: 191–203.

29.   Creanza N, Feldman MW. Complexity in models of cultural niche construction with selection and
homophily. Proc Natl Acad Sci U S A. 2014;111 Suppl 3: 10830–10837.

30.   Creanza N, Fogarty L, Feldman MW. Models of cultural niche construction with selection and
assortative mating. PLoS One. 2012;7: e42744.

31.   Gimelfarb A. Processes of Pair Formation Leading to Assortative Mating in Biological Populations:
Encounter-Mating Model. The American Naturalist. 1988. pp. 865–884. doi:10.1086/284827

32.   Centola D. An experimental study of homophily in the adoption of health behavior. Science.
2011;334: 1269–1272.

33. Salathé M, Bonhoeffer S. The effect of opinion clustering on disease outbreaks. J R Soc Interface.
2008;5: 1505–1508.

34. Yoder JS, Dworkin MS. Vaccination usage among an old-order Amish community in Illinois. Pediatr
Infect Dis J. 2006;25: 1182–1183.

35. Gastañaduy PA, Budd J, Fisher N, Redd SB, Fletcher J, Miller J, et al. A Measles Outbreak in an
Underimmunized Amish Community in Ohio. N Engl J Med. 2016;375: 1343–1354.

36. Bahta L, Ashkir A. Addressing MMR Vaccine Resistance in Minnesota's Somali Community. Minn Med.
2015;98: 33–36.

37. Wolff ER, Madlon-Kay DJ. Childhood vaccine beliefs reported by Somali and non-Somali parents. J Am
Board Fam Med. 2014;27: 458–464.

38. Laland K, Matthews B, Feldman MW. An introduction to niche construction theory. Evol Ecol.
2016;30: 191–202.

39. John Odling-Smee F, Laland KN, Feldman MW. Niche Construction: The Neglected Process in
Evolution (MPB-37). Princeton University Press; 2013.

40. O'Brien MJ, Laland KN. Genes, Culture, and Agriculture: An Example of Human Niche Construction.
Curr Anthropol. 2012;53: 434–470.

41. Fogarty L, Creanza N. The niche construction of cultural complexity: interactions between
innovations, population size and the environment. Philos Trans R Soc Lond B Biol Sci. 2017;372.
doi:10.1098/rstb.2016.0428

42. O'Brien MJ, Laland KN, Broughton JM, Cannon MD, Fuentes A, Gerbault P, et al. Genes, culture, and
agriculture: An example of human niche construction. Curr Anthropol. 2012;53: 000–000.

43. Fuentes A. Cooperation, conflict, and niche construction in the genus homo. War, peace, and human
nature. 2013; 78–94.

44. Perra N, Balcan D, Gonçalves B, Vespignani A. Towards a characterization of behavior-disease models.
PLoS One. 2011;6: e23084.

45. Mao L, Yang Y. Coupling infectious diseases, human preventive behavior, and networks--a conceptual
framework for epidemic modeling. Soc Sci Med. 2012;74: 167–175.

46. Chauhan S, Misra OP, Dhar J. Stability analysis of SIR model with vaccination. American journal of
computational and applied mathematics. 2014;4: 17–23.

47. Church KEM, Liu X. Analysis of a SIR model with pulse vaccination and temporary immunity: Stability,
bifurcation and a cylindrical attractor. Nonlinear Anal Real World Appl. 2019;50: 240–266.

48. May T, Silverman RD. "Clustering of exemptions" as a collective action threat to herd immunity.
Vaccine. 2003;21: 1048–1051.

49. Coe AB, Gatewood SBS, Moczygemba LR, Goode J-VKR, Beckner JO. The use of the health belief

model to assess predictors of intent to receive the novel (2009) H1N1 influenza vaccine. Innov Pharm.
2012;3: 1–11.

50. Reiter PL, Brewer NT, Gottlieb SL, McRee A-L, Smith JS. Parents' health beliefs and HPV vaccination of
their adolescent daughters. Soc Sci Med. 2009;69: 475–480.

51. Janz NK, Becker MH. The Health Belief Model: a decade later. Health Educ Q. 1984;11: 1–47.

52. Rosenstock IM. The Health Belief Model and Preventive Health Behavior. Health Educ Monogr.
1974;2: 354–386.

53. Jones CL, Jensen JD, Scherr CL, Brown NR, Christy K, Weaver J. The Health Belief Model as an
explanatory framework in communication research: exploring parallel, serial, and moderated
mediation. Health Commun. 2015;30: 566–576.

54. Wang E, Clymer J, Davis-Hayes C, Buttenheim A. Nonmedical exemptions from school immunization
requirements: a systematic review. Am J Public Health. 2014;104: e62–84.

55. Phadke VK, Bednarczyk RA, Salmon DA, Omer SB. Association Between Vaccine Refusal and Vaccine-
Preventable Diseases in the United States: A Review of Measles and Pertussis. JAMA. 2016;315:
1149–1158.

56. Pruitt RH, Kline PM, Kovaz RB. Perceived Barriers to Childhood Immunization Among Rural
Populations. Journal of Community Health Nursing. 1995. pp. 65–72.
doi:10.1207/s15327655jchn1202_1

57. Cavalli-Sforza LL, Feldman MW. Cultural transmission and evolution: a quantitative approach. Monogr
Popul Biol. 1981;16: 1–388.

58. Gangarosa EJ, Galazka AM, Wolfe CR, Phillips LM, Gangarosa RE, Miller E, et al. Impact of anti-vaccine
movements on pertussis control: the untold story. Lancet. 1998;351: 356–361.

59. Ozawa S, Mirelman A, Stack ML, Walker DG, Levine OS. Cost-effectiveness and economic benefits of
vaccines in low- and middle-income countries: a systematic review. Vaccine. 2012;31: 96–108.

60. Kennedy AM, Brown CJ, Gust DA. Vaccine beliefs of parents who oppose compulsory vaccination.
Public Health Rep. 2005;120: 252–258.

61. Leask J. Target the fence-sitters. Nature. 2011;473: 443–445.

62. Hill HA, Singleton JA, Yankey D, Elam-Evans LD, Cassandra Pingali S, Kang Y. Vaccination Coverage by
Age 24 Months Among Children Born in 2015 and 2016 — National Immunization Survey-Child,
United States, 2016–2018. MMWR. Morbidity and Mortality Weekly Report. 2019. pp. 913–918.
doi:10.15585/mmwr.mm6841e2

63. Hill HA, Elam-Evans LD, Yankey D, Singleton JA, Kang Y. Vaccination Coverage Among Children Aged
19–35 Months — United States, 2016. MMWR. Morbidity and Mortality Weekly Report. 2017. pp.
1171–1177. doi:10.15585/mmwr.mm6643a3

64. Omer SB, Salmon DA, Orenstein WA, Patricia deHart M, Halsey N. Vaccine Refusal, Mandatory

Immunization, and the Risks of Vaccine-Preventable Diseases. New England Journal of Medicine.
2009. pp. 1981–1988. doi:10.1056/nejmsa0806477

65. Gilkey MB, Calo WA, Marciniak MW, Brewer NT. Parents who refuse or delay HPV vaccine:
Differences in vaccination behavior, beliefs, and clinical communication preferences. Hum Vaccin
Immunother. 2017;13: 680–686.

66. Hanson KE, Koch B, Bonner K, McRee A-L, Basta NE. National Trends in Parental Human
Papillomavirus Vaccination Intentions and Reasons for Hesitancy, 2010–2015. Clin Infect Dis. 2018;67:
1018–1026.

67. Wong MCS, Wong ELY, Huang J, Cheung AWL, Law K, Chong MKC, et al. Acceptance of the COVID-19
vaccine based on the health belief model: A population-based survey in Hong Kong. Vaccine. 2021;39:
1148–1156.

68. Smith J, Lipsitch M, Almond JW. Vaccine production, distribution, access, and uptake. Lancet.
2011;378: 428–438.

69. Mathieu E, Ritchie H, Ortiz-Ospina E, Roser M, Hasell J, Appel C, et al. A global database of COVID-19
vaccinations. Nat Hum Behav. 2021;5: 947–953.

70. Burki T. Global COVID-19 vaccine inequity. Lancet Infect Dis. 2021;21: 922–923.

71. Solís Arce JS, Warren SS, Meriggi NF, Scacco A, McMurry N, Voors M, et al. COVID-19 vaccine
acceptance and hesitancy in low- and middle-income countries. Nat Med. 2021;27: 1385–1394.

72. Karthigesu SP, Chisholm JS, Coall DA. Do grandparents influence parents' decision to vaccinate their
children? A systematic review. Vaccine. 2018;36: 7456–7462.

73. Gowda C, Dempsey AF. The rise (and fall?) of parental vaccine hesitancy. Hum Vaccin Immunother.
2013;9: 1755–1762.

74. Ellithorpe ME, Adams R, Aladé F. Parents' Behaviors and Experiences Associated with Four
Vaccination Behavior Groups for Childhood Vaccine Hesitancy. Matern Child Health J. 2022;26: 280–
288.

75. Benin AL, Wisler-Scher DJ, Colson E, Shapiro ED, Holmboe ES. Qualitative Analysis of Mothers'
Decision-Making About Vaccines for Infants: The Importance of Trust. Pediatrics. 2006. pp. 1532–
1541. doi:10.1542/peds.2005-1728

76. Montgomery T, Berns JS, Braddock CH 3rd. Transparency as a Trust-Building Practice in Physician
Relationships With Patients. JAMA. 2020;324: 2365–2366.

1016