1

Title: The Cultural Evolution of Vaccine Hesitancy: Modeling the Interaction between Beliefs

and Behaviors

Authors: Kerri-Ann Anderson[a,b] and Nicole Creanza[a,b]

Affiliations:

[a] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA 37212

[b] Evolutionary Studies Initiative, Vanderbilt University, Nashville, TN, USA 37212

Correspondence: nicole.creanza@vanderbilt.edu

Abstract

Health perceptions and health-related behaviors can change at the population level as cultures evolve. In the last decade, despite the proven efficacy of vaccines, the developed world has seen a resurgence of vaccine-preventable diseases (VPDs) such as measles, pertussis, and polio. Vaccine hesitancy, an individual attitude influenced by historical, political, and socio-cultural forces, is believed to be a primary factor responsible for decreasing vaccine coverage, thereby increasing the risk and occurrence of VPD outbreaks. In recent years, mathematical models of disease dynamics have begun to incorporate aspects of human behavior, however they do not address how beliefs and motivations influence these health behaviors. Here, using a mathematical modeling framework, we explore the effects of cultural evolution on vaccine hesitancy and vaccination behavior. With this model, we shed light on facets of cultural evolution (vertical and oblique transmission, homophily, etc.) that promote the spread of vaccine hesitancy, ultimately affecting levels of vaccination coverage and VPD outbreak risk in a population. In addition, we present our model as a generalizable framework for exploring cultural evolution when humans' beliefs influence, but do not strictly dictate, their behaviors. This model offers a means of exploring how parents' potentially conflicting beliefs and cultural traits could affect their children's health and fitness. We show that vaccine confidence and vaccine-conferred benefits can both be driving forces of vaccine coverage. We also demonstrate that an assortative preference among vaccine-hesitant individuals can lead to increased vaccine hesitancy and lower vaccine coverage.

1. __Introduction__

Niche construction is a process in which organisms modify their local environment, thus altering selection pressures on themselves and the other organisms in that environment [1,2]. In *cultural* niche construction, humans modify their cultural environments—such as their beliefs, behaviors, preferences, and social contacts—in ways that subsequently alter evolutionary pressures on themselves and/or their culture [2]. Mathematical models of niche construction have traditionally been used in a biological and ecological context. More recently, this type of model has been expanded to explain the evolution of cultural behaviors, with applications to religion, fertility, and the evolution of large-scale human conflict [2–7]. Cultural niche construction theory recognizes that human evolution is in part directed by human behaviors. As such, using a niche construction framework allows for the exploration of a broader range of complex feedback scenarios resulting from selection pressures that may be caused by—or act upon—non-genetic traits. It may also provide insight into how otherwise deleterious traits become beneficial in certain environments and thus spread in a population [1]. Here, we propose a cultural niche construction model of the interactions between beliefs and behaviors, in which an individual's beliefs influence their behaviors, and these belief-behavior interactions can be affected by and shape the broader cultural landscape. We apply this model to the interactions between vaccine-related beliefs, such as vaccine hesitancy of individual parents, and vaccination behaviors, such as a pair of parents vaccinating their offspring. Modeling the belief-behavior interactions underlying vaccination coverage using a cultural niche construction framework allows us to better understand how vaccination "cultures" are formed and how they can be transformed to promote public health.

Understanding vaccination behaviors is a crucial aspect of preventing infectious disease outbreaks. The implementation of childhood vaccination policies has led to the eradication of smallpox and the elimination of poliomyelitis (polio) in the United States [8–10]. The high efficacy of the measles vaccine, combined with wide vaccine acceptance in developed countries, had resulted in measles previously being targeted for elimination by 2020 [11]. However, over the past decade, there has been a resurgence of vaccine-preventable diseases (VPDs) in developed countries despite the safety and efficacy of vaccines and high overall childhood vaccination rates [12–15]. Vaccine hesitancy, named one of the World Health Organization's ten threats to global health in 2019 [16], is believed to be responsible for decreasing vaccination coverage and thus

increasing the risk of vaccine-preventable disease outbreaks worldwide [17]. Vaccine hesitancy is

a complex and context-specific individual attitude influenced by multiple factors, such as

complacency (the belief that vaccination is unnecessary when the perceived risk of VPDs is low),

convenience (the accessibility and affordability of vaccines), and confidence (the level of trust in

the efficacy and safety of the vaccine, and in the healthcare system) [15,18]. Additionally, anti-

vaccine sentiments are still on the rise despite well-documented vaccine efficacy and safety,

including numerous studies debunking the spurious connection between vaccines and autism

[19] and other anti-vaccination arguments [20]. The spread of these sentiments and disease

outbreak risk are further exacerbated by homophily—the tendency of individuals to choose social

contacts and mates who are similar to themselves [6,7,21,22]. Network-based simulations

suggest that individuals with similar vaccine-hesitant opinions form groups that are more

susceptible to vaccine-preventable diseases, impeding the attainment of herd immunity and

substantially increasing the likelihood of disease outbreak in these clusters [23].

Even though some epidemiological models have begun to include aspects of human behavior

(e.g [24–26]), these models do not typically incorporate the effects of population beliefs and

changing cultural landscapes on disease transmission. For example, established epidemiological

models such as the Susceptible-Infected-Recovered (SIR) model have been modified to include a

vaccination component that can be useful in determining the intensity of intervention needed to

address an epidemic [27]; however these models do not generally address fluctuations in

vaccination rates or lower-than-expected rates of adoption based on cultural factors. One

notable exception is a recent study in which an SIR model of a contagious disease was paired with

another SIR model in which vaccine hesitancy is treated as a "cultural contagion"; this model

showed that the spread of anti-vaccine sentiment could cause epidemics that would otherwise

not have occurred [28]. However, it is still important that we understand how parents' beliefs,

which may differ from one another, interact with their perceptions of the relative risks of disease

and vaccines to shape the decision to vaccinate their children, which in turn affects the future risk

of vaccine-preventable disease outbreaks. Indeed, belief systems can act as the main barrier to

vaccination, as opposed to lack of vaccine access, particularly in wealthier countries [23,29]. For

example, increasing rates of non-medical exemption from vaccines (exemption on the basis of

religious, philosophical, and personal beliefs), have been observed in the United States [30,31].

Without these considerations, models commonly used in public health may be misleading; thus,

understanding and incorporating the underlying health cultures and their evolution, including the

interplay between beliefs and behaviors, will allow us to build more comprehensive and

representative models of vaccination dynamics and better support public health efforts.

In this study, we model the development and spread of vaccine hesitancy and childhood

vaccination through a cultural evolution framework, incorporating the transmission of vaccine

attitudes both from parents and from the community. We aim to assess the dynamic interactions

between beliefs (shaped by social interactions) and behaviors (influenced by these beliefs). Using

vaccine hesitancy and vaccination behaviors as a focal example of belief-behavior interactions,

we explore the situations in which vaccine hesitancy is most likely to spread, potentially reducing

childhood vaccination rates and leading to an increase in vaccine-preventable disease outbreaks.

In addition, we consider that the perception of the relative risks of a disease and its preventive

vaccine can fluctuate based on the prevalence of vaccination [26], such that the population's

vaccination coverage can influence the decision to vaccinate one's children. Finally, we take into

account that the decision to vaccinate a child is often the joint consideration of two individuals

who might have different vaccine attitudes, and we further incorporate homophily (assortative

mating) to understand how social subcultures might influence parental behaviors. Overall, we

propose that a generalizable modeling framework for belief-behavior interactions can help

inform public health strategies by improving our understanding of the cultural dynamics of

vaccine hesitancy.

2.   Methods

To model the evolution of vaccine beliefs and behaviors, we build on the cultural niche

construction framework of [6] to assess the effects of vaccine attitudes on vaccination behaviors

and on the resulting vaccination culture. We use this adapted model to explore how vaccination

patterns evolve in a population when a cultural trait, such as vaccine hesitancy, can influence but

not perfectly predict a behavior, such as vaccinating one's children.

We consider two cultural traits: **V**, a vaccination trait, and **A**, a vaccine attitude trait. Each

trait has two possible states, V$^+$ (vaccinated) or V$^-$ (unvaccinated) and A$^+$ (vaccine confident) or A$^-$

(vaccine hesitant), respectively. Thus, there are four possible phenotypes: V$^+$A$^+$ (type 1:

vaccinated and confident), V$^+$A$^-$ (type 2: vaccinated and hesitant), V$^-$A$^+$ (type 3: unvaccinated and

confident), and V$^-$A$^-$ (type 4: unvaccinated and hesitant), whose population frequencies are

denoted by x_1, x_2, x_3, and x_4, respectively, with $\sum_{i=1}^{4} x_i = 1$.

Table 1: List of parameters, their definitions, and default or initial values

Parameter	Meaning	Parameter	Meaning
V	Vaccination state (V$^+$ vaccinated, V$^-$ unvaccinated)	**A**	Vaccine attitude (A$^+$ confident, A$^-$ hesitant)
m_{ij}	Mating frequencies (given in **Table S1**)	α_k	Assortative mating parameter (homophily) **Default: α_1= 0, α_2= 0**
$B_{m,n}$	Probability that parental pairs vaccinate their children, which depends upon the parents' vaccination states (b_m) and vaccine attitude (c_n) (given in **Table S2**)	C_n	Probability that parental pairs transmit vaccine confidence to their children **Default: C_0= 0.01, C_1= C_2 = 0.5, C_3 = 0.99**
b_m	Probability that parental pairs support offspring vaccination given their vaccination states **Default: b_0= 0.01, b_1= b_2 = 0.5, b_3 = 0.99**	c_n	Probability that parental pairs support offspring vaccination given their vaccine attitude **Default: c_0= 0.01, c_1= c_2 = 0.5, c_3 = 0.99**
σ	Comprehensive selection coefficient for V$^+$, dependent on the population-wide vaccination rate (see **Figure 1**)	σ_{max}	The highest additional benefit that can be conferred by vaccination **Default: σ_{max}= 0.1**
Initial Phenotype Frequencies		x_1(V$^+$A$^+$) = 0.81, x_2 (V$^+$A$^-$) = 0.1, x_3 (V$^-$A$^+$) = 0.07, x_4 (V$^-$A$^-$) = 0.02	

134   The attitude trait (**A**) can influence the dynamics of the vaccination trait (**V**) in two ways:

135 by affecting the likelihood that couples vaccinate their offspring, and by determining with whom

136 each adult will preferentially pair in assortative interactions. The state of the vaccine attitude trait

137 (**A**) informs the value of an assortative mating parameter (α_k), which measures the departure

138 from random mating. We define a 'choosing parent', arbitrarily, as the first member of each

139 mating pair. The choosing parent's **A** state dictates the level of assortative mating, that is, the

140 degree to which an individual of a given **A** state will preferentially mate with another individual of

141 the same state, expressed by parameters α_k where k = {1, 2} and 0≤α_k≤1 (**Table S1**). If the

142 choosing parent is A$^+$, this individual mates preferentially with other A$^+$ individuals with

143 probability α_1, and mates randomly with probability 1−α_1, whereas if the choosing parent is A$^-$,

this individual mates preferentially with other A^- individuals with probability α_2, and mates

randomly with probability $1-\alpha_2$. There are sixteen possible mating pairs from the four

phenotypes described, and we use the notation $m_{i,j}$ to indicate the frequency of a mating

between a choosing parent of type i and the second parent of type j where $i, j = \{1, 2, 3, 4\}$ (**Table**

S1); for example, $m_{1,3}$ represents the mating frequency of V^+A^+ (x_1) and V^-A^+ (x_3).

Since the two traits (**A** and **V**) are transmitted vertically, for each phenotype we must specify

the probability that the mating produces an offspring of that phenotype. The vaccine confidence

trait (A^+) is transmitted with probability C_n, and the vaccine hesitancy trait (A^-) is transmitted with

probability $1-C_n$ (for $n = \{0, 1, 2, 3\}$ as shown in **Tables 2** and **Table S2**). If $C_0 = 0$, two A^- parents

will always produce A^- offspring, and if $C_3 = 1$, two A^+ parents will always produce A^+ offspring.

However, if $C_0 > 0$, two A^- parents can produce A^+ offspring at some probability, and similarly if C_3

< 1, two A^+ parents can produce A^- offspring with some probability.

Transmission of vaccination (V^+ with probability $B_{m,n}$ for $m, n = \{0, 1, 2, 3\}$; **Table 1**) is more

complex, since parents' vaccine attitudes (**A**), in addition to their own vaccination states (**V**), can

influence their behavior in vaccinating their offspring via a set of "influence parameters" that

inform vaccination probabilities. The probability that each mating pair produces an offspring with

the V^+ trait (i.e. vaccinates their offspring) is a scaled product of the influence of parental

attitudes (c_n for $n = \{0, 1, 2, 3\}$) and the influence of parental vaccination states (b_m for $m = \{0, 1, 2,$

$3\}$) (**Tables 2 and Table S2**). For example, for mating pair $V^+A^+ \times V^+A^-$, the combined vaccination

states ($V^+ \times V^+$) will influence vaccination behavior by b_3, and the combined attitude states, ($A^+ \times$

A^-), will influence vaccination behavior by c_2. Therefore, a $V^+A^+ \times V^+A^-$ mating will produce a V^+

offspring with probability $B_{3,2} = c_2\left(\frac{1+b_3}{2}\right)$; this pair will also produce an A^+ offspring with

probability C_2 based on their combined attitude states. Thus, according to the model, this pairing

will produce a V^+A^+ offspring with probability $B_{3,2}C_2$ and a V^+A^- offspring with probability

$B_{3,2}(1-C_2)$. We note that assortative mating ($\alpha_k > 0$) will increase the frequency of matings between

individuals that share an attitude trait, with these non-random interactions in turn skewing

vaccination outcomes toward those of same-state couples (via c_0 and c_3).

Transmission and influence probabilities are constant throughout a single simulation, with

values ranging from 0 to 1. At default settings, the influence parameters b_m and c_n, and the

transmission parameter C_n would take the following values: $C_0, b_0, c_0 = 0.01$; $C_1, C_2, b_1, b_2, c_1,$

$c_2 = 0.5$; and $C_3, b_3, c_3 = 0.99$. In our model, the influence of parental vaccine beliefs (c_n) is greater

than the influence of their own vaccination status (b_n) on their likelihood of vaccinating their

offspring, so offspring vaccination is guaranteed at some probability only if $c_n > 0$.

Table 2: Presence (+) and absence (−) subscript assignments. Demonstrating the trait presence (+) and
absence (−) combinations associated with m, n subscripts. For example, the + × − combinations is
associated with m and n subscript value 2: an $A^+ \times A^-$ pairing transmits A^+ at probability C_2. This rule applies
to parameters C_n, b_m, $B_{m,n}$, c_n, as shown in **Table S2**.

Subscript Value (m, n; e.g. b_m, C_n)	Associated Pairing (e.g. V × V, A × A)
0	− × −
1	− × +
2	+ × −
3	+ × +

Transmission and influence probabilities are constant throughout a single simulation, with

values ranging from 0 to 1. At default settings, the influence parameters b_m and c_n, and the

transmission parameter C_n would take the following values: C_0, b_0, $c_0 = 0.01$; C_1, C_2, b_1, b_2, c_1,

$c_2 = 0.5$; and C_3, b_3, $c_3 = 0.99$. In our model, the influence of parental vaccine beliefs (c_n) is greater

than the influence of their own vaccination status (b_n) on their likelihood of vaccinating their

offspring, so offspring vaccination is guaranteed at some probability only if $c_n > 0$.

The cultural selection pressure on vaccination is given by the parameter σ, such that the

frequency of the V^+A^+ and V^+A^- phenotypes are multiplied by 1+σ after vertical cultural

transmission has occurred. At the end of each timestep, the frequency of each phenotype is

divided by the sum of all four frequencies, ensuring that the frequencies sum to 1. This cultural

selection coefficient is implemented in the same way as a selection coefficient in a population-

genetic model, but unlike the latter, it is structured to encompass both biological fitness and

cultural selection pressures, including perceived risks or benefits of the vaccine itself, personal

cost-benefit analyses of preventative health behaviors, and the structural or societal-level factors

influencing vaccination rates [32,33]. This parameter modulates whether there are more or fewer

vaccinated individuals than expected: in other words, when σ>0, vaccinated individuals are more

common in a set of offspring than would be expected strictly based on the beliefs and vaccination

statuses of their parents. We calculate σ in each timestep as a function of the current vaccination

coverage (frequency of V$^+$, i.e. $x_1 + x_2$), and in each simulation we specify σ_{max} as the maximum

cultural selection pressure of getting vaccinated ($-1\leq\sigma_{max}\leq1$) (see the cultural selection coefficient

function in **Figure 1**). This function was constructed by fitting a curve to pre-specified conditions

that incorporate assumptions from evolutionary game theory (e.g. that herd immunity decreases

the incentive to vaccinate [34]): we assume that when vaccination coverage is low, the real and

perceived benefits of vaccination are highest, and thus, the cultural selection pressure is near

σ_{max}, however, as vaccination coverage increases, the perceived benefits of vaccination decrease

and the cultural selection pressure is reduced (**Figure 1**).

Figure 1: Cultural selection coefficient function. The cultural selection coefficient function was

constructed by fitting a curve to specified conditions, and considers both health and non-health related

effects. The selection coefficient (σ; vertical axis) is dependent on the frequency of vaccinated individuals

(V$^+$) in the population (horizontal axis). σ_{max} is the maximum cultural selection coefficient associated with

being vaccinated. Perceived vaccine benefit is reduced as vaccination coverage increases, since the

negative effects of the disease will be less apparent. Note: Of the σ_{max} values shown, only $\sigma_{max} = 0.1$ allows

the cultural selection pressure to be either positive or negative at a given timepoint depending on the

frequency of vaccination.

Thus far, we have described vertical cultural transmission from parent to offspring. The

model also incorporates a second phase with oblique cultural transmission (i.e. influence from

non-parental adults), in which individuals can change their inherited vaccine attitudes (**A**) due to

influence from other adults in the population. There are two probabilities associated with

attitude modulation: the probability that a vaccine hesitant (A⁻) individual adopts the vaccine

confident (A⁺) state (A⁻ to A⁺ transition probability, given by $A_{\rightarrow Confident}$ in **Figure 2**), and the

probability that an A⁺ individual adopts the A⁻ state (A⁺ to A⁻ transition probability, given by

$A_{\rightarrow Hesitant}$ in **Figure 2**). As with the strength of cultural selection (σ) described previously (**Figure**

1), the probability that offspring change their vaccine attitude is a function of the V⁺ frequency in

the population, constructed according to similar assumptions given in **Figure 2**. As the frequency

of vaccinated individuals (V⁺) increases in the population, vaccine-confident individuals (A⁺) are

more likely to become hesitant ($A_{\rightarrow Hesitant}$ probability increases) and vaccine-hesitant individuals

(A⁻) are less likely to become confident ($A_{\rightarrow Confident}$ probability decreases).

Figure 2: Attitude Transition Probability Function. Attitude transition probability functions were
constructed by fitting a curve to specified values based on the assumptions pictured. Attitude transition
probability (vertical axis) is a function of the vaccination frequency in the population (V⁺; horizontal axis).
The probability that a vaccine hesitant individual adopts vaccine confidence (A⁻ to A⁺ transition probability,
shown in black) is determined by the function $A_{\rightarrow Confident}$, and the probability that a vaccine confident
individual adopts vaccine hesitancy (A⁺ to A⁻ transition probability, shown with a blue dashed line) is
determined by the function $A_{\rightarrow Hesitant}$.

To compute the frequency of a given phenotype in the next iteration, we sum the

probability that each mating pair produces offspring of that phenotype over each of the sixteen

possible mating pairs. Cultural selection (σ), described above, then operates on offspring with the

V$^+$ trait. The full recursions, giving x_i' phenotype frequencies in the next iteration in terms of x_i in

the current iteration, are given in **Text S1**. If x_i' is equal to x_i, the system is at equilibrium. As we

do not incorporate a birth-death process or population asynchrony in this model, iterations in the

discrete-time format of our model should not be strictly interpreted as years or generations. We

instead interpret each iteration broadly as a timeframe in which the specified cultural

interactions could occur, which varies among individuals, populations, and cultures. Unless

otherwise stated, the model is initialized with phenotypic frequencies based on United States

data: x_1 (frequency of V$^+$A$^+$) = 0.81, x_2 (V$^+$A$^-$) = 0.1, x_3 (V$^-$A$^+$) = 0.07, x_4 (V$^-$A$^-$) = 0.02. These

frequencies were estimated using reports of Measles-Mumps-Rubella (MMR) vaccination rates

and estimates of vaccine attitude frequencies obtained from various sources in the literature

[35,36] and the Centers of Disease Control ChildVax database [37,38].

3.   Results

261   To test our model, we first initialized a population with a set of phenotype frequencies

and examined the changes in these frequencies over time with a given set of parameters (**3.1**).

Then, we evaluated the effects of each parameter by running simulations at multiple parameter

combinations and recording the population frequencies of each phenotype once the system

approached an equilibrium (**3.2–3.4**). In our first set of simulations (**3.1–3.3**), we include only

vertical transmission dynamics, i.e. only parent-to-offspring transmission, varying parameter

values in turn to test their effects on population vaccination behavior and attitudes. In the

vertical transmission phase of the model, parents choose whether to vaccinate their offspring

(i.e., transmit V$^+$) or to not vaccinate (V$^-$), and parents also transmit a vaccine attitude

(confidence, A$^+$, or hesitancy, A$^-$), each with a specified probability given the phenotypes of the

parents. The parental attitude state, vaccination status, assortative mating levels, and cultural

selection parameters interact to affect vaccination coverage (frequency of V$^+$ in the population)

and vaccine confidence (frequency of A$^+$).

3.1.   *Temporal dynamics of vaccine-related beliefs and behaviors*

To test whether the equilibrium phenotype frequencies were sensitive to starting

frequencies, we plotted the dynamics of each phenotype over time at default parameters (given

in **Table 1**). For each set of initial phenotype proportions tested, the phenotype frequencies in

the population quickly adjusted to approach equilibrium values and then gradually plateaued to a

stable equilibrium (vertical transmission: **Figure 3 and Figure S1**, vertical+oblique transmission:

Figure S2. This demonstrates that equilibrium frequencies of vaccination coverage and vaccine

confidence are determined by the parameter conditions rather than by the initial frequencies

themselves.

Figure 3: Equilibrium frequencies are determined by the parameter space, not by initial frequencies. The

change in each of the four phenotype frequencies and the total V^+ and A^+ frequencies (vertical axis) over

100 iterations of the model **with vertical transmission only** (horizontal axis). **Top Row:** Initial frequencies

are varied, such that we begin each simulation with a different phenotype at an initial high frequency

(0.81): V^+A^+ in panel **A**, V^+A^- in panel **B**, V^-A^+ in panel **C**, V^-A^- in panel **D**; the remaining phenotypes are set

to lower frequencies (0.1, 0.07, 0.02). See **Figure S1** for a full listing of these initial frequencies. **Bottom

Row:** The maximum cultural selection coefficient (σ_{max}) is varied: **E**. $\sigma_{max} = -0.1$; **F**. $\sigma_{max} = 0$; **G**. $\sigma_{max} = 0.1$; **H**.

$\sigma_{max} = 0.5$. Cultural selection against vaccinated individuals increases the frequency of V^-A^-, decreasing the

other frequencies (**E**), whereas increased cultural selection favoring vaccinated individuals increases V^+A^+

frequencies while decreasing the other frequencies (**F, G, H**). In all panels, the remaining parameters are

held at default values (**Table 1**).

When two parameters in particular are varied—maximum cultural selection (σ_{max}) or

confidence transmission ($C_1 = C_2$)—we observe a trade-off between the V^-A^- phenotype, which

dominates at lower values of these parameters, and the V^+A^+ phenotype, which dominates at

higher values (**Figures 3-4**). Interestingly, the "conflicting" phenotypes (when an individual's

attitude toward vaccinating their children does not match their own vaccination state: V^-A^+ and

V^+A^-) are present at their highest frequencies at neutral cultural selection (σ_{max} = 0, **Figure 3F**)

and/or neutral confidence transmission ($C_1 = C_2$ = 0.5, **Figure 4B**). Vaccinated individuals have the

same fitness regardless of their attitude (i.e. V^+A^+ bears the same selection pressure as V^+A^-), so it

is worth noting that at higher levels of confidence transmission and cultural selection, V^+A^+

increases in frequency but V^+A^- decreases in frequency (compare **Figure 3F-G, Figure 4B-C**). This

pattern seems to reflect their differing likelihoods of vaccinating their offspring: across all

possible partners, vaccinated but vaccine-hesitant parents (V^+A^-) are less likely to vaccinate their

offspring than vaccinated and vaccine-confident parents (V^+A^+), resulting in more V^- offspring.

Thus, when V^+ is favored by cultural selection, there is indirect selection against the vaccinated

but vaccine-hesitant (V^+A^-) phenotype (**Figure 3E-H**). Similarly, indirect selection against V^-A^+

occurs when V^- is favored by cultural selection (**Figure 3E**): compared to **Figure 3F**, we observe an

increase in V^-A^- individuals but a decrease in V^-A^+ individuals, who because of their vaccine

confidence have more V^+ offspring, which are culturally disfavored in this environment. When

cultural transmission from non-parental adults (oblique transmission) was included, described in

following sections, we observed similar patterns, but the final equilibria were more likely to be

polymorphic, with vaccinated, unvaccinated, confident, and hesitant phenotypes stabilizing at

more moderate frequencies than they would have with only vertical transmission (Compare

Figures 3-4 to Figures S3-S4).

Low confidence transmission ($C_1 = C_2$ = 0.1, **Figure 4A**) increases the frequency of vaccine

hesitancy (A^-) in the population over time, increasing the probability that more couples choose

not to vaccinate their offspring. However, the increase in vaccine hesitancy does not occur

equally in vaccinated and unvaccinated individuals: A^- frequency may increase overall in this

environment, but V^+A^- frequencies are lower and V^-A^- frequencies are higher (compared to

Figure 4B-C and **Figure S4**). At neutral confidence transmission probabilities (i.e. when couples

with one confident and one hesitant parent are equally likely to transmit either attitude), there is

a higher chance that the vaccinated but vaccine-hesitant (V^+A^-) phenotype is replenished.

However, if vaccine confidence is highly transmitted ($C_1 = C_2$ = 0.8), the V^+A^- frequency will be

reduced, as this phenotype is more likely to produce A^+ offspring than A^-, thus increasing V^+A^+

phenotype frequencies in the population (**Figure 4** and **Figure S4**). If we turn to the other

conflicting phenotype, unvaccinated but vaccine-confident (V^-A^+) individuals become more

common when A$^+$ increases in frequency in the population as $C_1 = C_2$ increases from 0.1 to 0.5

(**Figure 4** and **Figure S4**). In contrast, higher vaccine confidence transmission ($C_1 = C_2 = 0.8$) can

lead to a vaccination-promoting environment in which V$^-$ frequencies are reduced over time; thus

the V$^-$A$^+$ phenotype becomes rare and V$^+$A$^+$ predominates (**Figure 4** and **Figure S4**).

Figure 4: **Temporal Effects of Confidence Transmission**. The change in phenotype frequencies over 50
iterations as vaccine confidence transmission in mixed-attitude couples ($C_1 = C_2$) is varied (**A**. $C_1 = C_2 = 0.1$;
B. $C_1 = C_2 = 0.5$; **C**. $C_1 = C_2 = 0.8$) **with vertical transmission only,** while other parameters are held at default
values (**Table 1**). The population equilibrates at over 90% A$^-$V$^-$ at low confidence transmission (A).
Increasing the probability of confidence transmission results in less vaccine hesitancy and, in turn, higher
vaccination frequencies (V$^+$A$^+$).

3.2.   *Parent-to-Offspring Interactions (Simulations with vertical transmission only)*

Since our assessment of the temporal dynamics (**3.1**)demonstrated that our simulations

approach stable equilibria, we then modulated different sets of parameters and recorded the

phenotype frequencies at equilibrium, generating heat maps showing the results across a range

of parameters. In the first of these, we tested the relationship between vaccination probability

and vaccine confidence transmission. To directly alter vaccination probabilities while still

accounting for the couple's vaccine attitudes, we set ranges of values for $B_{m,n}$ that vary along the

horizontal axis of **Figure 5**, with the vaccination probability for two hesitant parents (e.g. $B_{0,0}$) on

the lower end of the range and the vaccination probability for two confident parents (e.g. $B_{3,3}$) on

the higher end of the range (**Table S3**). Confidence transmission probabilities are also structured

in this "range shift" manner (**Figure 5A-B, Table S3**). If we vary both confidence transmission

parameters and vaccination probability parameters by implementing range shifts in both C_n and

$B_{m,n}$, we observe a positive interaction between confidence transmission and vaccination

probability: vaccination coverage increases as either of these parameters are increased (**Figure**

5A).

Figure 5: Vaccination coverage levels are determined by an interaction between confidence transmission and vaccination probability. Heatmaps showing final vaccination coverage (**A, C**) and corresponding vaccine confidence (**B, D**) after 100 time-steps with **no oblique transmission**. In **A** and **B**, confidence transmission probabilities (C_n) are set within the range indicated on the vertical axis, and vaccination probabilities ($B_{m,n}$) are set within the range indicated on the horizontal axis with $B_{0,0}$, $B_{1,0}$, $B_{2,0}$ and $B_{3,0}$ taking the lowest value and $B_{3,3}$ taking the highest value (**Table S3**). In **C** and **D**, confidence transmission in mixed-attitude couples ($C_1 = C_2$) is varied along the vertical axis, while the vaccination probabilities ($B_{m,n}$) are set within the range indicated on the horizontal axis as in **A** and **B**. (**Table S3**). We show increased equilibrium vaccination coverage with increasing vaccination probability and confidence transmission probability ranges, while confidence levels are primarily dictated by proportion of the population transmitting confidence or hesitancy.

However, couples with mixed vaccination and/or attitude states ($V^+ \times V^-$, $A^+ \times A^-$) are

assumed to be more variable in their decision to vaccinate their offspring than parents who share

the same state. Thus, in the simulations that follow, we primarily modulated the specific

probabilities associated with these mixed-state pairings. In **Figure 5C-D**, we varied vaccination

probabilities ($B_{m,n}$) across the full range of individuals but modulated confidence transmission

probabilities only for mixed-attitude couples ($C_1 = C_2$), i.e. those with one vaccine-hesitant parent

and one vaccine-confident parent. In these tests, we observe increasing equilibrium vaccination

coverage as $B_{m,n}$ probabilities increase, with higher coverage in high-confidence transmission

environments (**Figure 5C-D**).

In both aforementioned simulations (**Figure 5**), we confirm vaccination coverage levels are

determined by an interaction between confidence transmission and vaccination probability,

whereas confidence levels are dictated primarily by levels of confidence transmission. In sum, the

degree to which parents with mixed vaccine-hesitant and vaccine-confident attitudes transmit

vaccine confidence instead of vaccine hesitancy to their offspring is a key factor in determining

population trait majorities which can drastically shift population dynamics.

We compared the effects of varying the confidence transmission probabilities for mixed-

attitude couples (C_1 and C_2) in combination with multiple factors: 1) the maximum cultural

selection coefficient (σ_{max}) (**Figure 6A-B**), 2) the vaccination influence parameters b_1 and b_2

(**Figure 6C-D**), 3) the attitude influence parameters c_1 and c_2 (**Figure 6E-F**), and 4) the vaccination

probabilities of couples with mixed states, $B_{1,1}$, $B_{1,2}$, $B_{2,1}$, $B_{2,2}$ (**Figure 6G-H**). In each examination,

we observed a C_n threshold: there is a mid-range value of C_n at which vaccination coverage and

vaccine confidence traits are polymorphic (i.e. both forms of each trait coexist in the population),

separating definitive high ($\gtrapprox 80\%$) and low ($\lessapprox 30\%$) levels of vaccination coverage and confidence.

This C_n threshold value is more sensitive to σ_{max} than to b_m, c_n, or $B_{m,n}$: the threshold value is

lowered as σ_{max} increases (diagonal line in **Figure 6A-B**). Although vaccination probability ($B_{m,n}$) is

dependent on both c_n, the influence of parental vaccine attitude, and b_m, the influence of

parental vaccination state (**Table S2**), modulating either type of influence of mixed-state parents

has little effect on the level of vaccination coverage and negligible effects on confidence levels at

each non-threshold C_n (**Figure 6C-F**).

Interestingly, direct modulation of the mixed-state couple vaccination probability ($B_{1,1} =$

$B_{1,2} = B_{2,1} = B_{2,2}$) also has little power in affecting coverage and confidence levels at equilibrium

(**Figure 6G-H**). We hypothesize that predominantly high or predominantly low confidence

transmission within a population reduces the occurrence of "mixed-state" pairings, i.e. if the

majority of the population becomes confident or hesitant, there are fewer confident-hesitant and

vaccinated-unvaccinated pairings. Thus, the effect of modulating mixed-state vaccination

probabilities ($B_{1,1}$, $B_{1,2}$, $B_{2,1}$, $B_{2,2}$) is significantly minimized as these couples approach low

frequencies in the population, and confidence transmission dominates the vaccination patterns.

Figure 6: **Vaccine Confidence Transmission Dictates Vaccination Coverage and Confidence Levels**
Heatmaps showing final vaccination coverage and vaccine confidence after 100 time-steps with **no**
oblique transmission, only parent-to-offspring transmission. The top row (**A, C, E, G**) shows vaccination
coverage (i.e. frequency of V^+ in the population) with low coverage in blue and high coverage in yellow;
the bottom row (**B, D, F, H**) shows the corresponding final vaccine confidence (i.e. frequency of A^+), with
low confidence in red and high confidence in yellow. Unless varied on the horizontal or vertical axis, other
parameters are set to the default values given in **Table 1**. In our model, parents' likelihood of vaccinating
their children depends on both their vaccination state and their attitude state. This figure shows that the
strength of parental transmission of vaccine confidence (C_n) has a much stronger effect on the equilibrium
levels of both vaccine coverage (V^+) and confidence (A^+) than other parameters: the maximum cultural
selection coefficient, σ_{max} (**A,B**), the influence of parental vaccination state, b_m (**C, D**), the level of influence
of parental vaccine attitudes on their vaccination behaviors, c_n (**E,F**), and the probability that mixed-state
parents vaccinate their offspring $B_{m,n}$ (**G,H**).

Next, we hold vaccine confidence transmission (C_n) at default probabilities, reminiscent of

Mendelian transmission, such that two vaccine confident or two vaccine hesitant parents

predictably transmit their vaccine attitude, and parents with differing vaccine attitudes each have

a ~50% chance of transmitting their own state, e.g. C_0 near 0, C_1 and C_2 at 0.5, C_3 near 1 **(Table 1)**.

Figure 7: **Cultural Selection Influences Vaccination Coverage and Vaccine Confidence.** Heatmaps showing final vaccination coverage (**A, C, E**) and final vaccination confidence (**B, D, F**) after 100 time-steps with **no oblique transmission**, only parent-to-offspring transmission. As in previous figures, parameters not varied here are given in **Table 1**. Parents' likelihood of vaccinating their children depends on both their vaccination state and their attitude state. At default probabilities of vaccine confidence transmission (C_n values in **Table 1**), these figures show that modulating the maximum cultural selection coefficient affects the equilibrium levels of vaccination coverage and vaccine confidence across the range of specified parameters: parental vaccination state influence, b_m (**A, B**), parental attitude state influence, c_n (**C,D**), and offspring vaccination probability, $B_{m,n}$ (**E,F**). Unless directly modulated (as in panels **E-F**), $B_{m,n}$ varies with b_m and c_n: $B_{m,n} = c_n\left(\frac{1+b_m}{2}\right)$.

We then varied cultural selection in combination with vaccination-associated probabilities (b_m, c_n, $B_{m,n}$). With C_n held constant, cultural selection (σ_{max}) is the primary factor determining vaccination coverage and confidence levels (**Figure 7**). Raising the maximum cultural selection coefficient increases the equilibrium level of vaccination coverage and vaccine confidence across various levels of vaccination state influence (b_m) (**Figure 7A-B**), vaccination attitude influence (c_n) (**Figure 7C-D**), and vaccination probability ($B_{m,n}$) (**Figure 7E-F**). Unlike in **Figure 6**, vaccine confidence does

not always mirror vaccination coverage across all levels of attitude influence (c_n) or vaccination

probabilities. Instead, vaccine confidence levels decline with increased c_n and increased $B_{m,n}$ for

$\sigma_{max} \lessgtr 0.3$ (**Figure 7D, F**), as well as for both increased c_n and increased b_m (**Figure S5**). This

dynamic is interesting as these parameters inform vaccination behavior, hinting that high

vaccination rates could reduce a populations' expected vaccine confidence. Vaccination coverage

and vaccine confidence remain low when cultural selection does not favor vaccination ($\sigma_{max} \lessgtr 0$),

i.e. parents vaccinate their children at or below the levels expected based on cultural

transmission rates.

458
3.3.   *Offspring can Change their Inherited Hesitancy State (Vertical and Oblique Dynamics)*

Increased exposure to the attitudes of the broader community (i.e. oblique cultural

transmission from non-parental adults in the population) could influence and change vaccination

beliefs inherited in childhood. Therefore, we next included these oblique effects in our model to

understand how they might modulate vaccine confidence and vaccination coverage levels. In the

oblique transmission phase of the model, offspring can change their vaccine attitude with some

probability based on the frequency of vaccination in the population (**Figure 2**). Thus, in addition

to the vertical transmission of attitudes and behaviors, phenotype frequencies are further

affected by the probability that adult offspring change their attitude (i.e. transition from vaccine

confident (A^+) to hesitant (A^-) and vice versa). By modulating the attitude transition probabilities

according to the vaccination coverage, we assume that when vaccine coverage (V^+ frequency, $x_1 +$

x_2) is low, disease occurrence is high and the negative effects of the disease are experienced

widely, thus the benefits of being vaccinated (and the costs of not being vaccinated) are more

evident [39,40]. As vaccination coverage (V^+) increases in the population, and thus disease

occurrence is low, the benefits to being vaccinated are less obvious, while low-probability costs

such as adverse reactions become more apparent and could be perceived as being riskier than

the disease itself.

The addition of oblique dynamics produces a pattern of vaccination coverage and vaccine

confidence similar to that of simulations run with solely vertical transmission (**Figure 6 and Figure**

7 compared to Figure 8 and Figure 9, and Figures 3-4 compared to **Figures S2-4**)—the level of

(vertical) vaccine confidence transmission still largely determines the level of vaccination

coverage and vaccine confidence (**Figure 8**). However, oblique cultural influences expanded the

polymorphic space, resulting in a wider range of intermediate C_n in which the different states of

each trait (vaccinated, unvaccinated, confident, and hesitant) are present in the population in

roughly equal proportions. In other words, there is a wider horizontal stripe of moderate values

between the definitively high and definitively low equilibrium frequencies in **Figure 8** than in

Figure 6). The addition of oblique transmission appears to lead to less polarized results overall,

moving the equilibrium levels of vaccination coverage and vaccine confidence away from extreme

values.

Figure 8: Vaccine confidence transmission dictates vaccination coverage and confidence levels (with

oblique transmission). Heatmaps showing final vaccination coverage (i.e. frequency of V^+ in the

population, with low coverage in blue and high coverage in yellow (**A, C, E, G**)) and final vaccine

confidence (i.e. frequency of A^+, with low confidence in red and high confidence in yellow (**B, D, F, H**))

after 100 time-steps in which oblique transmission of vaccine attitude can occur after parent-to-offspring

transmission. The likelihood that individuals change their vaccine beliefs depends on the current

vaccination coverage of the population (**Figure 2**). Unless varied on the horizontal or vertical axes, other

parameters are set to the default values given in **Table 1**. Parents' likelihood of vaccinating their children

depends on both their vaccination state and their attitude state; these figures show that the strength of

parental transmission of vaccine confidence (C_n) has a much stronger effect on the equilibrium levels of

both vaccine coverage (V^+) and confidence (A^+) than other tested parameters do: the maximum cultural

selection coefficient, σ_{max}, (**A,B**), the influence of parental vaccination state, b_m, (**C, D**), the level of

influence of parental attitudes on their vaccination behaviors, c_n, (**E,F**), and offspring vaccination

probability, $B_{m,n}$ (**G,H**).

With neutral confidence transmission ($C_1 = C_2 = 0.5$), we also observe an expansion of the

polymorphic space when we modulate cultural selection (σ_{max}) alongside the influence and

transmission parameters (**Figure 9**). Interestingly, in the cultural environment defined by this

parameter space, we observe a pattern that deviates from the expected association between

high vaccine confidence and high vaccination coverage: as the influence of vaccine attitudes (c_n)

and vaccination probabilities ($B_{m,n}$) increase (**Figure 9C-F**), the population's equilibrium

vaccination coverage increases while its vaccine confidence decreases. This pattern persisted

across all tested levels of maximum cultural selection (σ_{max}) (**Figure 9C-F**). In other words, we

observe higher levels of confidence when the influence of vaccine attitude is low (**Figure 9D**) and

vaccination probabilities are low (**Figure 9F**) than we do at higher values.

Figure 9: Cultural selection influences vaccination coverage and vaccine confidence (with oblique
transmission). Heatmaps showing final vaccination coverage (**A, C, E**) and final vaccination confidence (**B,**
D, F) after 100 time-steps **with oblique transmission**. As in previous figures, parameters not varied are
given in **Table 1**. Parents' likelihood of vaccinating their children depends on both their vaccination state
and their attitude state. At default probabilities of vaccine confidence transmission (C_n), these figures
show that modulating the maximum cultural selection coefficient affects the equilibrium levels of
vaccination coverage and vaccine confidence across the range of specified parameters: parental
vaccination state influence, b_m (**A, B**), parental attitude state influence, c_n (**C,D**), and offspring vaccination
probability, $B_{m,n}$ (**E,F**). Unless directly varied (as in panels **E-F**), $B_{m,n}$ varies as b_m and c_n are varied, as shown
in **Table 1**.

Figure 10: **The influence of parental traits on vaccination coverage and vaccine confidence at different**

levels of cultural selection. Heatmaps showing final vaccination coverage (**A, C**) and final vaccination

confidence (**B, D**) after 100 timesteps **with oblique transmission**. We modulate the interaction between

vaccination state influence (b_m; vertical axis) and attitude influence (c_n; horizontal axis) at various

maximum cultural selection coefficients: σ_{max} = 0 (**A, B**) and σ_{max} = 0.1 (**C, D**). As in previous figures,

unvaried parameters are given in **Table 1**. Vaccination frequency increases as both influence probabilities

increase and vaccination confidence decreases as both influence probabilities increase.

We explored the interaction between the influence parameters, b_m and c_n, at various

maximum cultural selection coefficients (σ_{max}) (**Figure 10**). Vaccination coverage and vaccine

confidence equilibrate at mid-range frequencies (between 0.3 and 0.8) across the range of b_m and

c_n, indicating that these trait frequencies are not particularly sensitive to either parameter.

Cultural selection favoring vaccination increases the equilibrium level of vaccination coverage and

vaccine confidence (**Figure 10 and Figure S6**). The most notable deviation between equilibrium

confidence and vaccination frequencies occurs at the intersection of the highest influence

parameter values (b_m and c_n), circumstances in which the parents' vaccination states and vaccine

attitudes overwhelmingly support offspring vaccination. In this top right section of the heat maps,

vaccination coverage is high while vaccine confidence is lower, indicating a behavioral pattern in

which mixed-trait couples are more inclined to vaccinate their offspring than transmit vaccine

confidence. Overall, the addition of oblique transmission to a population that would otherwise

equilibrate at high vaccination coverage (**Figure S5**) leads to increased attitude transition to

vaccine hesitancy and subsequently lower vaccine coverage.

3.4.   _Mating Preferences_

We hypothesized that mating preference (assortative mating) could modulate belief and

behavior dynamics and thus the vaccination coverage and confidence levels in the population. If

individuals are more likely to pair with individuals of the same vaccine attitude, such that same-

attitude couples become more common and mixed-attitude couples are less common, the

parameter values for mixed-attitude couples may have less impact on vaccination coverage and

confidence dynamics. Therefore, we analyzed the interaction between A^+ homophily (with α_1

indicating the preference of A^+ individuals for other A^+ individuals) and A^- homophily (with α_2

indicating the preference of A^- individuals for other A^- individuals) at various σ_{max} levels. When

vaccine attitudes are transmitted both from parent to offspring and between unrelated

individuals (vertical and oblique transmission) and there is neither cultural selection for nor

against being vaccinated ($\sigma_{max} = 0$), we observe a threshold region at roughly equal mating

preferences ($\alpha_1 \approx \alpha_2$; diagonal lines in **Figure 11C-D**); above this boundary (when $\alpha_1 > \alpha_2$)

vaccination coverage and confidence are much higher than below this boundary (when $\alpha_1 < \alpha_2$) .

When cultural selection explicitly does not favor vaccination (e.g. $\sigma_{max} = -0.1$, **Figure 11A-B**), low

vaccination coverage and confidence can occur even when there are more vaccine confident

couples in the population than hesitant couples ($\alpha_1 > \alpha_2$). Likewise, if cultural selection favors

being vaccinated ($\sigma_{max} > 0$, **Figure 11E-H**), the threshold between high and low equilibrium values

is shifted, such that high coverage and high confidence levels can potentially be attained even

when vaccine hesitant individuals preferentially pair with each other more than vaccine confident

individuals do ($\alpha_1 < \alpha_2$). We observe qualitatively similar patterns when vaccine attitudes are only

transmitted from parent to offspring (**Figure S7**); as we have previously observed in **Figures 8-10**,

the addition of oblique transmission leads to a broader polymorphic region than vertical

transmission alone. These patterns illustrate two overarching themes: 1) preferential interactions

between individuals with similar vaccine beliefs can dramatically shift the equilibrium levels of

vaccination coverage and confidence with all other parameters remaining equal, and 2) the actual

and perceived quality and efficacy of the vaccine are important to determining vaccine

acceptance.

583
Figure 11: Homophily between individuals with similar vaccine beliefs can shift equilibrium frequencies
of both vaccination coverage and confidence. Heatmaps showing final vaccination coverage (**A, C, E, G**)
and final vaccine confidence (**B, D, F, H**) after 100 timesteps **with oblique transmission**. As in previous
figures, unspecified parameters are given in **Table 1**. As vaccine-hesitant individuals (A⁻) increasingly
prefer to pair with one another (α_2; horizontal axis), vaccine-confident individuals (A⁺) must also
preferentially interact to maintain high vaccine coverage (α_1; vertical axis); this tradeoff is modulated by
the cultural selection pressures on vaccination (σ_{max} = -0.1 (**A, B**), σ_{max} = 0 (**C, D**) and σ_{max} = 0.1 (**E, F**), σ_{max} =
0.5 (**G, H**)).

4. **Discussion**

In this manuscript, we present a simplified model of a complex process: the spread of vaccine

attitudes and their effects on childhood vaccination frequency in a population. Increasing and

maintaining sufficient vaccination coverage to combat disease is more complex than simply

increasing vaccine availability or providing accurate information. A number of factors affect a

person's vaccine-related beliefs and parents' decision to vaccinate their children, including their

history with vaccinations and perception of the disease and vaccine effects. As such, it is

important that we understand how these personal factors can shape vaccination cultures and

thus affect public health. Using a cultural niche construction framework, we modeled the

transmission of vaccine attitudes and vaccination behavior in a variety of circumstances and

measured the resulting levels of vaccination coverage and vaccine confidence in the population.

Using this novel approach of modeling dynamically interacting beliefs and behaviors, we are able

to explore the interplay of cultural factors that drive vaccine attitudes and vaccination behavior,

providing insight into how vaccination cultures are formed, maintained, and evolve.

Our model demonstrates that the cultural landscape—current policies, transmission patterns

of beliefs and behaviors, etc.—can be more predictive of future levels of vaccine coverage and

confidence than current coverage and confidence levels in the population. Our simulations each

approached a stable equilibrium, and in general we could infer that a population with high

vaccination coverage will have low rates of vaccine hesitancy and vice versa. Further, our model

shows vaccine confidence transmission (C_n) to be the parameter that most strongly determines

vaccination coverage and confidence levels. That is, even though parents' decision to vaccinate

their children is based on both their level of confidence in vaccines and a consideration of their

own vaccination status, the probability of transmitting vaccine-positive attitudes is a stronger

predictor of future vaccination coverage than the probability of vaccination itself (**Figures 6 and**

8). Finally, our simulations also suggest that a pro-vaccination health culture can be undermined

by a vaccine hesitancy "echo chamber", possibly formed by a higher degree of preferential

assortment (homophily) among vaccine-hesitant individuals, who then form pairs more likely to

transmit vaccine-hesitancy to their children. Taken together, our results support the importance

of considering the cultural factors that have shaped current health-related beliefs and behaviors

if health policies aim to maintain or change the current conditions.

This model also shows that the perceived value and efficacy of a vaccine are important to

maintaining sufficient levels of vaccination coverage, especially if vaccine confidence is not being

robustly transmitted (or maintained in adulthood). Individuals essentially perform an internal

cost-benefit analysis based on their circumstances and interpretation of accessible information

when deciding to vaccinate. We aimed to be inclusive of their various considerations via our

comprehensive cultural selection coefficient. Increasing positive public perception through

honest and effective communication and reducing public concern about vaccines and increasing

vaccine safety could together drive increased vaccination trust and acceptance. Achieving the

optimal vaccination coverage lies not only in the hands of the public by vaccinating themselves

and their children, but also in the efforts of health officials and leaders in creating an

environment that fosters confidence by assuring the public of vaccine efficacy, safety, and value,

while providing convenient avenues to attain vaccines.

As with any model, we cannot fully capture the complex reality of the relationship

between vaccine hesitancy and vaccination behavior. First, though vaccination frequency data is

available for numerous vaccines, frequency data for vaccine attitudes are much less common,

with the two traditionally not surveyed together. Thus, there is no dataset that exactly estimates

the phenotypes presented here, for example, the number of vaccinated but hesitant (V^+A^-)

individuals in a population. The goals of vaccination attitude surveys have been primarily to

identify themes of vaccine hesitancy, and to a lesser degree, the themes of vaccination. However,

they do not report parent vaccination states or whether the child was actually vaccinated (on

schedule). With data presenting parent vaccination states alongside their vaccine attitudes and

vaccination decisions, we would be able to more accurately inform phenotype frequencies,

possibly extending the model to incorporate various types of hesitancy. We note, however, that

our results did not depend on the initial proportions of vaccination status or vaccine hesitancy, so

these data would primarily be for comparison to our equilibrium outcomes.

We were also constrained by limited data to inform our cultural transmission and

transition probabilities. In our model, baseline confidence transmission and influence

probabilities are structured according to a simple pattern of inheritance, such that each parent is

equally likely to influence an offspring's phenotype. However, cultural traits and vaccination

attitudes may not strictly follow this pattern: one parent might have more influence, or one

variant of a trait might be more likely to be transmitted. In addition, transmission probabilities

are constant in our model, remaining unaffected by changing cultural conditions throughout each

simulation, but in reality, these probabilities may fluctuate in response to a variety of factors

including vaccine type or family structure. Future developments of the model could include

modulating the probability of vaccine confidence transmission according to other aspects of the

cultural environment, such as the attitude frequencies in the population. We could also use the

current frequency of these cultural traits across different populations to generate more specific

hypotheses about their underlying cultural transmission processes [41,42] Our cultural selection

coefficient and attitude transition probabilities did vary with the frequency of vaccination

coverage, but the exact relationships could not be informed by existing data. Modulating both

the attitude transition probabilities and the cultural selection coefficient according to the level of

vaccination coverage in a population, however, reflects that perceptions about the vaccine and

its associated effects on health could be meaningfully different in a population with high

vaccination coverage than in one with low coverage.

Though vaccination coverage and vaccine confidence stabilized in our simulations, in

reality vaccination rates fluctuate over time in response to changing population dynamics,

perhaps never arriving at a stable equilibrium. For example, the increasingly rapid spread of

information [43] may cause attitudes and behaviors to change frequently over short periods of

time. In our model, most of the phenotype frequency fluctuations occur in the first few iterations

before quickly adjusting to an equilibrium. Unlike some models of population dynamics, this

model has a discrete-time format and does not consider asynchrony in population turnover.

Thus, the timescale of our model might not translate directly to years or generations, and we

avoid interpreting the number of iterations in literal terms. It is possible that if more realistic

birth and death processes were incorporated, the cultural dynamics would occur at different

timescales and would continue to fluctuate instead of approaching a stable equilibrium. In

addition, the grandparents of the children to be vaccinated also influence the parents'

vaccination decisions [44]. A restructuring of the timescale or the incorporation of population

asynchrony in our model could allow for consideration of these influences.

In this model, we constructed the offspring vaccination probability to be informed

primarily by parents' vaccine attitudes and secondarily by their own vaccination status. Though it

is understood that there is an interaction between parents' beliefs and their own experiences

with vaccines regarding their decision to vaccinate their children, accurately modeling the relative

contribution of these two factors could benefit from empirical studies on parental willingness to

vaccinate based on their beliefs and vaccination status. With our current formula ($B_{m,n}$, **Table S2**),

vaccine-confident parents who did not themselves receive childhood vaccines have a reduced

likelihood of vaccinating their offspring than vaccinated parents. In reality, parental vaccine

attitudes might even further outweigh their own vaccination status in their decision-making

process than we model here.

Finally, future developments of this model could include homophily of oblique

interactions, that is, if vaccine-related beliefs influenced not only one's mate choice but also

one's choice of social groups or information sources. On one hand, individuals who

disproportionately interact with vaccine-hesitant contacts would have a biased perspective that

vaccine hesitancy is more prevalent in the population than it actually is, which can reduce their

likelihood of vaccinating their children [45]; on the other hand, a high degree of homophily in

oblique interactions has been hypothesized to hinder the transmission of vaccine hesitancy to

vaccine confident individuals, reducing the spread of the belief overall [28]. Another potential

further exploration of the model includes modeling preferential assortment based on vaccination

status rather than vaccine attitude, which has been shown to occur in an empirical contact-

network study of high school students [46].

Our findings suggest that broad efforts to encourage and inform the public about vaccine

safety and efficacy will foster higher vaccine coverage, and thus points toward several

recommendations for public health policy and outreach. We recommend that accurate

information about vaccines be readily accessible through a variety of means, be easily

understood, and be supported by personal anecdotes since individuals who are skeptical about

vaccines might invest more time in seeking out information about them [47–49], and that

dialogue between people with different beliefs and attitudes be encouraged as it can help to

break the "echo chambers" of homophily, encouraging individuals to communicate and

empathize with one another. Therefore, to address vaccine hesitancy, our results underscore the

importance of considering the cultural beliefs and influences that underpin health behaviors.

5.   Acknowledgements

We are grateful for feedback on this model from Glenn Webb, Ann Tate, Tony Capra,

Kathy Friedman, Buddy Creech, and members of the Creanza lab. We thank the John Templeton

Foundation for funding this work.

References:

1. Laland K, Matthews B, Feldman MW. An introduction to niche construction theory. Evol Ecol. 2016;30: 191–202.

2. John Odling-Smee F, Laland KN, Feldman MW. Niche Construction: The Neglected Process in Evolution (MPB-37). Princeton University Press; 2013.

3. Fogarty L, Creanza N. The niche construction of cultural complexity: interactions between innovations, population size and the environment. Philos Trans R Soc Lond B Biol Sci. 2017;372. doi:10.1098/rstb.2016.0428

4. O'Brien MJ, Laland KN, Broughton JM, Cannon MD, Fuentes A, Gerbault P, et al. Genes, culture, and agriculture: An example of human niche construction. Curr Anthropol. 2012;53: 000–000.

5. Fuentes A. Cooperation, conflict, and niche construction in the genus homo. War, peace, and human nature. 2013; 78–94.

6. Creanza N, Fogarty L, Feldman MW. Models of cultural niche construction with selection and assortative mating. PLoS One. 2012;7: e42744.

7. Creanza N, Feldman MW. Complexity in models of cultural niche construction with selection and homophily. Proc Natl Acad Sci U S A. 2014;111 Suppl 3: 10830–10837.

8. Fenner F. A successful eradication campaign. Global eradication of smallpox. Rev Infect Dis. 1982;4: 916–930.

9. Kim-Farley R, Schonberger L, Nkowane B, Kew O, Bart K, Orenstein W, et al. POLIOMYELITIS IN THE USA: VIRTUAL ELIMINATION OF DISEASE CAUSED BY WILD VIRUS. The Lancet. 1984. pp. 1315–1317. doi:10.1016/s0140-6736(84)90829-8

10. Salk D. Eradication of Poliomyelitis in the United States. I. Live Virus Vaccine-Associated and Wild Poliovirus Disease. Clinical Infectious Diseases. 1980. pp. 228–242. doi:10.1093/clinids/2.2.228

11. Thompson KM, Strebel PM, Dabbagh A, Cherian T, Cochi SL. Enabling implementation of the Global Vaccine Action Plan: developing investment cases to achieve targets for measles and rubella prevention. Vaccine. 2013;31 Suppl 2: B149–56.

12. Atwell JE, Salmon DA. Pertussis resurgence and vaccine uptake: implications for reducing vaccine hesitancy. Pediatrics. 2014. pp. 602–604.

13. Kubin L. Is There a Resurgence of Vaccine Preventable Diseases in the U.S.? Journal of Pediatric Nursing. 2019. pp. 115–118. doi:10.1016/j.pedn.2018.11.011

14. Falagas ME, Zarkadoulia E. Factors associated with suboptimal compliance to vaccinations in children in developed countries: a systematic review. Curr Med Res Opin. 2008;24: 1719–1741.

15. Dubé E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. Vaccine hesitancy: an overview. Hum Vaccin Immunother. 2013;9: 1763–1773.

16. Scheres J, Kuszewski K. The Ten Threats to Global Health in 2018 and 2019. A welcome and informative communication of WHO to everybody. Zdrowie Publiczne i Zarządzanie. 2019. pp. 2–8. doi:10.4467/20842627oz.19.001.11297

17. Glanz JM, McClure DL, Magid DJ, Daley MF, France EK, Salmon DA, et al. Parental refusal of pertussis vaccination is associated with an increased risk of pertussis infection in children. Pediatrics. 2009;123: 1446–1451.

18. MacDonald NE, SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: Definition, scope and

determinants. Vaccine. 2015;33: 4161–4164.

19. Eggertson L. Lancet retracts 12-year-old article linking autism to MMR vaccines. CMAJ. 2010;182:
E199–200.

20. Rao TSS, Andrade C. The MMR vaccine and autism: Sensation, refutation, retraction, and fraud.
Indian J Psychiatry. 2011;53: 95–96.

21. Burley N. The meaning of assortative mating. Ethol Sociobiol. 1983;4: 191–203.

22. Gimelfarb A. Processes of Pair Formation Leading to Assortative Mating in Biological Populations:
Encounter-Mating Model. The American Naturalist. 1988. pp. 865–884. doi:10.1086/284827

23. Salathé M, Bonhoeffer S. The effect of opinion clustering on disease outbreaks. J R Soc Interface.
2008;5: 1505–1508.

24. Perra N, Balcan D, Gonçalves B, Vespignani A. Towards a characterization of behavior-disease models.
PLoS One. 2011;6: e23084.

25. Mao L, Yang Y. Coupling infectious diseases, human preventive behavior, and networks--a conceptual
framework for epidemic modeling. Soc Sci Med. 2012;74: 167–175.

26. Bauch CT. Imitation dynamics predict vaccinating behaviour. Proc Biol Sci. 2005;272: 1669–1675.

27. Chauhan S, Misra OP, Dhar J. Stability analysis of SIR model with vaccination. American journal of
computational and applied mathematics. 2014;4: 17–23.

28. Mehta RS, Rosenberg NA. Modelling anti-vaccine sentiment as a cultural pathogen. Evolutionary
Human Sciences. 2020;2: e21.

29. May T, Silverman RD. "Clustering of exemptions" as a collective action threat to herd immunity.
Vaccine. 2003;21: 1048–1051.

30. Wang E, Clymer J, Davis-Hayes C, Buttenheim A. Nonmedical exemptions from school immunization
requirements: a systematic review. Am J Public Health. 2014;104: e62–84.

31. Phadke VK, Bednarczyk RA, Salmon DA, Omer SB. Association Between Vaccine Refusal and Vaccine-
Preventable Diseases in the United States: A Review of Measles and Pertussis. JAMA. 2016;315:
1149–1158.

32. Pruitt RH, Kline PM, Kovaz RB. Perceived Barriers to Childhood Immunization Among Rural
Populations. Journal of Community Health Nursing. 1995. pp. 65–72.
doi:10.1207/s15327655jchn1202_1

33. Cavalli-Sforza LL, Feldman MW. Cultural transmission and evolution: a quantitative approach. Monogr
Popul Biol. 1981;16: 1–388.

34. Bauch CT, Bhattacharyya S. Evolutionary game theory and social learning can determine how vaccine
scares unfold. PLoS Comput Biol. 2012;8: e1002452.

35. Kennedy AM, Brown CJ, Gust DA. Vaccine beliefs of parents who oppose compulsory vaccination.
Public Health Rep. 2005;120: 252–258.

36. Leask J. Target the fence-sitters. Nature. 2011;473: 443–445.

37. Hill HA, Singleton JA, Yankey D, Elam-Evans LD, Cassandra Pingali S, Kang Y. Vaccination Coverage by
Age 24 Months Among Children Born in 2015 and 2016 — National Immunization Survey-Child,
United States, 2016–2018. MMWR. Morbidity and Mortality Weekly Report. 2019. pp. 913–918.
doi:10.15585/mmwr.mm6841e2

38. Hill HA, Elam-Evans LD, Yankey D, Singleton JA, Kang Y. Vaccination Coverage Among Children Aged

19–35 Months — United States, 2016. MMWR. Morbidity and Mortality Weekly Report. 2017. pp.
1171–1177. doi:10.15585/mmwr.mm6643a3

39. Gangarosa EJ, Galazka AM, Wolfe CR, Phillips LM, Gangarosa RE, Miller E, et al. Impact of anti-vaccine
movements on pertussis control: the untold story. Lancet. 1998;351: 356–361.

40. Ozawa S, Mirelman A, Stack ML, Walker DG, Levine OS. Cost-effectiveness and economic benefits of
vaccines in low- and middle-income countries: a systematic review. Vaccine. 2012;31: 96–108.

41. Kandler A, Powell A. Generative inference for cultural evolution. Philos Trans R Soc Lond B Biol Sci.
2018;373. doi:10.1098/rstb.2017.0056

42. Kandler A, Wilder B, Fortunato L. Inferring individual-level processes from population-level patterns
in cultural evolution. R Soc Open Sci. 2017;4: 170949.

43. Hornik J, Satchi RS, Cesareo L, Pastore A. Information dissemination via electronic word-of-mouth:
Good news travels fast, bad news travels faster! Computers in Human Behavior. 2015. pp. 273–280.
doi:10.1016/j.chb.2014.11.008

44. Karthigesu SP, Chisholm JS, Coall DA. Do grandparents influence parents' decision to vaccinate their
children? A systematic review. Vaccine. 2018;36: 7456–7462.

45. Brunson EK. The Impact of Social Networks on Parents' Vaccination Decisions. Immunization
Strategies and Practices. 2018. pp. 123–133. doi:10.1542/9781610022774-the_impact

46. Barclay VC, Smieszek T, He J, Cao G, Rainey JJ, Gao H, et al. Positive network assortativity of influenza
vaccination at a high school: implications for outbreak risk and herd immunity. PLoS One. 2014;9:
e87042.

47. Gowda C, Dempsey AF. The rise (and fall?) of parental vaccine hesitancy. Hum Vaccin Immunother.
2013;9: 1755–1762.

48. Ellithorpe ME, Adams R, Aladé F. Parents' Behaviors and Experiences Associated with Four
Vaccination Behavior Groups for Childhood Vaccine Hesitancy. Matern Child Health J. 2022;26: 280–
288.

49. Benin AL, Wisler-Scher DJ, Colson E, Shapiro ED, Holmboe ES. Qualitative Analysis of Mothers'
Decision-Making About Vaccines for Infants: The Importance of Trust. Pediatrics. 2006. pp. 1532–
1541. doi:10.1542/peds.2005-1728