Epidemiology of adverse events attributed to airway management in pediatric

anesthesia: A protocol for a Japan Pediatric Difficult Airway in Anesthesia study

(J-PEDIA)

Taiki Kojima[1,2*], Yusuke Yamauchi[1¶], Fumio Watanabe[1¶], Shogo Ichiyanagi[1¶], Yasuma

Kobayashi[3&], Yu Kaiho[4&], Shugo Kasuya[5&], Kevin Y. Urayama[6&], Norifumi Kuratani[7&,]

Yasuyuki Suzuki[5&], on behalf of J-PEDIA study collaborators

[1] Department of Anesthesiology, Aichi Children's Health and Medical Center, Aichi,

Japan

[2] Division of Comprehensive Pediatric Medicine, Nagoya University Graduate School of

Medicine, Aichi, Japan

[3] Children's Heart Center, Saitama Children's Medical Center, Saitama, Japan

[4] Department of Anesthesiology, Tohoku University Hospital, Sendai, Japan

[5] Department of Critical Care and Anesthesiology, National Center for Child Health and

Development, Tokyo, Japan

[6] Graduate School of Public Health, St. Luke's International University, Tokyo, Japan

[7] Department of Anesthesiology, Saitama Children's Medical Center, Saitama, Japan

*Corresponding author

Taiki Kojima

E-mail: daiki_kojima@sk00106.achmc.pref.aichi.jp (TK)

[¶] These authors contributed equally to this work.

[&] These authors also contributed equally to this work.

<u>Abbreviations</u>

AE: Adverse event

NEAR4KIDS: National Emergency Airway Registry for children

Abstract

Introduction

Failure to secure an airway during general anesthesia is a major cause of adverse events

(AEs) in children. The safety of peddiatric anesthesia may be improved by identifying the

incidence of AEs and their attributed risk factors. However, previous large cohort studies

have not been appropriately designed to explore such aspects. Thus, this study aims to

determine the incidence of AEs and the risk factors attributed to airway management under

general anesthesia in children.

Methods and design

This prospective, multi-center, registry-based, observational study will be conducted in

four tertiary care hospitals in Japan from June 2022 to May 2025. Children younger

than 18 years of age undergoing surgical and/or diagnostic test procedures under general

anesthesia or sedation by anesthesiologists will be enrolled in this study. Data on patient

characteristics, discipline of anesthesia providers, and methodology of airway

management will be collected through a standardized verification system under

monitoring by institutional research leaders among the recruited institutions to minimize

the loss of collected data. The primary outcome and exposure are AEs and presence of

difficult airway features with potential confounders, which are related to the failure to

secure the airway, and the variability in the anesthesia providers' levels, adjusted using

hierarchical multivariable regression models with mixed effects.

Discussion

This study will determine the incidence of AEs and the risk factors related to airway

management under general anesthesia in children.

Trial registration

This study is registered as a prospective observational study in the University hospital

Medical Information Network (UMIN) (registration number UMIN000047351, April 1,

2022).

Introduction

The incidence of perioperative life-threatening adverse events (AEs) in children is higher than that in adults due to their unique physiological and anatomical characteristics [1]. These AEs are related to hypoxia in children commonly occur following failure to secure the airway since children are less tolerant to apnea [1-4]. Therefore, identifying the incidence of AEs and the risk factors in securing the airway is desired for safe and strategic airway management in pediatric anesthesia.

Epidemiological data on AEs during general anesthesia in children have been reported in previous large prospective cohort studies. The APRICOT study, which was the largest multi-center prospective study conducted in Europe, reported an incidence rate of 5.2% for severe perioperative events in children [5]. In addition, Subramanyam reported prediction models for perioperative respiratory adverse events based on preoperative patient comorbidities [6]. However, the APRICOT study was not designed to explore AEs specifically attributed to airway management during general anesthesia. Therefore, it did not include detailed information regarding the difficult airway features of patients, anesthesia methods, devices utilized for securing the airway, and disciplines of anesthesia providers. The PeDI registry study, a multi-center cohort study conducted in children's hospitals in the US, reported that the incidence rate of intraoperative AEs

was high in children with difficult airway features [2]. However, approximately 80% of

the study cohort were patients with difficult airway features; therefore, the results did

not represent the entire pediatric population. In addition, these major studies did not

adjust for the variability in the techniques and skill level of anesthesia providers (e.g.,

each anesthesia provider could have a specific methodology for securing the airway and

anesthesia). Therefore, the incidence of AEs and the related risk factors in securing the

airway under general anesthesia in children remain unclear. In addition, to our

knowledge, there is a lack of well-designed pediatric prospective studies using registry-

based, real-world data in the Asian region regarding the incidence of AEs and the risk

factors attributed to airway management.

This study aims to investigate the incidence of AEs attributed to airway

management during general anesthesia in children and to identify their risk factors.

Methods

Study design and setting

This is a prospective, registry-based, real-world, multicenter, observational study that

will initiate data registration from June 1, 2022, to May 31, 2025, in five tertiary care

hospitals in Japan: Aichi Children's Health and Medical Center (Aichi), National Center

for Child Health and Development (Tokyo), Saitama Prefectural Children's Medical

Center (Saitama), Tohoku University Hospital (Sendai), and Hokkaido University

Hospital (Sapporo).

Target population

Inclusion criteria. We will enroll cases that meet all of the following criteria.

- Children younger than 18 years of age who receive advanced airway management

under general anesthesia or sedation for surgical and/or test procedures conducted by

anesthesiologists or anesthesia providers under the supervision of anesthesiologists.

- General anesthesia or sedation is performed in operating suites, catheterization

laboratory rooms, rooms for radiological imaging and procedures (e.g., computed

tomography, magnetic resonance imaging, radiation therapy), or general ward.

Exclusion criteria. We will exclude cases that meet at least one of the following

criteria.

- Cases outside operating suites where anesthesiologists are consulted for airway

management.

- Cases where airway management is performed in the emergency department and

intensive care unit.

- Duplicate cases during the study period.

- Patients or their families who refuse to participate in the study for any reason.

119   Data collection

This registry-based observational study will prospectively collect data regarding the

characteristics of patients and surgery, disciplines of anesthesia providers, practice of

airway management, occurrence of AEs, and treatments for AEs.

Data quality control. Data will be initially collected by anesthesia providers for each

attempt to secure the airway with paper-based data collection forms. The collected

initial paper-form-based data will be verified by each site-specific research leader

through a standardized process among the four recruited institutions. Site-specific

research leaders will confirm that the capture rate is ≥95% of cases at each institution.

To standardize the definitions of the terms for data collection, they are predetermined

and described in the operational research manual created by the primary investigator

(TK). The research operational committee will be convened regularly to confirm the

definitions of the terms used in the data collection process. Site-specific research leaders

instructed the anesthesia providers to increase the accuracy of the collected data at each

recruited institution, aiming to minimize misclassification bias (Figure 1).

Data storage and process. The initially collected paper-form-based data will include

information that can identify the study participants (i.e., medical record number and

name initials). Thus, only anonymous information will be registered in the Research

Electronic Data Capture (REDCap®, Tokyo, Japan) [7] system hosted by the National

Center for Child Health and Development. Site-specific research leaders will utilize a

standardized participant's correspondence file, which will include the participant's

initials, medical record number, and REDCap registration number for a potential review

(Figure 1).

Endpoints. The primary and secondary endpoints are any AEs attributed to securing

the airway and decreased values of peripheral capillary oxygen saturation (SpO_2) from

the time of initiation to completion of securing the airway. The criteria for AEs were

predetermined [i.e., cardiac arrest (status of dead or alive at 24 hours after the critical

event happened), laryngospasm, upper airway obstruction, severe cough, bronchial

intubation, esophageal intubation, vomiting, hypotension, hypertension, tooth injury,

pneumothorax, mediastinal emphysema, bronchospasm, atelectasis, pulmonary edema,

stridor, airway trauma, arrhythmia, agitation, dislodgement of airway securing devices]

based on the National Emergency Airway Registry for children (NEAR4KIDS), which

is a national registry database to evaluate AEs during emergency tracheal intubation in

pediatric intensive care units mostly located in North America [8]. The AEs have been

classified as severe and non-severe before data collection. Severe AEs include cardiac

arrest, esophageal intubation with a $\geq 5\%$ decrease in SpO_2, vomiting with aspiration,

hypotension requiring treatments, laryngospasm, pneumothorax, mediastinal

emphysema, bronchospasm, pulmonary edema, and direct airway trauma. Non-severe

AEs include bronchial intubation, esophageal intubation with a $< 5\%$ decrease in SpO_2,

vomiting without aspiration, hypertension requiring treatments, upper airway

obstruction, severe cough, tooth injury, atelectasis, stridor, arrhythmia, and

dislodgement of airway securing devices.

Exposures and potential confounders. Exposure is defined as the presence of any

of the difficult airway features (i.e., preoperative recognition of difficult airway by

anesthesiologists, history of difficult airway, restricted cervical flexion, restricted mouth

opening, narrow thyromental distance, upper airway obstruction, or anatomical barrier

to visualize glottic opening, midfacial hypoplasia, large tongue, micrognathia, and large

head circumference) that were predetermined based on previous studies [9]. Potential

confounders were selected before initiation of data collection based on a review of

previous literature and clinical experience of experienced anesthesiologists among the

research team members. The selected potential confounders were age, sex, prematurity,

obesity (weight-for-age ≥95 percentile), preoperative comorbidities, families smoking

status, provider discipline, location, institution, fiscal year, Cormack-Lehan grade,

method of anesthesia induction, status of nothing per os and a full stomach, type of

device for securing the airway, and application of external laryngeal manipulation or

cricoid pressure [9-14].

179   Statistical considerations

Sample size estimation. The reported incidence rate of critical events occurring

during general anesthesia induction in children was 5.2%, including an incidence of

1.9% for cardiovascular instability in the general pediatric population who received

general anesthesia [5]. Therefore, we estimated the sample size with the assumption of

the lowest incidence of critical adverse events during general anesthesia in children as

2.0%. The estimated sample size was approximately 16,000 assuming a 99% probability

of obtaining a 95% Wilson CI with ± 0.3% of the half-width for the incidence of critical

adverse events.

Statistical analysis. For summary statistics, categorical variables will be described as

numbers and percentages, whereas normally and non-normally distributed continuous

variables will be described as means and standard deviations or medians and

interquartile ranges. Univariable and multivariable analyses will be performed to

evaluate risk factors associated with the primary outcome. Baseline differences between

subjects with and without difficult airway features will be evaluated using the Student's

t-test for normally distributed continuous variables, Wilcoxon rank-sum test for

continuous variables with skewed distributions, and chi-square test or Fisher's exact test

for categorical variables. Multilevel logistic regression models with mixed effects will

be used to evaluate the association between the occurrence of adverse events while

securing the airway and the presence of difficult airway features [15]. Hierarchical

relative risk regression models will be developed using robust error variances or

hierarchical logistic regression models for multivariable analysis. Potential confounders

for patient, anesthesia, provider characteristics, and institutions will be selected based

on clinical relevance and adjusted for fixed effects. The variance contributed by the

clustering effects based on the different clinical practices among anesthesia providers

will be adjusted (each subject as level 1, each anesthesia provider as level 2) as random

effects [15]. Cases with missing values will be removed from the analysis. Data will be

analyzed using STATA 17.0 (StataCorp, College Station, TX, USA), with a two-sided

p-value of <0.05, serving as the criterion for assessing the null hypothesis for each

analysis.

212 Ethics statement

This study will be conducted in accordance with the declaration of Helsinki. The study

protocol was reviewed and approved by the institutional review board at Aichi

Children's Health and Medical Center (approval number 2021051, September 29, 2021)

and local ethical approval will be obtained from each recruited institution before

initiating data collection. The institutional review board considered that written consent

could be waived by performing the opt-out procedure in the current clinical study.

220 Dissemination

221   The results of this study will be reported in a peer-reviewed journal and a relevant

222   cademic conference.

Harms

No harmful events will occur during data collection because of the nature of an

observational study without interventions.

Discussion

This is a prospective, registry-based, multicenter, observational study that aims to

describe real-world data on the incidence of AEs and the risk factors attributed to

securing the airway in children under general anesthesia. Investigating real-world data

regarding patient characteristics (including the prevalence of difficult airway features)

and characteristics of pediatric anesthesia practice can provide anesthesia providers with

useful information to improve the safety and effectiveness of airway management.

This study has several important limitations owing to its observational nature

of the study. First, unmeasured confounders could remain unadjusted owing to the

observational nature of the study. Second, there might be a selection bias due to the

sampling method. Third, reporting bias (i.e., misclassification bias) can occur due to

incorrect reports during data collection. To address these potential limitations, we

carefully designed this study to collect detailed information regarding the exposure

(presence of difficult airway features), potential confounders, and institutions based on

previous literature and meetings with experienced board-certified anesthesiologists

during the development of the study protocol. In addition, we structured a standardized

system to confirm the data collected by local research leaders among the recruited

institutions. This standard data confirmation system has been used in NEAR4KIDS,

which achieved a data capture rate of ≥95% [11]. Our standardized data collection

system allows our study to maintain the quality of the collected data and minimize

potential information biases (e.g., misclassification). In addition, the REDCap® data

registration system does not allow data registration to proceed if there are missing data.

We will use this system for data registration to minimize missing data for future

analysis.

Previous registry-based prospective studies (i.e., APRICOT study, PeDI

registry study) were performed mainly in Europe and North America [2,5]. The

evidence regarding the incidence of AEs related to securing the airway is insufficient in

Asian settings. The incidence and risks of AEs attributed to airway management in the

Japanese pediatric population might be different from those in Europe and North

America owing to the differences in the characteristics of the patient population and

anesthesia practices. This prospective, registry-based, real-world observational study in

Japan can be a stepping stone for further research to identify the specific risk factors for

AEs attributed to failure to secure the airway in the Asian region.

Ethics statements

Patient consent for publication

Not required.

Acknowledgments

None.

Author Contributions

Conceptualization: Taiki Kojima, Yusuke Yamauchi, Fumio Watanabe, Shogo

Ichiyanagi, Yasuma Kobayashi, Yu Kaiho, Shugo Kasuya, Norifumi Kuratani and

Yasuyuki Suzuki

Data curation: Taiki Kojima, Yusuke Yamauchi

Funding acquisition: Taiki Kojima [Grants-in-Aid for Scientific Research (Kakanhi

Grant) #22K09085, April 1, 2022].

Methodology: Taiki Kojima, Yusuke Yamauchi, Fumio Watanabe, Shogo Ichiyanagi

Writing - original draft: Taiki Kojima

Writing – review & editing: Taiki Kojima, Yusuke Yamauchi, Fumio Watanabe,

Shogo Ichiyanagi, Yasuma Kobayashi, Yu Kaiho, Shugo Kasuya, Kevin Y. Urayama,

Norifumi Kuratani, Yasuyuki Suzuki

References

1. Else SDN, Kovatsis PG. A narrative review of oxygenation During pediatric intubation

and airway procedures. Anesth Analg. 2020;130: 831-840. doi:

10.1213/ANE.0000000000004403.

2. Fiadjoe JE, Nishisaki A, Jagannathan N, Hunyady AI, Greenberg RS, Reynolds PI, et

al. Airway management complications in children with difficult tracheal intubation from

the Pediatric Difficult Intubation (PeDI) registry: A prospective cohort analysis. Lancet

Respir Med. 2016;4: 37-48. doi: 10.1016/S2213-2600(15)00508-1.

3. Schibler A, Hall GL, Businger F, Reinmann B, Wildhaber JH, Cernelc M, et al.

Measurement of lung volume and ventilation distribution with an ultrasonic flow meter

in healthy infants. Eur Respir J. 2002;20: 912-918. doi: 10.1183/09031936.02.00226002.

4. Humphreys S, Pham TM, Stocker C, Schibler A. The effect of induction of anesthesia

and intubation on end-expiratory lung level and regional ventilation distribution in

cardiac children. Paediatr Anaesth. 2011;21: 887-893. doi: 10.1111/j.1460-

9592.2011.03547.x.

5. Habre W, Disma N, Virag K, Becke K, Hansen TG, Jöhr M, et al. Incidence of severe

critical events in paediatric anaesthesia (apricot): A prospective multicentre observational

study in 261 hospitals in Europe. Lancet Respir Med. 2017;5: 412-425. doi:

10.1016/S2213-2600(17)30116-9.

6. Subramanyam R, Yeramaneni S, Hossain MM, Anneken AM, Varughese AM.

Perioperative respiratory adverse events in pediatric ambulatory anesthesia: Development

and validation of a risk prediction tool. Anesth Analg. 2016;122: 1578-1585. doi:

10.1213/ANE.0000000000001216.

7. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research Electronic

Data Capture (REDCap)--A metadata-driven methodology and workflow process for

providing translational research informatics support. J Biomed Inform. 2009;42: 377-381.

doi: 10.1016/j.jbi.2008.08.010.

8. Branca A, Tellez D, Berkenbosch J, Rehder KJ, Giuliano JS, Gradidge E, et al. The

New Trainee Effect in Tracheal Intubation Procedural Safety Across PICUs in North

America: A Report From National Emergency Airway Registry for Children. Pediatr Crit

Care Med. 2020;21: 1042-1050. doi: 10.1097/PCC.0000000000002480.

9. Heinrich S, Birkholz T, Ihmsen H, Irouschek A, Ackermann A, Schmidt J. Incidence

and predictors of difficult laryngoscopy in 11,219 pediatric anesthesia procedures.

Paediatr Anaesth. 2012;22: 729-736. doi: 10.1111/j.1460-9592.2012.03813.x.

10. Valois-Gómez T, Oofuvong M, Auer G, Coffin D, Loetwiriyakul W, Correa JA.

Incidence of difficult bag-mask ventilation in children: A prospective observational study.

Paediatr Anaesth. 2013;23: 920-926. doi: 10.1111/pan.12144.

11. Kojima T, Laverriere EK, Owen EB, Harwayne-Gidansky I, Shenoi AN, Napolitano

N, et al. Clinical impact of external laryngeal manipulation during laryngoscopy on

tracheal intubation success in critically ill children. Pediatr Crit Care Med. 2018;19: 106-

114. doi: 10.1097/PCC.0000000000001373.

12. Mirghassemi A, Soltani AE, Abtahi M. Evaluation of laryngoscopic views and related

influencing factors in a pediatric population. Paediatr Anaesth. 2011;21: 663-667. doi:

10.1111/j.1460-9592.2011.03555.x.

13. Akpek EA, Mutlu H, Kayhan Z. Difficult intubation in pediatric cardiac anesthesia. J

Cardiothorac Vasc Anesth. 2004;18: 610-612. doi: 10.1053/j.jvca.2004.07.003.

14. Graciano AL, Tamburro R, Thompson AE, Fiadjoe J, Nadkarni VM, Nishisaki A.

Incidence and associated factors of difficult tracheal intubations in pediatric ICUs: A

report from National Emergency Airway Registry for Children: NEAR4KIDS. Intensive

Care Med. 2014;40: 1659-1669. doi: 10.1007/s00134-014-3407-4.

15. Austin PC. A tutorial on multilevel survival analysis: Methods, models and

applications. Int Stat Rev. 2017;85: 185-203. doi: 10.1111/insr.12214.

Data are collected in paper-based forms by assigned anesthesia providers in each case of advanced airway management

Site-specific research leaders verify the initial paper-form-based data and the capture rate is ≥95% of cases at each institution

Collected data in paper-based forms will be anonymized, and registered in REDCap® system in each institution

Site-specific research leaders utilize a standardized participant's correspondence file for a potential future review

Participant's correspondence file will be stored in each institution

Figure 1. Flow of collection, quality control, and storage of data

Figure1