Title

How is the COVID-19 pandemic impacting our life, mental health, and well-being? Design and preliminary findings of the pan-Canadian longitudinal COHESION Study

Authors

Stephan Gabet, PhD, PharmD,[1,2] Benoit Thierry, MSc,[2] Rania Wasfi, PhD,[3] Margaret De Groh, PhD,[3] Guido Simonelli, MD, MSc,[4,5] Catherine Hudon, MD, PhD,[6] Lily Lessard, PhD,[7,8] Ève Dubé, PhD,[9] Bouchra Nasri, PhD,[1,2] Yan Kestens, PhD,[1,2] Grégory Moullec, PhD[1,5]

Affiliations

[1] Département de Médecine Sociale et Préventive, École de Santé publique de l'Université de Montréal (ESPUM), Québec (QC), Canada.

[2] Centre de Recherche en Santé Publique (CReSP), Université de Montréal (UdeM), Québec (QC), Canada.

[3] Centre for Surveillance and Applied Research, Health Promotion and Chronic Disease Prevention Branch, Public Health Agency of Canada / Government of Canada, Ottawa, Ontario (ON), Canada.

[4] Département de Médecine, Université de Montréal (UdeM), Québec (QC), Canada

[5] Centre de Recherche du Centre intégré universitaire de santé et de services sociaux du Nord-de-l'Île-de-Montréal (CIUSSS-NIM), Québec (QC), Canada.

[6] Département de Médecine de Famille et Médecine d'urgence, Université de Sherbrooke (UdeS), Québec (QC), Canada.

24 [7] Département des sciences de la santé, Université du Québec à Rimouski, Québec (QC),

25 Canada.

26 [8] Centre de recherche du CISSS de Chaudière-Appalaches, Québec (QC), Canada.

27 [9] Centre de recherche du CHU de Québec, Université de Laval, Québec (QC), Canada.

Corresponding author

E-mail: gregory.moullec@umontreal.ca (GM)

Abstract (255 words)

With the advent of the COVID-19 pandemic, in-person social interactions and opportunities

for accessing resources that sustain health and well-being have drastically reduced. We

therefore designed the pan-Canadian population-based prospective *COVID-19: HEalth and*

Social Inequities across Neighbourhoods (COHESION) cohort to provide deeper

understanding of how the COVID-19 pandemic context affects mental health and well-being,

key determinants of health, and health inequities.

This paper presents the design of the two-phase COHESION Study, and descriptive

results from the first phase conducted between May 2020 and September 2021. During that

period, the COHESION research platform collected monthly data linked to COVID-19 such

as infection and vaccination status, perceptions and attitudes regarding pandemic-related

measures, and information on participants' physical and mental health, well-being, sleep,

loneliness, resilience, substances use, living conditions, social interactions, activities, and

mobility.

The 1,268 people enrolled in the Phase 1 COHESION Study are for the most part from

Ontario (47%) and Quebec (33%), aged 48 ± 16 years [mean ± standard deviation (SD)], and

mainly women (78%), White (85%), with a university degree (63%), and living in large urban

centers (70%). According to the 298 ± 68 (mean ± SD) prospective questionnaires completed

each month in average, the first year of follow-up reveals significant temporal variations in

standardized indexes of well-being, loneliness, anxiety, depression, and psychological distress.

The COHESION Study will allow identifying trajectories of mental health and well-

being while investigating their determinants and how these may vary by subgroup, over time,

and across different provinces in Canada, in the unique context of the COVID-19 pandemic.

Keywords

COVID-19; mental health; prospective cohort; social interactions; study design; well-being.

Abbreviations

COHESION: *COVID-19: HEalth and Social Inequities across Neighbourhoods*

Word count

3,748 words (from the beginning of the Introduction to the end of the Results)

1. Introduction

On January 10, 2020, WHO announced the identification of a new strain of coronavirus, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), causing a mild to severe respiratory illness – which may progress to pneumonia and respiratory failure – named coronavirus disease 2019 (COVID-19) [1]. The first positive case of COVID-19 in Canada was reported on January 25, 2020 and a state of health emergency was declared across the country between March 13 and March 22, 2020, depending on the province [2,3]. Mid-August 2021, there were just over 1.4 million cases in Canada and more than 26,700 deaths [2].

Public health measures to control transmission have brought vast sectors of economic activity to a halt, leading to massive unemployment and reductions in income, while reducing people's daily movements and opportunities for in-person social interactions [4]. This has affected people's opportunities and access to resources that sustain mental health and well-being. The COVID-19 conditions reinforce the role of various social and environmental health determinants, with differential impacts on the mental health and well-being of populations, depending on age [5], gender [6], housing conditions [7], education [8], job type [9], income [10], or ethnic background [11–13] and, more generally, social and environmental contexts [14,15].

COVID-19 conditions including lockdown, curfew and physical distancing measures reduce social contacts, increase social isolation and feelings of loneliness, and decrease levels of social support; these dimensions directly influence well-being and mental health [16,17]. Canadian data shows pandemic-related increases in social isolation, domestic violence and anxiety [18], with certain groups such as teens, older adults, women and racialized communities particularly at-risk [19]. Daily mobility and related physical activity are reduced through confinement measures and reduced activity spaces have been linked to depressive symptoms [20,21] and sleep troubles [22]. Walkable environments and access to green space

are key environmental conditions linked to positive health outcomes including physical

activity and well-being [23]. With shrinking activity spaces and policy constraints curbing

daily mobility, the role of residential living conditions is further amplified.

Priority populations often bear the burden of poor social and environmental living

conditions and have been shown to be disproportionately affected by COVID-19 [24,25].

Overcrowding or living in inadequate dwellings are important determinants of mental health

and well-being; moreover, prolonged exposure to home environments during lockdown

conditions further exacerbate these impacts [26,27]. Economic hardship, which is linked to

income reduction and job instability, has heavily contributed to the mental health burden of

Canadians, and is also related to housing instability and food insecurity [28,29]. In turn,

detrimental health behavior such as alcohol or other substance use are increasing and are

linked to poorer mental health outcomes [30].

In total, there is an urgent need to better understand, in particular, how the unintended

long-term consequences of COVID-19 pandemic and mitigation measures, contextual

conditions (e.g., housing conditions, neighborhood characteristics), and behavior (e.g.,

mobility, social interaction, sleep) are linked to mental health and well-being trajectories.

Furthermore, impacts are likely to vary between population groups (for instance, according to

gender, age, racialized communities, or deprivation level).

2. Two-phase COHESION Study objectives

We developed the pan-Canadian *COVID-19: HEalth and Social Inequities across Neighbourhoods* (COHESION) Study to better understand how the COVID-19 pandemic affects health, key determinants of health, and health inequities, with a focus on mental health and well-being. This study will provide longitudinal evidence of how these change over time and across different provinces in Canada.

In the context of the COVID-19 pandemic, the COHESION Study aims (Fig 1):

- evaluating the direct impact of physical and social neighborhood characteristics on mental health and well-being trajectories, while controlling for individual-level health behaviors and socio-demographics;
- evaluating if and how physical and social neighborhood characteristics may modify the associations between individual-level predictors and mental health and well-being trajectories.

This paper presents the design and methods used in the two phases of the COHESION study. It also reports descriptive statistics from the longitudinal follow-up of the first phase of the study and lessons learned from its implementation.

3. Two-phase COHESION Study design

3.1. Study design and recruitment

The two-phase COHESION Study is a pan-Canadian population-based prospective cohort study. COHESION Phase 1 was conducted between May 2020 and September 2021. COHESION Phase 2 will be launched in May 2022. All Phase 1 participants will be invited to participate in Phase 2, and additional recruitments (n = 10,000) will be done.

After consenting to participate to the study, participants are invited to fill a short eligibility questionnaire. Inclusion criteria are being aged 15 years or above, currently residing in Canada, and reading or speaking English or French. Eligible participants are then invited to complete a baseline questionnaire.

The recruitment for COHESION Phase 1 was launched on May 11 2020. We used a combination of methods that have proven successful from previous experience in recruiting population-based research project samples [31], including media communication (e.g., newspaper articles, radio interviews), social media promotion (e.g., Facebook, Twitter, Instagram, and LinkedIn), and outreach through partners' local networks (e.g., local health authorities that relayed the study on their website or through their newsletters).

For COHESION Phase 2, we will further use quota sampling at the health region level, based on 2016's Census data (i.e., age composition, gender, income, educational attainment, and ethnic background). We engaged Potloc Inc., a tech-enabled consumer research company that conducts survey sampling through social networks (Facebook, Twitter, Instagram, LinkedIn), for geo-targeting of respondents based on the sampling quotas. Potloc Inc.'s algorithm will push sociodemographic and geographically targeted online ads to our study until local quotas are attained (targets are monitored daily) and that 10,000 participants have thereby been enrolled.

Enrollment and participation in the two-phase COHESION Study are voluntary, and a

raffle of three prizes ($100 gift cards) is drawn every month for active participants for the

duration of the study.

3.2. Data collection

3.2.1. Involvement options

Two levels of involvement are proposed. Participants can opt for: i) participating in the online

self-administered questionnaires only (baseline + invitation to follow up questionnaires); or ii)

participating in the questionnaires and downloading a mobile phone application to provide

additional active and passive data (cf. 3.2.5).

3.2.2. Baseline questionnaires

Through the COHESION Study Phase 1 baseline questionnaire (35-45 min), participants

report on: their sociodemographic characteristics (e.g., gender, ethnic or cultural background,

education level, employment status, income, relationship status, and household composition);

housing condition (e.g., house tenure, dwelling type and size, outdoor space) and satisfaction;

physical (e.g., chronic disease diagnosis) and mental health conditions (e.g., anxiety and

depression), and substance use (alcohol, tobacco, vape, and cannabis); COVID-19 infection

status, vulnerability towards COVID-19, perception of- and compliance with COVID-19

mitigation measures, and pre-pandemic employment, income, substances use, loneliness,

resilience, and social interactions. Additionally, a map-based questionnaire collects data on

activity locations and social contacts (cf. 3.2.4). For parents, a supplementary module on

mental health and perceptions of COVID-19 mitigation measures concerning adolescents and

children who live in the household is administered (participants answer on behalf of children
in their household).

The COHESION Study Phase 2 baseline questionnaire will consist in one core (10
min) and two optional complementary sections (15 min each). The core baseline
questionnaire includes: sociodemographic characteristics and key housing conditions;
COVID-19 vaccination status, perceptions of COVID-19 pandemic and related mitigation
measures; standardized modules assessing general health (first item of the 12-item Short-
Form Health Survey) [32], well-being (5-item World Health Organization Well-Being Index,
WHO-5 Index) [33], sleep credit (Pittsburgh Sleep Quality Index, PSQI, 4 items only) [34],
and psychological distress (Psychological Distress Scale, Kessler-6) [35]; and an outdoor
mobility and social interactions module. Optional complementary baseline sections include
the map-based questionnaire on activity locations and social contacts (cf. 3.2.4) and additional
in-depth questions on relevant themes: additional items retrieved from the Phase 1 baseline
questionnaire (it means all items not already included in the Phase 2 core baseline
questionnaire); standardized modules on loneliness (University of California Los Angeles,
UCLA, 3-item loneliness score) [36], anxiety symptoms (7-item Generalized Anxiety
Disorder, GAD-7) [37], depression symptoms (9-item Patient Health Questionnaire, PHQ-9)
[38], resilience (6-item Brief Resilience Scale) [39], sense of belonging (Canada Community
Health Survey, CCHS) [40], and physical activity (Godin Leisure-Time Exercise) [41]; items
for assessing conspiracy beliefs.

Slightly shorten versions of the Phase 2 baseline questionnaires (i.e., excluding
questions about birth, ethnics, etc.) have been provided for participants coming from Phase 1
and wishing going on Phase 2.

3.2.3. Follow-up questionnaires

Follow-up questionnaires of the COHESION Study are short questionnaires (15 min). For

Phase 1, they were first offered biweekly until August 2020, then monthly. COVID-19-related

topics cover infection and/or vaccination status, perceived vulnerability, perception of and

compliance with mitigation measures, position regarding vaccination, and conspiracy beliefs.

Health-related questions focus on general health (SF-12, first item) [32], well-being (WHO-5

Index) [33], sleep credit (PSQI, 4 items only) [34], loneliness (UCLA 3-item loneliness score)

[36], anxiety symptoms (GAD-7) [37], depression symptoms (PHQ-9) [38], psychological

distress (Kessler-6) [35], and sense of belonging (CCHS) [40] (Table 1). Employment status,

household income, and substance use are also documented. Additionally, participants are

asked at each follow-up about any changes concerning their place of residence and housing

conditions. Supplementary module on mental health and perception of COVID-19 mitigation

measures concerning children or adolescents living at home is administered when applicable.

For Phase 2, follow-up questionnaires will be split in a core ('light') and an optional

complementary ('complete') sections, and will be offered every two months. It will include

the same questions as for Phase 1, adding a standardized module for assessing physical

activity (Godin Leisure-Time Exercise Questionnaire) [41].

Questions can vary between follow-ups, depending on the situation and priorities

raised by the research team and our public health partners. Thus, these additional themes can

focus on health insecurity (i.e., access to health care, prescriptions and medicine) [42], sleep

troubles (PSQI) [34], food insecurity (10-item Health Canada Household Food Security

Module) [43], and children's difficulties (if any) [44].

3.2.4. Use of VERITAS-Social to collect daily mobility and social interaction data

One of the specificities of COHESION is that it integrates, in baseline and follow-up

questionnaires of the two phases, the *Visualization and Evaluation of Route Itineraries,*

Travel destinations, Activity spaces and Social interactions (VERITAS-Social) questionnaire.

For COHESION, it was adapted to locate a possible list of up to 20 activities carried out

during the past seven days (Fig 2). As previously described, VERITAS-Social is an

interactive map-based questionnaire that jointly collects an individual's social network and

activity locations [44]; it is an adaptation of the VERITAS tool, an interactive questionnaire

for geo-locating places, and related information of interest (e.g., frequency of visit,

transportation modes used) [45]. It uses a Google Map module to facilitate the location of

activity places. The social module asks if an activity location is generally visited alone or with

someone else; participants can identify one or more individuals, or a group of people (*see*

S1a-b Figs). In other words, it is a name generator that identifies network members based on

their co-presence at reported destinations [46]. Data on network members include age, gender,

type of relationship (e.g., friend, acquaintance), frequency of interactions, and duration of the

relationship; for groups, data includes the number of people in the group, and the duration of

the relationship.

Once all activities, corresponding locations, and all people with whom these activities

are carried out have been reported, participants are asked to identify members of their social

network from whom they receive support or with whom they enjoy spending time with,

including members who may not have been reported among the people seen at usual activity

venues. Finally, participants were able to indicate interpersonal relationships between network

members (i.e., who knows whom), and whether some specific individuals belonged to

documented groups.

3.2.5. *Optional mobile application*

Participants enrolling in the mobile phone arm of the study were invited to download Ethica

Data mobile application (cf. 3.2.1), used successfully in previous research projects [47]. Once

installed and launched, the application passively monitors location and mobility (e.g.,

stationary, in vehicle, walking, or biking) using GPS data (1 minute of data collected every 5

minutes), physical activity (step counter), and social contacts (other smartphones), derived

from listings of other Bluetooth discoverable devices in the participant's surroundings [48].

The app further prompts participants to complete Ecological Momentary Assessment (EMA)

questionnaires three times a day for one week every month. EMA questionnaires include short

self-reports (<1 min) on well-being, sleep, mood (Short Mood Scale, 6 items) [49], and social

interactions (*see* S2 Fig).

3.2.6. *Environmental data*

Because some of the data collected is spatial (e.g. place of residence, VERITAS-Social

locations, mobile app GPS data), we are able to add area-level environmental data to our

dataset. This allows to explore links between individual-level measures and social and

environmental contextual conditions. Numerous environmental datasets covering Canada are

available through the *Canadian Urban Environmental Health Research Consortium*

(CANUE).

265   Contextual variables of interest include measures of neighborhood deprivation, such as

266   the Material and Social Deprivation Indices (MSDI; data 2016) available at the Canadian

267   Census dissemination area level (i.e., the smallest Canadian standard geographic area, with a

268   population of 400 to 700 persons) [50]. These two composite indexes combine Census data on

269   educational attainment, employment ratio, average income, and household composition.

270   The urbanization degree is measured using *Statistics Canada* classification (at the

four-digit code area level; data 2016), based on the number of inhabitants in population

centers [51]. "Small", "medium" and "large" urban population centers correspond to areas

embracing between 1,000 and 29,999, between 30,000 and 99,999, and 100,000 and more

inhabitants, respectively, while "rural area" is a residual value gathering all areas located

outside population centers.

Greenness is evaluated using the growing season Normalized Difference Vegetation

Index (NDVI) (at the six-digit code area level; data 2019), based on Landsat 8 satellite data

[52,53].

Area-level walkability is measured using the *Canadian Active Living Environments*

(CAN-ALE) database [54].

Additional localized and temporalized data on COVID-19 pandemic conditions,

including lockdowns, curfews, vaccination plan, number of cases, and emergence of variants

is also available through the *Canadian Institute for Health Information* (CIHI) compilation

[55].

3.3. Ethics and data management plan

Before completing the eligibility questionnaire, potential participants are first invited to read

the consent form and provide an electronic written consent to participate in the study. In

Phase 2, a second consent will be required from participant wishing to register for the

prospective follow-ups after having completed baseline questionnaire. Phase 1 COHESION

was approved by the ethics board of the *Centre de Recherche du Centre Hospitalier de*

l'Université de Montréal (CRCHUM; MP-02-2021-8924) and by the *Public Health Agency of*

Canada (PHAC) ethical review board (REB 2020-016P). Phase 2 COHESION was approved

by both the ethics board of the *Centre Intégré Universitaire de Santé et de Services Sociaux*

du Nord-de-l'Île-de-Montréal (CIUSSS-NIM ; 2022-2327) and PHAC (REB 2020-016P).

For data security and privacy proposal, all data are hosted on *Compute Canada* servers

with secondary backups on hospital-grade internal servers. All data has been stripped from

any personally identifying information, with only the principal investigator being able to track

the records back to one participant. Researchers can access anonymized individual level

records by connecting directly through a Secure Shell (SSH) to the database hosted on

Compute Canada (a SSH allows remote machines accessing data in a secure way since the

connection is encrypted). They can also access aggregated data for analysis through a secure

online platform hosted by Tableau [56]. A document listing all the data management policies

governing data access and storage has been submitted to the ethic board for approval.

4. COHESION Study Phase 1 first results

4.1. Recruitment and participants

Among the 2,557 people who completed the COHESION Phase 1 eligibility questionnaire, 2,346 (92%) met the selection criteria (Fig 3). Among these ones, 1,268 (54%) completed the baseline questionnaire and have been enrolled in the COHESION Study, and among the latter, 557 (44%) opted for the Ethica mobile application in addition to the online self-administered questionnaires. Although recruitment was continuously open during the Phase 1 timespan, the major part of participants joined the study during the first months following the study launch: 832 (66%) recruitments after three months of follow-up, 1,136 (90%) after six months (Fig 4).

COHESION Phase 1 participants are distributed all across Canada; they live mainly in Ontario (597, 47%) and Quebec (417, 33%) (Fig 5), and 910 (72%) declared English as their favorite language. They are in average 48 ± 16 (mean ± standard deviation, SD) and are mainly women (78%), White (85%), born in Canada (85%), with university or post-graduate level degree (63%), and in a relationship (67%) (Table 2a). The most part is owner (62%) and lives in a house (66%), with their partner or family (74%), without children at home (72%), with private outside space access (97%), with pets (56%), and in large urban centers (i.e., in areas including more than 100,000 inhabitants); though participants show wide contrasts in terms of home surroundings' greenness and neighborhood's material and social deprivation (Table 2b). Participants are, by decreasing order, employed (58%), retired (19%), unemployed (14%), on leave or disabled (5%), or students (2%); they are mostly satisfied about their household annual income (77%) (Table 2a). Concerning health, they are 44% and 35% being affected by a physical (i.e., heart disease, lung disease, cancer, high blood pressure, diabetes, severe obesity, and/or autoimmune disease) or a mental (i.e., depressive disorder and/or anxiety disorder) chronic disease, respectively. A monthly consumption of alcohol, cigarettes

and/or vape, and cannabis was reported by 72%, 14%, and 18% of participants at baseline, respectively. Lastly, they are 27% and 38% considering their selves and/or someone in their household at a high risk of being infected by- or of complications of COVID-19, respectively.

4.2. Prospective follow-up and attrition

Throughout the first year of follow-up, i.e., from May 2020 to July 2021, up to 17 follow-up questionnaires were administered to participants (Fig 4), and 758 (60%) participants completed at least one (Fig 3). Depending on their date of recruitment, participants were contacted between one and seventeen times as part of the prospective follow-up waves (Fig 4), and they filled out in average 29% $\pm$ 36% (mean $\pm$ SD) of the follow-up questionnaires they received. Among the first year of follow-up, in average 298 $\pm$ 68 participants completed the follow-up questionnaire by wave (Fig 6, *see* S1 Table).

Attrition rate throughout the Phase 1 follow-up appeared statistically distinct in some specific groups (Table 3). The older the participants were and the higher the education level was, the less the attrition was (p <0.001 and p = 0.069, respectively); average age was 50 $\pm$ 16 and 45 $\pm$ 16 (mean $\pm$ SD) in people participating or not in the longitudinal follow-up, respectively (p <0.001). Attrition appeared higher in employed people and lower in retired ones (p <0.001) (Table 3), and also higher in people living in family (p = 0.005), with children (p = 0.013), with pets (p = 0.001), in a house (p = 0.001), and not owner (p = 0.001) (*see* S2a Table). Lastly, people with a chronic physical disease (p = 0.039) or at a high risk of being infected by the COVID-19 (p = 0.001) were less prone to participate to the follow-up. No contrast was noticed according to substance use or residential social and environmental components (*see* S2a-b Tables).

4.3. Well-being and mental health

From June 2020 to July 2021, we administered to participants well-being, sleep credit, loneliness, anxiety, depression and psychological distress -related standardized modules 5, 9, 13, 8, 8, and 11 times, respectively (Table 1). During this period, each thematic module was completed at least once by 481 (38%), 630 (50%), 658 (52%), 612 (48%), 612 (48%), and 692 (55%) Phase 1 participants, respectively (*see* S3a-b Tables).

The median of the 5-WHO well-being Index ranged from 44% (interval interquartile, IQR: 24-65; wave 21/27) to 52% (IQR: 32-72; wave 20/52) depending on the wave. Well-being appeared decreasing waves after waves, the median index value from the last two administered waves (21/23 and 21/27, it means 2021, June and July) being significantly lower than the one from the first three ones (20/52, 21/15, and 21/19, it means 2020, December, 2021, April, and 2021, May, respectively) (Fig 7a; *see* S4a Table). The sleep duration appeared stable throughout the first year of the Phase 1 follow-up (mean ± SD: from 7.9 ± 1.3 to 8.0 ± 1.4 hours; waves 21/23 and 20/30, respectively); no change was statistically found. During the prospective follow-up, the median of the UCLA 3-item score of loneliness oscillated between 5 (IQR: 3-6; wave 21/27) and 6 (IQR: 4-7; wave 21/15). These changes were statistically significant, with a feeling of loneliness higher during the period from 2020, October (wave 20/44) to 2021, April (wave 21/15), yet for the wave 20/52, corresponding to the festive season. Moreover, all UCLA 3-item score median values through the follow-up were significantly higher than the UCLA 3-item score median assessed retrospectively for the period prior to the COVID-19 pandemic (4; IQR: 3-6).

Concerning mental health, the first year of follow-up of the Phase 1 showed statistically significant changes between questionnaire waves regarding reporting of anxiety symptoms, depression symptoms, and psychological distress (Fig 7b; *see* S4b Table). Thus, GAD-7 median score ranged between 4 (IQR: 1-7; wave 21/27) and 5 (IQR: 2-10; wave 21/15) and PHQ-9 median score varied from 4 (IQR: 1-7; wave 21/27) and 6 (IQR: 3-9.3;

wave 20/30). For these both composite indexes, the median value from the wave 21/27 (2021,

July) was the lowest among all waves administered during the first year of follow-up, and

some statistically significant contrasts were observed. Lastly, Kessler-6 psychological distress

median score oscillated from 4 (IQR: 1-9; wave 20/52) and 5 (IQR: 2-10; wave 20/44);

several comparisons between psychological distress median score from the wave 20/52 (2020,

December) and other waves were statistically significant.

5. Discussion

In total, more than 1,200 people enrolled in the Phase 1 of the pan-Canadian population-based COHESION cohort, and about 60% of them participated in the prospective follow-up. The COHESION Study overrepresents women, White and graduated people. The first year of follow-up reveals significant temporal variations in standardized indices of well-being, loneliness, anxiety, depression, and psychological distress. Thus, well-being appeared decreasing waves after waves over the follow-up whereas sleep duration appeared stable throughout. Feeling of loneliness was significantly higher for each follow-up wave compared to the period prior to the COVID-19 pandemic while oscillations in mental health indices were observed over the follow-up timespan. Indices of loneliness and psychological distress were significantly better around the festive period.

Challenges. lessons learned from Phase 1, and proposed adaptations for Phase 2

Loss to follow-up is an issue in any longitudinal study. In Phase 1, attrition was related to age, education level, employment, housing conditions, and some health conditions, similar to other longitudinal studies [57,58]. To try to reduce attrition in Phase 2 and better retain those that are at greatest risk of loss to follow-up, we have devised the following strategies: 1) Reminders: Our questionnaire platform provides automated reminders for online follow-up survey completion including for EMA; 2) Flexibility: Phase 2 will offer easier participation options, relying on a multi-stage on-boarding process, which includes a short initial 10-min questionnaire and easy-to-sign-up longitudinal follow-ups, with possible 'light' or 'complete' choices. To increase accessibility, participation and retention (notably with younger participants), we have also optimized the online questionnaire for ease of use through smartphones. 3) Gifts: While enrollment and participation to the COHESION Study are free

and voluntary, a lottery of 3 monthly prizes ($100 gift cards) is organized for active

participants. In Phase 2, participants who choose the full participation (by completing the

optional follow-up questionnaire modules) will be entered twice in the raffle to increase their

chance of winning. 4) Improved feedback: In December 2020, an opinion survey was sent to

participants to gather feedback on the experience of COHESION Study Phase 1, including on

perceived complexity, length, and usability of the questionnaires, interest in surveyed topics,

interactions with the mobile application. Suggestions for improvement included better

messaging on the purpose of the study, including through regular updates and better

recognition of their contribution, and sharing stories to increase the sense of belonging. These

elements were considered when re-designing the study for Phase 2, in collaboration with the

Center of Excellence on Partnership with Patients and the Public [59].

It is important to include marginalized populations better. We have worked in

collaboration with public health partners to identify best recruitment strategies, including by

building connections with local partners (e.g., Médecins du Monde, Red Cross, Food banks)

that are working directly with various priority groups and have established trust with these

individuals and communities [60]. The use of high-precision targeted social media campaigns,

provided by partnering consumer research company Potloc Inc., along with continuous

monitoring of stratified targets by region, age and gender, should also contribute to optimize

sample representation. Now, because the survey uses online technologies that may be a

barrier for participation in remote communities and for more marginalized groups, we will

also offer computer-assisted phone interviews during working hours through our study

helpline.

Finally, as suggested by Fumagalli et al. (2013), promising approaches based on

persona principles as those used in marketing strategies might help. Thus, we plan to tailor the

content of newsletters, to strengthen the sense of belonging of those most at risk of dropping

out, a strategy that has proved effective for re-engaging young and busy people [61]. Despite

all these methodological efforts to minimize it, the risk of attrition can hardly be eliminated.

One way to account for the potential effect of residual attrition bias is through various

analytical strategies, including the use of inverse probability weights for trajectory analyses

(e.g., growth curve models) on mental health and well-being.

One critical issue with our cohort that was launched in mid-spring 2020 is the lack of a

true pre-pandemic baseline. This can partly be circumvented with retrospective questions, but

as time goes by, the recall bias increases, particularly for subjective mental health measures

[62]. After first using specific 'pre-pandemic' reference questions, a strategy we adopted,

from fall 2020 onwards, to minimize the cognitive effort and bias, was to use a combination

of broader questions about psychological changes since the pandemic outbreak (improvement,

deterioration, no change) and very factual questions for which recall bias is expected lower

(i.e., use of mental health services and substance use in the pre-pandemic year). Despite these

adaptations, the risk of recall bias remains, meaning that interpretation about change from

pre-pandemic measures should be done with caution.

An advantage of a prospective cohort design is the potential for adapting surveys to

new or unforeseen circumstances, including evolving priorities of public health authorities.

As months passed by and the situation evolved, we developed and administered new modules

to address such needs, including on substance use, a priority identified by our partner the

Public Health Agency of Canada, or on parent-related stress during the back-to-school periods.

Repeated surveys also provide the opportunity for timely dissemination of findings. To

facilitate dissemination and use of data by our public health partners, we developed an online

dashboard for real-time monitoring of key indicators (www.cohesionstudy.ca/dashboard).

Throughout phase 2 of COHESION, we will continue adapting our survey content with timely

themes, helping uncovering key pathways linking individual trajectories, environmental

contexts, and health and equity outcomes.

An important contribution of this project is to offer both a pan-Canadian portrait while

allowing local oversampling and providing very contextualized information. An example of

local adaptation is planned for public health territory of CIUSSS Nord-de-l'Île-de-Montréal

(NIM) in Québec [63], with a target local sample of 1,000 participants. Data will be used to

support the co-construction of tailored local intersectoral interventions towards mental health

and more broadly to support activities towards health equity as part of a longer-term

pandemic recovery strategy.

While COHESION Study Phase 1 aimed to understand the mechanisms linking

residential living conditions (built environment, surrounding greenness, neighborhood

deprivation) to differential trajectories of mental health and well-being since the outbreak of

the pandemic, Phase 2 may also provide decision support for public health authorities across

Canada. This contextualized information on a potential representative sample of general

population would be particularly important for policy makers in the pandemic recovery period

where it will be crucial to address health inequities related to income, housing, daily mobility

and social interactions, intimate partner violence, childhood, access to food and health care,

and racism exacerbated over the past two years.

6. Conclusion

Real-time monitoring and evaluation of the unintended consequences on mental health and health inequities of the pandemic is essential for shaping and adapting effective public health policies and programs targeting contextual living conditions (e.g., pedestrianization of streets, securing access to parks, housing renovation programs, permanent supportive housing programs, neighborhood greening program). We will benefit from the support of the Uni-Cité Collaboratory [64] which specializes in science to policy approaches – to equip research teams and cities with tools to better incorporate science into urban public policy. Our study will provide a comprehensive portrait of the key pathways of the COVID-19 impacts on mental health and well-being across Canada. Our flexible infrastructure will ensure that we can adjust to local needs and to the evolving situation from pandemic to post-pandemic recovery.

7. Acknowledgments

The authors would like to warmly thank Dre. Kate Zinszer (*Centre de Recherche en Santé*

Publique, Montreal, QC, Canada) for her crucial involvement in the foundation of this project;

Alexandra Otis (*CIUSSS Nord-de-l'Ile-de-Montréal*, Montreal, QC, Canada) and Marina

Najjar (*Centre de Recherche en Santé Publique*, Montreal, QC, Canada) for their decisive

coordination work in the Phase 1 COHESION study; Alexie Kim, Salma Sahil, and

Andréanne Fortin for their assistance in study's communication; Alexandre Naud for his help

with the VERITAS questionnaire; and Catherine Grenon for her help in analyzing the

verbatims of the completed surveys. NDVI and MSDI metrics, indexed to DMTI Spatial Inc.

postal codes, were provided by the *Canadian Urban Environmental Health Research*

Consortium (CANUE).

8. Funding

514 The COHESION Study is supported by the *Public Health Agency of Canada* (PHAC, Ref #

515 4500416825 – 450041483) and the *Fonds de Recherche du en Santé du Québec en Santé*

516 *(FRQ-S)* and the *Ministère de l'Économie et de l'Innovation du Québec* (Ref # 52266), and

517 has benefited from infrastructure support from the *Canadian Foundation for Innovation* (CFI,

518 Ref # 41072).

9. Conflict of interest

521 YK holds shares in Polygon Research Inc., the data collection platform that is used for the

522 COHESION study.

523 The content and views expressed in this article are those of the authors and do not necessarily

524 reflect those of the Government of Canada.

10. References

1.   World Health Organization. Rolling updates on coronavirus disease (COVID-19). 2020
[cited 25 Aug 2021]. Available: https://www.who.int/emergencies/diseases/novel-
coronavirus-2019/events-as-they-happen
2.   Government of Canada. COVID-19 daily epidemiology update. 2021 [cited 25 Aug
2021]. Available: https://health-infobase.canada.ca/covid-19/epidemiological-
summary-covid-19-cases.html
3.   Bernier C-A, Boire-Schwab D, Bélanger C, Castonguay J-S, Fimiani M, Fitz-Simon N,
et al. COVID-19: Emergency Measures Tracker. 2021 [cited 25 Aug 2021]. Available:
https://www.mccarthy.ca/fr/node/63791
4.   Statistics Canada. Canadian Economic Dashboard and COVID-19. 2021 [cited 30 Mar
2021]. Available: https://www150.statcan.gc.ca/n1/pub/71-607-x/71-607-x2020009-
eng.htm
5.   Bruine de Bruin W. Age Differences in COVID-19 Risk Perceptions and Mental
Health: Evidence From a National U.S. Survey Conducted in March 2020. Isaacowitz
D, editor. The Journals of Gerontology: Series B. 2021;76: e24–e29.
doi:10.1093/geronb/gbaa074
6.   Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. The
Lancet. 2020. doi:10.1016/S0140-6736(20)30526-2
7.   Tsai J, Wilson M. COVID-19: a potential public health problem for homeless
populations. The Lancet Public Health. 2020. doi:10.1016/S2468-2667(20)30053-0
8.   Jenkins EK, McAuliffe C, Hirani S, Richardson C, Thomson KC, McGuinness L, et al.
A portrait of the early and differential mental health impacts of the COVID-19
pandemic in Canada: Findings from the first wave of a nationally representative cross-
sectional survey. Preventive Medicine. 2021;145: 106333.
doi:10.1016/j.ypmed.2020.106333
9.   The New York Times. The Workers Who Face the Greatest Coronavirus Risk. 15 Mar
2020.
10. Chen JT, Krieger N. Revealing the unequal burden of COVID-19 by income,
race/ethnicity, and household crowding: US county versus zip code analyses. Journal
of Public Health Management and Practice. 2021;27: S46–S56.
doi:10.1097/PHH.0000000000001263
11. Van Lancker W, Parolin Z. COVID-19, school closures, and child poverty: a social
crisis in the making. The Lancet Public Health. 2020. doi:10.1016/S2468-
2667(20)30084-0
12. Yancy CW. COVID-19 and African Americans. JAMA - Journal of the American
Medical Association. 2020. doi:10.1001/jama.2020.6548
13. Pareek M, Bangash MN, Pareek N, Pan D, Sze S, Minhas JS, et al. Ethnicity and
COVID-19: an urgent public health research priority. The Lancet. 2020.
doi:10.1016/S0140-6736(20)30922-3
14. Samuelsson K, Barthel S, Colding J, Macassa G. Urban nature as a source of resilience
during social distancing amidst the coronavirus pandemic. 2020.
doi:10.31219/osf.io/3wx5a
15. Varshney LR, Socher R. COVID-19 Growth Rate Decreases with Social Capital. 2020.
doi:10.1101/2020.04.23.20077321

16. Courtet P, Olie E, Debien C, Vaiva G. Keep socially (but not physically) connected and
carry on: Preventing suicide in the age of COVID-19. Journal of Clinical Psychiatry.
2020. doi:10.4088/JCP.20com13370
17. Monteith LL, Holliday R, Brown TL, Brenner LA, Mohatt N V. Preventing Suicide in
Rural Communities During the COVID-19 Pandemic. Journal of Rural Health.
Blackwell Publishing Ltd; 2021. pp. 179–184. doi:10.1111/jrh.12448
18. Dawkins G. Mental health survey shows Manitobans feeling more anxious due to
social isolation. Winnipeg Sun. 5 May 2020.
19. Mcmaster G. Why COVID-19 loneliness can be especially hard on teens. 2020.
20. Vallée J, Cadot E, Roustit C, Parizot I, Chauvin P. The role of daily mobility in mental
health inequalities: The interactive influence of activity space and neighbourhood of
residence on depression. Social Science & Medicine. 2011;73: 1133–1144.
doi:10.1016/j.socscimed.2011.08.009
21. Chung MH, Leung SF, Välimäki M. Use of tracking technology to examine life-space
mobility among people with depression: a systematic review protocol. BMJ Open.
2020;10: e034208. doi:10.1136/bmjopen-2019-034208
22. Astell-Burt T, Feng X, Kolt GS. Does access to neighbourhood green space promote a
healthy duration of sleep? Novel findings from a cross-sectional study of 259 319
Australians. BMJ Open. 2013;3: e003094. doi:10.1136/bmjopen-2013-003094
23. Public Health Agency of Canada. The Chief Public Health Officer's Report on the
State of Public Health in Canada 2017 – Designing Healthy Living. 2017.
24. Abedi V, Olulana O, Avula V, Chaudhary D, Khan A, Shahjouei S, et al. Racial,
Economic, and Health Inequality and COVID-19 Infection in the United States. Journal
of Racial and Ethnic Health Disparities. 2021;8: 732–742. doi:10.1007/s40615-020-
00833-4
25. Turcotte M, Hango D. Incidence des conséquences économiques de la COVID-19 sur
les préoccupations sociales des Canadiens. 2020.
26. Pevalin DJ, Reeves A, Baker E, Bentley R. The impact of persistent poor housing
conditions on mental health: A longitudinal population-based study. Preventive
Medicine. 2017;105. doi:10.1016/j.ypmed.2017.09.020
27. Schellenberg G, Fonberg J. Housing characteristics and staying at home during the
COVID-19 pandemic. 2020.
28. Deaton J, Deaton B. Food security and Canada's agricultural system challenged by
COVID-19. Canadian Journal of Agricultural Economics. 2020;68.
doi:10.1111/cjag.12227
29. Loopstra R. Vulnerability to food insecurity since the COVID-19 lockdown. 2020.
30. Rotermann M. Canadians who report lower self-perceived mental health during the
COVID-19 pandemic more likely to report increased use of cannabis, alcohol and
tobacco. 2020.
31. Wasfi R, Poirier Stephens Z, Sones M, Laberee K, Pugh C, Fuller D, et al. Recruiting
Participants for Population Health Intervention Research: Effectiveness and Costs of
Recruitment Methods for the INTERACT (INTErventions, Research, and Action in
Cities Team) study. Journal of Medical Internet Research. 2021. doi:10.2196/21142
32. Turner-Bowker D, Hogue SJ. Short Form 12 Health Survey (SF-12). Encyclopedia of
Quality of Life and Well-Being Research. Dordrecht: Springer Netherlands; 2014. pp.
5954–5957. doi:10.1007/978-94-007-0753-5_2698
33. Bech P. Measuring the dimension of psychological general well-being by the WHO-5.
In: Quality of life newsletter. 2004.
34. Smyth C. The Pittsburgh Sleep Quality Index (PSQI). Journal of Gerontological
Nursing. 1999;25: 10–10. doi:10.3928/0098-9134-19991201-10

35. Kessler RC, Green JG, Gruber MJ, Sampson NA, Bromet E, Cuitan M, et al. Screening
for serious mental illness in the general population with the K6 screening scale: results
from the WHO World Mental Health (WMH) survey initiative. International Journal of
Methods in Psychiatric Research. 2010;19: 4–22. doi:10.1002/mpr.310

36. Russell DW. UCLA Loneliness Scale (Version 3): Reliability, validity, and factor
structure. Journal of Personality Assessment. 1996;66: 20–40.
doi:10.1207/s15327752jpa6601_2

37. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing
generalized anxiety disorder: The GAD-7. Archives of Internal Medicine. 2006;166.
doi:10.1001/archinte.166.10.1092

38. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: Validity of a brief depression
severity measure. Journal of General Internal Medicine. 2001;16. doi:10.1046/j.1525-
1497.2001.016009606.x

39. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief
resilience scale: Assessing the ability to bounce back. International Journal of
Behavioral Medicine. 2008;15: 194–200. doi:10.1080/10705500802222972

40. Statistics Canada. Canadian Community Health Survey (CCHS) - Mental Health. [cited
1 May 2020]. Available:
https://www23.statcan.gc.ca/imdb/p3Instr.pl?Function=assembleInstr&a=1&&lang=en
&Item_Id=119788

41. Godin G, Shephard RJ. A simple method to assess exercise behavior in the community.
Canadian journal of applied sport sciences. 1985;10: 141–6.

42. Canadian Income Survey - 2019 Questionnaire [Unmet health care needs module].
[cited 16 Mar 2022]. Available:
https://www23.statcan.gc.ca/imdb/p3Instr.pl?Function=assembleInstr&a=1&&lang=en
&Item_Id=1270841

43. Office of Nutrition Policy and Promotion, Health Products and Food Branch. Canadian
Community Health Survey, Cycle 2.2, Nutrition (2004) - Income-Related Household
Food Security in Canada. Ottawa, CA; 2007.

44. Naud A, Sueur C, Basile C, Kestens Y. Combining social network and activity space
data for health research: tools and methods. Health and Place. 2020;66: 102454.
doi:10.1016/j.healthplace.2020.102454

45. Chaix B, Kestens Y, Perchoux C, Karusisi N, Merlo J, Labadi K. An interactive
mapping tool to assess individual mobility patterns in neighborhood studies. American
Journal of Preventive Medicine. 2012;43. doi:10.1016/j.amepre.2012.06.026

46. Campbell KE, Lee BA. Name generators in surveys of personal networks. Social
Networks. 1991;13: 203–221. doi:10.1016/0378-8733(91)90006-F

47. Fuller D, Bell S, Firth CL, Muhajarine N, Nelson T, Stanley K, et al. Wave 1 results of
the INTerventions, Research, and Action in Cities Team (INTERACT) cohort study:
Examining spatio-temporal measures for urban environments and health. Health &
Place. 2021. doi:10.1016/j.healthplace.2021.102646

48. Yang YS, Ryu GW, Han I, Oh S, Choi M. Ecological Momentary Assessment Using
Smartphone-Based Mobile Application for Affect and Stress Assessment. Healthcare
Informatics Research. 2018;24: 381. doi:10.4258/hir.2018.24.4.381

49. Wilhelm P, Schoebi D. Assessing Mood in Daily Life. European Journal of
Psychological Assessment. 2007;23: 258–267. doi:10.1027/1015-5759.23.4.258

50. Pampalon R, Hamel D, Gamache P, Philibert MD, Raymond G, Simpson A. An Area-
based Material and Social Deprivation Index for Public Health in Québec and Canada.
Canadian Journal of Public Health. 2012;103: S17–S22.

51.   Statistics Canada. Population centre size class values by province and territory, 2016
Census. [cited 2 Aug 2021]. Available: https://www12.statcan.gc.ca/census-
recensement/2016/ref/dict/tab/t1_13-eng.cfm

52.   Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth
Engine: Planetary-scale geospatial analysis for everyone. Remote Sensing of
Environment. 2017;202: 18–27. doi:10.1016/j.rse.2017.06.031

53.   Google Earth Engine. USGS Landsat 8 TOA Reflectance (Orthorectified) - 2013 to
2017. 2017. Available:
https://explorer.earthengine.google.com/detail/LANDSAT/LC8_L1T_TOA

54.   Herrmann T, Gleckner W, Wasfi RA, Thierry B, Kestens Y, Ross NA. A pan-Canadian
measure of active living environments using open data. Health Reports. 2019;30.
doi:10.25318/82-003-x201900500002-eng

55.   Canadian Institute for Health Information. COVID-19 Intervention Timeline in Canada.
2021 [cited 3 Aug 2021]. Available: https://www.cihi.ca/en/covid-19-intervention-
timeline-in-canada

56.   Fields E. Tableau Online Security in the Cloud. 2021.

57.   Jacobsen E, Ran X, Liu A, Chang CCH, Ganguli M. Predictors of attrition in a
longitudinal population-based study of aging. International Psychogeriatrics. 2021;33:
767–778. doi:10.1017/S1041610220000447

58.   Young AF, Powers JR, Bell SL. Attrition in longitudinal studies: Who do you lose?
Australian and New Zealand Journal of Public Health. 2006;30: 353–361.
doi:10.1111/J.1467-842X.2006.TB00849.X

59.   CEPPP – Center of Excellence for Patient and Public Partnership. [cited 4 Apr 2022].
Available: https://ceppp.ca/en/

60.   Wasfi R, Stephens ZP, Sones M, Laberee K, Pugh C, Fuller D, et al. Recruiting
Participants for Population Health Intervention Research: Effectiveness and Costs of
Recruitment Methods for a Cohort Study. J Med Internet Res. 2021;23.
doi:10.2196/21142

61.   Fumagalli L, Laurie H, Lynn P. Experiments with methods to reduce attrition in
longitudinal surveys. Journal of the Royal Statistical Society: Series A (Statistics in
Society). 2013;176: 499–519. doi:10.1111/J.1467-985X.2012.01051.X

62.   Hipp L, Bünning M, Munnes S, Sauermann A. Problems and pitfalls of retrospective
survey questions in COVID-19 studies. Survey Research Methods. 2020;14: 109–114.
doi:10.18148/SRM/2020.V14I2.7741

63.   CIUSSS du Nord-de-l'île-de-Montréal | Portrait de l'établissement. [cited 4 Apr 2022].
Available: https://www.ciusssnordmtl.ca/a-propos/qui-sommes-nous/portrait-de-
letablissement/

64.   Home — Collaboratoire Uni-Cité. [cited 4 Apr 2022]. Available: https://www.uni-
cite.ca/home

65.   Wasfi R, Kestens Y. Built environment and health. Urban Form and Accessibility.
Elsevier; 2021. pp. 187–205. doi:10.1016/b978-0-12-819822-3.00005-5

Tables and figures

3 tables and 7 figures (in color).

Table 1. Main information collected in the framework of the COHESION Study Phase 1 prospective follow-up (June 2020 to July 2021).

Theme	Assessment tool	Assessment period	20/26	20/30	20/32	20/34	20/36	20/38	20/40	20/44	20/48	20/52	21/03	21/07	21/11	21/15	21/19	21/23	21/27
COVID-19																			
COVID-19 infection status		At the questionnaire administration point	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
COVID-19-pandemic-related perceptions		At the questionnaire administration point	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Anti-COVID-19 measures compliance		At the questionnaire administration point				X	X	X	X	X	X	X	X	X	X	X	X	X	X
Position towards anti-COVID-19 vaccine		At the questionnaire administration point							X			X					X	X	X
Anti-COVID-19 vaccination status		At the questionnaire administration point													X	X	X	X	X
Health / Well-being																			
General health	SF-12 (first item only)	During the past 2 weeks	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Well-being	WHO-5 Index	During the past 2 weeks										X				X	X	X	X
Loneliness	UCLA 3-item loneliness score	During the past 2 weeks						X	X	X	X	X	X	X	X	X	X	X	X
Anxiety symptoms	7-item General Anxiety Disorder (GAD-7)	During the past 2 weeks	X	X				X				X				X	X	X	X
Depression symptoms	9-item Patient Health Questionnaire (PHQ-9)	During the past 2 weeks	X	X				X				X				X	X	X	X
Psychological distress symptoms	6-item Psychological Distress Scale (Kessler-6)	During the past 2 weeks				X	X	X	X	X	X	X	X						
Behaviors																			
Outdoor mobility	VERITAS	During the past 7 days	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Social interactions	VERITAS	During the past 7 days	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Sleep credit	Pittsburgh Sleep Quality Index	During the past month	X	X				X				X		X		X	X	X	X
Substance use [a]	Frequency questionnaire	During the past 2 weeks			X		X		X	X		X		X		X	X	X	X

[a] Four items: alcohol, cigarettes, vape, and cannabis.

Follow-up questionnaire waves are named according to their week and year of release (for instance, "20/26" for the follow-up questionnaire proposed to participants in the 26th week of 2020).

Table 2a. Description of COHESION Study Phase 1 participants, at recruitment ($N = 1,268$).

Characteristics	N	Percent
Demographics		
Age category		
15-24	103	8.1 %
25-34	231	18.2 %
35-44	227	17.9 %
45-54	230	18.1 %
55-64	282	22.2 %
$\geq$65	195	15.4 %
Gender		
Woman	985	77.7 %
Man	243	19.2 %
Other identity	40	3.2 %
Ethnic or cultural group		
White	1,080	85.2 %
Other group	188	14.8 %
Born in Canada		
Yes	1,073	84.6 %
No	195	15.4 %
Education level		
School degree	179	14.1 %
College degree	278	21.9 %
University degree	408	32.2 %
Post-graduate degree	393	31.0 %
Unknown / Not answered	10	0.8 %
Employment		
Employed, full-time	545	43.0 %
Employed, part-time	148	11.7 %
Self-employed	36	2.8 %
Unemployed	183	14.4 %
Retired	245	19.3 %
On leave / Disabled	59	4.7 %
Student	24	1.9 %
Other	28	2.2 %
Relationship		
In a relationship / Married	847	66.8 %
Single	282	22.2 %
Separated / Divorced / Widowed	135	10.6 %
Other	4	0.3 %
Household characteristics		
Household composition		
Alone	274	21.6 %
With my family	482	38.0 %
With my partner	451	35.6 %
Other	61	4.8 %
Children (living at home)		
0	918	72.4 %
1	155	12.2 %
2	137	10.8 %
$\geq$3	58	4.6 %
Household annual income		
<$21,000	153	12.1 %
$21,000-$36,000	173	13.6 %
$36,000-$48,000	141	11.1 %
$48,000-$96,000	337	26.6 %
$\geq$$96,000	279	22.0 %
Unknown / Not answered	185	14.6 %

Table 2a. To be continued.

Characteristics	N	Percent
Satisfaction regarding income/needs		
Very well	453	35.7 %
Well	522	41.2 %
Not well	271	21.4 %
Unknown / Not answered	22	1.7 %
Pet ownership		
Yes	693	56.4 %
No	536	43.6 %
Housing		
House tenure		
Owner	786	62.0 %
Renter	382	30.1 %
Other	69	5.4 %
Unknown / Not answered	31	2.4 %
Dwelling type		
House	833	65.7 %
Apartment / Condo	419	33.0 %
Other	12	0.9 %
Unknown / Not answered	4	0.3 %
Private outside space		
Yard or nature	963	78.4 %
Balcony or terrace	225	18.3 %
No	41	3.3 %
General health		
Chronic physical disease		
Yes	557	43.9 %
No	688	54.3 %
Unknown / Not answered	23	1.8 %
Chronic mental disease		
Yes	441	34.8 %
No	794	62.6 %
Unknown / Not answered	33	2.6 %
Substance use		
Alcohol [a]		
No	352	27.8 %
At least once a month	245	19.3 %
At least once a week	465	36.7 %
At least once a day	206	16.2 %
Cigarettes and/or vape [a]		
No	1,095	86.4 %
At least once a month	16	1.3 %
At least once a week	11	0.9 %
At least once a day	146	11.5 %
Cannabis [a]		
No	1,036	81.7 %
At least once a month	88	6.9 %
At least once a week	57	4.5 %
At least once a day	87	6.9 %
COVID-19 [b]		
At a high risk of being infected		
Yes	327	26.6 %
No	902	73.4 %
At a high risk of complications		
Yes	465	37.8 %
No	764	62.2 %

[a] During the month before completing baseline questionnaire.
[b] Participant and/or someone in his/her household.

Table 2b. Residential social and environmental measures (*N* = 1,268).

Indexes	*N*	Percent
Urbanization degree [a]		
Large center	891	70.4 %
Medium center	102	8.1 %
Small center	138	10.9 %
Rural	135	10.7 %
Surrounding greenness (NDVI) [b]		
<0.28	254	20.1 %
0.28-0.36	259	20.5 %
0.36-0.43	266	21.0 %
0.43-0.51	233	18.4 %
≥0.51	253	20.0 %
Material deprivation index [c]		
Quintile 1	295	24.5 %
Quintile 2	279	23.2 %
Quintile 3	257	21.4 %
Quintile 4	203	16.9 %
Quintile 5	169	14.0 %
Social deprivation index [c]		
Quintile 1	162	13.5 %
Quintile 2	162	13.5 %
Quintile 3	199	16.5 %
Quintile 4	251	20.9 %
Quintile 5	429	35.7 %

[a] According to *Statistics Canada* classification, "small", "medium" and "large" centers correspond to areas including between 1,000 and 29,999, between 30,000 and 99,999, and 100,000 and more inhabitants, respectively, while "rural area" is a residual value gathering all areas located outside population centers (at the four-digit code area level; data 2016) [51].

[b] Quintiles of the growing season Normalized Difference Vegetation Index (NDVI) in the COHESION Study cohort (at the six-digit code area level; data 2019) [52,53].

[c] Quintiles of the distribution over the whole Canadian territory (at the Canadian Census dissemination area level; data 2016) [50].

Table 3. Attrition in the COHESION Study Phase 1, and comparison of participants enrolling or not in the prospective follow-up according to demographics (*N* = 1,268).

Characteristics	Participating in follow-ups (*N* = 758)		Participated in baseline only (*N* = 510)		*p*-value [a]
	N	Percent	*N*	Percent	
Age category					<0.001
15-24	47	6.2 %	56	11.0 %	
25-34	120	15.8 %	111	21.8 %	
35-44	134	17.7 %	93	18.2 %	
45-54	131	17.3 %	99	19.4 %	
55-64	183	24.1 %	99	19.4 %	
≥65	143	18.9 %	52	10.2 %	
Gender					0.63
Woman	591	78.0 %	394	77.3 %	
Man	146	19.3 %	97	19.0 %	
Other identity	21	2.8 %	19	3.7 %	
Ethnic or cultural group					0.64
White	649	85.6 %	431	84.5 %	
Other group	109	14.4 %	79	15.5 %	
Born in Canada					0.12
Yes	631	83.2 %	442	86.7 %	
No	127	16.8 %	68	13.3 %	
Education level					0.069
School degree	94	12.4 %	85	16.7 %	
College degree	169	22.3 %	109	21.4 %	
University degree	235	31.0 %	173	33.9 %	
Post-graduate degree	254	33.5 %	139	27.3 %	
Unknown / Not answered	6	0.8 %	4	0.8 %	
Employment					<0.001
Employed, full-time	300	39.6 %	245	48.0 %	
Employed, part-time	85	11.2 %	63	12.4 %	
Self-employed	23	3.0 %	13	2.5 %	
Unemployed	99	13.1 %	84	16.5 %	
Retired	192	25.3 %	53	10.4 %	
On leave / Disabled	29	3.8 %	30	5.9 %	
Student	13	1.7 %	11	2.2 %	
Other	17	2.2 %	11	2.2 %	
Relationship					0.59
In a relationship / Married	511	67.4 %	336	65.9 %	
Single	160	21.1 %	122	23.9 %	
Separated / Divorced / Widowed	85	11.2 %	50	9.8 %	
Other	2	0.3 %	2	0.4 %	

[a] Chi-square test (or Fisher exact test for low numbers).

Figure 1. Conceptual model of the COHESION Study.
Adapted from Wasfi & Kestens (2021) [65].

Can you locate and name the places you have gone to in the past 7 days?

These places could include work, school, shops, services, leisure, cafes, runs, walks, parks, visiting someone's home, etc.

Figure 2. Collecting data on living and activity places with VERITAS-Social.
Screenshot for a fictional participant; the house icon locates the participant's residential address and the pin icons symbolize the visited places located by the participant.

Figure 3. Flow chart of participants' recruitment to the COHESION Study Phase 1 (June 2020 to July 2021).

Figure 4. Timeline of recruitment of the COHESION Study Phase 1 participants ($N = 1,268$).

Follow-up questionnaire waves are named according to their week and year of release (for instance, "20/26" for the follow-up questionnaire proposed to participants in the 26th week of 2020).

All participants	1,268		
Alberta (AB)	61	4.8 %	
British Columbia (BC)	123	9.7 %	
Manitoba (MB)	31	2.4 %	
New Brunswick (NB)	13	1.0 %	
Newfoundland and Labrador (NL)	0	0 %	
Nova Scotia (NS)	12	0.9 %	
Ontario (ON)	597	47.1 %	
Prince Edward Island (PE)	2	0.2 %	
Quebec (QC)	417	32.9 %	
Saskatchewan (SK)	11	0.9 %	
Northwest Territories (NT)	1	0.1 %	
Nunavut (NU)	0	0 %	
Yukon (YT)	0	0 %	

Figure 5. Geographical distribution of COHESION Study Phase 1 participants across Canada (*N* = 1,268).

Figure 6. Participation in Phase 1 of the COHESION Study, baseline and follow-ups (June 2020 to July 2021; $N = 1,268$).

Follow-up questionnaire waves are named according to their week and year of release (for instance, "20/26" for the follow-up questionnaire proposed to participants in the 26th week of 2020). Green flow shows participants responding for the first time to a follow-up questionnaire, blue flow shows participants responding again to a follow-up questionnaire, and grey flow shows participants not responding to follow-ups.

Figure 7a. (A) Well-being (WHO-5 Index) level, (B) sleep duration, and (C) feeling of loneliness (UCLA 3-item loneliness score) throughout the first year of prospective follow-up of the COHESION Study Phase 1.
Significance of comparison tests, after Bonferroni correction: * <0.05, ** <0.01, *** <0.001. Green: baseline questionnaire; blue: follow-up questionnaire. Follow-up questionnaire waves are named according to their week and year of release (for instance, "20/26" for the follow-up questionnaire proposed to participants in the 26th week of 2020).

Figure 7b. Symptoms of (A) anxiety (7-item General Anxiety Disorder, GAD-7, score), (B) depression (9-item Patient Health Questionnaire, PHQ-9, score), and (C) psychological distress (6-item Kessler Psychological Distress Scale) throughout the first year of prospective follow-up of the COHESION Study Phase 1.

Significance of comparison tests, after Bonferroni correction: * <0.05, ** <0.01, *** <0.001. Follow-up questionnaire waves are named according to their week and year of release (for instance, "20/26" for the follow-up questionnaire proposed to participants in the 26th week of 2020