Emotional dysregulation in childhood and disordered eating and self-harm in adolescence: Prospective associations and mediating pathways

Naomi Warne[1,2], Jon Heron[1,2], Becky Mars[1,3], Francesca Solmi[4], Lucy Biddle[1,5], David Gunnell[1,3], Gemma Hammerton[1,2], Paul Moran[1,3,5], Marcus Munafò[2,3,6], Ian Penton-Voak[3,6], Andy Skinner[2,7], Anne Stewart[8,9], Helen Bould[1,2,10]

[1]Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School, University of Bristol

[2]Medical Research Council Integrative Epidemiology Unit, University of Bristol

[3]NIHR Biomedical Research Centre, University Hospitals Bristol and Weston NHS Foundation Trust, University of Bristol

[4]Division of Psychiatry, University College London

[5]NIHR Applied Research Collaboration West at University Hospitals Bristol and Weston NHS Foundation Trust

[6]School of Psychological Science, University of Bristol

[7]Integrative Cancer Epidemiology Programme, Bristol Medical School, University of Bristol

[8]Department of Psychiatry, University of Oxford

[9]Oxford Health NHS Foundation Trust

[10]Gloucestershire Health and Care NHS Foundation Trust

Word count: 5827

Running head: Emotional dysregulation, disordered eating, and self-harm

Abstract

Background: Emotional dysregulation may be a risk factor for disordered eating and self-harm in young people, but few prospective studies have assessed these associations long-term, or considered potential mediators. We examined prospective relationships between childhood emotional dysregulation and disordered eating and self-harm in adolescence; and social cognition, emotional recognition and being bullied as mediators.

Methods: We analysed Avon Longitudinal Study of Parents and Children data on 3453 males and 3481 females. We examined associations between emotional dysregulation at 7 years and any disordered eating and any self-harm at 16 years with probit regression models. We also assessed whether social cognition (7 years), emotional recognition (8 years) and bullying victimisation (11 years) mediated these relationships.

Results: Emotional dysregulation at age 7 years was associated with disordered eating (fully adjusted probit B (95% CI) = 0.082 (0.029, 0.134)) and self-harm (fully adjusted probit B (95% CI) = 0.093 (0.036, 0.150)) at age 16 years. There was no evidence of sex interactions or difference in effects between self-harm and disordered eating. Mediation models found social cognition was a key pathway to disordered eating (females 51.2%; males 27.0% of total effect) and self-harm (females 15.7%; males 10.8% of total effect). Bullying victimisation was an important pathway to disordered eating (females 17.1%; males 10.0% of total effect), but only to self-harm in females (15.7% of total effect). Indirect effects were stronger for disordered eating than self-harm.

Conclusions: In males and females, emotional dysregulation in early childhood is associated with disordered eating and self-harm in adolescence and may be a useful target for prevention and treatment. Mediating pathways appeared to differ by sex and outcome, but social cognition was a key mediating pathway for both disordered eating and self-harm.

Keywords: ALSPAC; self-harm; disordered eating; emotional regulation; prospective; mediation

Abbreviations: ALSPAC – Avon Longitudinal Study of Parents and Children

Introduction

Eating disorders affect up to 6% of female adolescents (Smink, Van Hoeken, Oldehinkel, & Hoek,

2014) and are associated with significant morbidity and mortality (Arcelus, Mitchell, Wales, &

Nielsen, 2011). Disordered eating behaviours (including fasting, binge-eating and purging) are

common (Warne et al., 2021), and associated with negative health and social outcomes, including

eating disorders (Kärkkäinen, Mustelin, Raevuori, Kaprio, & Keski-Rahkonen, 2018; Tabler & Utz,

2015). Self-harm in young people is also common and associated with significant morbidity and

mortality (Bergen et al., 2012; McManus et al., 2019). Critically, there is high comorbidity between

disordered eating and self-harm. Prevalence of self-harm in patients with eating disorders is 14-64%

(Cucchi et al., 2016; Svirko & Hawton, 2007); in the community, almost two-thirds of 16-year-old and

24-year-old females reporting self-harm also report disordered eating (Warne et al., 2021). This

observed co-occurrence may reflect common aetiological pathways and shared risk factors (Svirko &

Hawton, 2007).

Research indicates that emotional dysregulation - the inability to be aware of, accept, regulate, and

modify emotional reactions and subsequent behaviours (Gratz & Roemer, 2004) - is linked to both

eating disorders and self-harm. Emotional dysregulation is cross-sectionally associated with eating

disorders (Lavender et al., 2015; Oldershaw, Lavender, Sallis, Stahl, & Schmidt, 2015) and self-harm

(Colmenero-Navarrete, García-Sancho, & Salguero, 2021; Wolff et al., 2019). Some prospective

studies have found associations between emotional dysregulation and eating disorders (Henderson

et al., 2021; McLaughlin, Hatzenbuehler, Mennin, & Nolen-Hoeksema, 2011; Nolen-Hoeksema, Stice,

Wade, & Bohon, 2007) and self-harm in adolescence (Garisch & Wilson, 2015; Robinson et al., 2019;

Voon, Hasking, & Martin, 2014; You, Lin, & Leung, 2014; Zhou, Daukantaitė, Lundh, Wångby-Lundh,

& Ryde, 2020). However, most of these studies follow-up for a year or less and focus on individual

eating disorders/behaviours, or single maladaptive emotional regulation strategies, rather than

global deficits. If prospective relationships for global emotional regulation deficits are established

over longer periods, then it would enable more time to intervene on modifiable mediators, and

potentially enable earlier interventions and effective prevention.

The mechanisms underlying the potential prospective associations of emotional dysregulation with

disordered eating and self-harm are unknown. Three potential mediators of the association between

emotional dysregulation and subsequent disordered eating and self-harm are social cognition,

emotion recognition and bullying victimisation. Problems with social cognition (cognitive processes

underlying social interaction such as perception, interpretation, and generation of responses to

other people's behaviours and mental states (Green et al., 2008)) in childhood are associated with

higher subsequent rates of adolescent suicide attempts (Culpin et al., 2018), and disordered eating

(Solmi et al., 2020). Cross-sectional studies find that people who self-harm (Seymour et al., 2016)

and who have eating disorders (Caglar-Nazali et al., 2014) have increased difficulties with facial

emotion recognition, and adolescents with dysregulated mood, a concept similar to emotional

dysregulation, are less accurate than controls at identifying facial emotions (Rich et al., 2008).

Emotional dysregulation is prospectively associated with bullying victimisation (Winsper, Hall,

Strauss, & Wolke, 2017) and being bullied is associated with increased risk of subsequent self-harm

(Bowes, Wolke, Joinson, Lereya, & Lewis, 2014; Lereya et al., 2013) and disordered eating (Copeland

et al., 2015). Difficulties with social cognition and emotion recognition may therefore represent

distinct mediating pathways from emotional dysregulation to eating disorders and/or self-harm, by,

in turn, increasing a child's susceptibility to bullying victimisation. Investigation of these factors as

mediators has the potential to identify early modifiable targets and facilitate development of

interventions to prevent both self-harm and eating disorders.

Our aims were to investigate the (1) association between emotional dysregulation in childhood and

disordered eating and self-harm in adolescence in a population-based cohort and (2) extent to which

social cognition, emotion recognition of others' faces and being bullied mediate these associations.

Methods

The research protocol was preregistered on Open Science Framework (https://osf.io/43kcg/).

Deviations from the protocol are listed in Appendix S1.

Study design and participants

We used data from a prospective birth cohort: the Avon Longitudinal Study of Parents and Children

(ALSPAC) (Boyd et al., 2013; Fraser et al., 2013). Pregnant women with an expected delivery date

between 1st April 1991 and 31st December 1992 living in the Avon area of Bristol, UK, were invited to

take part. Of the 14,541 pregnancies, there were 13,988 children alive at 1 year of age. Ethical

approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local

Research Ethics Committees. Informed consent for the use of data collected via questionnaires and

clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law

Committee at the time. The study website contains details of all the available data through a fully

searchable data dictionary and variable search tool:

http://www.bristol.ac.uk/alspac/researchers/our-data/. We used a sample of 6,934 children (3,453

males, 3,481 females) from the core sample (Figure S1).

Exposure

Mothers reported on their child's emotional dysregulation (mean age 6 years 9 months) over the

past 6 months using the Strengths and Difficulties Questionnaire-Dysregulation Profile (SDQ-DP)

(Holtmann, Becker, Banaschewski, Rothenberger, & Roessner, 2011). This is a reliable, valid, and

stable measure of emotional dysregulation (Deutz et al., 2018) composed of emotional, hyperactivity

and conduct SDQ subscales (Table S1) (Goodman, 2001). To avoid small parameter estimates, and

additional decimal places in quoted values, this variable was rescaled by dividing all values by 4,

close to the standard deviation of the SDQ-DP in this sample (possible range 0-7.5). We chose not to

standardize the variable as the SD would be expected to differ between complete case and imputed

samples.

Outcomes

Past-year self-harm and disordered eating were assessed via self-report questionnaire during

adolescence (mean age 16 years 8 months).

Disordered eating

Questions were adapted from the Youth Risk Behaviour Surveillance System (Kann et al., 1996).

Participants were asked about the presence and frequency of fasting (not eating for at least a day to

lose weight/avoid gaining weight), purging (vomiting or taking laxatives/other medications to lose

weight/avoid gaining weight), binge-eating (eating a large amount of food in a short amount of time,

with a sense of loss of control), and excessive exercise (exercise in order to lose weight/avoid gaining

weight that routinely interfered with daily routine or work) (Table S2). Our outcome was *any*

disordered eating (fasting, purging, binge-eating, or excessive exercise at any frequency in the past

year). Previous analysis of this data highlights that binge-eating and fasting are the more frequently

reported behaviours (binge-eating: 16.2% of females, 4.3% of males; fasting: 20.7% of females, 3.4%

of males), whereas smaller proportions of participants reported purging (9.5% of females, 1.9% of

males) and excessive exercise (2.1% of females, 0.8% of males) (Warne et al., 2021).

Self-harm

We used two questions adapted from the Child and Adolescent Self-Harm in Europe study (Madge et

al., 2008): whether the adolescent had hurt themselves on purpose (e.g., taking an overdose of pills

or cutting themselves) and the last time they did so. Our outcome was *any self-harm* (regardless of

suicidal intent) in the last year (Table S2) as suicidal intent was not measured.

Mediators

Social cognition was measured using maternal reports on the Social and Communication Disorders

Checklist (Skuse, Mandy, & Scourfield, 2005) at mean age 7 years 8 months (possible range 0-24).

Emotion recognition was measured using the Diagnostic Assessment of Non-Verbal Accuracy

(Nowicki & Duke, 1994) at mean age 8 years 8 months. We used the total number of facial emotions

incorrectly identified (Reed et al., 2021) (possible range 0-24). For bullying victimisation, we used the

summed score of bullying behaviours (Table S3) reported by participants at mean age 12 years 10

months on the Bullying and Friendship Interview Schedule (Wolke, Woods, Stanford, & Schulz, 2001)

(possible range 0-27). Further details available in Appendix S2.

Confounders

Potential confounders of the paths between exposure, mediators and outcomes included sex

(recorded at birth), socioeconomic disadvantage, maternal mental health, and child general

cognitive ability. We also considered Body Mass Index (BMI), at age 12 years 10 months, to be an

intermediate confounder in mediation analyses. Details on confounder measurement and coding are

available in Appendix S2.

Statistical methods

We used a Structural Equation Modelling framework and the Weighted Least Squares estimator

(WLSMV) in Mplus v8.3 (Muthén & Muthén, 2017).

Prospective associations

First, we used bivariate probit regression to determine associations between emotional

dysregulation and the pair of correlated binary outcomes: disordered eating and self-harm. Wald

tests were used to test whether 1) there was an overall association between emotional

dysregulation and both outcomes, 2) the magnitude of the exposure-outcome effect differed

between outcomes, and 3) interactions with sex. We sequentially adjusted for potential

confounders. Probit coefficients can be interpreted as the effect that a one-unit increase in exposure

has on the predicted z-score of the probability of the outcome. To aid interpretation, we derived

marginal effects from the fully adjusted model and produced a margins plot showing how predicted

probability of each outcome would be expected to vary across the observed range of emotional

dysregulation.

Mediating pathways

Secondly, we extended the model, incorporating social cognition, emotion recognition and bullying

victimisation as potential mediators (Figure 1). For this model, the sample was stratified to permit all

paths to vary by sex. We adjusted all direct and indirect paths for confounders and additionally

adjusted for the intermediate confounding effect of low or high BMI. As paths through BMI were not

of substantive interest, indirect effect estimates for these specific paths were combined with those

related to the three mediators of interest, or the direct effect, as appropriate (Figure S2).

Indirect paths were quantified with the product-of-coefficients method, facilitated using probit

regression for all binary/categorical dependent variables (MacKinnon, Lockwood, Brown, Wang, &

Hoffman, 2007). We employed bootstrapping with 1000 bootstrap samples and the bootstrap SEs

were used when pooling across imputed samples ("Rubin's rules"). We express indirect effects as

percentages of total effect. Finally, we explored associations for (a) each mediator/outcome pairing,

and (b) the exposure and each mediator.

Missing data

We sought to address potential bias due to missing data by using a Fully Conditional Specification

approach with -mi impute chained- in Stata version 16 (StataCorp, 2019). We used 25 cycles of

regression-switching and produced 250 imputed datasets with imputation sample stratified by sex

(more information in Appendix S3).

Results

Descriptive results

Descriptive statistics for the key variables are presented in Table 1. The sample had low levels of

socioeconomic disadvantage and child ethnicity was predominantly white (95.9%) with a small

proportion of children from non-white ethnic groups (4.1%). In the sample, maternal ethnicity

(3.88% missing) was reported as white (94.4%), Black Caribbean (0.37%), Other Black (0.23%), Indian

(0.37%), Pakistani (0.10%), Chinese (0.43%), and Other (0.43%). We were unable to report

proportion of mothers reporting Black African ethnicity as ALSPAC stipulates censoring small cell

sizes to preserve anonymity. At age 16, 15.3% females and 5.6% males reported past year self-harm,

whereas 33.9% females and 9.4% males reported past year disordered eating. A comparison of

observed and imputed samples for all model variables is available in Table S4.

Prospective associations

Table 2 shows probit estimates for the imputed sample. There was modest evidence (p=0.019) for a

positive association between emotional dysregulation and both outcomes, however estimates

increased markedly in magnitude following adjustment for sex. In the fully adjusted model, a one

unit increase in our emotional dysregulation scale was associated with a 0.082 (95% CI 0.029, 0.134)

increase in z-score for probability of disordered eating and a 0.093 (95% CI 0.036, 0.150) increase in

z-score for probability of self-harm (Figure 2). As shown in Table S5, there was little evidence that

emotional dysregulation had a differential impact on disordered eating and self-harm (difference in

estimates p=0.745) and little evidence that the magnitude of associations varied by sex (interaction

p=0.600). Complete case sample results were consistent with imputed sample results (Table S5).

Mediating pathways

Estimated indirect effects (Table 3) show differences in the proportion of the total effect mediated

for each combination of sex and outcome which ranged from 15.7% total indirect effect for self-

harm in males, to 62.2% for disordered eating in females.

Social cognition was the dominant mediator between emotional dysregulation and both self-harm

and disordered eating, with substantial pathways via social cognition alone, and via both social

cognition and bullying. In females, social cognition accounted for 51.2% (46.3%+4.9%) of the total

effect of emotional dysregulation on disordered eating, and 15.7% (11.5%+4.2%) for self-harm;

whilst in males these figures were lower, at 27.0% (21.0%+6.0%) and 10.8% (8.8%+2.0%)

respectively.

In females, mediation via bullying victimisation accounted for 17.1% (12.2%+4.9%+0.0%) of the total

effect on disordered eating and 15.7% (11.5%+4.2%+0.0%) of the total effect on self-harm. For

males, bullying victimisation accounted for 10.0% (6.0%+4.0%+0.0%) of the total effect on

disordered eating but only 4.0% (2.0%+2.0%+0.0%) of the total effect for self-harm.

Finally, emotion recognition accounted for a negligible proportion of the total effect for both

outcomes in both females and males. Additional analyses (Tables S6 and S7) found that whilst

emotion recognition was strongly related to emotional dysregulation (particularly for males) the

association between emotion recognition and both self-harm and disordered eating was modest.

Discussion

Children with higher emotional dysregulation were more likely to report disordered eating and self-

harm during adolescence. Mediating pathways appeared to differ by sex (females had two dominant

indirect effects via social cognition and bullying, whereas males had these two dominant pathways

for disordered eating, but only one dominant indirect effect (social cognition) for self-harm).

Pathways also differed by outcome, with stronger indirect effects found for disordered eating than

self-harm. These results suggest that emotional dysregulation, social cognition, and potentially

bullying, may be useful targets to prevent disordered eating and self-harm.

Our findings are consistent with previous literature which has found positive prospective

associations between emotional dysregulation and self-harm (Garisch & Wilson, 2015; Robinson et

al., 2019; Voon et al., 2014; You et al., 2014; Zhou et al., 2020) and disordered eating (Henderson et

al., 2021; McLaughlin et al., 2011; Nolen-Hoeksema et al., 2007). Our study extends these findings to

a longer follow up (9 years) from childhood to adolescence, and considers both self-harm and

disordered eating, which may share some aetiological factors (Svirko & Hawton, 2007). In

combination with previous literature, our findings suggest that interventions targeting emotional

dysregulation may be useful for prevention and treatment of eating disorders and self-harm, as it is

modifiable at the individual level (Gratz, Weiss, & Tull, 2015). Improving emotional regulation skills

may be useful as a universal transdiagnostic preventative intervention for adolescents (Volkaert,

Wante, Loeys, Boelens, & Braet, 2021). However, further longitudinal research is needed to

understand potential preventative effects for disordered eating and self-harm. Training in emotional

regulation skills can be helpful for adults with eating disorders (Davies et al., 2012) and self-harm

(DeCou, Comtois, & Landes, 2019) and preliminary evidence suggests it may also be useful in

adolescents (Giombini et al., 2021; Kothgassner et al., 2021).

Social cognition was an important mediating pathway for both disordered eating and self-harm,

consistent with literature on prospective links between autism spectrum traits and disordered eating

(Solmi et al., 2020) and self-harm (Culpin et al., 2018). Our work extends previous findings and

suggests that emotional dysregulation may precede these associations. However, we cannot rule out

the possibility that social cognition problems measured at age 7 represent earlier differences in

social cognition that precede emotional dysregulation. One way to sever the link between emotional

dysregulation and subsequent disordered eating and self-harm could be to develop interventions

that de-stigmatise atypical social cognition in children. Alternatively, interventions to strengthen

social cognition (Fernández-Sotos et al., 2018), or strongly related concepts such as mentalisation

(the capacity to interpret and understand behaviour from oneself and others based on underlying

mental states (Simonsen, Jakobsen, Grøntved, & Kjaersdam Telléus, 2020)) may have a place in the

treatment of eating disorders and self-harm. Previous research has found lower mentalisation ability

is associated with eating disorders (Jewell et al., 2016; Simonsen et al., 2020), deficits in

mentalisation (excessive certainty about mental states) predict poorer treatment outcomes for

adolescents with anorexia nervosa (Jewell et al., 2021). Mentalisation-based therapies may also be

effective in improving core body image psychopathology in adults with eating disorders (Robinson et

al., 2016), and reducing self-harm in adolescents (Griffiths et al., 2019; Rossouw & Fonagy, 2012).

We found stronger indirect effects in relation to disordered eating than for self-harm, suggesting

that addressing difficulties with social cognition or mentalisation may be more effective in reducing

or preventing disordered eating than self-harm. Further research on associations between

difficulties in social cognition, mentalisation and eating patterns is warranted.

Bullying victimisation was also a key mediating pathway for disordered eating and self-harm in

females, but only for disordered eating in males. The lack of association in males may have been

driven by very weak associations between bullying victimisation and self-harm (Table S6), and it is

possible that bullying victimisation led to alternative negative outcomes in males that were not

assessed. Nevertheless, our findings suggest that effective anti-bullying interventions (e.g. in schools

(Fraguas et al., 2021)) may be beneficial in preventing disordered eating regardless of sex, and in

preventing self-harm in females.

In the context of our analysis, emotion recognition was not an important mediator. Emotional

dysregulation was associated with emotion recognition, but we did not find strong associations

between emotion recognition and either disordered eating or self-harm. It may be that difficulties in

recognising specific emotions (e.g. anger and fear) are more important than overall accuracy in

emotion recognition. It is also important to acknowledge that our emotion recognition task focused

on facial emotions of others, but difficulties associated with recognising one's own emotions (i.e.

alexithymia) may help explain the relationship between emotion dysregulation and subsequent

disordered eating and self-harm. Indeed associations have previously been reported between

alexithymia and eating disorders, symptoms of eating disorders (Nowakowski, McFarlane, & Cassin,

2013; Shank et al., 2019) and also with self-harm (Norman, Oskis, Marzano, & Coulson, 2020). We

did not have the data to explore this in the current study, but future research would benefit from

considering recognition of one's own emotions.

Strengths and limitations

Strengths include using a large prospective cohort with considerable information on potential

confounders and including both self-harm and disordered eating. Limitations include measuring any

frequency of disordered eating rather than diagnostic-level frequency or a research diagnosis of

eating disorders. This was so we could assess the relationship between emotional dysregulation

across a spectrum of potentially impairing eating behaviours that are not always picked up in clinical

samples. Our results may not generalise to individuals with eating disorder diagnoses. We were

unable to differentiate self-harm with and without suicidal intent in the last year as this data was not

available at age 16. The sample was mainly white and less socioeconomically disadvantaged, so

results may not generalise to other ethnic and socioeconomic groups. Furthermore, we were unable

to list all of the races and ethnicities present in the sample (i.e. list the "non-white" child ethnicity

groups separately) as, due to small numbers, this data was censored to preserve anonymity. Findings

on social cognition as a potential mediator should be interpreted cautiously as social cognition was

measured close in time to the exposure, and by the same informant, which may have inflated

associations. The measure of bullying did not include items specifically pertaining to bullying in

relation to appearance or weight, so the strength of association of bullying in relation to subsequent

disordered eating in both boys and girls may be an underestimate. Although we adjusted for several

important confounders, it is possible there may be residual confounding. Another limitation is that

although we measured our putative mediators at a later time point than emotional dysregulation,

this does not mean that deficits in social cognition and emotion recognition are necessarily a

consequence of emotional dysregulation. Indeed, it is possible that that difficulties with emotion

regulation, social cognition, and emotion recognition are all indicators of a broader deficit in social

and emotional processing, rather than causal of each other. Further work should determine whether

these risk factors represent distinct phenomena. Finally, we did not statistically examine the

difference in proportion-mediated for males and females. Consequently, sex differences should be

interpreted with some caution.

Conclusion

Emotional dysregulation in childhood may be an important precursor for adolescent disordered

eating and self-harm. Social cognition was a key mediating pathway for both outcomes and sexes

and being bullied was important for disordered eating in both males and females, and self-harm in

females. Indirect effects were stronger for disordered eating than for self-harm. Findings suggest

that emotional dysregulation, social cognition and bullying may be useful targets in the prevention

and treatment of disordered eating and self-harm.

Key points and relevance

Key points and relevance

- Few studies have examined associations between childhood emotional dysregulation and adolescent disordered eating/self-harm
- We found emotional dysregulation (7 years) was associated with both self-harm and disordered eating (16 years), but associations did not differ by sex or outcome
- Measuring emotional dysregulation may be a useful way of identifying young people at high risk of developing disordered eating/self-harm and a potential target for interventions
- Social cognition and bullying were important mediating pathways and may be targets for interventions

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

Funding

This work was supported by funding from the Medical Research Council/Medical Research Foundation (MR/S020292/1).

The UK Medical Research Council and Wellcome (217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and NW, JH and HB will serve as guarantors for the contents of this paper. A comprehensive list of grants funding is available on the ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf). This research was specifically funded by NIH (MH087786-01) and Wellcome Trust (GR067797MA). Additional funding is detailed in Appendix S4.

Correspondence to
Dr Helen Bould

University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS82BN

+441173313353

helen.bould@bristol.ac.uk

References

Arcelus, J., Mitchell, A. J., Wales, J., & Nielsen, S. (2011). Mortality rates in patients with anorexia nervosa and other eating disorders: A meta-analysis of 36 studies. *Archives of General Psychiatry*, *68*(7), 724–731.

Bergen, H., Hawton, K., Waters, K., Ness, J., Cooper, J., Steeg, S., & Kapur, N. (2012). Premature death after self-harm: A multicentre cohort study. *The Lancet*, *380*(9853), 1568–1574.

Bowes, L., Wolke, D., Joinson, C., Lereya, S. T., & Lewis, G. (2014). Sibling bullying and risk of depression, anxiety, and self-harm: A prospective cohort study. *Pediatrics*, *134*(4), e1032–e1039.

Boyd, A., Golding, J., Macleod, J., Lawlor, D. A., Fraser, A., Henderson, J., … Smith, G. D. (2013). Cohort profile: The 'Children of the 90s'-The index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*, *42*(1), 111–127.

Caglar-Nazali, H. P., Corfield, F., Cardi, V., Ambwani, S., Leppanen, J., Olabintan, O., … Treasure, J. (2014). A systematic review and meta-analysis of "Systems for Social Processes" in eating disorders. *Neuroscience and Biobehavioral Reviews*, *42*, 55–92.

Colmenero-Navarrete, L., García-Sancho, E., & Salguero, J. M. (2021). Relationship Between Emotion Regulation and Suicide Ideation and Attempt in Adults and Adolescents: A Systematic Review. *Archives of Suicide Research*, 1–34.

Copeland, W. E., Bulik, C. M., Zucker, N., Wolke, D., Lereya, S. T., & Costello, E. J. (2015). Does childhood bullying predict eating disorder symptoms? A prospective, longitudinal analysis. *International Journal of Eating Disorders*, *48*(8), 1141–1149.

Cucchi, A., Ryan, D., Konstantakopoulos, G., Stroumpa, S., Kaçar, A. S., Renshaw, S., … Kravariti, E. (2016). Lifetime prevalence of non-suicidal self-injury in patients with eating disorders: A

systematic review and meta-analysis. *Psychological Medicine*, *46*(7), 1345–1358.

Culpin, I., Mars, B., Pearson, R. M., Golding, J., Heron, J., Bubak, I., … Rai, D. (2018). Autistic Traits

and Suicidal Thoughts, Plans, and Self-Harm in Late Adolescence: Population-Based Cohort

Study. *Journal of the American Academy of Child and Adolescent Psychiatry*, *57*(5), 313-320.e6.

Davies, H., Fox, J., Naumann, U., Treasure, J., Schmidt, U., & Tchanturia, K. (2012). Cognitive

remediation and emotion skills training for anorexia nervosa: An observational study using

neuropsychological outcomes. *European Eating Disorders Review*, *20*(3), 211–217.

DeCou, C. R., Comtois, K. A., & Landes, S. J. (2019). Dialectical Behavior Therapy Is Effective for the

Treatment of Suicidal Behavior: A Meta-Analysis. *Behavior Therapy*, *50*(1), 60–72.

Deutz, M. H. F., Shi, Q., Vossen, H. G. M., Huijding, J., Prinzie, P., Deković, M., … Woltering, S. (2018).

Evaluation of the Strengths and Difficulties Questionnaire-Dysregulation Profile (SDQ-DP).

Psychological Assessment, *30*(9), 1174–1185.

Fernández-Sotos, P., Torio, I., Fernández-Caballero, A., Navarro, E., González, P., Dompablo, M., &

Rodriguez-Jimenez, R. (2018). Social cognition remediation interventions: A systematic

mapping review. *PLoS ONE*, *14*(6), e0218720.

Fraguas, D., Diáz-Caneja, C. M., Ayora, M., Durán-Cutilla, M., Abregú-Crespo, R., Ezquiaga-Bravo, I., …

Arango, C. (2021). Assessment of School Anti-Bullying Interventions: A Meta-analysis of

Randomized Clinical Trials. *JAMA Pediatrics*, *175*(1), 44–45.

Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey smith, G., … Lawlor, D. A.

(2013). Cohort profile: The avon longitudinal study of parents and children: ALSPAC mothers

cohort. *International Journal of Epidemiology*, *42*(1), 97–110.

Garisch, J. A., & Wilson, M. S. (2015). Prevalence, correlates, and prospective predictors of non-

suicidal self-injury among New Zealand adolescents: Cross-sectional and longitudinal survey

data. *Child and Adolescent Psychiatry and Mental Health*, *9*(1), 1–11.

Giombini, L., Nesbitt, S., Kusosa, R., Fabian, C., Easter, A., & Tchanturia, K. (2021). Adapted emotion skills training group for young people with anorexia nervosa: A mixed-methods evaluation. *Neuropsychiatrie*, *35*(2), 76–83.

Goodman, R. (2001). Psychometric properties of the strengths and difficulties questionnaire. *Journal of American Academy of Child and Adolescent Psychiatry*, *40*(11), 1337–1345.

Gratz, K. L., & Roemer, L. (2004). Multidimensional Assessment of Emotion Regulation and Dysregulation: Development, Factor Structure, and Initial Validation of the Difficulties in Emotion Regulation Scale. *Journal of Psychopathology and Behavioral Assessment*, *26*(1), 41–54.

Gratz, K. L., Weiss, N. H., & Tull, M. T. (2015). Examining emotion regulation as an outcome, mechanism, or target of psychological treatments. *Current Opinion in Psychology*, *3*, 85–90.

Green, M. F., Penn, D. L., Bentall, R., Carpenter, W. T., Gaebel, W., Gur, R. C., … Heinssen, R. (2008). Social cognition in schizophrenia: An NIMH workshop on definitions, assessment, and research opportunities. *Schizophrenia Bulletin*, *34*(6), 1211–1220.

Gregg, P., Propper, C., & Washbrook, E. (2007). Understanding the Relationship between Parental Income and Multiple Child Outcomes: A Decomposition Analysis. *LSE STICERD Research Paper No. CASE129*. Retrieved from https://ssrn.com/abstract=1159360

Griffiths, H., Duffy, F., Duffy, L., Brown, S., Hockaday, H., Eliasson, E., … Schwannauer, M. (2019). Efficacy of Mentalization-based group therapy for adolescents: The results of a pilot randomised controlled trial. *BMC Psychiatry*, *19*(1), 1–13. https://doi.org/10.1186/s12888-019-2158-8

Henderson, M., Bould, H., Flouri, E., Harrison, A., Lewis, G., Lewis, G., … Solmi, F. (2021). Association of Emotion Regulation Trajectories in Childhood with Anorexia Nervosa and Atypical Anorexia Nervosa in Early Adolescence. *JAMA Psychiatry*, *78*(11), 1249–1257.

Holtmann, M., Becker, A., Banaschewski, T., Rothenberger, A., & Roessner, V. (2011). Psychometric validity of the strengths and difficulties questionnaire-dysregulation profile. *Psychopathology*, *44*(1), 53–59.

Jewell, T., Collyer, H., Gardner, T., Tchanturia, K., Simic, M., Fonagy, P., & Eisler, I. (2016). Attachment and mentalization and their association with child and adolescent eating pathology: A systematic review. *International Journal of Eating Disorders*. https://doi.org/10.1002/eat.22473

Jewell, T., Herle, M., Serpell, L., Eivors, A., Simic, M., Fonagy, P., & Eisler, I. (2021). Attachment and mentalization as predictors of outcome in family therapy for adolescent anorexia nervosa. *European Child and Adolescent Psychiatry*, *1*, 1–11. https://doi.org/10.1007/s00787-021-01930-3

Kann, L., Warren, C. W., Harris, W. A., Collins, J. L., Williams, B. I., Ross, J. G., & Kolbe, L. J. (1996). Youth Risk Behavior Surveillance- United States, 1995. *Journal of School Health*, *66*(10), 365–377.

Kärkkäinen, U., Mustelin, L., Raevuori, A., Kaprio, J., & Keski-Rahkonen, A. (2018). Do Disordered Eating Behaviours Have Long-term Health-related Consequences? *European Eating Disorders Review*, *26*(1), 22–28.

Kothgassner, O. D., Goreis, A., Robinson, K., Huscsava, M. M., Schmahl, C., & Plener, P. L. (2021). Efficacy of dialectical behavior therapy for adolescent self-harm and suicidal ideation: A systematic review and meta-analysis. *Psychological Medicine*, *51*(7), 1057–1067.

Lavender, J. M., Wonderlich, S. A., Engel, S. G., Gordon, K. H., Kaye, W. H., & Mitchell, J. E. (2015). Dimensions of emotion dysregulation in anorexia nervosa and bulimia nervosa: A conceptual review of the empirical literature. *Clinical Psychology Review*, *40*, 111–122.

Lereya, S. T., Winsper, C., Heron, J., Lewis, G., Gunnell, D., Fisher, H. L., & Wolke, D. (2013). Being

bullied during childhood and the prospective pathways to self-harm in late adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, *52*(6), 608-618.e2.

MacKinnon, D. P., Lockwood, C. M., Brown, C. H., Wang, W., & Hoffman, J. M. (2007). The intermediate endpoint effect in logistic and probit regression. *Clinical Trials*, *4*(5), 499–513.

Madge, N., Hewitt, A., Hawton, K., Wilde, E. J. De, Corcoran, P., Fekete, S., … Ystgaard, M. (2008). Deliberate self-harm within an international community sample of young people: Comparative findings from the Child & Adolescent Self-harm in Europe (CASE) Study. *Journal of Child Psychology and Psychiatry*, *49*(6), 667–677.

McLaughlin, K. A., Hatzenbuehler, M. L., Mennin, D. S., & Nolen-Hoeksema, S. (2011). Emotion dysregulation and adolescent psychopathology: A prospective study. *Behaviour Research and Therapy*, *49*, 544–554.

McManus, S., Gunnell, D., Cooper, C., Bebbington, P. E., Howard, L. M., Brugha, T., … Appleby, L. (2019). Prevalence of non-suicidal self-harm and service contact in England, 2000–14: repeated cross-sectional surveys of the general population. *The Lancet Psychiatry*, *6*(7), 573–581.

Muthén, L. K., & Muthén, B. . (2017). Mplus User's Guide. Los Angeles, CA: Muthén & Muthén.

Nolen-Hoeksema, S., Stice, E., Wade, E., & Bohon, C. (2007). Reciprocal relations between rumination and bulimic, substance abuse, and depressive symptoms in female adolescents. *Journal of Abnormal Psychology*, *116*(1), 198–207.

Norman, H., Oskis, A., Marzano, L., & Coulson, M. (2020). The relationship between self-harm and alexithymia: A systematic review and meta-analysis. *Scandinavian Journal of Psychology*, *61*(6), 855–876. https://doi.org/10.1111/SJOP.12668

Nowakowski, M. E., McFarlane, T., & Cassin, S. (2013). Alexithymia and eating disorders: A critical review of the literature. *Journal of Eating Disorders*, *1*(1), 1–14. https://doi.org/10.1186/2050-2974-1-21/TABLES/1

Nowicki, S., & Duke, M. P. (1994). Individual differences in the nonverbal communication of affect: The diagnostic analysis of nonverbal accuracy scale. *Journal of Nonverbal Behavior*, *18*(1), 9–35.

Oldershaw, A., Lavender, T., Sallis, H., Stahl, D., & Schmidt, U. (2015). Emotion generation and regulation in anorexia nervosa: A systematic review and meta-analysis of self-report data. *Clinical Psychology Review*, *39*, 83–95.

Reed, Z. E., Mahedy, L., Jackson, A., Davey Smith, G., Penton-Voak, I., Attwood, A. S., & Munafò, M. R. (2021). Examining the bidirectional association between emotion recognition and social autistic traits using observational and genetic analyses. *Journal of Child Psychology and Psychiatry*, *62*(11), 1330–1338.

Rich, B. A., Grimley, M. E., Schmajuk, M., Blair, K. S., Blair, J. R., & Leibenluft, E. (2008). Face emotion labeling deficits in children with bipolar disorder and severe mood dysregulation. *Development and Psychopathology*, *20*(2), 529–546.

Robinson, K., Garisch, J. A., Kingi, T., Brocklesby, M., O'Connell, A., Langlands, R. L., … Wilson, M. S. (2019). Reciprocal Risk: the Longitudinal Relationship between Emotion Regulation and Non-suicidal Self-Injury in Adolescents. *Journal of Abnormal Child Psychology*, *47*(2), 325–332.

Robinson, P., Hellier, J., Barrett, B., Barzdaitiene, D., Bateman, A., Bogaardt, A., … Fonagy, P. (2016). The NOURISHED randomised controlled trial comparing mentalisation-based treatment for eating disorders (MBT-ED) with specialist supportive clinical management (SSCM-ED) for patients with eating disorders and symptoms of borderline personality disorder. *Trials*, *17*(1), 1–15. https://doi.org/10.1186/S13063-016-1606-8/FIGURES/2

Rossouw, T. I., & Fonagy, P. (2012). Mentalization-based treatment for self-harm in adolescents: A randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, *51*(12), 1304-1313.e3. https://doi.org/10.1016/j.jaac.2012.09.018

Seymour, K. E., Jones, R. N., Cushman, G. K., Galvan, T., Puzia, M. E., Kim, K. L., … Dickstein, D. P.

(2016). Emotional face recognition in adolescent suicide attempters and adolescents engaging in non-suicidal self-injury. *European Child and Adolescent Psychiatry*, *25*(3), 247–259.

Shank, L. M., Tanofsky-Kraff, M., Kelly, N. R., Jaramillo, M., Rubin, S. G., Altman, D. R., … Yanovski, J. A. (2019). The Association Between Alexithymia and Eating Behavior in Children and Adolescents. *Appetite*, *142*, 104381. https://doi.org/10.1016/J.APPET.2019.104381

Simonsen, C. B., Jakobsen, A. G., Grøntved, S., & Kjaersdam Telléus, G. (2020). The mentalization profile in patients with eating disorders: a systematic review and meta-analysis. *Nordic Journal of Psychiatry*. https://doi.org/10.1080/08039488.2019.1707869

Skuse, D. H., Mandy, W. P. L., & Scourfield, J. (2005). Measuring autistic traits : heritability , reliability and validity of the Social and Communication Disorders Checklist. *British Journal of Psychiatry*, *187*, 568–572.

Smink, F. R. E., Van Hoeken, D., Oldehinkel, A. J., & Hoek, H. W. (2014). Prevalence and severity of DSM-5 eating disorders in a community cohort of adolescents. *International Journal of Eating Disorders*, *47*(6), 610–619.

Solmi, F., Bentivegna, F., Bould, H., Mandy, W., Kothari, R., Rai, D., … Lewis, G. (2020). Trajectories of autistic social traits in childhood and adolescence and disordered eating behaviours at age 14 years: A UK general population cohort study. *Journal of Child Psychology and Psychiatry*.

StataCorp. (2019). Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC.

Svirko, E., & Hawton, K. (2007). Self-Injurious Behavior and Eating Disorders: The Extent and Nature of the Association. *Suicide and Life-Threatening Behavior*, *37*(4), 409–421.

Tabler, J., & Utz, R. L. (2015). The influence of adolescent eating disorders or disordered eating behaviors on socioeconomic achievement in early adulthood. *International Journal of Eating Disorders*, *48*(6), 622–632.

Volkaert, B., Wante, L., Loeys, T., Boelens, E., & Braet, C. (2021). The Evaluation of Boost Camp: A

Universal School-Based Prevention Program Targeting Adolescent Emotion Regulation Skills.

School Mental Health.

Voon, D., Hasking, P., & Martin, G. (2014). Change in emotion regulation strategy use and its impact

on adolescent nonsuicidal self-injury: A three-year longitudinal analysis using latent growth

modeling. *Journal of Abnormal Psychology*, *123*(3), 487–498.

Warne, N., Heron, J., Mars, B., Moran, P., Stewart, A., Munafò, M., … Bould, H. (2021). Comorbidity

of self-harm and disordered eating in young people: Evidence from a UK population-based

cohort. *Journal of Affective Disorders*, *282*, 386–390.

Winsper, C., Hall, J., Strauss, V. Y., & Wolke, D. (2017). Aetiological pathways to Borderline

Personality Disorder symptoms in early adolescence: Childhood dysregulated behaviour,

maladaptive parenting and bully victimisation. *Borderline Personality Disorder and Emotion

Dysregulation*, *4*(1), 1–10.

Wolff, J. C., Thompson, E., Thomas, S. A., Nesi, J., Bettis, A. H., Ransford, B., … Liu, R. T. (2019).

Emotion dysregulation and non-suicidal self-injury: A systematic review and meta-analysis.

European Psychiatry, *59*, 25–36.

Wolke, D., Woods, S., Stanford, K., & Schulz, H. (2001). Bullying and victimization of primary school

children in England and Germany: Prevalence and school factors. *British Journal of Psychology*,

92(4), 673–696.

You, J., Lin, M. P., & Leung, F. (2014). A Longitudinal Moderated Mediation Model of Nonsuicidal

Self-injury among Adolescents. *Journal of Abnormal Child Psychology*, *43*(2), 381–390.

Zhou, Y., Daukantaitė, D., Lundh, L. G., Wångby-Lundh, M., & Ryde, A. (2020). Adolescents' Emotion

Regulation Strategies Questionnaire: Initial Validation and Prospective Associations With

Nonsuicidal Self-Injury and Other Mental Health Problems in Adolescence and Young

Adulthood in a Swedish Youth Cohort. *Frontiers in Psychiatry*, *11*, 1–14.

Tables

Table 1. Descriptive statistics (imputed sample, n=6,934)

		Total sample (n=6,934)	Males (n=3,453)	Females (n=3,481)
		Mean or % (SE)	Mean or % (SE)	Mean or % (SE)
	Demographic characteristics			
Ethnicity (non-white)[a]		4.1% (0.25)	4.2% (0.36)	4.0% (0.35)
Parental social class[b]				
	Lowest parental social class codes iv and v ("manual" partly skilled and unskilled occupations)	4.4% (0.26)	3.8% (0.35)	4.9% (0.39)
	Code iii ("manual" and "non-manual" skilled occupations)	35.4% (0.60)	35.6% (0.84)	35.2% (0.84)
	Code ii ("non-manual" managerial and technical occupations)	45.0% (0.62)	45.1% (0.88)	44.9% (0.87)
	Highest parental social class code i ("non-manual" professional occupations)	15.2% (0.44)	15.5% (0.63)	15.0% (0.62)
Lowest parental income quintile[c]		15.8% (0.46)	16.2% (0.66)	15.5% (0.65)
Maternal education[d]				
	No high school qualification (vocational or less)	22.0% (0.51)	22.2% (0.72)	21.9% (0.71)
	High school (O level)	34.9% (0.58)	35.6% (0.83)	34.3% (0.82)
	Beyond high school (A level or degree)	43.0% (0.60)	42.2% (0.85)	43.8% (0.85)
Father absence (any)[e]		21.5% (0.51)	21.5% (0.72)	21.5% (0.73)
	Exposure			

Emotional dysregulation (rescaled)[f] (range 0-6)	1.60 (0.013)	1.69 (0.018)	1.52 (0.018)
Mediators			
Social cognition (range 0-23)	2.76 (0.045)	3.16 (0.065)	2.36 (0.062)
Emotion recognition (range 0-21)	4.60 (0.034)	4.80 (0.048)	4.40 (0.048)
Bully victimisation (range 0-22)	1.81 (0.036)	1.78 (0.051)	1.84 (0.049)
Outcomes			
Disordered eating	21.7% (0.68)	9.4% (0.80)	33.9% (1.04)
Self-harm	10.4% (0.49)	5.6% (0.60)	15.3% (0.76)

SE = standard error of the estimated mean or proportion

[a] Reported by maternal questionnaire in pregnancy

[b] Defined as highest parental occupation on 1991 British Office of Population and Census Statistics job codes reported during pregnancy

[c] Defined as average weekly household disposable income measured via maternal questionnaire at 33 and 37 months, divided into quintiles and re-scaled to account for estimated housing benefits, family size and composition (Gregg, Propper, & Washbrook, 2007). The lowest category indicates proportions for the lowest quintile

[d] Highest maternal educational qualification reported by mothers during pregnancy

[e] Any absence of child's father reported by mothers at ages 1, 2, 3, 7, 8 and 10 years

[f] Emotional dysregulation was rescaled (by a factor of 4) for the analyses

Table 2. Prospective association between childhood emotional dysregulation and adolescent disordered eating and self-harm (imputed sample, n=6,934)

	Outcome		
	Disordered eating Probit B (95% CI)	Self-harm Probit B (95% CI)	Wald test (2 d.f.), p-value
Unadjusted	0.048 (0.003, 0.094)	0.061 (0.011, 0.111)	7.91, 0.0192
Adjusted for sex	0.095 (0.045, 0.145)	0.086 (0.034, 0.138)	20.10, <0.001
Additionally adjusted for socioeconomic disadvantage	0.086 (0.036, 0.136)	0.086 (0.033, 0.140)	17.57, <0.001
Additionally adjusted for maternal mental health	0.076 (0.024, 0.127)	0.075 (0.019, 0.131)	12.47, 0.002
Fully adjusted for all baseline confounders	0.082 (0.029, 0.134)	0.093 (0.036, 0.150)	15.95, <0.001

Wald test = Omnibus test of the association between exposure and both outcomes

Table 3. Mediating pathways from emotional dysregulation to disordered eating and self-harm (imputed sample, n=6,934)

	Males				Females			
	Disordered eating		Self-harm		Disordered eating		Self-harm	
	Probit B (SE)	% of total	Probit B (SE)	% of total	Probit B (SE)	% of total	Probit B (SE)	% of total
Total effect	0.100 (0.053)		0.102 (0.055)		0.082 (0.034)		0.096 (0.036)	
Direct effect	0.076 (0.063)	76.0%	0.085 (0.065)	83.3%	0.030 (0.039)	36.6%	0.069 (0.042)	71.9%
Total indirect effect	0.024 (0.034)	24.0%	0.016 (0.034)	15.7%	0.051 (0.020)	62.2%	0.027 (0.022)	28.1%
Emotional dysregulation → social cognition → outcome	0.021 (0.033)	21.0%	0.009 (0.033)	8.8%	0.038 (0.020)	46.3%	0.011 (0.022)	11.5%
Emotional dysregulation → emotion recognition → outcome	-0.007 (0.004)	-7.0%	0.003 (0.004)	2.9%	-0.001 (0.001)	-1.2%	0.001 (0.002)	1.0%
Emotional dysregulation → bullying victimisation → outcome	0.004 (0.003)	4.0%	0.002 (0.003)	2.0%	0.010 (0.004)	12.2%	0.011 (0.005)	11.5%
Emotional dysregulation → social cognition → bullying victimisation → outcome	0.005 (0.004)	6.0%	0.002 (0.004)	2.0%	0.004 (0.002)	4.9%	0.004 (0.002)	4.2%
Emotional dysregulation → emotion recognition → bullying victimisation → outcome	0.000 (0.000)	0.0%	0.000 (0.000)	0.0%	0.000 (0.000)	0%	0.000 (0.000)	0.0%

% of total = percentage of the total effect explained by the pathway. Fully adjusted for all confounders. SE incorporates bootstrapping (1000 iterations) and

pooling across 250 imputed datasets.

Figures

Figure 1. Hypothesised model linking emotional dysregulation with bivariate outcomes of self-harm and disordered eating

Direct effects are depicted with lines of double thickness. Confounders and pathways involving confounders are depicted by dashed lines. For clarity, some paths between baseline confounders and mediators are not shown.

Figure 2. The predicted probability of self-harm and disordered eating across the range of emotional dysregulation.

All baseline confounders, other than sex, have been held at their mean levels. Dotted lines indicate 95% CI.