Comparison of the burnout among medical residents before and during the pandemic

Hüseyin KÜÇÜKALİ[1], Sezanur Nazlı TÜRKOĞLU[2], Shams HASANLI[3], Fatma Nur DAYANIR ÇOK[4], Hazal Cansu CULPAN[5], Osman HAYRAN[6]

[1] Queen's University Belfast, Centre for Public Health, Belfast, UK. ORCID: 0000-0003-1669-3107

[2] Bezmialem Vakif University, School of Medicine, Department of Public Health, Istanbul, TR. ORCID: 0000-0003-4638-2596

[3] University of Health Sciences, Hamidiye International School of Medicine, Istanbul, TR. ORCID: 0000-0001-6148-8431

[4] Dicle University, School of Medicine, Department of Public Health, Diyarbakir, TR. ORCID: 0000-0002-1554-1892

[5] Karaman Central Community Health Center, Karaman, TR. ORCID: 0000-0001-9244-3818

[6] Istanbul Medipol University, School of Medicine, Department of Public Health, Istanbul, TR. ORCID: 0000-0002-9994-5033

* Corresponding author: Hüseyin Küçükali (h.kucukali[at]qub.ac.uk)

ABSTRACT

Objective: This study aims to compare the level of burnout syndrome in medical residents before and during the COVID-19 pandemic and identify potential risk factors.

Methods: This cross-sectional study was conducted on medical residents from three different university hospitals in Turkey in March 2021, one year after the pandemic hit Turkey. Burnout is measured by the Maslach Burnout Inventory which assesses three dimensions of it: emotional exhaustion, depersonalization, and personal accomplishment. Collected data were combined and compared with data from a previous study held in the same hospitals in December 2019, three months before the pandemic.

Results: 412 medical residents from three universities participated. The mean age was 27.8±2.4 and half of them were female. Compared to pre-pandemic levels, no significant differences in emotional exhaustion (pre:19.0±7.6 post:18.8±7.8), depersonalization (pre:7.3±4.3 post:7.2±4.4), and personal accomplishment (pre:20.8±5.1 post:21.1±5) scores were observed one year after the pandemic. Adjusting for confounders, multiple linear regression models indicated that those who are female, are in a surgical speciality, have vulnerable cohabitants, and have more night shifts face higher emotional exhaustion. Depersonalisation is higher among those who spent more years in residency, have more night shifts, or have COVID-19 outpatient duty. Females and those who have vulnerable cohabitants have lower levels of Personal Achievement.

Conclusion: This study does not support the hypothesis that pandemic increases the burnout levels. Yet it identifies a couple of pandemic-related factors that are associated with burnout and confirms the association of several previously known factors.

Keywords: Burnout, COVID-19, mental health, resident physician

INTRODUCTION

Resident physicians in Türkiye are more precarious than ever. Physician identity was wounded before the pandemic (Koytak, 2021). The recent decrease in purchasing power and public disputes on physician wages did not do better. Türkiye is already one of the countries with the lowest doctor per population (1.8/1000) among OECD countries (average of 3.5) (OECD, 2021). Considering "The Great Resignation" (Sheather & Slattery, 2021) following the COVID-19 pandemic, a possible wave of resignations may intensify the problems arising from inadequate human resources, such as short consultation times and long waiting times. Because the residency program also works as a tool for distributing doctors after the training more equally among regions, this may also play a role in health inequalities. Lower completion of the residency program would mean fewer specialists in the following years. This may risk the achievements of universal health coverage in Türkiye, particularly for some specialist care. Burnout is known to be a major driver of dissatisfaction and withdrawal from job among medical doctors and studies are alarming about an increase during the pandemic (Sheather & Slattery, 2021).

Burnout has been a mental and physical syndrome characterized by emotional distress that usually occurs in individuals who need to use emotional resources as part of their work (Maslach & Jackson, 1981). It represents with emotional exhaustion, depersonalization, and loss of sense of accomplishment due to prolonged exposure to work-related stress (Agha et al., 2019; Biksegn et al., 2016; Leiter & Maslach, 2014; Moreira et al., 2018). In particularly, during residency period, stress is elongated in terms of time and intensity, and insufficient adaptation leads to burnout or psychobiological exhaustion of physicians (Ironside et al., 2019; Navinés et al., 2016). Burnout affects physicians' clinical judgments, their ability to communicate and the provision of quality care for patients (Dyrbye et al., 2017; Elbarazi et al., 2017; Sanfilippo et al., 2017; Shah et al., 2019; Kane, 2019). In addition, burnout causes higher medical errors due to fatigue and reduced attention, which lead to neglect of personal safety precautions and patient safety (Dimitriu et al., 2020).

It is vital to address burnout in the context of the COVID-19 pandemic. In addition to the economic crisis and high unemployment rate, pandemic stress also affected the physical and psychological health of people (Blustein et al., 2020; Xiong et al., 2020). Countries confronted this period differently, so the impact of a pandemic on the quality of life varies according to the country and the individual (Suryavanshi et al., 2020). While the primary focus of frontline healthcare workers is to minimize transmission and treat patients with COVID-19, the impact, and consequences of the pandemic on mental health are significant (Fiest et al., 2021). During this period, mental health problems such as insomnia, anxiety, depression, burnout syndrome were observed in many study on healthcare workers (Barello et al., 2020; Brito-Marques et al., 2021; Pappa et al., 2020; Çevik & Ungan, 2021).

Several recent studies have found an increased level of burnout in healthcare professionals and medical assistants exposed to the pandemic. A study on medical residents during COVID-19 pandemic indicated a burnout prevalence of 76% (Osama et al., 2020) which is higher than the studies that are conducted before the pandemic (Malik et al., 2016; Serenari et al., 2019; Mion et al., 2013).

In the context of Türkiye, students graduate from medical school after 6 years as medical doctors. Who wants to get specialization chooses a department in a state or foundation owned university according to the score they got in the 'Medical Specialization Exam' (Mert et al., 2022; Sahin & Akcicek, 2005). Duration of medical residency training varies between 3-5 years, depending on the specialty area. In addition to theoretical training, residents take a central role in inpatient and outpatient care and night shifts (Küçükali & Akgöğ, 2021).

In 2020, there were a total of 26,181 residents in 68 university hospitals and 86 training and research hospitals in Türkiye. While there has been a significant transformation in many aspects of Türkiye's health system in the last two decades, an adequate improvement has not been achieved in the work conditions of medical residents. Being overwhelmed by the heavy workload in the hospital, lack of time, energy and support for specialty training, low wages, mobbing, violence from patients and their relatives are among the ongoing problems (Küçükali & Akgöğ, 2021). According to the results of a survey conducted with 1069 medical residents from 11 provinces in Türkiye in 2007, the most common complaints during the residency training period are; excessive workload, excessive number of shifts and economic problems. In addition, medical residents stated that they found the training they received insufficient and that they had communication problems with their trainers (Aysan et al., 2008).

Although it is generally assumed that the incidence of burnout among physicians increased during the COVID-19 pandemic, we wanted to test this hypothesis on medical residents in a context where pre-pandemic level of burnout is already high. It was our chance to have a study (Küçükali et al., 2022) that was conducted in three university hospitals in Türkiye just before the pandemic. Using that study as a point of reference, the current study aims to compare the level of burnout in medical residents before and during the COVID-19 pandemic and identify potential risk factors.

MATERIALS AND METHODS

This cross-sectional study is a replication of a previous study (Küçükali et al., 2022) which investigated the burnout level of medical residents in three university hospitals in Türkiye in December 2019, just before the COVID-19 pandemic emerged. The current study was conducted in the same hospitals in March 2021 (from March 5th to May 5th, 2021) – one year after the pandemic hit Türkiye. There were approximately eight hundred medical residents in three hospitals combined. The study aimed to reach all active medical residents in hospitals excluding non-clinical departments (basic medical sciences, medical pathology, and public health).

Data was collected via a questionnaire that includes new questions regarding COVID-19 in addition to inherited the questions on sociodemographic and occupational factors, and Turkish adaptation of the Maslach Burnout Inventory (Ergin, 1992) from the previous study. Sociodemographic factors included age, sex, and marital status. Occupational factors included professional experience in years after obtaining medical degree, area of medical specialty, years spent in residency program, number of night shifts assigned last month. COVID-19 related factors include having a COVID-19 related duty, having infected by COVID-19 and living with someone at risk of severe COVID-19.

Well-established Maslach Burnout Inventory (MBI) measures burnout in 3 dimensions: emotional exhaustion (EE), depersonalization (DP), and lack of personal achievement (PA) via 22 questions. Questions used to be scored in 5-level Likert style ranging from "never" to "every day". The inventory yields three scores from three sub-scales where burnout syndrome presented by high scores in emotional exhaustion and depersonalization, and low scores in personal accomplishment.

MBI is not a stand-alone diagnostic tool (Amosun & Dantile, 1996) . Since there is no cut-off value, there is no distinction as having burnout or not (Amosun & Dantile, 1996; Pehlevan, 2017). The result is the probability of having burnout (Kleijweg et al., 2013). Although the three sub-dimensions of the burnout scale are related to each other, they are different concepts. For this reason, a total burnout score is not calculated during the evaluation (Pehlevan, 2017). While evaluating burnout in accordance with Maslach's definition, the scores of each sub-dimension are collected and interpreted separately (Pehlevan, 2017; Williamson et al., 2018). In addition, as the burnout increases, the scores of the

emotional exhaustion and depersonalization sub-dimensions increase, while the personal achievement score decreases (Pehlevan, 2017).

Researchers approached participants in their work environment (usually in doctors' rooms in their clinics), informed them about the study, obtained their written consent and handed them a hard copy of the questionnaire. In the beginning, questionnaires were applied in a self-answered manner. However, researchers couldn't reach all residents due to their unforseable shift schedules. For those who cannot be reached after two rounds of visits in all departments, we sent online versions of the questionnaires via instant messaging applications. 69 residents participated to study via online form.

Figure 1. Directed acyclic graph showing causal assumptions

To compare our results with the results of the aforementioned pre-pandemic study, we requested and obtained the data of the previous study. The combined dataset was analysed in Python (numpy v1.21.6, scipy v1.7.3, statsmodels v0.12.2). Mean total scores and standard deviations for each of the three subscales were calculated. Mean scores were compared between study periodsusing the independent samples t-test. Causal assumptions are presented in Figure 1. We build separate multiple linear regression models to estimate total effect of each variable on each of the three burnout sub-scale scores while avoiding "Table 2 Fallacy" (Westreich and Greenland, 2012). To eliminate possible confounding effects, we included confounder variables for given variable into equation according to Figure 1. Normality of distribution of the residuals are assessed with P-P plots. Statistical significance was assumed for the probabilities below 0.05 in statistical tests.This study was reviewed and approved by İstanbul Medipol University's Non-invasive Clinical Studies Ethics Committee on 4.3.2021 (No: 284) and by the Ministry of Health on 5.3.2021.

RESULTS

A total of 412 medical residents from three university hospitals participated in the current study. Participants' characteristics including demographics and COVID-19 related variables were given alongside the previous study in Table 1. The mean age of the residents was 27.8±2.4 and half of them were female.

Table 1. Characteristics of the participants

	Before the pandemic (December 2019)*		During the pandemic (March 2021)	
	n	**%**	**n**	**%**
Sex				
Female	248	50.9	206	50.0
Male	239	49.1	206	50.0
Marital status				
Single	310	63.7	294	71.4
Married	175	35.9	118	28.6
Widowed	1	0.2	0	0.0
Divorced	1	0.2	0	0.0
Hospital type				
Foundation owned	196	40.2	258	62.6
State owned	291	59.8	154	37.4
Specialty group				
Internal	291	59.8	265	64.3
Surgical	196	40.2	142	34.5
Unknown	0	0.0	5	1.2
Vulnerable cohabitant				
No	487	100.0	341	82.8
Yes	0	0.0	71	17.2
COVID-19 diagnosis				
No	487	100.0	267	64.8
Yes	0	0.0	145	35.2
COVID-19 duty				
Only Inpatient	0	0.0	145	35.2
Only outpatient	0	0.0	32	7.8
Both	0	0.0	174	42.2
None	487	100.0	60	14.6
Unknown	0	0.0	1	0.2
Total	487	100.0	412	100.0
	Mean	**Std.**	**Mean**	**Std.**
Age	27.9	2.4	27.8	2.4
Experience (years)	3.4	2.2	3.9	2.3
Residency year	2.3	1.1	2.2	1.3
Night shifts per month	6.0	2.9	6.2	2.7

* Different from published paper (Küçükali et. al. 2022), participants from pathology department (n=3) are excluded in here.

The mean sub-scale scores of two studies compared in Table 2. Point estimates and distributions for each subscale score was similar in both study.

Table 2. Comparison of mean subscale scores before and during pandemic

	Before the pandemic (December 2019)		During the pandemic (March 2021)		
	Mean	**Std.**	**Mean**	**Std.**	**p**
Emotional Exhaustion	19.0	7.6	18.8	7.8	0.707
Depersonalization	7.3	4.3	7.2	4.4	0.792
Personal Accomplishment	20.8	5.1	21.1	5.0	0.310

Table 3 presents the total effect of each variable on burnout subscale scores. We built a separate linear regression model to estimate the total effect of each variable on each subscale score. In each model we adjusted for confounders specific to relationship of focus. Pandemic, i.e. study period, did not show a significant difference in any of the subscales. Emotional Exhaustion score is higher among who are female (1.77 [95%CI 0.76-2.77]), having trained in a surgical specialty (1.16 [95%CI 0.10-2.21]), have a vulnerable cohabitant (2.79 [95%CI 0.78-4.80]), and have more night shifts per month (0.46 [95%CI 0.25-0.66]). Depersonalisation score is higher among who spent more time in residency (0.29 [95%CI 0.00-0.58]), had a COVID-19 related outpatient duty (1.71 [95%CI 0.82-2.60]), and have more night shifts per month (0.23 [95%CI 0.12-0.35]). Personal Achievement score is lower among who are female (0.71 [95%CI 0.04-1.37]) and have a vulnerable cohabitant (1.66 [95%CI 0.28-3.03]).

Table 3. Total effect of each variable on burnout subscale scores

	B*	95% C.I.		p
		Lower	Upper	
Emotional Exhaustion				
Age	0.03	-0.18	0.25	0.755
Sex (Male)	-1.77	-2.77	-0.76	0.001
Marital status (Married)	0.07	-1.10	1.23	0.912
Experience years	0.01	-0.38	0.41	0.943
Hospital type (State)	-0.46	-1.47	0.55	0.371
Specialty type (Surgical)	1.16	0.10	2.21	0.031
Residency year	0.03	-0.49	0.54	0.922
Study period (During pandemic)	-0.19	-1.21	0.82	0.707
Vulnerable cohabitant	2.79	0.78	4.80	0.007
COVID-19 diagnosis	0.38	-1.23	1.99	0.643
COVID-19 duty	1.66	-0.48	3.81	0.129
COVID-19 duty – outpatient	1.21	-0.37	2.79	0.134
COVID-19 duty – inpatient	1.65	-0.18	3.49	0.078
Night shifts per month	0.46	0.25	0.66	0.000
Depersonalisation				
Age	-0.09	-0.21	0.03	0.152
Sex (Male)	-0.18	-0.75	0.39	0.539
Marital status (Married)	0.30	-0.35	0.95	0.366
Experience years	-0.00	-0.23	0.22	0.968
Hospital type (State)	0.28	-0.29	0.85	0.338
Specialty type (Surgical)	0.29	-0.31	0.88	0.347
Residency year	0.29	0.00	0.58	0.049
Study period (During pandemic)	-0.08	-0.65	0.49	0.792
Vulnerable cohabitant	-0.08	-1.22	1.07	0.897
COVID-19 diagnosis	-0.24	-1.15	0.66	0.599
COVID-19 duty	0.54	-0.69	1.76	0.391
COVID-19 duty – outpatient	1.71	0.82	2.60	0.000
COVID-19 duty – inpatient	0.83	-0.21	1.87	0.119
Night shifts per month	0.23	0.12	0.35	0.000
Personal Achievement				
Age	0.08	-0.06	0.22	0.270
Sex (Male)	0.71	0.04	1.37	0.039
Marital status (Married)	-0.16	-0.93	0.62	0.690
Experience years	0.19	-0.07	0.45	0.156
Hospital type (State)	-0.07	-0.74	0.60	0.847
Specialty type (Surgical)	-0.53	-1.23	0.18	0.142
Residency year	0.04	-0.30	0.39	0.801
Study period (During pandemic)	0.35	-0.33	1.02	0.310
Vulnerable cohabitant	-1.66	-3.03	-0.28	0.018
COVID-19 diagnosis	-0.98	-2.05	0.10	0.074
COVID-19 duty	0.63	-0.80	2.07	0.387
COVID-19 duty – outpatient	-0.38	-1.44	0.68	0.485
COVID-19 duty – inpatient	0.89	-0.34	2.12	0.155
Night shifts per month	-0.10	-0.23	0.04	0.167

* Coefficients in each line obtained by a separate regression model that is adjusted for confounders of given variable according to assumptions presented in Figure 1. Confounder adjustment for each variable was as follow:

- age : none
- sex_male : none
- marital_married : age
- experience : age
- hospital_state : none
- specialty_surgical : sex_male
- residency_year : age
- pandemic : none
- vulnerable_cohabitant : pandemic, age, sex_male, marital_married
- covid_diagnosis : pandemic, age, sex_male, marital_married, hospital_state, specialty_surgical, residency_year
- covid_duty : pandemic, age, sex_male, hospital_state, specialty_surgical, residency_year, covid_duty_outpatient, covid_duty_inpatient
- covid_duty_outpatient : pandemic, age, sex_male, hospital_state, specialty_surgical, residency_year
- covid_duty_inpatient : pandemic, age, sex_male, hospital_state, specialty_surgical, residency_year
- night_shift_pm : pandemic, age, sex_male, hospital_state, specialty_surgical, residency_year

DISCUSSION

The main objectives of our study were to compare the level of burnout in medical residents before and during the COVID-19 pandemic and identify risk factors that may be associated with burnout. Our findings suggest that during pandemic medical residents have similar but high levels of burnout as before the pandemic. We have identified that having a vulnerable cohabitant and outpatient COVID-19 duty are risk factors for burnout during pandemic alongside known factors such as female sex, surgical specialty, residency year and number of night shifts.

In the literature a range of causes is suggested for increased burnout during the pandemic. New tasks due to the pandemic, fear of illness, lack of knowledge on using personal protective equipment (PPE), inadequate PPE, faulty infection control measures, faulty communication and directives, lack of emotional support and preparation, low autonomy, lack of management support, low hospital management support, lack of appreciation from supervisors, job insecurity, perceived fatality, not being able to spare enough time for the patient, fear of mistakes, medical neglect, dissatisfaction with patient care, fear of carrying a disease to the family, staying in a place other than the family home for security measures, not being able to see the family, being young, isolation and stigma are some of those(Aba, 2022; Dumea et al., 2022; Kılıç et al., 2021; Sengul et al., 2021; Soares et al., 2022).

Seda-Gombau et. al. conducted series of cross-sectional studies in November 2016, January 2019, and October 2020 to see the change of burnout levels among primary care physicians in Catalonia. Although the study has quite small sample size, it showed increase in burnout as a result of upward trend in EE and DP, and downward trend in PA subscales by the pandemic (Seda-Gombau et al., 2021). Contrary to that study and our expectations, our study does not support the hypothesis of higher burnout during the pandemic. This may be due to already high levels of burnout among medical residents in Türkiye. Another explanation would be that exacerbating effects of the pandemic may be balanced by the relative decrease in demand of other (elective) health services and imposed flexible work schedules during pandemic. Anecdotally, some of our participants commented that except COVID-19 specific duties, their overall workload is actually decreased.

On the other hand, our findings regarding burnout subscale scores are consistent with many studies that are conducted in different regions of Türkiye throughout the pandemic. For example, in April 2020 Kılıç et. al. measured burnout levels of 748 physicians from 59 cities of Türkiye via online convenience survey where EE was 19.7 ± 8, DP was 6.9 ± 4.5 and, PA was 8.9 ± 5.1 (Kılıç et al., 2021). It is worth noting that this study reverse-coded PA subscale and if our study was coded same way our estimation of PA score would be 10.9 ± 5.0. In May 2020 Sevinç et. al. estimated median EE as 22 (IQR: 15-28.7), DP as 6 (IQR: 3-9), and PA as 23 (IQR: 17-26) among 104 physicians, residents, and nurses in an intensive care unit in Istanbul (Sevinc et al., 2022). Similarly, in December 2020, Aba estimated mean EE as 20.3 ± 8.8, DP as 7.9 ± 4.4, and PA as 20.5 ± 4.6 among 350 healthcare workers in Türkiye (Aba, 2022). Those studies provide not only similar point estimates but also similar dispersion for three subscale scores. Of course, heterogeneity in some degree between regions and professional should be expected.

It is understandable that emotional exhaustion, which is evaluated as feeling emotionally distant from work, feeling tired at the beginning and end of the workday, feeling nervous, stressed and overworked due to working with people all day, exhausted and disappointed, helpless due to work, is high in healthcare workers living with a person at risk of COVID-19.

In studies, EE scores were found to be higher in women, single people, those with less professional years, with longer working hours, those who experienced sleep deprivation, and who worked in shifts (Kılıç et al., 2021; Seda-Gombau et al., 2021; Sengul et al., 2021). The effects of work-home conflicts

and maternal responsibilities on female health workers are known (Kılıç et al., 2021; Soares et al., 2022). In our study, multiple regression analysis showed that woman have higher the EE score. It is known that the high number of night shifts that cause sleep deprivation and poor-quality sleep which increases burnout (Seda-Gombau et al., 2021; Sevinc et al., 2022). In our study, multiple regression analysis also showed that higher number of night shifts per month is associated with higher the EE score.

Again, in some studies conducted during the pandemic period, the emotional exhaustion scores of healthcare workers who were in direct contact with patients due to the high infectivity of COVID-19, and who were afraid of infecting their family members with COVID-19 were found to be high (Jose et al., 2020; Liu et al., 2020; Luceño-Moreno et al., 2020; Serrão et al., 2021; Soto-Cámara et al., 2021).

In our study, analysis showed that more time spent in residencyand higher number of night shifts per month is associated with higher DP score.

Some studies show that the residency year does not make a difference in DP (Passos et al., 2022; Wilson et al., 2017). It is known that exposure to criticism and violence increases DP scores (Passos et al., 2022; Wilson et al., 2017). In our study, the fact that the DP score was high in the younger ones but the increase in residency year was a positive predictor of DP may be evaluated in this context. It was found that the burnout levels (EE, DP, and PA) of night shift workers were significantly higher than those working the day shift (Hacimusalar et al., 2021; Sengul et al., 2021). In our study, it was reported that DP scores were higher in shift workers, which was consistent with the literature.

In the early period of the pandemic, that is, at a time when there are many uncertainties, the burnout frequency of healthcare professionals working on the front line was found to be lower. (Wu et al., 2020). However in our study, residents who worked in COVID-19 outpatient services has the higher depersonalisation score. This difference can be related to duration of the exposure and prolonged exposure may lead to depersonalisation.

Different relationships were found between gender and PA (Aba, 2022; Sengul et al., 2021). Although in some studies, as ours, men had a higher average of PA than women, in other studies, it was the opposite (Aba, 2022; Kılıç et al., 2021; Sengul et al., 2021). In a study examining the causal ranking of the three sub-dimensions of burnout among male and female physicians in 2011; It was found that personal accomplishment for men developed independently of the other two sub-dimensions. Despite an increase in emotional exhaustion and depersonalization, a sense of personal accomplishment has increased among male physicians. For men, it was thought that the decrease in personal accomplishment was not a dimension of burnout and they had a sense of self-efficacy that was not affected by emotional exhaustion and depersonalization. For women, it was observed that the sense of personal accomplishment was affected by the other two sub-dimensions of burnout. It has been stated that women who experience emotional exhaustion and become desensitized to their patients may feel guilty and think that they are personally unsuccessful(Houkes et al., 2011).

COVID-19 is, undoubtedly, the biggest distruption that health systems have faced in last century. It is fair to expect it to increase stress and, by extension, burnout among healthcare workers. However, our study does not support this hypothesis. We could not find a significant difference burnout levels of medical residents before and during pandemic. Could it simply be explained by selection bias or small sample size? Despite many others our study did not used convenience sampling with online surveys and tried to reach most of the residents in investigated hospitals. We were able to reach approximately half of the population and distribution of participants were quite similar with reference study. Regarding the medical residents, to our knowledge, we included more participants than any other studies conducted during pandemic in Türkiye.

We would like to raise four possible explanations and hypotheses for further research. First, burnout levels were already high before pandemic and pandemic did not / could not increase them further. Second, in contrary to induced demand beforehand, decrease in demand to other (elective) health services and relatively flexible work schedules during pandemic created a space where medical residents can relax. Third, capacity of health system in Türkiye was able to manage stress on the system. Fourth, pandemic management was successful to protect health system from excess workload. Nevertheless, medical residents are facing serious levels of burnout. As central players in the health system their problems should be always a priority in health management until we reach better working conditions for medical residents.

Limitations

It is important to note that this study was not longitudinal, i.e., data is not necessarily collected from the same residents in the reference study. There is a constant change in residents in each hospital. Thus, this study does not show a causal relationship between the pandemic and burnout among medical residents. Still, the comparison in this study provides better evidence than existing studies in the Turkish context for both local decision-makers and global researchers. Some may argue that multiple comparisons require adjustment for inflation in familywise type 1 error. We did not see a need for adjustment for multiple comparisons. Because we only tested familywise hypotheses while we were comparing burnout levels of two groups and none of the three was significant even at 0.05 level (Table 2). Although the study has achieved an approximate 50% participation rate (exact number of all residents in the hospitals was not provided to us), selection bias should be always kept in mind while interpreting the findings. Residents who experience burnout may be more or less willing to participate in the study than others. For example, percentage of residents from state owned hospital was lower in the current study than previous study. Their participation may be related with their burnout. Lastly, 69 participants answered the online form, and they have a significantly higher mean EE score (21.9±7.5) than those who answered the paper form (18.2±7.7).

Acknowledgements: Authors would like to thank our interns Abidin Fatih Emhan, Ahmet Burak Börekçi, Kadir Kolçak, and Serdar Yıldız for their support in data collection.

Ethical approval: This study is reviewed and approved by İstanbul Medipol University's Non-invasive Clinical Studies Ethics Committee on 4.3.2021 (No: 284) and by the Ministry of Health on 5.3.2021.

Conflict of interest: Authors declare no conflict of interest.

Funding: This study did not receive any external funding.

Data availability: Data used in this study is available on request from the corresponding author.

REFERENCES

Aba, M. (2022). *COVID-19 salgınında görev alan sağlık çalışanlarında travma sonrası stres belirtileri, depresif belirtiler ve tükenmişlik belirtilerinin yordayıcısı olarak belirsizliğe tahammülsüzlük, algılanan tehdit boyutu ve duygusal emek değişkenlerinin rolünün incelenmesi* Hacettepe Üniversitesi, Sosyal Bilimler Enstitüsü]. http://hdl.handle.net/11655/25827

Abdelghani, M., El-Gohary, H. M., Fouad, E., & Hassan, M. S. (2020). Addressing the relationship between perceived fear of COVID-19 virus infection and emergence of burnout symptoms in a sample of Egyptian physicians during COVID-19 pandemic: a cross-sectional study. *Middle East Current Psychiatry*, *27*(1), 70. https://doi.org/10.1186/s43045-020-00079-0

Agha, R., Abdall-Razak, A., Crossley, E., Dowlut, N., Iosifidis, C., & Mathew, G. (2019). STROCSS 2019 guideline: strengthening the reporting of cohort studies in surgery. *International Journal of Surgery*, *72*, 156-165. https://doi.org/10.1016/j.ijsu.2019.11.002

Amosun, S. L., & Dantile, N. D. (1996). Are we graduating physiotherapists with the burnout syndrome? *South African Journal of Physiotherapy*, *52*(3), 69-72.

Aysan, E., Köroğlu, G., Türkeli, V., Özgönül, A., Özyaşar, A., GÜLÜMser, Ç., Kilic, K., Ertürk, T., Özdemir, Ö., & Alpaslan, A. H. (2008). Resident physicians in Turkey: results of a survey of 1069 residents from 11 provinces. *Turkish Journal of Medical Sciences*, *38*(1), 35-42.

Barello, S., Palamenghi, L., & Graffigna, G. (2020). Burnout and somatic symptoms among frontline healthcare professionals at the peak of the Italian COVID-19 pandemic. *Psychiatry research*, *290*, 113129-113129. https://doi.org/10.1016/j.psychres.2020.113129

Biksegn, A., Kenfe, T., Matiwos, S., & Eshetu, G. (2016). Burnout status at work among health care professionals in a tertiary hospital. *Ethiopian Journal of Health Sciences*, *26*(2), 101-108. https://doi.org/10.4314/ejhs.v26i2.3

Blustein, D. L., Duffy, R., Ferreira, J. A., Cohen-Scali, V., Cinamon, R. G., & Allan, B. A. (2020). Unemployment in the time of COVID-19: a research agenda. *Journal of Vocational Behavior*, *119*, 103436. https://doi.org/https://doi.org/10.1016/j.jvb.2020.103436

Borusiak, P., Mazheika, Y., Bauer, S., Haberlandt, E., Krois, I., Fricke, C., Simon, L., Beschoner, P., Jerg-Bretzke, L., & Geiser, F. (2022). The impact of the COVID-19 pandemic on pediatric developmental services: a cross-sectional study on overall burden and mental health status. *Archives of Public Health*, *80*(1), 1-13. https://doi.org/10.1186/s13690-022-00876-5

Brito-Marques, J., Franco, C. M. R., Brito-Marques, P. R., Martinez, S. C. G., & Prado, G. F. D. (2021). Impact of COVID-19 pandemic on the sleep quality of medical professionals in Brazil. *Arquivos de Neuropsiquiatria*, *79*(2), 149-155. https://doi.org/10.1590/0004-282X-anp-2020-0449

Çevik, H., & Ungan, M. (2021). The impacts of the COVID-19 pandemic on the mental health and residency training of family medicine residents: findings from a nationwide cross-sectional survey in Turkey. *BMC Family Practice*, *22*(1), 226. https://doi.org/10.1186/s12875-021-01576-9

de Wit, K., Mercuri, M., Wallner, C., Clayton, N., Archambault, P., Ritchie, K., Gérin-Lajoie, C., Gray, S., Schwartz, L., & Chan, T. (2020). Canadian emergency physician psychological distress and burnout during the first 10 weeks of COVID-19: a mixed-methods study. *Journal of the American College of Emergency Physicians Open*, *1*(5), 1030-1038. https://doi.org/10.1002/emp2.12225

Dimitriu, M. C. T., Pantea-Stoian, A., Smaranda, A. C., Nica, A. A., Carap, A. C., Constantin, V. D., Davitoiu, A. M., Cirstoveanu, C., Bacalbasa, N., Bratu, O. G., Jacota-Alexe, F., Badiu, C. D., Smarandache, C. G., & Socea, B. (2020). Burnout syndrome in Romanian medical residents in time of the COVID-19 pandemic. *Medical Hypotheses*, *144*, 109972. https://doi.org/10.1016/j.mehy.2020.109972

Dumea, E., Efrim, N. D., Petcu, A., Anghel, L., & Puscasu, C. G. (2022). Burnout syndrome in personnel of an infectious diseases hospital, one year after the outbreak of the COVID-19 pandemic.

BRAIN. Broad Research in Artificial Intelligence and Neuroscience, *13*(1Sup1), 230-246. https://doi.org/https://doi.org/10.18662/brain/13.1Sup1/316

Dyrbye, L., Shanafelt, T., Sinsky, C., Cipriano, P., Bhatt, J., Ommaya, A., West, C., & Meyers, D. (2017). Burnout among health care professionals: a call to explore and address this underrecognized threat to safe, high-quality care. *NAM Perspectives*, *7*. https://doi.org/10.31478/201707b

Elbarazi, I., Loney, T., Yousef, S., & Elias, A. (2017). Prevalence of and factors associated with burnout among health care professionals in Arab countries: a systematic review. *BMC Health Services Research*, *17*(1), 491. https://doi.org/10.1186/s12913-017-2319-8

Elhadi, M., Msherghi, A., Elgzairi, M., Alhashimi, A., Bouhuwaish, A., Biala, M., Abuelmeda, S., Khel, S., Khaled, A., Alsoufi, A., Elmabrouk, A., Alshiteewi, F. B., Hamed, T. B., Alhadi, B., Alhaddad, S., Elhadi, A., & Zaid, A. (2020). Burnout syndrome among hospital healthcare workers during the COVID-19 pandemic and civil war: a cross-sectional study [Original Research]. *Frontiers in Psychiatry*, *11*. https://www.frontiersin.org/article/10.3389/fpsyt.2020.579563

Ergin, C. (1992). *Doktor ve hemsirelerde tukenmislik ve Maslach tukenmislik olceginin uyarlanmasi* VII. Ulusal Psikoloji Kongresi, 22th September 1992 Ankara (Turkey).

Fiest, K. M., Parsons Leigh, J., Krewulak, K. D., Plotnikoff, K. M., Kemp, L. G., Ng-Kamstra, J., & Stelfox, H. T. (2021). Experiences and management of physician psychological symptoms during infectious disease outbreaks: a rapid review. *BMC Psychiatry*, *21*(1), 91. https://doi.org/10.1186/s12888-021-03090-9

Hacimusalar, Y., Misir, E., Kahve, A. C., Hacimusalar, G. D., Guclu, M. A., & Karaaslan, O. (2021). The effects of working and living conditions of physicians on burnout level and sleep quality. *La Medicina del lavoro*, *112*(5), 346. https://doi.org/10.23749/mdl.v112i5.11268

Houkes, I., Winants, Y., Twellaar, M., & Verdonk, P. (2011). Development of burnout over time and the causal order of the three dimensions of burnout among male and female GPs. A three-wave panel study. *BMC Public Health*, *11*(1), 240. https://doi.org/10.1186/1471-2458-11-240

Ironside, K., Becker, D., Chen, I., Daniyan, A., Kian, A., Saheba, N., & Hollander, R. (2019). Resident and faculty perspectives on prevention of resident burnout: a Focus group study. *The Permanente Journal*, *23*. https://doi.org/10.7812/tpp/18-185

Jose, S., Dhandapani, M., & Cyriac, M. C. (2020). Burnout and Resilience among Frontline Nurses during COVID-19 Pandemic: A Cross-sectional Study in the Emergency Department of a Tertiary Care Center, North India. *Indian J Crit Care Med*, *24*(11), 1081-1088. https://doi.org/10.5005/jp-journals-10071-23667

Kane, L. (2019). *Medscape national physician burnout, depression & suicide report 2019*. https://www.medscape.com/slideshow/2019-lifestyle-burnout-depression-6011056#5

Kılıç, O. H. T., Murat, A., Varol, U., Sofuoğlu, Z., Çoban, İ., Gülmez, H., Güvendi, G., & Mete, B. D. (2021). Factors affecting burnout in physicians during COVID-19 pandemic. *EgeJournal of Medicine*, *60*(2), 136-144. https://doi.org/10.19161/etd.951043

Kleijweg, J. H. M., Verbraak, M. J. P. M., & Van Dijk, M. K. (2013). The clinical utility of the Maslach Burnout Inventory in a clinical population. *Psychological Assessment*, *25*(2), 435.

Koytak, E. (2021, October). The Changing Social Position of the Medical Profession in the Pandemic. *Tıbbi ve Toplumsal Açıdan Pandemi*, Istanbul, Turkey.

Küçükali, H., & Akgöğ, A. (2021). We do not know enough about burnout among medical residents in Turkey: a rapid review of the literature. *Erciyes Medical Journal*. https://doi.org/10.14744/etd.2021.25564

Küçükali, H., Hayran, O., Ankaralı, Z. C., Cengiz, C., Karakaya, R. N., & Yüce, F. Z. (2022). Medical residents are facing burnout: who are at stake? *Journal of Basic and Clinical Health Sciences*, *6*(2).

Leiter, M., & Maslach, C. (2014). *Burnout at work: a psychological perspective (an edited volume)*. https://doi.org/10.4324/978131589416

Levin, J. (2019). Mental health care for survivors and healthcare workers in the aftermath of an outbreak. In *Psychiatry of pandemics* (pp. 127-141). Springer. https://doi.org/https://doi.org/10.1007/978-3-030-15346-5_11

Liu, X., Chen, J., Wang, D., Li, X., Wang, E., Jin, Y., Ma, Y., Yu, C., Luo, C., & Zhang, L. (2020). COVID-19 outbreak can change the job burnout in health care professionals. *Frontiers in Psychiatry*, *11*, 563781.

López-Núñez, M. I., Rubio-Valdehita, S., Diaz-Ramiro, E. M., & Aparicio-García, M. E. (2020). Psychological capital, workload, and burnout: what's new? the impact of personal accomplishment to promote sustainable working conditions. *Sustainability*, *12*(19), 8124. https://doi.org/https://doi.org/10.3390/su12198124

Luceño-Moreno, L., Talavera-Velasco, B., García-Albuerne, Y., & Martín-García, J. (2020). Symptoms of Posttraumatic Stress, Anxiety, Depression, Levels of Resilience and Burnout in Spanish Health Personnel during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, *17*(15). https://doi.org/10.3390/ijerph17155514

Malik, A. A., Bhatti, S., Shafiq, A., Khan, R. S., Butt, U. I., Bilal, S. M., Khan, H. S., Kashif Malik, M., & Ayyaz, M. (2016). Burnout among surgical residents in a lower-middle income country - Are we any different? *Annals of Medicine and Surgery*, *9*, 28-32. https://doi.org/10.1016/j.amsu.2016.05.012

Maslach, C., & Jackson, S. E. (1981). The measurement of experienced burnout. *Journal of organizational behavior*, *2*(2), 99-113. https://doi.org/https://doi.org/10.1002/job.4030020205

Maslach, C., Schaufeli, W. B., & Leiter, M. P. (2001). Job burnout. *Annual review of psychology*, *52*(1), 397-422. https://doi.org/https://doi.org/10.1146/annurev.psych.52.1.397

Mert, L., Demirel, M., Kendirci, A. Ş., Akgül, T., & Kılıçoğlu, Ö. İ. (2022). Reasons for resident resignations from Orthopedic Residency Programs in Turkey: A cross-sectional survey from residents' perspectives.

Mion, G., Libert, N., & Journois, D. (2013). [Burnout-associated factors in anesthesia and intensive care medicine. 2009 survey of the French Society of anesthesiology and intensive care]. *Annales Françaises d'Anesthésie et de Réanimation*, *32*(3), 175-188. https://doi.org/10.1016/j.annfar.2012.12.004 (Facteurs associés au burnout en anesthésie-réanimation. Enquête 2009 de la Société française d'anesthésie et de réanimation.)

Moreira, L., Govêia, C., Miranda, D., & Amorim, M. (2018). Burnout in anesthesiology: when exhaustion overcomes the duty. *Journal of Anaesthesia and Therapeutics*, *1*(2).

Navinés, R., Martín-Santos, R., Olivé, V., & Valdés, M. (2016). Work-related stress: Implications for physical and mental health. *Medicina Clínica*, *146*(8), 359-366. https://doi.org/10.1016/j.medcli.2015.11.023 (Estrés laboral: implicaciones para la salud física y mental.)

Nituica, C., Bota, O. A., Blebea, J., Cheng, C.-I., & Slotman, G. J. (2021). Factors influencing resilience and burnout among resident physicians - a national survey. *BMC medical education*, *21*(1), 514-514. https://doi.org/10.1186/s12909-021-02950-y

Osama, M., Zaheer, F., Saeed, H., Anees, K., Jawed, Q., Syed, S. H., & Sheikh, B. A. (2020). Impact of COVID-19 on surgical residency programs in Pakistan; A residents' perspective. Do programs need formal restructuring to adjust with the "new normal"? A cross-sectional survey study. *International Journal of Surgery*, *79*, 252-256. https://doi.org/10.1016/j.ijsu.2020.06.004

Pappa, S., Ntella, V., Giannakas, T., Giannakoulis, V. G., Papoutsi, E., & Katsaounou, P. (2020). Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Brain, behavior, and immunity*, *88*, 901-907. https://doi.org/10.1016/j.bbi.2020.05.026

Passos, L. S., Alves, V. d. S., Rodrigues, L. A. R. L., & Saraiva, L. V. (2022). Burnout syndrome in resident physicians of a Federal University. *Revista da Associação Médica Brasileira*, *68*, 24-30. https://doi.org/10.1590/1806-9282.20210241

Pehlevan, O. (2017). 112 Acil sağlık hizmetlerinde çalışan personelin tükenmişlik ve iş doyumları arasındaki ilişkinin incelenmesi. *Gazi Üniversitesi, Sosyal Bilimler Enstitüsü*.

Sahin, H., & Akcicek, F. (2005). Internal medicine residency training in Turkey. *European Journal of Internal Medicine*, *16*(8), 590-595. https://doi.org/https://doi.org/10.1016/j.ejim.2005.08.002

Sanfilippo, F., Noto, A., Foresta, G., Santonocito, C., Palumbo, G. J., Arcadipane, A., Maybauer, D. M., & Maybauer, M. O. (2017). Incidence and factors associated with burnout in anesthesiology: a systematic review. *BioMed Research International*, *2017*, 8648925. https://doi.org/10.1155/2017/8648925

Seda-Gombau, G., Montero-Alía, J. J., Moreno-Gabriel, E., & Torán-Monserrat, P. (2021). Impact of the COVID-19 pandemic on burnout in primary care physicians in Catalonia. *International journal of environmental research and public health*, *18*(17), 9031. https://doi.org/10.3390/ijerph18179031

Sengul, H., Bulut, A., & Kahraman, B. (2021). The impact of changing processes in the COVID-19 pandemic on health care workers' burnout syndrome: web-based questionnaire study. *Medical Bulletin of Haseki/Haseki Tip Bulteni*. https://doi.org/10.4274/haseki.galenos.2021.7130

Serenari, M., Cucchetti, A., Russo, P. M., Fallani, G., Mattarozzi, K., Pinna, A. D., Colonnello, V., Poggioli, G., & Cescon, M. (2019). Burnout and psychological distress between surgical and non-surgical residents. *Updates in Surgery*, *71*(2), 323-330. https://doi.org/10.1007/s13304-019-00653-0

Serrão, C., Duarte, I., Castro, L., & Teixeira, A. (2021). Burnout and Depression in Portuguese Healthcare Workers during the COVID-19 Pandemic—The Mediating Role of Psychological Resilience. *International Journal of Environmental Research and Public Health*, *18*(2). https://doi.org/10.3390/ijerph18020636

Sevinc, S. A., Metin, S., Basi, N. B., Cinar, A. S., Ozkan, M. T., & Oba, S. (2022). Anxiety and burnout in anesthetists and intensive care unit nurses during the COVID-19 pandemic: a cross-sectional study. *Brazilian Journal of Anesthesiology*. https://doi.org/10.1016/j.bjane.2021.07.011

Shah, A., Wyatt, M., Gourneau, B., Shih, G., & De Ruyter, M. (2019). Emotional exhaustion among anesthesia providers at a tertiary care center assessed using the MBI burnout survey. *Psychology, Health & Medicine*, *24*(5), 620-624. https://doi.org/10.1080/13548506.2018.1546019

Shanafelt, T. D., Hasan, O., Dyrbye, L. N., Sinsky, C., Satele, D., Sloan, J., & West, C. P. (2015). Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014.

Shanafelt, T. D., West, C. P., Sinsky, C., Trockel, M., Tutty, M., Wang, H., Carlasare, L. E., & Dyrbye, L. N. (2022). Changes in burnout and satisfaction with work-life integration in physicians and the general US working population between 2011 and 2020.

Sheather, J., & Slattery, D. (2021). The great resignation—how do we support and retain staff already stretched to their limit? BMJ, n2533. https://doi.org/10.1136/bmj.n2533

Soares, J. P., Oliveira, N. H. S. d., Mendes, T. d. M. C., Ribeiro, S. d. S., & Castro, J. L. d. (2022). Burnout-related factors in health professionals during the COVID-19 pandemic: an integrative review. *Saúde em Debate*, *46*, 385-398. https://doi.org/10.1590/0103-11042022E126I

Soto-Cámara, R., García-Santa-Basilia, N., Onrubia-Baticón, H., Cárdaba-García, R. M., Jiménez-Alegre, J. J., Reques-Marugán, A. M., Molina-Oliva, M., Fernández-Domínguez, J. J., Matellán-Hernández, M. P., Morales-Sanchez, A., & Navalpotro-Pascual, S. (2021). Psychological Impact of the COVID-19 Pandemic on Out-of-Hospital Health Professionals: A Living Systematic Review. *Journal of Clinical Medicine*, *10*(23). https://doi.org/10.3390/jcm10235578

Suryavanshi, N., Kadam, A., Dhumal, G., Nimkar, S., Mave, V., Gupta, A., Cox, S. R., & Gupte, N. (2020). Mental health and quality of life among healthcare professionals during the COVID-19 pandemic in India. *Brain and Behavior*, *10*(11), e01837. https://doi.org/10.1002/brb3.1837

Şahin, Z. (2011). *Bir devlet hastanesinde çalışan hemşireler ile bir vakıf üniversitesi hastanesinde çalışan hemşirelerin iş stres ve iş doyumu düzeylerinin incelenmesi* Maltepe Üniversitesi, Sosyal Bilimler Enstitüsü]. https://hdl.handle.net/20.500.12415/2797

Türkmen, İ., & Aslan, E. Ç. *COVID-19 pandemisi sürecinde sağlık çalışanlarının algılanan stres ve duygusal tükenme düzeylerinin incelenmesi* International Black Sea Coastline Countries Scientific Research Symposium – VI,

Williamson, K., Lank, P. M., Cheema, N., Hartman, N., & Lovell, E. O. (2018). Comparing the Maslach Burnout Inventory to Other Well-Being Instruments in Emergency Medicine Residents. *J Grad Med Educ*, *10*(5), 532-536. https://doi.org/10.4300/jgme-d-18-00155.1

Wilson, W., Raj, J. P., Narayan, G., Ghiya, M., Murty, S., & Joseph, B. (2017). Quantifying burnout among emergency medicine professionals. *Journal of emergencies, trauma, and shock*, *10*(4), 199. https://doi.org/10.4103/JETS.JETS_36_17

Westreich, D., & Greenland, S. (2013). The Table 2 Fallacy: Presenting and Interpreting Confounder and Modifier Coefficients. *American Journal of Epidemiology, 177*(4), 292–298. https://doi.org/10.1093/aje/kws412

Wu, Y., Wang, J., Luo, C., Hu, S., Lin, X., Anderson, A. E., Bruera, E., Yang, X., Wei, S., & Qian, Y. (2020). A Comparison of Burnout Frequency Among Oncology Physicians and Nurses Working on the Frontline and Usual Wards During the COVID-19 Epidemic in Wuhan, China. *Journal of Pain and Symptom Management*, *60*(1), e60-e65. https://doi.org/10.1016/j.jpainsymman.2020.04.008

Xiong, J., Lipsitz, O., Nasri, F., Lui, L. M. W., Gill, H., Phan, L., Chen-Li, D., Iacobucci, M., Ho, R., Majeed, A., & McIntyre, R. S. (2020). Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *Journal of Affective Disorders*, *277*, 55-64. https://doi.org/10.1016/j.jad.2020.08.001

Zarei, E., Ahmadi, F., Sial, M. S., Hwang, J., Thu, P. A., & Usman, S. M. (2019). Prevalence of burnout among primary health care staff and its predictors: a study in Iran. *International journal of environmental research and public health*, *16*(12). https://doi.org/10.3390/ijerph16122249